



Universiteit
Leiden
The Netherlands

Structural dynamics of light harvesting proteins, photosynthetic membranes, and cells observed by spectral editing solid-state NMR
Pandit, A.

Citation

Pandit, A. (2022). Structural dynamics of light harvesting proteins, photosynthetic membranes, and cells observed by spectral editing solid-state NMR. *The Journal Of Chemical Physics*, 157(2). doi:10.1063/5.0094446

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3492087>

Note: To cite this publication please use the final published version (if applicable).

Structural dynamics of light harvesting proteins, photosynthetic membranes, and cells observed by spectral editing solid-state NMR

Cite as: J. Chem. Phys. **157**, 025101 (2022); <https://doi.org/10.1063/5.0094446>

Submitted: 03 April 2022 • Accepted: 14 June 2022 • Accepted Manuscript Online: 15 June 2022 •

Published Online: 08 July 2022

 Anjali Pandit

COLLECTIONS

Paper published as part of the special topic on [Photosynthetic Light-Harvesting and Energy Conversion](#)



[View Online](#)



[Export Citation](#)



[CrossMark](#)

ARTICLES YOU MAY BE INTERESTED IN

[Chlorophyll a de-excitation pathways in the LHCII antenna](#)

The Journal of Chemical Physics **156**, 070902 (2022); <https://doi.org/10.1063/5.0073825>

[Observation of conformational dynamics in single light-harvesting proteins from cryptophyte algae](#)

The Journal of Chemical Physics **157**, 035102 (2022); <https://doi.org/10.1063/5.0095763>

[Improved estimation for energy dissipation in biochemical oscillations](#)

The Journal of Chemical Physics **157**, 025102 (2022); <https://doi.org/10.1063/5.0092126>



[Learn More](#)

Special Topics Open for Submissions

Structural dynamics of light harvesting proteins, photosynthetic membranes, and cells observed by spectral editing solid-state NMR

Cite as: J. Chem. Phys. 157, 025101 (2022); doi: 10.1063/5.0094446

Submitted: 3 April 2022 • Accepted: 14 June 2022 •

Published Online: 8 July 2022



Anjali Pandit^{a)}

AFFILIATIONS

Department of Solid-State NMR, Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

Note: This paper is part of the JCP Special Topic on Photosynthetic Light-Harvesting and Energy Conversion.

^{a)}Author to whom correspondence should be addressed: a.pandit@lic.leidenuniv.nl

ABSTRACT

Photosynthetic light-harvesting complexes have a remarkable capacity to perform robust photo-physics at ambient temperatures and in fluctuating environments. Protein conformational dynamics and membrane mobility are processes that contribute to the light-harvesting efficiencies and control photoprotective responses. This short review describes the application of magic angle spinning nuclear magnetic resonance (NMR) spectroscopy for characterizing the structural dynamics of pigment, protein, and thylakoid membrane components related to light harvesting and photoprotection. I will discuss the use of dynamics-based spectral editing solid-state NMR for distinguishing rigid and mobile components and assessing protein, pigment, and lipid dynamics on sub-nanosecond to millisecond timescales. Dynamic spectral editing NMR has been applied to investigate light-harvesting complex II protein conformational dynamics inside lipid bilayers and in native membranes. Furthermore, we used the NMR approach to assess thylakoid membrane dynamics. Finally, it is shown that dynamics-based spectral editing NMR for reducing spectral complexity by filtering motion-dependent signals enabled us to follow processes in live photosynthetic cells.

Published under an exclusive license by AIP Publishing. <https://doi.org/10.1063/5.0094446>

INTRODUCTION

Photosynthetic light-harvesting and energy conversion processes take place in the photosynthetic membrane, the thylakoid, and encompass the interplay between pigments and protein complexes in a dynamic membrane environment.^{1–3} Inside the thylakoid membrane, robust photophysical and photochemical processes are carried out at ambient temperatures. Thermal fluctuations, protein conformational dynamics, and membrane transitions contribute to the efficiency of light harvesting and excitation transfer and allow for flexibility of these processes to respond to fluctuating environmental conditions.^{3–8} The assessment of photosynthetic protein, pigment, and membrane dynamics on nano- to millisecond timescales provides insight into the molecular and nanoscale mechanisms of photosynthesis regulation. This short review describes the application of Magic Angle Spinning solid-state (MAS ss) nuclear magnetic resonance (NMR) spectroscopy in combination with dynamics-based spectral editing to gain insight into the structural dynamics

of light-harvesting pigment–protein complexes and for characterizing protein and lipid dynamics in thylakoid membranes. MAS ssNMR is widely used as a tool for structural characterization of membrane proteins and biomembranes at the atomic level.^{9–12} In solution-state NMR spectroscopy, rapid tumbling of molecules averages out anisotropic interactions, such as dipolar coupling and chemical shift anisotropy, which results in long relaxation times for the magnetization and narrow peaks. Large structures, such as membrane proteins in lipid bilayers, polymers, crystals, and molecular assemblies, experience slow tumbling that is usually too slow for averaging out anisotropic interactions, resulting in broad peaks or NMR invisibility. In the solid state, NMR spectroscopy on solids or semi-solids is combined with MAS. Under MAS, the sample inside a small rotor is held under an angle of 54.7° with respect to the static magnetic field (Z-axis) and the XY plane and spun around its axis with frequencies typically ranging from 5 to 50 kHz depending on the rotor size and instrumentation (Fig. 1). Fast rotation of the sample at this fixed angle with respect to the magnetic field reduces




FIG. 1. The concept of MAS NMR. The sample inside an NMR rotor is held at an angle θ of 54.7° (magic angle) with respect to the magnetic field B_z and spun fast around its axis.

contributions of anisotropic interactions by averaging those, similar to rapid tumbling in the solution state.¹³ Due to strong ^1H – ^1H dipolar couplings and magnetic field inhomogeneities caused by structural disorder, ^1H line widths are broadened in the solid state, even with MAS applied. Solid-state NMR on biological samples typically involves ^{13}C or $^{13}\text{C}/^{15}\text{N}$ detection of isotope-labeled samples, sometimes requiring dedicated labeling strategies.¹⁴

DYNAMICS-BASED SPECTRAL EDITING NMR

A basic ^{13}C NMR experiment encompasses single pulse excitation, also termed direct polarization (DP), of the ^{13}C nuclei (90° pulse) followed by ^{13}C detection. A drawback of this type of NMR experiment is that ^{13}C longitudinal T_1 relaxation times for molecules with low internal motions are very long, requiring long recycle delay intervals in between repeated scans. ^{13}C ssNMR experiments commonly employ ^1H – ^{13}C heteronuclear cross-polarization (CP), where magnetization is transferred from protons to carbons via dipolar interactions, and ^{13}C is detected. Compared to DP, CP NMR provides enhancement of signal proportional to $\gamma^1\text{H}/\gamma^{13}\text{C}$ and shortens the relaxation times because of the faster

longitudinal relaxation of excited protons compared to ^{13}C . In contrast to DP, signal intensities of CP NMR are affected by internal motions that reduce the efficiency of dipolar polarization transfer, leading to the invisibility of molecules with sub-microsecond dynamics.¹⁵ Highly mobile residues can be detected by an alternative polarization transfer NMR scheme, Insensitive Nuclei Enhanced by Polarization Transfer (INEPT), where magnetization is transferred through bond via scalar couplings (J couplings).^{16,17} INEPT-based polarization transfer makes use of echo pulses and renders inefficient for rigid molecules of which the transverse T_2 relaxation times are very fast. As a consequence, INEPT-based NMR only detects signals of molecules with dynamical motions in the (sub)nanosecond range. The combination of CP and INEPT experiments provides a way to selectively detect rigid molecules (via CP) or very mobile molecules (via INEPT).^{15,18} Changes in the structural dynamics are detectable if they move signals in or out the CP or INEPT NMR detection window. DP signal intensities are motion-independent and DP spectra detect signals within a wide dynamics range. Figure 2 summarizes the dynamics ranges for CP, INEPT, and DP NMR detection under the experimental conditions that we used in 1D MAS NMR experiments for analyzing membrane and cell dynamics.

Dynamics-based spectral editing ssNMR has found a wide range of applications. Conventional CP-based ssNMR spectroscopy on membrane proteins or protein assemblies does not detect protein sites with fast or intermediate dynamics that are typical for unstructured loop and tail segments. The combination of ssNMR with spectral editing has been applied by van der Wel and co-workers to address the rigid and flexible regions of amyloid fibrils and their polymorphs and gain molecular insight into their structures and dynamic disorder.^{19,20} In addition, they could reveal the structure and dynamics of peripherally bound membrane proteins that contain flexible, water-exposed sites.²¹ Furthermore, J -coupling-based INEPT ssNMR spectroscopy has been applied to determine the unfolded extra-membrane structures of integral membrane proteins, in addition to the folded, rigid protein sites that are embedded in the membrane and accessible with CP-based ssNMR. The NMR dynamics filtering technique is not only limited to resolving protein structural dynamics but also could resolve aspects of hydration and water behavior in hydrogels and detect phase transitions in lipid membranes and surfactants.^{22–24} A novel technique based on the principles of dynamics-based spectral editing is comprehensive multi-phase (CMP) NMR.²⁵ The CMP NMR method involves a specialized NMR probe for simultaneous detection of solid, semi-solid, and liquid phases. CMP NMR has been applied in *in situ*




FIG. 2. Dynamics time scales for NMR detection via DP- (yellow), CP- (blue), or INEPT-based (red) MAS NMR experiments and processes occurring on the respective time scales.

NMR experiments on intact micro-organisms or heterogeneous biomaterials for distinction and analysis of rigid, semi-solid, and fluid constituents.

ssNMR SPECTROSCOPY APPLIED TO PHOTOSYNTHETIC ANTENNA SYSTEMS

In the past few decades, various structures of photosynthetic light-harvesting proteins have been resolved at high resolution by crystallography and cryo-electron microscopy (cryo-EM).^{26–30} Yet, open questions remain concerning the pigment structural and electronic environments and dynamics of light-harvesting protein and pigments that may control light-harvesting and photoprotective processes. ssNMR is a technique that has compelling advantages when it comes to resolving atomic details of pigment–protein interactions and dynamics in a flexible protein environment. Moreover, as crystallization is not a requirement, the molecular structure can be resolved within mesoscopic self-assemblies with substantial dynamics or inhomogeneity. ssNMR was applied to determine the ground-state electronic structures of chlorophylls (Chls) in purple non-sulfur bacterial antenna systems^{31–33} revealing molecular mechanisms by which mutual interactions with the protein matrix stabilize the pigment–protein quaternary structures. Alia, de Groot, and co-workers found that the BChls in LH2 antenna complexes modify the electronic structures of the ligating histidines with mutual charge transfer between the histidines and BChls.^{34,35} Furthermore, the group of Matysik has applied photochemically induced dynamic nuclear polarization (photo-CIDNP) to uncover the electronic structure and special properties of the reaction center special pair Chls in intact plants, algae, and diatoms.^{36–38} de Groot and Buda combined ssNMR data and further optimization by density functional theory (DFT) calculations for elucidation of the molecular structures (unit cells) of chlorosomes and man-made artificial light-harvesting tubular assemblies.^{39–41} In recent work from the lab of de Groot, Sevink, and co-workers, molecular dynamics simulations were applied to show how dynamic disorder drives excitation transfer in chlorosome assemblies.^{42,43} They created a protocol for *in silico* generation of atomically resolved plastic, tubular structures with defined dihedral angles, and used unit cells derived from ssNMR data on chlorosomes to generate MD models of chlorosome tubular assemblies, which now can be used as a platform to investigate the links between chlorosome structural mechanics, plasticity, and optical properties.⁴²

Light-Harvesting Complex II (LHCII) is the major light-harvesting protein in plants and green algae, which is responsible for capturing light and guiding the excitation energy to Photosystem II (PSII) and under high-light conditions also to PSI. An unresolved intriguing property of LHC-type light-harvesting antenna complexes of plants and algae is their capacity to switch their function between fluorescent, light-harvesting, photoprotective, and excitation-quenched states via reversible conformational transitions.^{6,44,45} X-ray structures of several LHC complexes have been obtained,^{29,30,46,47} which formed the basis for atomistic and coarse-grained MD simulations.^{48–54} ssNMR spectroscopy may bridge the gap between *in silico* protein dynamic models and crystallographic pigment–protein structures and highlight specific pigment and protein sites with high plasticity. Moreover, ssNMR spectroscopy is applicable to membrane-embedded proteins and

biological membranes and could shed light on protein–lipid interactions that may have importance for the dual function of LHC complexes.

This short review describes the application of dynamics-based spectral editing ssNMR for gaining insight into the conformational dynamics of the most abundant LHC complex, LHCII, in native-like membrane environments and inside native membranes. Furthermore, it is shown that dynamics-based spectral editing ssNMR is applicable for determining lipid membrane dynamics in intact photosynthetic cells and holds promise for simultaneous detection of structural dynamics and metabolic processes in live photosynthetic cells.

CONFORMATIONAL DYNAMICS OF LIGHT-HARVESTING COMPLEX II

The conformational dynamics of LHCII has been investigated by various techniques, including single-molecule fluorescence, electron paramagnetic resonance (EPR), and neutron scattering techniques.^{6,44,55,56} Single molecule fluorescence has revealed that individual pigment–protein complexes fluctuate between fluorescent and fluorescence-quenched states.⁶ Neutron scattering methods including (quasi-) inelastic neutron scattering (QENS) can measure directly protein dynamics extending ps–ns time scales. A combined optical and QENS low-temperature study on LHCII demonstrated that there is a correlation between conformational dynamics and the position of the LHCII Chl excited electronic states due to protein conformational fluctuations.⁵⁷ EPR studies revealed that the protein complex possesses a flexible N tail and rigid transmembrane interior.⁵⁵ Recently, it was found via neutron scattering techniques that LHCII proteins can adopt different quaternary structures, including small oligomers, depending on the detergent types.⁵⁸ We employed dynamics-based spectral editing NMR to assess the structure and dynamics of LHCII in native and native-like membrane environments.⁵⁹ ¹³C-labeled LHCII was obtained from *Chlamydomonas reinhardtii* (Cr) cells that were grown on ¹³C-acetate. The NMR INEPT spectrum of LHCII reconstituted in thylakoid lipid bilayers contained a moderate set of signals that were assigned to the highly flexible N-tail of LHCII and the tails of two chlorophylls [see LHCII structures in Figs. 3(a) and 3(b) and spectrum in Fig. 3(e)]. INEPT NMR detects highly dynamic and unstructured protein segments that are often not accessible for crystallization.^{60–63} Consistent with this, the protein N tail and the tails of two Chls (Chl 605 and 606) have not been resolved in the crystal structures of LHCII.^{30,46} Sites with intermediate dynamics were identified based on the invisibility of predicted NMR signals in the CP-based spectrum, indicated in the structure in Fig. 3(b). Those sites that undergo motions on microsecond timescales—the timescale for domain motions and protein allostery—are located at the membrane–water interface and part of the stromal loop and lumen helix edges connected to the amphipathic helices D and E. Most of the Chl macrocycle atom signals were invisible with CP detection, indicating that the LHCII Chl rings undergo motions on microsecond timescales. The Chl macrocycles become visible in CP-based NMR at temperatures below 240 K, suggesting strongly reduced pigment and protein dynamics below this temperature, correlating with the onset temperature for conformational motions as indicated in QENS measurements on LHCII.⁵⁷




FIG. 3. NMR-detected protein dynamics of LHCII. (a)–(c) LHCII structure with detected dynamic sites indicated. Highly mobile sites are in red and sites with intermediate (μs) dynamics in yellow. (a) Chl dynamics, showing for Chls 605 and 606; (b) protein dynamics, and (c) flexible sites in LHCII. The arrows with an S point to Ser sites that may convert to strand conformations in npq2 LHCII. The LHCII structures were modeled using Lhcbm structures from PDB-IDs 6KAC, 7DZ8, and 7E0H. (d) and (e) D, CP-MAS ^{13}C - ^{13}C spectrum with identified protein helix (h) and coil (c) amino acid types, Chl (green) and lipid (blue) signals indicated. (e) Overlaid spectra DP-MAS ^{13}C - ^{13}C spectrum (red) and INEPT-TOBSY ^{13}C - ^{13}C spectrum (blue) with Chl phytol signals indicated. (d) and (e) are modified with permission from Azadi-Chegeni *et al.*, *Biophys. J.* **120**(2), 270–283 (2021). Copyright 2021 Cell Press.

Influence of zeaxanthin

Notable changes in structure and dynamics were observed for LHCII from the xanthophyll-cycle impaired mutant npq2 that constitutively accumulates zeaxanthin (Zea) in its thylakoid membranes.⁶⁴ The npq2 mutant contained LHCII complexes binding Zea instead of Vio and had destabilized LHCII trimers so that the LHCII

complexes were isolated as monomers. Monomeric Zea-LHCII complexes from the npq2 mutant display a larger static disorder than wild-type LHCII trimers, shown as broader lines in CP-based NMR spectra, and lacked sites with intermediate dynamics. Distinctive spectral changes were observed in the stromal loop regions where Ser residues in strand conformation were identified [see Fig. 3(c)]. Protein signals in the INEPT spectrum were conserved, indicating

that the npq2 LHCII also contained a highly flexible N tail. Strong signals of monogalactosyldiacylglycerol (MGDG) and digalactosyldiacylglycerol (DGDG) lipids in the CP-based NMR spectrum of npq2 LHCII indicated that multiple lipids were associated with npq2 LHCII, which were strongly protein bound. In contrast, the CP spectrum of wild-type LHCII trimers only contained a lipid signal of phosphatidyl glycerol (PG), which is a structural lipid that is bound to each monomer in LHCII.³⁰ Additional coarse-grain MD simulations supported the NMR data of npq2 LHCII and demonstrated that monomerization of LHCII increases structural displacements and promotes lipid association, while substitution of Vio by Zea has an additional effect on the protein–lipid affinities.⁶⁴

Dynamic picture of LHCII and photoprotective states

From the NMR studies, a dynamic picture of LHCII emerges in which the protein has (1) a highly dynamical, disordered N tail, (2) flexible loops at the stromal and luminal membrane–water interface, which undergo microsecond local motions, and (3) stromal loop and N-terminal coil that can undergo conformational transitions that are stabilized by monomerization or Zea. The NMR-determined LHCII dynamic sites agree well with atomistic MD simulation studies that have shown that the N terminus is highly disordered and that the most flexible domains are exposed to the membrane–water interface.^{49,53} Reversible transitions have been proposed to occur in two LHCII pigment domains: one formed by lutein (Lut1) and the Chl *a* 610, 611, and 612 and another by Lut2 and Chls 602 and 603.^{6,49,65} Protein dynamics has been associated with transitions from active light-harvesting to excitation-quenched states: motions in the N terminus of LHCII correlated with fluctuations in Chl611–612 excitonic coupling⁴⁹ and LHC quenching mechanisms may involve short-range carotenoid interactions with multiple Chls that are fine-tuned by protein motions and modest conformational rearrangements.^{66–68} Similar effects were found for LHCII using QENS, where it was found that a variety of conformational substates with different spectral positions may be thermally accessible, with a special focus on Chl612, implying that dynamic pigment–protein interactions fine-tune electronic energy levels and electron–phonon coupling of LHC II.⁵⁷ An NMR comparison between LHCII in fully active and strongly quenched states showed characteristic changes in signals of Chl *a* macrocycle rings and heterogeneity of the stromal coil connected to the N terminus, while the protein core of intercrossing helices containing the binding sites of Chl610 and Chl602 was unchanged.^{69,70} Heterogeneity of the LHCII N-terminus has also been reported in electron paramagnetic resonance (EPR) spin-label studies.⁵⁵ The combined NMR results would be consistent with a hypothetical mechanism in which conformational transitions at the stromal loop and N terminus, together with local displacements at the lumen membrane–water interface, propagate into the LHC pigment domains where they are able to control quencher states.

LHCII in native membranes

How dynamic is LHCII in a native thylakoid membrane? To address this question, we performed an NMR analysis on purified ¹³C-isotope labeled thylakoid membranes from *Chlamydomonas reinhardtii* (Cr) cells that were grown in ¹³C-labeled media.⁶⁴ The results revealed that in the native membranes,

LHCII does not contain a highly dynamic N-tail, and both large fluctuations and microsecond local dynamics were reduced. Structures of photosynthetic Photosystem (PS)I- and PSII-LHCII supercomplexes have provided detailed information on LHCII protein–protein, protein–pigment, and protein–lipid interactions in thylakoid membranes.^{27,28,71–73} The stromal and luminal loops are involved in LHCII–LHCII interactions²⁷ and the N terminus of LHCII can interact with PSI core subunits.⁷³ In stacked membranes, LHCII stromal loops can be involved in interactions between adjacent membranes.⁷⁴ Such interactions are likely to reduce local motions of LHCII in native membranes and stabilize its N terminus. The absence of highly dynamic protein signals in the thylakoid spectrum, however, implies that also unbound LHCII complexes have stabilized N tails. To conclude, the internal dynamics of LHCII is significantly lower in native membranes than in *in vitro* systems demonstrating that the conformational dynamics of LHCs is tightly coupled with their local environments.

THYLAKOID MEMBRANE DYNAMICS

Thylakoid membranes containing the photosynthetic apparatus have evolved remarkable plasticity, forming complex architectures with stacked and unstacked appressed membrane regions and hexagonal phases.^{75–77} Changes in thylakoid structure and dynamics have been observed with respect to stacking, protein lateral diffusion, and membrane reorganization in response to light conditions.^{78–81} Inelastic neutron scattering has been applied to study the flexibility and motions of photosynthetic membranes inside leaves or whole cells, revealing revealed membrane undulatory motions coupled to photosynthetic activity.^{82,83}

Combined INEPT-DP- and CP-based 1D MAS NMR spectroscopy provides a usual means to qualify and quantify lipid dynamics and membrane phase transitions^{24,84} that we applied to assess the dynamics of protein and lipid thylakoid components.^{85–87} The ¹³C NMR spectrum of thylakoid membranes contains broad bands of the protein backbone and side chains with superimposed sharp peaks from lipids, where we distinguish the accumulated *n*CH₂ signals of the lipid acyl chain, CH atoms of unsaturated lipid tails, and the head atoms of galactosyl lipids (see Fig. 4). Furthermore, we could distinguish *all-trans* from *trans-gauche* lipid conformations.⁸⁸ The thylakoid CP spectrum is selective for signals of the protein backbone and immobilized lipids and the INEPT spectrum mainly contains signals of mobile lipids.

Influence of zeaxanthin

We addressed the influence of Zea on thylakoid membrane dynamics. For membranes of the npq2 mutant, a temperature-dependent increase in protein dynamics within the temperature range 3–25 °C was smaller than for the wild-type, while the increase in dynamic lipids was larger as depicted in Fig. 3.⁸⁵ Given the fact that npq2 LHCII monomers strongly bind lipid and have a greater content of immobilized lipids compared to wild-type LHCII trimers, this suggests that the npq2 membranes contain separate regions of immobilized protein–lipid aggregates and lipid pools with high mobility. The overall reduction in membrane fluidity that has been observed for Zea-containing membranes is explained by the fact that the majority of lipids are immobilized so that the overall lipid




FIG. 4. DP, CP, and INEPT ^{13}C MAS NMR spectra of stacked (black) and unstacked (red) thylakoid membranes and the difference spectrum of stacked minus unstacked (blue). Changes in protein backbone (CO and C_α) and side chain dynamics are observed in CP and changes in lipid dynamics in INEPT. The negative peak in the DP difference spectrum is attributed to lipid *trans-gauche* to *all-trans* transition. Sharp peaks between 60 and 105 ppm are attributed to sucrose present in the stacked membrane sample. The figure is modified with permission from Nami *et al.*, BBA Adv. 1, 100015 (2021). Copyright 2021 Elsevier.

dynamics inside npq2 membranes is lower than that in wild-type membranes [Fig. 5(b)].⁸⁹

Influence of membrane stacking

Light-dependent reorganization of thylakoid membranes involves unstacking of stacked grana in high light, which is controlled by phosphorylation. We analyzed the effect of cation-induced unstacking on thylakoid protein and lipid dynamics, causing loss of long-range ordering and migration of LHCII complexes from PSII to PSI.⁸⁷ Unstacking increased the microsecond dynamics of proteins seen as a loss of the protein signal in CP and increased (sub)-nanosecond dynamics of mobile lipids seen as an increase of the lipid signal in INEPT (Fig. 4). The latter was confirmed by additional EPR spin-label experiments performed in the same study that showed increased dynamics of both bulk and annular lipids without changes in their relative fractions. An unexpected

result was that, different from thermally induced dynamics, unstacking particularly affected the dynamics of protein backbone helices with relatively high loss of CP signal intensities at the left wing of the protein CO band (Fig. 4). The result suggests that the release of thylakoid stacking interactions and protein long-range ordering enhances microsecond fluctuations of membrane protein helices. The effect may be similar to a phenomenon that has been observed for purple membranes, where the loss of protein ordering increased microsecond rigid body motions of bacteriorhodopsin transmembrane helices.⁹⁰ Fluorescence decay-associated spectra of the stacked and unstacked membranes showed remarkable similarities with those of intact *Cr* cells kept in state I and state II,⁹¹ suggesting that changes in lipid dynamics and helix motions may occur during membrane state transitions in intact cells in a similar fashion as depicted in Fig. 5(d).

DYNAMIC PROCESSES IN LIVE *Cr* CELLS

NMR spectroscopy is a non-invasive technique and can be applied to endogenous cell compounds, giving access to monitor *in-cell* processes or determine protein structures inside intact cells.⁹² MAS NMR with spectral editing has the advantage that solid-, gel-, and liquid-like phases can be distinguished,²⁵ allowing simultaneous detection of metabolites, macromolecules, and solids in intact organisms. We demonstrated the application of NMR dynamics spectral editing for the investigation of ^{13}C -labeled *Cr* whole cells.^{86,93} Cell viability during NMR experiments was tested,⁹³ concluding that, at moderate rotor spinning frequencies, the *Cr* cells would survive inside the NMR rotors for several hours.

Membrane dynamics

Membrane dynamics of whole *Cr* cells was assessed at different equilibrated temperatures between 3 and 25 °C. For *Cr* algae, the majority of the lipids reside in the cellular and thylakoidal membranes, with about one-third found in the thylakoid membranes and two-third in the cellular membrane.⁹⁴ In contrast to the trend that was observed for isolated thylakoid membranes, in whole cells, the rigidification of lipids was observed at increased temperatures,⁸⁶ as illustrated in Fig. 5(c). Inspection of the DP NMR spectra that are insensitive to dynamics discarded any spectral changes associated with lipid conversion or degradation. Further experiments are required to interpret the results, which suggest that cell protective mechanisms against cold stress may rapidly counteract temperature-dependent changes in membrane fluidity.

Metabolic processes and structural changes

In recent work, we performed MAS NMR with dynamics spectral editing on *Cr* cells that were kept under concentrated-induced, dark anoxia conditions, which resulted in fermentation.⁹³ Whereas previous experiments on thylakoid membranes and cells did not show significant changes in DP spectra over time, here new peaks emerged over a time scale of 0–10 h, depending on the set temperature, which could be assigned to the production of ethanol, CO_2 , and glycerol, formation of glucose products, and slow accumulation of triacylglycerol (TAG) and fatty acids (FFA) (Fig. 6). CP and INEPT spectra provided further insight into structural changes,




FIG. 5. Influence of temperature and stacking conditions on the dynamics of thylakoid membranes and intact *Cr* membranes, as observed with CP- and INEPT-detected MAS NMR. Relative dynamics is illustrated by the lengths of the black arrows. (a) and (b) Temperature effect on protein (blue ovals) and lipid dynamics in wild-type (a) and npq2 (b) thylakoid membranes. (c) Temperature effect on protein and lipid dynamics in intact *Cr* cells. (d) Thylakoid protein and lipid dynamics in stacked and unstacked membranes. Conditions for INEPT and CP detection as described in Ref. 86 are given with S_H , lipid order parameter, and, τ_c , rotational correlation time.

involving the breakdown of starch and galactolipids and progressive enhancement of lipid dynamics over time that we attributed to destabilization or breakdown of cellular or thylakoidal membranes. After 6–12 h depending on the set temperature, the production of ethanol and CO_2 stopped while glycerol production continued. The

time scales correlated with measured cell viability under the same conditions, suggesting that continued glycerol production resulted from non-specific processes occurring in the lysates of death cells. This study demonstrates the proof of principle for the following molecular processes and macromolecular changes in live cells as one




FIG. 6. MAS NMR monitoring of lipid conversion and fermentation in live *Cr* cells. The figure is modified with permission from Nami *et al.*, *Angew. Chem., Int. Ed.* **61**(14), e202117521 (2022). Copyright 2021 Wiley.

of the exciting potential usages of spectral-editing ssNMR. Further applications will benefit from technical improvements to expand cell lifetimes inside MAS NMR rotors and protect them against mechanistic stresses, which are subject for further research.

ACKNOWLEDGMENTS

This work was financially supported, in part, by a Fundamental Research on Matter (FOM) Projectruimte grant of the Netherlands Organization of Scientific Research (NWO) (Grant No. 680.91.15.19) and a CW-VIDI grant of the Netherlands Organization of Scientific Research (NWO) (Grant No. 723.012.103).

AUTHOR DECLARATIONS

Conflict of Interest

The author has no conflicts to disclose.

Author Contributions

Anjali Pandit: Writing – original draft (lead); Writing – review and editing (lead).

DATA AVAILABILITY

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

REFERENCES

- 1 R. E. Blankenship, “The basic principles of photosynthetic energy storage,” in *Molecular Mechanisms of Photosynthesis* (Wiley, 2002), pp. 1–10.
- 2 M. Pribil, M. Labs, and D. Leister, “Structure and dynamics of thylakoids in land plants,” *J. Exp. Bot.* **65**(8), 1955–1972 (2014).
- 3 M. P. Johnson and E. Wientjes, “The relevance of dynamic thylakoid organization to photosynthetic regulation,” *Biochim. Biophys. Acta, Bioenerg.* **1861**(4), 148039 (2020).
- 4 R. Moya, A. C. Norris, T. Kondo, and G. S. Schlau-Cohen, “Observation of robust energy transfer in the photosynthetic protein allophycocyanin using single-molecule pump–probe spectroscopy,” *Nat. Chem.* **14**(2), 153–159 (2022).
- 5 Y.-C. Cheng and G. R. Fleming, “Dynamics of light harvesting in photosynthesis,” *Annu. Rev. Phys. Chem.* **60**(1), 241–262 (2009).
- 6 G. S. Schlau-Cohen, H.-Y. Yang, T. P. J. Krüger, P. Xu, M. Gwizdala, R. van Grondelle, R. Croce, and W. E. Moerner, “Single-molecule identification of quenched and unquenched states of LHCII,” *J. Phys. Chem. Lett.* **6**(5), 860–867 (2015).
- 7 H. Kirchhoff, S. Haferkamp, J. F. Allen, D. B. A. Epstein, and C. W. Mullineaux, “Protein diffusion and macromolecular crowding in thylakoid membranes,” *Plant Physiol.* **146**(4), 1571–1578 (2008).
- 8 E. A. Arsenault, Y. Yoneda, M. Iwai, K. K. Niyogi, and G. R. Fleming, “Vibronic mixing enables ultrafast energy flow in light-harvesting complex II,” *Nat. Commun.* **11**(1), 1460 (2020).
- 9 M. Baldus, A. T. Petkova, J. Herzfeld, and R. G. Griffin, “Cross polarization in the tilted frame: Assignment and spectral simplification in heteronuclear spin systems,” *Mol. Phys.* **95**(6), 1197–1207 (1998).
- 10 V. Ladizhansky, “Applications of solid-state NMR to membrane proteins,” *Biochim. Biophys. Acta, Proteins Proteomics* **1865**(11, Part B), 1577–1586 (2017).
- 11 B. J. Wylie, H. Q. Do, C. G. Borcik, and E. P. Hardy, “Advances in solid-state NMR of membrane proteins,” *Mol. Phys.* **114**(24), 3598–3609 (2016).
- 12 M. Bonaccorsi, T. Le Marchand, and G. Pintacuda, “Protein structural dynamics by Magic-Angle Spinning NMR,” *Curr. Opin. Struct. Biol.* **70**, 34–43 (2021).
- 13 M. J. Duer, “The basics of solid-state NMR,” in *Solid-State NMR Spectroscopy Principles and Applications* (Wiley, 2001), pp. 1–72.
- 14 J. Boisdouvier and L. E. Kay, “Advanced isotopic labeling for the NMR investigation of challenging proteins and nucleic acids,” *J. Biomol. NMR* **71**(3), 115–117 (2018).
- 15 J. Matlahov and P. C. A. van der Wel, “Hidden motions and motion-induced invisibility: Dynamics-based spectral editing in solid-state NMR,” *Methods* **148**, 123–135 (2018).
- 16 M. Baldus and B. H. Meier, “Total correlation spectroscopy in the solid state. The use of scalar couplings to determine the through-bond connectivity,” *J. Magn. Reson., Ser. A* **121**(1), 65–69 (1996).
- 17 B. Elena, A. Lesage, S. Steuernagel, A. Böckmann, and L. Emsley, “Proton to carbon-13 INEPT in solid-state NMR spectroscopy,” *J. Am. Chem. Soc.* **127**(49), 17296–17302 (2005).
- 18 A. Nowacka, N. A. Bongartz, O. H. S. Ollila, T. Nylander, and D. Topgaard, “Signal intensities in ^1H - ^{13}C CP and INEPT MAS NMR of liquid crystals,” *J. Magn. Reson.* **230**, 165–175 (2013).
- 19 J. C. Boatz, T. Piretra, A. Lasorsa, I. Matlahov, J. F. Conway, and P. C. A. van der Wel, “Protofilament structure and supramolecular polymorphism of aggregated mutant huntingtin exon 1,” *J. Mol. Biol.* **432**(16), 4722–4744 (2020).
- 20 H.-K. Lin, J. C. Boatz, I. E. Krabbendam, R. Kodali, Z. Hou, R. Wetzel, A. M. Dolga, M. A. Poirier, and P. C. A. van der Wel, “Fibril polymorphism affects immobilized non-amyloid flanking domains of huntingtin exon 1 rather than its polyglutamine core,” *Nat. Commun.* **8**(1), 15462 (2017).
- 21 M. Li, A. Mandal, V. A. Tyurin, M. DeLucia, J. Ahn, V. E. Kagan, and P. C. A. van der Wel, “Surface-binding to cardiolipin nanodomains triggers cytochrome *c* pro-apoptotic peroxidase activity via localized dynamics,” *Structure* **27**(5), 806–815.e4 (2019).
- 22 M. El Hariri El Nokab, A. Lasorsa, K. O. Sebakhy, F. Picchioni, and P. C. A. van der Wel, “Solid-state NMR spectroscopy insights for resolving different water pools in alginate hydrogels,” *Food Hydrocolloids* **127**, 107500 (2022).
- 23 A. Nowacka, P. C. Mohr, J. Norrman, R. W. Martin, and D. Topgaard, “Polarization transfer solid-state NMR for studying surfactant phase behavior,” *Langmuir* **26**(22), 16848–16856 (2010).
- 24 C. Galvagnion, D. Topgaard, K. Makasewicz, A. K. Buell, S. Linse, E. Sparr, and C. M. Dobson, “Lipid dynamics and phase transition within α -synuclein amyloid fibrils,” *J. Phys. Chem. Lett.* **10**(24), 7872–7877 (2019).
- 25 D. Courtier-Murias, H. Farooq, H. Masoom, A. Botana, R. Soong, J. G. Longstaffe, M. J. Simpson, W. E. Maas, M. Fey, B. Andrew, J. Struppe, H. Hutchins, S. Krishnamurthy, R. Kumar, M. Monette, H. J. Stronks, A. Hume, and A. J. Simpson, “Comprehensive multiphase NMR spectroscopy: Basic experimental approaches to differentiate phases in heterogeneous samples,” *J. Magn. Reson.* **217**, 61–76 (2012).
- 26 Y. L. Mobarhan, B. Fortier-McGill, R. Soong, W. E. Maas, M. Fey, M. Monette, H. J. Stronks, S. Schmidt, H. Heumann, W. Norwood, and A. J. Simpson, “Comprehensive multiphase NMR applied to a living organism,” *Chem. Sci.* **7**(8), 4856–4866 (2016).
- 27 L. Shen, Z. Huang, S. Chang, W. Wang, J. Wang, T. Kuang, G. Han, J.-R. Shen, and X. Zhang, “Structure of a $\text{C}_2\text{S}_2\text{M}_2\text{N}_2$ -type PSII-LHCII supercomplex from the green alga *Chlamydomonas reinhardtii*,” *Proc. Natl. Acad. Sci. U. S. A.* **116**(42), 21246–21255 (2019).
- 28 X. Sheng, Z. Liu, E. Kim, and J. Minagawa, “Plant and algal PSII-LHCII supercomplexes: Structure, evolution and energy transfer,” *Plant Cell Physiol.* **62**(7), 1108–1120 (2021).
- 29 M. Fan, M. Li, Z. Liu, P. Cao, X. Pan, H. Zhang, X. Zhao, J. Zhang, and W. Chang, “Crystal structures of the PsbS protein essential for photoprotection in plants,” *Nat. Struct. Mol. Biol.* **22**(9), 729–735 (2015).
- 30 Z. Liu, H. Yan, K. Wang, T. Kuang, J. Zhang, L. Gui, X. An, and W. Chang, “Crystal structure of spinach major light-harvesting complex at 2.72 Å resolution,” *Nature* **428**(6980), 287–292 (2004).
- 31 T. A. Egorova-Zachernyuk, J. Hollander, N. Fraser, P. Gast, A. J. Hoff, R. Cogdell, H. J. M. de Groot, and M. Baldus, “MAS NMR on a uniformly [^{13}C , ^{15}N] labeled LH2 light-harvesting complex from *Rhodospseudomonas Acidophila* 10050 at ultra-high magnetic fields,” in *Perspectives on Solid State NMR in Biology*, edited by S. R. Kiihne and H. J. M. de Groot (Springer Netherlands, Dordrecht, 2001), pp. 171–183.
- 32 A. Pandit, F. Buda, A. J. van Gammeren, S. Ganapathy, and H. J. M. de Groot, “Selective chemical shift assignment of bacteriochlorophyll *a* in uniformly

- [^{13}C - ^{15}N]-labeled light-harvesting 1 complexes by solid-state NMR in ultrahigh magnetic field," *J. Phys. Chem. B* **114**(18), 6207–6215 (2010).
- ³³S. Ganapathy, A. J. van Gammeren, F. B. Hulsbergen, and H. J. M. de Groot, "Probing secondary, tertiary, and quaternary structure along with protein-cofactor interactions for a helical transmembrane protein complex through ^1H spin diffusion with MAS NMR spectroscopy," *J. Am. Chem. Soc.* **129**(6), 1504–1505 (2007).
- ³⁴Alia, J. Matysik, C. Soede-Huijbregts, M. Baldus, J. Raap, J. Lugtenburg, P. Gast, H. J. van Gorkom, A. J. Hoff, and H. J. M. de Groot, "Ultrahigh field MAS NMR dipolar correlation spectroscopy of the histidine residues in light-harvesting complex II from photosynthetic bacteria reveals partial internal charge transfer in the B850/His complex," *J. Am. Chem. Soc.* **123**(20), 4803–4809 (2001).
- ³⁵Alia, J. Matysik, I. de Boer, P. Gast, H. J. van Gorkom, and H. J. de Groot, "Heteronuclear 2D (^1H - ^{13}C) MAS NMR resolves the electronic structure of coordinated histidines in light-harvesting complex II: Assessment of charge transfer and electronic delocalization effect," *J. Biomol. NMR* **28**(2), 157–164 (2004).
- ³⁶A. Diller, E. Roy, P. Gast, H. J. van Gorkom, H. J. M. de Groot, C. Glaubitz, G. Jeschke, J. Matysik, and A. Alia, " ^{15}N photochemically induced dynamic nuclear polarization magic-angle spinning NMR analysis of the electron donor of photosystem II," *Proc. Natl. Acad. Sci. U. S. A.* **104**(31), 12767–12771 (2007).
- ³⁷J. C. Zill, M. Kansy, R. Goss, A. Alia, C. Wilhelm, H. J. Matysik, " ^{15}N photo-CIDNP MAS NMR on both photosystems and magnetic field-dependent ^{13}C photo-CIDNP MAS NMR in photosystem II of the diatom *Phaeodactylum tricornutum*," *Photosynth. Res.* **140**(2), 151–171 (2019).
- ³⁸G. J. Janssen, E. Daviso, M. van Son, H. J. de Groot, A. Alia, and J. Matysik, "Observation of the solid-state photo-CIDNP effect in entire cells of cyanobacteria *Synechocystis*," *Photosynth. Res.* **104**(2), 275–282 (2010).
- ³⁹S. Ganapathy, G. T. Oostergetel, P. K. Wawrzyniak, M. Reus, A. Gomez Maqueo Chew, F. Buda, E. J. Boekema, D. A. Bryant, A. R. Holzwarth, and H. J. M. de Groot, "Alternating syn-anti bacteriochlorophylls form concentric helical nanotubes in chlorosomes," *Proc. Natl. Acad. Sci. U. S. A.* **106**(21), 8525–8530 (2009).
- ⁴⁰S. Ganapathy, S. Sengupta, P. K. Wawrzyniak, V. Huber, F. Buda, U. Baumeister, F. Würthner, and H. J. M. de Groot, "Zinc chlorins for artificial light-harvesting self-assemble into antiparallel stacks forming a microcrystalline solid-state material," *Proc. Natl. Acad. Sci. U. S. A.* **106**(28), 11472–11477 (2009).
- ⁴¹A. Pandit, K. Ocakoglu, F. Buda, T. van Marle, A. R. Holzwarth, and H. J. M. de Groot, "Structure determination of a bio-inspired self-assembled light-harvesting antenna by solid-state NMR and molecular modeling," *J. Phys. Chem. B* **117**(38), 11292–11298 (2013).
- ⁴²X. Li, F. Buda, H. J. M. de Groot, and G. J. A. Sevink, "Contrasting modes of self-assembly and hydrogen-bonding heterogeneity in chlorosomes of *Chlorobaculum tepidum*," *J. Phys. Chem. C* **122**(26), 14877–14888 (2018).
- ⁴³X. Li, F. Buda, H. J. M. de Groot, and G. J. A. Sevink, "Molecular insight in the optical response of tubular chlorosomal assemblies," *J. Phys. Chem. C* **123**(26), 16462–16478 (2019).
- ⁴⁴T. P. J. Krüger, C. Iliaia, M. P. Johnson, A. V. Ruban, E. Papagiannakis, P. Horton, and R. van Grondelle, "Controlled disorder in plant light-harvesting complex II explains its photoprotective role," *Biophys. J.* **102**(11), 2669–2676 (2012).
- ⁴⁵A. V. Ruban, "Light harvesting control in plants," *FEBS Lett.* **592**(18), 3030–3039 (2018).
- ⁴⁶J. Standfuss, A. C. Terwisscha van Scheltinga, M. Lamborghini, and W. Kühlbrandt, "Mechanisms of photoprotection and nonphotochemical quenching in pea light-harvesting complex at 2.5 Å resolution," *EMBO J.* **24**(5), 919–928 (2005).
- ⁴⁷X. Pan, M. Li, T. Wan, L. Wang, C. Jia, Z. Hou, X. Zhao, J. Zhang, and W. Chang, "Structural insights into energy regulation of light-harvesting complex CP29 from spinach," *Nat. Struct. Mol. Biol.* **18**(3), 309–315 (2011).
- ⁴⁸N. Liguori, R. Croce, S. J. Marrink, and S. Thallmair, "Molecular dynamics simulations in photosynthesis," *Photosynth. Res.* **144**(2), 273–295 (2020).
- ⁴⁹N. Liguori, X. Periole, S. J. Marrink, and R. Croce, "From light-harvesting to photoprotection: Structural basis of the dynamic switch of the major antenna complex of plants (LHCII)," *Sci. Rep.* **5**(1), 15661 (2015).
- ⁵⁰S. Thallmair, P. A. Vainikka, and S. J. Marrink, "Lipid fingerprints and cofactor dynamics of light-harvesting complex II in different membranes," *Biophys. J.* **116**(8), 1446–1455 (2019).
- ⁵¹F. J. van Eerden, D. H. de Jong, A. H. de Vries, T. A. Wassenaar, and S. J. Marrink, "Characterization of thylakoid lipid membranes from cyanobacteria and higher plants by molecular dynamics simulations," *Biochim. Biophys. Acta, Biomembr.* **1848**(6), 1319–1330 (2015).
- ⁵²V. Daskalakis, S. Papadatos, and U. Kleinekathöfer, "Fine tuning of the photosystem II major antenna mobility within the thylakoid membrane of higher plants," *Biochim. Biophys. Acta, Biomembr.* **1861**(12), 183059 (2019).
- ⁵³V. Daskalakis, S. Papadatos, and T. Stergiannakos, "The conformational phase space of the photoprotective switch in the major light harvesting complex II," *Chem. Commun.* **56**(76), 11215–11218 (2020).
- ⁵⁴N. Liguori, S. R. R. Campos, A. M. Baptista, and R. Croce, "Molecular anatomy of plant photoprotective switches: The sensitivity of PsbS to the environment, residue by residue," *J. Phys. Chem. Lett.* **10**(8), 1737–1742 (2019).
- ⁵⁵C. Dockter, A. H. Müller, C. Dietz, A. Volkov, Y. Polyhach, G. Jeschke, and H. Paulsen, "Rigid core and flexible terminus: Structure of solubilized light-harvesting chlorophyll a/b complex (LHCII) measured by EPR," *J. Biol. Chem.* **287**(4), 2915–2925 (2012).
- ⁵⁶M. Golub, L. Rusevich, K.-D. Irrgang, and J. Pieper, "Rigid versus flexible protein matrix: Light-harvesting complex II exhibits a temperature-dependent phonon spectral density," *J. Phys. Chem. B* **122**(28), 7111–7121 (2018).
- ⁵⁷K. Vrandečić, M. Rätsep, L. Wilk, L. Rusevich, M. Golub, M. Reppert, K.-D. Irrgang, W. Kühlbrandt, and J. Pieper, "Protein dynamics tunes excited state positions in light-harvesting complex II," *J. Phys. Chem. B* **119**(10), 3920–3930 (2015).
- ⁵⁸M. Golub, H. Lokstein, D. Soloviov, A. Kuklin, D. C. F. Wieland, and J. Pieper, "Light-harvesting complex II adopts different quaternary structures in solution as observed using small-angle scattering," *J. Phys. Chem. Lett.* **13**(5), 1258–1265 (2022).
- ⁵⁹F. Azadi-Chegeni, M. E. Ward, G. Perin, D. Simionato, T. Morosinotto, M. Baldus, and A. Pandit, "Conformational dynamics of light-harvesting complex II in a native membrane environment," *Biophys. J.* **120**(2), 270–283 (2021).
- ⁶⁰T. Gopinath and G. Veglia, "Probing membrane protein ground and conformationally excited states using dipolar- and J-coupling mediated MAS solid state NMR experiments," *Methods* **148**, 115–122 (2018).
- ⁶¹M. E. Ward, E. Ritz, M. A. M. Ahmed, V. V. Bamm, G. Harauz, L. S. Brown, and V. Ladizhansky, "Proton detection for signal enhancement in solid-state NMR experiments on mobile species in membrane proteins," *J. Biomol. NMR* **63**(4), 375–388 (2015).
- ⁶²Y. Luo, S. Xiang, P. J. Hooikaas, L. van Bezouwen, A. S. Jijumon, C. Janke, F. Förster, A. Akhmanova, and M. Baldus, "Direct observation of dynamic protein interactions involving human microtubules using solid-state NMR spectroscopy," *Nat. Commun.* **11**(1), 18 (2020).
- ⁶³P. C. A. van der Wel, "New applications of solid-state NMR in structural biology," *Emerging Top. Life Sci.* **2**(1), 57–67 (2018).
- ⁶⁴F. Azadi-Chegeni, S. Thallmair, M. E. Ward, G. Perin, S. J. Marrink, M. Baldus, T. Morosinotto, and A. Pandit, "Protein dynamics and lipid affinity of monomeric, zeaxanthin-binding LHCII in thylakoid membranes," *Biophys. J.* **121**(3), 396–409 (2022).
- ⁶⁵T. Kondo, A. Pinnola, W. J. Chen, L. Dall'Osto, R. Bassi, and G. S. Schlau-Cohen, "Single-molecule spectroscopy of LHCSR1 protein dynamics identifies two distinct states responsible for multi-timescale photosynthetic photoprotection," *Nat. Chem.* **9**(8), 772–778 (2017).
- ⁶⁶E. Cignoni, M. Lapillo, L. Cupellini, S. Acosta-Gutiérrez, F. L. Gervasio, and B. Mennucci, "A different perspective for nonphotochemical quenching in plant antenna complexes," *Nat. Commun.* **12**(1), 7152 (2021).
- ⁶⁷L. Cupellini, D. Calvani, D. Jacquemin, and B. Mennucci, "Charge transfer from the carotenoid can quench chlorophyll excitation in antenna complexes of plants," *Nat. Commun.* **11**(1), 662 (2020).
- ⁶⁸C. Gray, T. Wei, T. Polivka, V. Daskalakis, and C. D. P. Duffy, "Trivial excitation energy transfer to carotenoids is an unlikely mechanism for non-photochemical quenching in LHCII," *Front. Plant Sci.* **12**, 797373 (2022).
- ⁶⁹K. Sunku, H. J. M. de Groot, and A. Pandit, "Insights into the photoprotective switch of the major light-harvesting complex II (LHCII): A preserved core of arginine-glutamate interlocked helices complemented by adjustable loops," *J. Biol. Chem.* **288**(27), 19796–19804 (2013).

- ⁷⁰A. Pandit, M. Reus, T. Morosinotto, R. Bassi, A. R. Holzwarth, and H. J. M. de Groot, "An NMR comparison of the light-harvesting complex II (LHCII) in active and photoprotective states reveals subtle changes in the chlorophyll *a* ground-state electronic structures," *Biochim. Biophys. Acta, Bioenerg.* **1827**(6), 738–744 (2013).
- ⁷¹X. Sheng, X. Liu, P. Cao, M. Li, and Z. Liu, "Structural roles of lipid molecules in the assembly of plant PSII–LHCII supercomplex," *Biophys. Rep.* **4**(4), 189–203 (2018).
- ⁷²E. Kim, A. Watanabe, C. D. P. Duffy, A. V. Ruban, and J. Minagawa, "Multimeric and monomeric photosystem II supercomplexes represent structural adaptations to low- and high-light conditions," *J. Biol. Chem.* **295**(43), 14537–14545 (2020).
- ⁷³Z. Huang, L. Shen, W. Wang, Z. Mao, X. Yi, T. Kuang, J.-R. Shen, X. Zhang, and G. Han, "Structure of photosystem I-LHCI-LHCII from the green alga *Chlamydomonas reinhardtii* in state 2," *Nat. Commun.* **12**(1), 1100 (2021).
- ⁷⁴P. Albanese, S. Tamara, G. Saracco, R. A. Scheltema, and C. Pagliano, "How paired PSII–LHCII supercomplexes mediate the stacking of plant thylakoid membranes unveiled by structural mass-spectrometry," *Nat. Commun.* **11**(1), 1361 (2020).
- ⁷⁵S. B. Krumova, C. Dijkema, P. de Waard, H. Van As, G. Garab, and H. van Amerongen, "Phase behavior of phosphatidylglycerol in spinach thylakoid membranes as revealed by ³¹P-NMR," *Biochim. Biophys. Acta* **1778**(4), 997–1003 (2008).
- ⁷⁶G. Garab, B. Ughy, P. d. Waard, P. Akhtar, U. Javornik, C. Kotakis, P. Šket, V. Karlický, Z. Materová, V. Špunda, J. Plavec, H. van Amerongen, L. Vígh, H. V. As, and P. H. Lambrev, "Lipid polymorphism in chloroplast thylakoid membranes—As revealed by ³¹P-NMR and time-resolved merocyanine fluorescence spectroscopy," *Sci. Rep.* **7**(1), 13343 (2017).
- ⁷⁷H. Kirchhoff, "Architectural switches in plant thylakoid membranes," *Photosynth. Res.* **116**(2–3), 481–487 (2013).
- ⁷⁸E. Janik, W. Maksymiec, W. Grudziński, and W. I. Gruszecki, "Strong light-induced reorganization of pigment–protein complexes of thylakoid membranes in rye (spectroscopic study)," *J. Plant Physiol.* **169**(1), 65–71 (2012).
- ⁷⁹G. Nagy, R. Ünnep, O. Zsiros, R. Tokutsu, K. Takizawa, L. Porcar, L. Moyet, D. Petroutsos, G. Garab, G. Finazzi, and J. Minagawa, "Chloroplast remodeling during state transitions in *Chlamydomonas reinhardtii* as revealed by noninvasive techniques in vivo," *Proc. Natl. Acad. Sci. U. S. A.* **111**(13), 5042–5047 (2014).
- ⁸⁰H. Kirchhoff, C. Hall, M. Wood, M. Herbstová, O. Tsaabari, R. Nevo, D. Charuvi, E. Shimoni, and Z. Reich, "Dynamic control of protein diffusion within the granal thylakoid lumen," *Proc. Natl. Acad. Sci. U. S. A.* **108**(50), 20248–20253 (2011).
- ⁸¹H. Kirchhoff, "Diffusion of molecules and macromolecules in thylakoid membranes," *Biochim. Biophys. Acta, Bioenerg.* **1837**(4), 495–502 (2014).
- ⁸²L.-R. Stingaciu, H. O'Neill, M. Liberton, V. S. Urban, H. B. Pakrasi, and M. Ohl, "Revealing the dynamics of thylakoid membranes in living cyanobacterial cells," *Sci. Rep.* **6**(1), 19627 (2016).
- ⁸³R. Ünnep, G. Nagy, M. Markó, and G. Garab, "Monitoring thylakoid ultrastructural changes in vivo using small-angle neutron scattering," *Plant Physiol. Biochem.* **81**, 197–207 (2014).
- ⁸⁴T. M. Ferreira, O. H. S. Ollila, R. Pigliapochi, A. P. Dabkowska, and D. Topgaard, "Model-free estimation of the effective correlation time for C–H bond reorientation in amphiphilic bilayers: ¹H–¹³C solid-state NMR and MD simulations," *J. Chem. Phys.* **142**(4), 044905 (2015).
- ⁸⁵F. Azadi Chegeni, G. Perin, K. B. Sai Sankar Gupta, D. Simionato, T. Morosinotto, and A. Pandit, "Protein and lipid dynamics in photosynthetic thylakoid membranes investigated by *in-situ* solid-state NMR," *Biochim. Biophys. Acta* **1857**(12), 1849–1859 (2016).
- ⁸⁶F. Azadi-Chegeni, C. Schiphorst, and A. Pandit, "In vivo NMR as a tool for probing molecular structure and dynamics in intact *Chlamydomonas reinhardtii* cells," *Photosynth. Res.* **135**(1–3), 227–237 (2018).
- ⁸⁷F. Nami, L. Tian, M. Huber, R. Croce, and A. Pandit, "Lipid and protein dynamics of stacked and cation-depletion induced unstacked thylakoid membranes," *BBA Adv. I*, 100015 (2021).
- ⁸⁸R. N. Purusottam, L. Sénicourt, J.-J. Lacapère, and P. Tekely, "Probing the gel to liquid-crystalline phase transition and relevant conformation changes in liposomes by ¹³C magic-angle spinning NMR spectroscopy," *Biochim. Biophys. Acta, Biomembr.* **1848**(12), 3134–3139 (2015).
- ⁸⁹M. Havaux, "Carotenoids as membrane stabilizers in chloroplasts," *Trends Plant Sci.* **3**(4), 147–151 (1998).
- ⁹⁰S. Tuzi, A. Naito, and H. Saitō, "Local protein structure and dynamics at kinked transmembrane α -helices of [^{1-¹³C}]Pro-labeled bacteriorhodopsin as revealed by site-directed solid-state ¹³C NMR," *J. Mol. Struct.* **654**(1–3), 205–214 (2003).
- ⁹¹L. M. Włodarczyk, E. Dinc, R. Croce, and J. P. Dekker, "Excitation energy transfer in *Chlamydomonas reinhardtii* deficient in the PSI core or the PSII core under conditions mimicking state transitions," *Biochim. Biophys. Acta, Bioenerg.* **1857**(6), 625–633 (2016).
- ⁹²E. Luchinat and L. Banci, "In-cell NMR: A topical review," *IUCr* **4**(Pt. 2), 108–118 (2017).
- ⁹³F. Nami, M. J. Ferraz, T. Bakkum, J. M. F. G. Aerts, and A. Pandit, "Real-time NMR recording of fermentation and lipid metabolism processes in live microalgae cells," *Angew. Chem., Int. Ed.* **61**(14), e202117521 (2022).
- ⁹⁴D. R. Janero and R. Barrnett, "Cellular and thylakoid-membrane glycolipids of *Chlamydomonas reinhardtii* 137+," *J. Lipid Res.* **22**(7), 1119–1125 (1981).