



Universiteit
Leiden
The Netherlands

Gene-educational attainment interactions in a multi-ancestry genome-wide meta-analysis identify novel blood pressure loci

Fuentes, L. de las; Sung, Y.J.; Noordam, R.; Winkler, T.; Feitosa, M.F.; Schwander, K.; ... ; Lifelines Cohort Study

Citation

Fuentes, L. de las, Sung, Y. J., Noordam, R., Winkler, T., Feitosa, M. F., Schwander, K., ... Fornage, M. (2020). Gene-educational attainment interactions in a multi-ancestry genome-wide meta-analysis identify novel blood pressure loci. *Molecular Psychiatry*, 26, 2111-2125. doi:10.1038/s41380-020-0719-3

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3185192>

Note: To cite this publication please use the final published version (if applicable).



Gene-educational attainment interactions in a multi-ancestry genome-wide meta-analysis identify novel blood pressure loci

Lisa de las Fuentes ^{1,2} et al.

Received: 21 August 2019 / Revised: 19 March 2020 / Accepted: 24 March 2020 / Published online: 5 May 2020
© The Author(s), under exclusive licence to Springer Nature Limited 2020

Abstract

Educational attainment is widely used as a surrogate for socioeconomic status (SES). Low SES is a risk factor for hypertension and high blood pressure (BP). To identify novel BP loci, we performed multi-ancestry meta-analyses accounting for gene-educational attainment interactions using two variables, “Some College” (yes/no) and “Graduated College” (yes/no). Interactions were evaluated using both a 1 degree of freedom (DF) interaction term and a 2DF joint test of genetic and interaction effects. Analyses were performed for systolic BP, diastolic BP, mean arterial pressure, and pulse pressure. We pursued genome-wide interrogation in Stage 1 studies ($N = 117\,438$) and follow-up on promising variants in Stage 2 studies ($N = 293\,787$) in five ancestry groups. Through combined meta-analyses of Stages 1 and 2, we identified 84 known and 18 novel BP loci at genome-wide significance level ($P < 5 \times 10^{-8}$). Two novel loci were identified based on the 1DF test of interaction with educational attainment, while the remaining 16 loci were identified through the 2DF joint test of genetic and interaction effects. Ten novel loci were identified in individuals of African ancestry. Several novel loci show strong biological plausibility since they involve physiologic systems implicated in BP regulation. They include genes involved in the central nervous system-adrenal signaling axis (*ZDHH17*, *CADPS*, *PIK3C2G*), vascular structure and function (*GNB3*, *CDON*), and renal function (*HAS2* and *HAS2-AS1*, *SLIT3*). Collectively, these findings suggest a role of educational attainment or SES in further dissection of the genetic architecture of BP.

Introduction

Educational attainment is among the most widely used indices of socioeconomic status (SES) in epidemiologic studies [1,2]. Multiple studies have demonstrated a step-wise decline in all-cause mortality with increasing levels of education [1]. Compared with other measures of SES, such as income and occupation, the use of educational attainment has several advantages: it is stable after young adulthood, simple to capture, has a low nonresponse rate, and is not affected by poor health in adulthood [1,3]. Furthermore, the relationship between educational attainment and cardiovascular disease traits tend to be more consistent and

stronger [4]. Higher educational attainment is related to improved health efficacy (such as preventive health behaviors and problem-solving capacity), improved access to health care, and more favorable socio-psychological conditions (such as personal control and social support) [2,3].

Several variables of educational attainment investigated in epidemiologic studies in relation to cardiovascular risk traits include continuous variables (such as years of education) and various partitions (such as completing high school or completing college degree) [5–11]. Low educational attainment is related to high blood pressure (BP) and increased hypertension (HTN) risk as evidenced in a meta-analysis of 51 studies across 20 countries [3]. Educational attainment is also related to coronary artery disease [12], coronary calcification [13], and other cardiovascular risk traits including metabolic syndrome [10], lipid levels [9,10,14], smoking behavior [12,15], salt intake [16,17], and leisure-time physical activity [18]. Furthermore, the genetic effects on HTN may vary as a function of educational attainment. For example, a heritability study of EUR-ancestry male twins showed higher heritability of HTN at higher education levels ($h^2 = 0.63$ with >14 years of education compared with $h^2 = 0.46$ with ≤ 14

Supplementary information The online version of this article (<https://doi.org/10.1038/s41380-020-0719-3>) contains supplementary material, which is available to authorized users.

✉ Lisa de las Fuentes
lfuentes@wustl.edu

Extended author information available on the last page of the article

years of education) [19], suggesting interactions between genes and educational attainment.

While genome-wide association studies have investigated the genetic contributions to educational attainment [6], there has been no comprehensive effort to examine the role played by gene-environment interactions in BP using educational attainment as the environmental exposure. Within the CHARGE Gene-Lifestyle Interactions Working Group [20], we performed genome-wide meta-analysis of systolic BP (SBP), diastolic BP (DBP), mean arterial pressure (MAP), and pulse pressure (PP), accounting for gene-educational attainment interactions. Based on the availability of data across participating studies, we considered two educational attainment variables, “Some College” (yes/no, for any education beyond high school) and “Graduated College” (yes/no, for completing a 4-year college degree). Herein, we report our findings based on up to 411,225 individuals from five ancestry groups.

Subjects and methods

Participating studies

We performed our analysis in two stages (Fig. 1). A total of 42 cohorts including 117 438 men and women (aged 18–80 years) from European (EUR), African (AFR), Asian (ASN), Hispanic (HIS), and Brazilian (BRZ) ancestries contributed to Stage 1 genome-wide interaction analyses (Table S1). An additional 49 cohorts including 293 787 individuals contributed to Stage 2 analyses of top single nucleotide variants (SNVs, also including a small number of insertion and deletion [indels] variants) selected from Stage 1 (Table S2). Participating studies are described in Supplementary

Material. Since discoveries to date are largely from EUR populations, considerable effort was made to recruit most of the available non-EUR cohorts into Stage 1. Each study obtained informed consent from participants and approval from the appropriate institutional review boards.

Phenotype and lifestyle variables

We performed our analysis for SBP, DBP, MAP, and PP. After computing SBP and DBP when multiple measurements were available, we adjusted for antihypertensive medication use by adding 15 mmHg and 10 mmHg to SBP and DBP, respectively [21]. After medication adjustment, MAP was computed as $(SBP + 2DBP)/3$, and PP was computed as $SBP - DBP$ (SBP–DBP). To reduce the influence of possible outliers, Winsorizing was performed for each BP value that was more than six standard deviations away from the mean. Descriptive statistics for these four BP traits are presented in Tables S3 and S4. For educational attainment, two dichotomous variables were created. The first variable, “Some College” (SomeCol), was coded as 1 if the subject received any education beyond high school, including vocational school (and as 0 if no education beyond high school). The second variable, “Graduated College” (GradCol), was coded as 1 if the subject completed at least a 4-year college degree (and as 0 for any education less than a 4-year degree). Subjects with missing data for BP, education attainment, or any covariates were excluded from analysis.

Genotype data

Genotyping was performed by each participating study using Illumina (San Diego, CA, USA) or Affymetrix (Santa Clara, CA, USA) genotyping arrays. Imputation was performed using the 1000 Genomes Project [22] Phase I Integrated Release Version 3 Haplotypes (2010-11 data freeze, 2012-03-14 haplotypes) as a reference panel, in most cohorts. Information on genotype and imputation for each study is presented in Tables S5 and S6.

Analysis methods

Each study performed association analyses using the following model:

$$E(Y) = \beta_0 + \beta_G G + \beta_E \text{Educ} + \beta_{GE} G \times \text{Educ} + \beta_C C,$$

where Y is the BP variable (SBP, DBP, MAP, or PP value), Educ is the educational variable (SomeCol or GradCol), and G is the dosage of the imputed genetic variant coded additively from 0 to 2. C is the vector of covariates, including age, sex, field center (for multi-center studies), and principal components. In addition, studies in Stage 1 performed




Fig. 1 Study design with summary of data included in this study. Educ: education status (considering either SomeCol or GradCol status separately); PC principal component, EUR European, AFR African, ASN Asian, HIS Hispanic, BRZ Brazilian, SNV single nucleotide variant, TRANS trans-ancestry (i.e., combining all ancestry groups through meta-analysis).

association analysis using the following genetic main-effect model with education attainment:

$$E(Y) = \beta_0 + \beta_G G + \beta_E \text{Educ} + \beta_C C.$$

Each study provided the estimated SNV effect (β_G), estimated SNV-educational attainment interaction effect (β_{GE}), their robust standard errors (SEs), and a robust estimate of the covariance between β_G and β_{GE} . We performed meta-analysis using the 1 degree of freedom (DF) test of the interaction effect (β_{GE}) and 2DF test of both SNV (β_G) and interaction effects (β_{GE}). Inverse-variance weighted meta-analysis was performed for the 1DF test and the joint meta-analysis of Manning et al. [23] for the 2DF test, both using METAL [24]. In Stage 1 EUR, AFR, ASN meta-analyses, variants were included if they were available in more than 5,000 samples or at least three cohorts (these filters were not applied to BRZ or HIS because of the fewness of cohorts included in these meta-analyses). We applied genomic control correction [25] twice in Stage 1, first for study-specific GWAS results and again for meta-analysis results. Genome-wide significant ($P < 5 \times 10^{-8}$) and suggestive ($P < 1 \times 10^{-6}$) variants in Stage 1 were taken forward into Stage 2 analysis. Genomic control correction was not applied to the Stage 2 results as association test was performed for select variants. Results presented reflect meta-analyses combining Stages 1 and 2. Loci were defined by physical distance (± 1 Mb around the index SNV of the respective locus).

Quality control (QC)

Each participating cohort in Stage 1 excluded variants with minor allele frequency (MAF) $< 1\%$. We performed extensive QC using the R package EasyQC [26] for all cohort-specific and meta-analysis results. For Stages 1 and 2, we excluded all variants with imputation quality measure < 0.5 . In addition, to remove unstable study-specific results that reflected small sample size, low minor allele count (MAC), or low imputation quality, we excluded variants for which the minimum of (MAC0, MAC1) \times imputation quality < 20 , where MAC0 and MAC1 are the MAC in the two education strata (Educ = 0 and Educ = 1). The allele frequencies provided by each cohort were compared against those from the relevant ancestry-specific 1000 Genomes reference panel. Marker names were harmonized to ensure consistency across cohorts. In addition, we visually compared summary statistics (e.g., mean, median, standard deviation, inter-quartile range) of all effect estimates, SEs, and P values. We examined SE-N plots [26] and quantile-quantile (QQ) plots to reveal issues with trait transformation or other analytical problems. Any problems encountered during QC steps were resolved through communication with the analysts from the participating studies. More detailed information about the QC steps, including major

QC problems encountered and how they were resolved, are described elsewhere [20,27].

Characterization of functional roles

A suite of tools implemented in FUMA GWAS [28] was used to identify functional roles for the index variants and nearby variants in linkage disequilibrium (LD; $r^2 \geq 0.2$) in each of the novel BP loci. LD information was obtained from the 1000 Genomes Project reference genome for the ancestry with the most significant ancestry-specific association. If the most significant association was in trans-ancestry analyses, the reference genome for the ancestry with the next most significant association was used instead [29]. One index insertion/deletion locus was not identified in any of the reference genomes by FUMA and therefore not detailed. Nearest gene annotations were limited to protein coding, long non-coding RNAs, and non-coding RNAs within 10 kb of index variants and variants in LD ($r^2 \geq 0.2$) with the index variant [30].

For the index and LD variants, we used the RegulomeDB score [31], which reflects a summary of annotations with known and predicted regulatory elements such as DNAase hypersensitivity, binding sites of transcription factors, and promoter regions, and Combined Annotation Dependent Depletion (CADD) [32] scores, which predict deleteriousness of variants. The 15-core chromatin state (ChromHMM) [33,34] was assessed for 129 epigenomes (labeled E001-E129) to identify histone modifications consistent with epigenetic regulation of gene expression. Expression quantitative trait loci (eQTLs) were determined using the GTEx_v7 database [35] for index and LD variants. Using nearest-gene annotations, FUMA GWAS was used to generate tissue-specific gene expression data (GTEx V7 dataset, 53 tissue types); significance was determined as a Benjamini–Hochberg false discovery rate (FDR) < 0.05 .

Results

Overview

We performed a meta-analysis of gene-education interactions on SBP, DBP, MAP, and PP in two stages (Fig. 1). In Stage 1, we pursued genome-wide interrogation in 117 438 individuals of EUR, AFR, ASN, HIS, and BRZ ancestries (summary information, Table 1). After extensive quality control (QC), we performed genome-wide meta-analyses at ~ 18.8 million SNVs and a small number of insertion and deletion (indels) variants imputed using the 1000 Genomes Project reference panel (QQ plots, Fig. S1). Through the 1DF test of the interaction effect and the 2DF joint test of the SNV and interaction effects, we identified 1481

Table 1 Basic characteristics of cohorts in Stages 1 and 2 in each ancestry.

Stage	Max. N	Some College? (Yes/No)		Grad. College? (Yes/No)	% Male	% HT	% HT Meds	Age	SBP		DBP		MAP		PP		
		N	% Yes						Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Stage 1																	
EUR	84,276	47,870	57.7	79,040	32.9	28.7	17.0	54.7	7.8	130.3	19.1	78.3	11.2	95.6	12.6	52.0	13.1
AFR	16,665	16,665	48.8	16,665	40.6	52.4	40.3	51.5	9.2	134.4	23.2	81.2	13.4	98.9	15.5	53.2	16.1
ASN	11,352	11,352	10.0	748	53.0	49.6	29.5	56.8	9.4	139.3	16.1	80.2	11.1	99.9	13.5	59.1	16.2
BRZ	3689	3689	36.8	3689	45.6	29.0	7.6	34.5	3.8	123.4	15.3	76.6	10.0	92.2	11.1	46.8	10.2
HIS	1456	1456	35.2	1456	48.4	41.4	33.3	60.8	9.73	131.2	24.8	74.9	11.7	93.6	14.9	56.4	18.5
Stage 1 Total	117,438	81,032	47.8	101,598	36.5	34.2	21.4	53.9	8.1	131.6	19.4	78.8	11.5	96.4	13.1	52.8	13.8
Stage 2																	
EUR	264,052	242,524	54.3	257,037	47.6	48.5	21.7	54.9	8.9	138.5	20.2	83.3	11.4	101.7	13.4	55.2	13.6
AFR	7198	7198	20.8	7198	40.4	56.8	44.1	54.7	9.5	137.6	21.8	83.8	12.9	101.7	14.8	53.8	14.9
ASN	10,906	10,906	21.4	5947	40.1	37.4	24.9	55.8	9.0	134.2	23.1	80.7	12.8	98.6	15.4	53.5	14.8
HIS	11,631	11,631	35.5	11,631	41.1	25.4	15.0	45.3	13.6	123.6	19.7	75.0	11.8	91.2	13.6	48.6	13.1
Stage 2 Total	293,787	272,259	51.3	281,813	46.6	47.4	22.1	54.6	9.1	137.7	20.3	82.9	11.5	101.2	13.6	54.8	13.7
TOTAL	411,225	353,291	50.5	383,411	43.7	43.6	21.9	54.4	8.8	136.0	20.1	81.7	11.5	99.8	13.4	54.2	13.7

The cell entries for the covariates and BP traits corresponds to sample-size weighted averages across all cohorts in each category.

genome-wide significant ($P < 5 \times 10^{-8}$) and 3309 suggestive ($P < 1 \times 10^{-6}$) variants associated with any BP trait in any ancestry and/or in trans-ancestry analysis. All significant and suggestive variants were tested for association in 293 787 additional individuals of EUR, AFR, ASN, and HISP ancestries in Stage 2.

We performed meta-analyses combining Stages 1 and 2 (Manhattan Plots, Fig. S2). We identified 84 known BP loci. This includes 82 loci identified through main-effect only analyses [36–41], including 18 recently reported by Hoffmann et al. [42], Evangelou et al. [43], and Giri et al. [44]; and two loci (*TFAP2A* and *PCDH9*) recently reported by our consortium through gene-smoking and gene-alcohol interaction analyses [27,45], which suggest the inter-correlated nature of the various lifestyle traits.

We identified 18 novel genome-wide significant loci ($P < 5 \times 10^{-8}$) located at least 1 Mb away from any known BP loci (Table 2). Nine loci were identified through the combined analyses of Stage 1 and 2; the remaining nine loci were identified in Stage 1 but not available in Stage 2 for combined analyses (Tables S7). The LocusZoom plots of these novel loci are presented in Fig. S3. Two loci (*SLIT3* and *HRH4*) were identified through the 1DF test of interaction effects. At both loci, the genetic effect on DBP was stronger and beneficial in higher education and weaker and detrimental in lower education. For example, at *SLIT3*, the minor allele A was associated with a 4.82 mmHg lower DBP in higher education (GradCol = 1), whereas it was associated with a 2.25 mmHg higher DBP in GradCol = 0. The remaining 16 loci were identified through the 2DF joint test of the SNV and interaction effects; twelve loci were identified considering ‘Some College’ (SomeCol) and four loci were identified considering ‘Graduated College’ (GradCol).

Ancestry-specific and trans-ancestry analyses

Novel loci were identified through separate analyses of AFR (12 loci), EUR (1 locus), trans-ancestry (4 loci), and in both AFR and trans-ancestry (1 locus). This highlights the importance of including non-EUR populations to identify novel BP loci. By nature, AFR populations carry more rare and low-frequency variants that may be very rare or monomorphic in other ancestral groups [22]; the MAF for the novel index SNVs range from 0.02 to 0.04 in AFR. The enhanced discovery of novel loci in AFR ancestry may be attributable to the relatively higher MAF in this population versus in EUR. For example, rs141962517 (*CAPDS*) with a MAF = 0.02 in AFR was significantly associated with SBP (2DF $P = 3.07 \times 10^{-10}$; 1DF Interaction $P = 1.99 \times 10^{-7}$); this variant was not present in other ancestries after filtering.

Among the 18 novel loci, three loci were identified only through trans-ancestry analyses, as none of the

Table 2 Eighteen new loci associated with BP traits that are at least 1 Mb away from any known BP locus.

Locus	Nearest Gene	rsID	Chr:Pos	EA	EAF (AFR/ASN/EUR/HIS)	Genetic Effect		GxE Interaction		P value		Trait	Educ.	Anc.
						Effect	SE	Effect	SE	Interaction	2DF Joint			
1	<i>CDC14A</i>	rs114558965	1:100829685	a	0.97/. /. /1.00	-0.23	1.02	5.02	1.27	2.54E-03	2.86E-09	SBP	SC	AFR
2	<i>LIN01249</i>	rs9308788	2:4711095	a	0.02/0.22/0.06/0.06	2.97	2.13	-11.93	2.64	5.04E-05	4.47E-08	SBP	GC	AFR
3	<i>ARL4C</i>	rs145586115	2:235604646	t	0.97/. /. /0.99	4.80	0.87	-2.57	1.29	1.16E-01	1.92E-08	SBP	SC	AFR
4	<i>CADPS</i>	rs141962517	3:62514061	t	0.98/. /. /.	-2.46	1.54	12.85	2.23	1.99E-07	3.07E-10	SBP	GC	AFR
5	<i>PCDH7</i>	rs74458816	4:31363388	a	0.03/. /. /0.00	-0.18	0.70	-3.46	0.91	2.26E-03	4.90E-09	MAP	SC	AFR
6	<i>E1F4E</i>	rs2141284	4:99704167	a	. /. /0.01/0.01	-1.35	0.22	1.01	0.38	2.16E-02	6.73E-09	MAP	GC	TRANS
7	<i>SPEF2</i>	rs115523707	5:35756623	t	0.03/. /0.00/0.00	-0.83	0.64	-2.73	0.88	5.91E-02	7.22E-09	PP	SC	AFR
8	<i>ANKRD34B</i>	rs66907226	5:79863455	d	0.46/0.81/0.58/0.69	0.48	0.14	0.04	0.18	9.86E-01	4.43E-08	SBP	SC	EUR
9	<i>SLC04C1</i>	rs114175587	5:100994204	t	0.02/. /. /0.00	-4.84	0.83	3.84	1.29	3.55E-03	2.40E-08	PP	SC	TRANS
10	<i>SLIT3</i>	rs142385399	5:168166731	a	0.04/. /0.00/0.01	2.25	0.74	-7.07	1.22	2.79E-08	4.76E-08	DBP	GC	AFR
11	<i>THSD7A</i>	rs200612978	7:11493906	d	0.03/0.03/. /0.01	-0.40	1.27	-4.94	1.57	5.71E-02	4.81E-08	SBP	SC	AFR
12	<i>HAS2-AS1</i>	rs112332671	8:122673983	a	0.02/. /. /0.00	-1.37	1.18	-4.55	1.52	6.03E-02	1.89E-09	PP	SC	AFR
13	<i>CDON</i>	rs12295584	11:125841078	a	0.95/. /0.99/0.99	2.87	0.57	-1.07	0.84	2.20E-01	4.71E-08	SBP	SC	TRANS
14	<i>DSTNP2</i>	rs75535814	12:6996683	t	0.02/. /. /0.00	-8.78	1.45	7.24	1.95	3.17E-04	5.79E-09	SBP	SC	AFR
15	<i>PIK3C2G</i>	rs189555401	12:18443065	t	0.03/. /. /0.01	-3.23	0.57	1.82	1.15	8.92E-02	4.10E-08	DBP	GC	TRANS
16	<i>E2F7; ZDHC17</i>	rs17043233	12:77648216	t	0.02/. /. /0.00	-4.07	0.58	4.26	0.90	1.72E-05	1.39E-11	MAP	SC	TRANS
17	<i>HRH4</i>	rs8099516	18:22108763	t	0.04/. /. /0.01	1.71	0.61	-5.39	0.95	7.83E-09	5.04E-08	DBP	GC	AFR
18	<i>LOC100289473</i>	rs113809930	20:1711768	a	0.98/0.88/0.97/0.94	-0.69	0.79	3.67	0.94	1.70E-03	3.48E-08	DBP	SC	AFR

Positions are based on build 37. Effect is in mmHg unit.

BP blood pressure, DBP diastolic BP, EA effect allele, EAF effect allele frequency observed in our cohorts, MAP mean arterial pressure, P P value of the joint test with 2 degrees of freedom of genetic main and interaction effects, PP pulse pressure, SBP systolic BP, SE standard error, 2DF joint IDF interaction, P P value of the interaction test with 1 degree of freedom. The smallest P values between IDF interaction test and the joint 2DF test are in boldface, SC SomeCol, EUR European, AFR African, TRANS, trans-ancestry (i.e., combining all ancestry groups through meta-analysis).




Fig. 2 LocusZoom plots for 2 BP loci related to CNS-adrenal signaling. **a** MAP-associated locus adjacent to *ZDHHC17*, identified in AFR and in trans-ancestry, shows roles in CNS-adrenal signaling. In neurons, *ZDHHC17* encodes a membrane protein that mediates fusion of synaptic vesicles to the plasma membrane, enabling the release of neurotransmitters. Murine *zdhhc17* knockout models show impaired hippocampal memory and reduced synaptic plasticity, providing potential biological links to working memory and subsequent educational attainment. **b** A locus intragenic to *CADPS*, identified in AFR, is

of potential biologic relevance given this gene's expression in CNS tissue and role in regulating the fusion of neuroendocrine vesicles and release of vasoactive catecholamines from both adrenal and neural tissue. Three LD SNVs have CADD scores >10, and four LD SNVs have ChromHMM state signals consistent with strong evidence of transcription regulation. SBP systolic blood pressure, DBP diastolic blood pressure, MAP mean arterial pressure, PP pulse pressure. The plots were created using LocusZoom (<http://locuszoom.sph.umich.edu/>).

ancestry-specific analyses reached genome-wide significance. For example, the index SNV (rs189555401) representing the four-variant locus within *PIK3C2G* was suggestively associated with DBP ($P = 1.31 \times 10^{-7}$) in AFR and not even nominally associated in HIS ($P = 9.67 \times 10^{-2}$). However, in trans-ancestry analysis combining these two ancestral groups, the association reached genome-wide significance ($P = 4.10 \times 10^{-8}$).

Functional annotation and eQTL evidence

To obtain functional annotations for the index variants and nearby variants in LD ($r^2 \geq 0.2$), we used FUMA GWAS [28]. Among the 18 index variants representing our novel loci, two variants were intronic to a non-coding RNA (ncRNA), six variants were intronic, nine variants were intergenic, and the remaining variant (rs66907226) was an indel without available annotation. Among the 499 variants that include both the index variants and nearby variants in LD, there were four exonic, four exonic-ncRNA, 119 intronic, 67 intronic-ncRNA, two 3' untranslated region (UTR), seven up/downstream flanking, and 296 intergenic variants (Table S8). Of the 499 variants, 13 had RegulomeDB [31] scores better than or equal to 3a, suggesting at least moderate evidence for involvement in transcription regulation (Table S9). In total, 32 SNVs have CADD [32] scores ≥ 10 ,

representing the top 10% of predicted deleteriousness for SNVs genome-wide. A single SNV (rs112332671) ~20 kb upstream of *HAS2* and 16 kb downstream of the ncRNA *HAS2-AS1* had a CADD score of 20.1, placing it in the top 1% of predicted deleteriousness.

The 15-core chromatin state (ChromHMM) [33] was assessed for 127 epigenomes in the 17 index variants (Table S9). Two had histone chromatin markers in regions flanking the transcription start site and one in a region associated with strong transcription in relevant tissues including brain. Among all 499 LD variants, 45 had histone chromatin markers characteristic of a transcription start site, 64 had markers consistent with strong transcription, and 25 were in enhancer regions. One LD variant (rs555713705) was identified as *cis*-acting eQTLs [46,47] for heart tissue in the GTEx_v7 database (FDR P value ranging from 3.90×10^{-3}).

Biologic plausibility of the new BP loci

Three novel BP loci are related to the central nervous system (CNS)-adrenal signaling axis that is critical for BP regulation. A locus (Fig. 2a) adjacent to *ZDHHC17*, identified in AFR and in trans-ancestry analyses, encodes a membrane protein that mediates fusion of synaptic vesicles to the plasma membrane. *CADPS* (Fig. 2b), identified in

AFR, is expressed in CNS tissue. Three variants in LD have CADD scores >10, and four SNVs have ChromHMM state signals consistent with strong evidence of transcription regulation. *PIK3C2G*, identified in trans-ancestry analyses, also shows roles in CNS-adrenal signaling. Three variants in LD in this locus have CADD scores >10, including one with a CADD score of 18 that is predicted to reside in an enhancer region in fetal adrenal cells.

Two novel BP loci are related to renal fibrosis and cation exchange. A locus, which includes a variant intragenic to *SLIT3*, showed interaction evidence with educational attainment in AFR (rs142385399, $P = 2.79 \times 10^{-8}$). A locus also identified in AFR includes *HAS2* and *HAS2-AS1*, which play roles in renal fibrosis. In addition, we identified two novel BP loci related to pathways involved in vascular smooth muscle cell structure and function. A locus identified by trans-ancestry analyses included a variant intragenic to *CDON*, which is expressed in vascular smooth muscle cells. A locus identified in AFR includes *GNB3*, which encodes a subunit critical for signal transduction of several vasoactive peptide G protein-coupled receptors involved in BP regulation. A SNV in this locus shows ChromHMM chromatin states consistent with strong transcription regulation in multiple tissues, and three SNVs have strong *cis*-eQTL association with *GNB3* expression in nerve, artery, and skeletal muscle tissue (minimum FDR P value 1.20×10^{-43}).

Discussion

A relationship between educational attainment and BP has been well demonstrated [48–51]. Furthermore, AFR ancestry individuals have been shown to have a higher burden of HTN than EUR-ancestry [52]. However, higher-educated AFR ancestry individuals bear approximately twice the burden of HTN as compared with their EUR-ancestry counterparts [48,51], demonstrating that educational differences did not fully account for this observed racial disparity. Herein, we reported genome-wide meta-analyses for SBP, DBP, MAP, and PP accounting for gene-educational attainment interactions across five ancestry groups. We pursued a genome-wide interrogation in 117 438 individuals (in Stage 1) and follow-up analysis at selected variants in 293,787 additional individuals (in Stage 2). Through the combined meta-analysis of stages 1 and 2, we identified 84 known and 18 novel loci at genome-wide significance. As known loci have been discussed elsewhere, this report highlights several novel loci show biologic plausibility by involving physiologic systems implicated in BP regulation.

The CNS-adrenal signaling axis is critical for BP regulation. Three novel BP loci (*ZDHHC17*, *CADPS*, and *PIK3C2G*) are related to these pathways. In neurons, *ZDHHC17* encodes a membrane protein that mediates fusion

of synaptic vesicles to the plasma membrane, enabling the release of neurotransmitters [53]. Murine *zdhhc17* knockout models show impaired hippocampal memory and reduced synaptic plasticity, providing potential biological links to working memory and subsequent educational attainment [54]. Although a biological connection between *ZDHHC17* and BP traits is not well established, *zdhhc17* expression induces neurite outgrowth in a rodent adrenal-derived cell line [55]. *Cadps* plays a role in regulating the fusion of neuroendocrine vesicles and release of vasoactive catecholamines in calf adrenal and neural tissue [56]. *Pik3c2g* encodes a phosphoinositide kinases that are expressed in a sexually dimorphic pattern specifically in a zone of the mouse adrenal cortex believed to play a role in steroid sex hormone production [57]. Furthermore, *PIK3C2G* is under-expressed in human hypertensive kidneys, providing a potential biological link between the expression of adrenal mineralocorticoid hormones and their target organ [58]. Among alcohol-preferring rats, *pik3c2g* expression is also increased in the cerebral periaqueductal gray, a region involved in pain, fear, and anxiety responses [59], possibly providing a link to drivers of SES in humans [60]. Notably, the loci including *ZDHHC17* and *CADPS* demonstrated some evidence of interaction with educational attainment ($P = 1.72 \times 10^{-5}$ and 1.99×10^{-7} , respectively).

Two new BP loci (*HAS2* and *HAS2-AS1*, *SLIT3*) show potential roles in renal function. A locus that includes a variant intragenic to *SLIT3* had a significant interaction term with educational attainment. *SLIT3* encodes a cell-cell adhesion molecule that binds its receptor, ROBO4, in human-derived endothelial stem cells directing the formation of vascular networks [61]. *SLIT3* also plays a role in directing neuronal growth in the brain [62,63], and in renal and cardiac development [64]. A locus including *HAS2* and *HAS2-AS1* is also of interest for roles played in renal fibrosis. *HAS2-AS1* is an antisense ncRNA simultaneously expressed and thought to stabilize the *HAS2* transcript [65].

Two new BP loci (*GNB3*, *CDON*) have been shown to regulate pathways involved in vascular smooth muscle cell structure and function. We identified a locus in *GNB3*, which encodes a G protein-coupled receptor subunit involved in BP regulation. Several candidate gene association studies have identified the synonymous *GNB3* variant C825T (rs5443), resulting in a splice variant of the $\beta 3$ subunit, as significantly associated with essential HTN [66,67], with BP response to diuretic [68] and β -adrenergic receptor blockade [69], and other cardiovascular traits [70]. Another locus identified by trans-ancestry analyses included a variant intragenic to *CDON*; this gene is expressed in vascular smooth muscle cells [71], and encodes a cell-surface receptor complex that regulates myocyte differentiation in rodents [72].

This large-scale multi-ancestry study has several limitations. First, the practice of adjusting SBP and DBP by

adding 15 and 10 mmHg for antihypertensive use is based on a method derived from a EUR-ancestry cohort [21]. While this approach is common among GWAS of BP traits [36], we acknowledge that this practice may not be equally appropriate and/or justified in all ancestry groups. Second, while the sample sizes in diverse ancestries are a strength, resulting in the identification of several novel BP loci particularly in AFR ancestry, several identified loci included low-frequency variants that require further validation. Third, main effect only analysis without educational attainment was not performed, and this limits our ability to resolve if novel loci identified through the 2DF joint test could be found without considering educational attainment. Fourth, the use of educational attainment as a proxy for SES can present some challenges. The socioeconomic impact of education has changed over time and may differ according to birth cohort, as well as in other subgroups defined by gender, ancestry, region, and/or country [1,49]. Even with similar levels of educational attainment, social and environmental experiences were different between AFR and EUR individuals in the United States, especially those educated in the 1960s and 1970s, resulting in residual confounding inequities between the ancestral groups [9,73]. This additional source of heterogeneity may have reduced power for trans-ancestry analyses.

In summary, this multi-ancestry study that used gene-education interactions on BP traits identified 18 novel loci and validated 84 known BP loci. Ten novel loci were identified in individuals of AFR ancestry, demonstrating the need for pursuing genetic studies in diverse populations. Several novel loci involve physiologic systems implicated in BP regulation including genes involved in CNS-adrenal signaling, vascular structure and function, and renal function. Two loci showed interaction evidence with educational attainment. These findings may identify a role for educational attainment and SES in further dissection of the genetic architecture of BP.

Acknowledgements This project was largely supported by a grant from the U.S. National Heart, Lung, and Blood Institute (NHLBI), the National Institutes of Health, R01HL118305. A Career Development Award (K25HL121091), also from the NHLBI, enabled Dr. Sung to play a major role on this project. Dr. Kilpeläinen was supported by the Novo Nordisk Foundation (NNF18CC0034900 and NNF17OC0026848). Full set of study-specific funding sources and acknowledgments appear in the Supplementary Material. These authors constitute the writing group: L.d.l.F., Y.J.S., R.N., T.W., M.F.F., K.S., P.B.M., P.W.F., D.C.R., M.F.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Kaplan GA, Keil JE. Socioeconomic factors and cardiovascular disease: a review of the literature. *Circulation*. 1993;88:1973–98.
- Liberatos P, Link BG, Kelsey JL. The measurement of social class in epidemiology. *Epidemiol Rev*. 1988;10:87–121.
- Leng B, Jin Y, Li G, Chen L, Jin N. Socioeconomic status and hypertension: a meta-analysis. *J Hypertens*. 2015;33:221–9.
- Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health*. 1992;82:816–20.
- Basson J, Sung YJ, Schwander K, Kume R, Simino J, de las Fuentes L, et al. Gene-education interactions identify novel blood pressure loci in the Framingham Heart Study. *Am J Hypertens*. 2014;27:431–44.
- Lee JJ, Wedow R, Okbay A, Kong E, Maghziyan O, Zacher M, et al. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet*. 2018;50:1112–21.
- Hertz RP, Unger AN, Cornell JA, Saunders E. Racial disparities in hypertension prevalence, awareness, and management. *Arch Intern Med*. 2005;165:2098–104.
- Stephoe A, Hamer M, Butcher L, Lin J, Brydon L, Kivimaki M, et al. Educational attainment but not measures of current socioeconomic circumstances are associated with leukocyte telomere length in healthy older men and women. *Brain Behav Immun*. 2011;25:1292–8.
- Metcalfe PA, Sharrett AR, Folsom AR, Duncan BB, Patsch W, Hutchinson RG, et al. African American-white differences in lipids, lipoproteins, and apolipoproteins, by educational attainment, among middle-aged adults: the atherosclerosis risk in communities study. *Am J Epidemiol*. 1998;148:750–60.
- Matthews KA, Kelsey SF, Meilahn EN, Kuller LH, Wing RR. Educational attainment and behavioral and biologic risk factors for coronary heart disease in middle-aged women. *Am J Epidemiol*. 1989;129:1132–44.
- Rietveld CA, Medland SE, Derringer J, Yang J, Esko T, Martin NW, et al. GWAS of 126,559 individuals identifies genetic variants associated with educational attainment. *Science*. 2013;340:1467–71.
- Smith GD, Hart C, Watt G, Hole D, Hawthorne V. Individual social class, area-based deprivation, cardiovascular disease risk factors, and mortality: the Renfrew and Paisley Study. *J Epidemiol Community Health*. 1998;52:399–405.
- Gallo LC, Matthews KA, Kuller LH, Sutton-Tyrrell K, Edmondowicz D. Educational attainment and coronary and aortic calcification in postmenopausal women. *Psychosom Med*. 2001;63:925–35.
- Jacobsen BK, Thelle DS. Risk factors for coronary heart disease and level of education. The Tromso Heart Study. *Am J Epidemiol*. 1988;127:923–32.
- Pierce JP, Fiore MC, Novotny TE, Hatziandreu EJ, Davis RM. Trends in cigarette smoking in the United States. Educational differences are increasing. *JAMA*. 1989;261:56–60.
- Stamler J, Elliott P, Appel L, Chan Q, Buzzard M, Dennis B, et al. Higher blood pressure in middle-aged American adults with less education-role of multiple dietary factors: the INTERMAP study. *J Hum Hypertens*. 2003;17:655–775.
- Tian HG, Hu G, Dong QN, Yang XL, Nan Y, Pietinen P, et al. Dietary sodium and potassium, socioeconomic status and blood pressure in a Chinese population. *Appetite*. 1996;26:235–46.
- Kaplan GA, Lazarus NB, Cohen RD, Leu DJ. Psychosocial factors in the natural history of physical activity. *Am J Prev Med*. 1991;7:12–17.

19. McCaffery JM, Papandonatos GD, Lyons MJ, Niaura R. Educational attainment and the heritability of self-reported hypertension among male Vietnam-era twins. *Psychosom Med.* 2008;70:781–6.
20. Rao DC, Sung YJ, Winkler TW, Schwander K, Borecki I, Cupples LA et al. Multiancestry study of gene-lifestyle interactions for cardiovascular traits in 610 475 individuals from 124 cohorts: design and rationale. *Circ Cardiovasc Genet.* 2017;10:e001649.
21. Tobin MD, Sheehan NA, Scurren KJ, Burton PR. Adjusting for treatment effects in studies of quantitative traits: antihypertensive therapy and systolic blood pressure. *Stat Med.* 2005;24:2911–35.
22. 1000 Genomes Project Consortium, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature.* 2012;491:56–65.
23. Manning AK, LaValley M, Liu CT, Rice K, An P, Liu Y, et al. Meta-analysis of gene-environment interaction: joint estimation of SNP and SNP x environment regression coefficients. *Genet Epidemiol.* 2011;35:11–18.
24. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics.* 2010;26:2190–1.
25. Devlin B, Roeder K. Genomic control for association studies. *Biometrics.* 1999;55:997–1004.
26. Winkler TW, Day FR, Croteau-Chonka DC, Wood AR, Locke AE, Magi R, et al. Quality control and conduct of genome-wide association meta-analyses. *Nat Protoc.* 2014;9:1192–212.
27. Sung YJ, Winkler TW, de Las Fuentes L, Bentley AR, Brown MR, Kraja AT, et al. A large-scale multi-ancestry genome-wide study accounting for smoking behavior identifies multiple significant loci for blood pressure. *Am J Hum Genet.* 2018;102:375–400.
28. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun.* 2017;8:1826.
29. Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res.* 2012;40:D930–934.
30. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* 2010;38:e164.
31. Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, et al. Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res.* 2012;22:1790–7.
32. Kircher M, Witten DM, Jain P, O’Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet.* 2014;46:310–5.
33. Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenyk M, Yen A, et al. Integrative analysis of 111 reference human epigenomes. *Nature.* 2015;518:317–30.
34. Ernst J, Kellis M. ChromHMM: automating chromatin-state discovery and characterization. *Nat Methods.* 2012;9:215–6.
35. GTEx Consortium. Genetic effects on gene expression across human tissues. *Nature.* 2017;550:204–13.
36. Newton-Cheh C, Johnson T, Gateva V, Tobin MD, Bochud M, Coin L, et al. Genome-wide association study identifies eight loci associated with blood pressure. *Nat Genet.* 2009;41:666–76.
37. Ehret GB, Munroe PB, Rice KM, Bochud M, Johnson AD, Chasman DI, et al. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature.* 2011;478:103–9.
38. Ehret GB, Ferreira T, Chasman DI, Jackson AU, Schmidt EM, Johnson T, et al. The genetics of blood pressure regulation and its target organs from association studies in 342,415 individuals. *Nat Genet.* 2016;48:1171–84.
39. Liu C, Kraja AT, Smith JA, Brody JA, Franceschini N, Bis JC, et al. Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat Genet.* 2016;48:1162–70.
40. Surendran P, Drenos F, Young R, Warren H, Cook JP, Manning AK, et al. Trans-ancestry meta-analyses identify rare and common variants associated with blood pressure and hypertension. *Nat Genet.* 2016;48:1151–61.
41. Levy D, Ehret GB, Rice K, Verwoert GC, Launer LJ, Dehghan A, et al. Genome-wide association study of blood pressure and hypertension. *Nat Genet.* 2009;41:677–87.
42. Hoffmann TJ, Ehret GB, Nandakumar P, Ranatunga D, Schaefer C, Kwok PY, et al. Genome-wide association analyses using electronic health records identify new loci influencing blood pressure variation. *Nat Genet.* 2017;49:54–64.
43. Evangelou E, Warren HR, Mosen-Ansorena D, Mifsud B, Pazoki R, Gao H, et al. Genetic analysis of over 1 million people identifies 535 new loci associated with blood pressure traits. *Nat Genet.* 2018;50:1412–25.
44. Giri A, Hellwege JN, Keaton JM, Park J, Qiu C, Warren HR, et al. Trans-ethnic association study of blood pressure determinants in over 750,000 individuals. *Nat Genet.* 2019;51:51–62.
45. Feitosa MF, Kraja AT, Chasman DI, Sung YJ, Winkler TW, Ntalla I, et al. Novel genetic associations for blood pressure identified via gene-alcohol interaction in up to 570K individuals across multiple ancestries. *PLoS ONE.* 2018;13:e0198166.
46. The GTEx Consortium. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science.* 2015;348:648–60.
47. Westra HJ, Peters MJ, Esko T, Yaghootkar H, Schurmann C, Ketunen J, et al. Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nat Genet.* 2013;45:1238–43.
48. Hypertension Detection and Follow-up Program Cooperative Group. Race, education and prevalence of hypertension. *Am J Epidemiol.* 1977;106:351–61.
49. Sorel JE, Ragland DR, Syme SL, Davis WB. Educational status and blood pressure: the Second National Health and Nutrition Examination Survey, 1976–1980, and the Hispanic Health and Nutrition Examination Survey, 1982–1984. *Am J Epidemiol.* 1992;135:1339–48.
50. Steffen PR. The cultural gradient: culture moderates the relationship between socioeconomic status (SES) and ambulatory blood pressure. *J Behav Med.* 2006;29:501–10.
51. Vargus CM, Ingram DD, Gillum RF. Incidence of hypertension and educational attainment: the NHANES I epidemiologic followup study. First National Health and Nutrition Examination Survey. *Am J Epidemiol.* 2000;152:272–8.
52. Benjamin EJ, Virani SS, Callaway CW, Chamberlain AM, Chang AR, Cheng S, et al. Heart disease and stroke statistics-2018 update: a report from the American Heart Association. *Circulation.* 2018;137:e67–e492.
53. Ohyama T, Verstreken P, Ly CV, Rosenmund T, Rajan A, Tien AC, et al. Huntingtin-interacting protein 14, a palmitoyl transferase required for exocytosis and targeting of CSP to synaptic vesicles. *J Cell Biol.* 2007;179:1481–96.
54. Milnerwood AJ, Parsons MP, Young FB, Singaraja RR, Franciosi S, Volta M, et al. Memory and synaptic deficits in Hip14/DHHC17 knockout mice. *Proc Natl Acad Sci USA.* 2013;110:20296–301.
55. Shi W, Wang F, Gao M, Yang Y, Du Z, Wang C, et al. ZDHHC17 promotes axon outgrowth by regulating TrkA-tubulin complex formation. *Mol Cell Neurosci.* 2015;68:194–202.
56. Elhamedani A, Martin TF, Kowalchuk JA, Artalejo CR. Ca(2+) dependent activator protein for secretion is critical for the fusion of dense-core vesicles with the membrane in calf adrenal chromaffin cells. *J Neurosci.* 1999;19:7375–83.
57. El Wakil A, Mari B, Barhanin J, Lalli E. Genomic analysis of sexual dimorphism of gene expression in the mouse adrenal gland. *Horm Metab Res.* 2013;45:870–3.
58. Marques FZ, Campain AE, Tomaszewski M, Zukowska-Szczzechowska E, Yang YH, Charchar FJ, et al. Gene expression

- profiling reveals renin mRNA overexpression in human hypertensive kidneys and a role for microRNAs. *Hypertension*. 2011;58:1093–8.
59. McClintick JN, McBride WJ, Bell RL, Ding ZM, Liu Y, Xuei X, et al. Gene expression changes in glutamate and GABA-A receptors, neuropeptides, ion channels, and cholesterol synthesis in the periaqueductal gray following binge-like alcohol drinking by adolescent alcohol-preferring (P) rats. *Alcohol Clin Exp Res*. 2016;40:955–68.
 60. McEwen BS, Gianaros PJ. Central role of the brain in stress and adaptation: links to socioeconomic status, health, and disease. *Ann NY Acad Sci*. 2010;1186:190–222.
 61. Paul JD, Coulombe K, Toth PT, Zhang Y, Marsboom G, Bindokas VP, et al. SLIT3-ROBO4 activation promotes vascular network formation in human engineered tissue and angiogenesis in vivo. *J Mol Cell Cardiol*. 2013;64:124–31.
 62. Ypsilanti AR, Zagar Y, Chedotal A. Moving away from the midline: new developments for Slit and Robo. *Development*. 2010;137:1939–52.
 63. Blockus H, Chedotal A. Slit-Robo signaling. *Development*. 2016;143:3037–44.
 64. Liu J, Zhang L, Wang D, Shen H, Jiang M, Mei P, et al. Congenital diaphragmatic hernia, kidney agenesis and cardiac defects associated with Slit3-deficiency in mice. *Mech Dev*. 2003;120:1059–70.
 65. Michael DR, Phillips AO, Krupa A, Martin J, Redman JE, Altafer A, et al. The human hyaluronan synthase 2 (HAS2) gene and its natural antisense RNA exhibit coordinated expression in the renal proximal tubular epithelial cell. *J Biol Chem*. 2011;286:19523–32.
 66. Siffert W, Roskopf D, Siffert G, Busch S, Moritz A, Erbel R, et al. Association of a human G-protein beta3 subunit variant with hypertension. *Nat Genet*. 1998;18:45–48.
 67. Bagos PG, Elefsinioti AL, Nikolopoulos GK, Hamodrakas SJ. The GNB3 C825T polymorphism and essential hypertension: a meta-analysis of 34 studies including 14,094 cases and 17,760 controls. *J Hypertens*. 2007;25:487–500.
 68. Turner ST, Schwartz GL, Chapman AB, Boerwinkle E. C825T polymorphism of the G protein beta(3)-subunit and anti-hypertensive response to a thiazide diuretic. *Hypertension*. 2001;37:739–43.
 69. Filigheddu F, Reid JE, Troffa C, PinnaPargaglia P, Argiolas G, Testa A, et al. Genetic polymorphisms of the beta-adrenergic system: association with essential hypertension and response to beta-blockade. *Pharmacogenomics J*. 2004;4:154–60.
 70. Bojic T, Milovanovic B, Cupic SJ. Genetic polymorphisms of neurocardiovascular disorders. *Arch Med*. 2015;7:1–22.
 71. Aoki T, Kataoka H, Ishibashi R, Nozaki K, Hashimoto N. Gene expression profile of the intima and media of experimentally induced cerebral aneurysms in rats by laser-microdissection and microarray techniques. *Int J Mol Med*. 2008;22:595–603.
 72. Takaesu G, Kang JS, Bae GU, Yi MJ, Lee CM, Reddy EP, et al. Activation of p38alpha/beta MAPK in myogenesis via binding of the scaffold protein JLP to the cell surface protein Cdo. *J Cell Biol*. 2006;175:383–8.
 73. Tyroler HA. Socioeconomic status in the epidemiology and treatment of hypertension. *Hypertension*. 1989;13 5 Suppl:194–97.

Affiliations

Lisa de las Fuentes ^{1,2} · Yun Ju Sung² · Raymond Noordam³ · Thomas Winkler⁴ · Mary F. Feitosa ⁵ · Karen Schwander² · Amy R. Bentley ⁶ · Michael R. Brown⁷ · Xiuqing Guo⁸ · Alisa Manning^{9,10} · Daniel I. Chasman^{11,12} · Hugues Aschard^{13,14} · Traci M. Bartz¹⁵ · Lawrence F. Bielak¹⁶ · Archie Campbell ¹⁷ · Ching-Yu Cheng ^{18,19} · Rajkumar Dorajoo ²⁰ · Fernando P. Hartwig ^{21,22} · A. R. V. R. Horimoto²³ · Changwei Li²⁴ · Ruifang Li-Gao²⁵ · Yongmei Liu²⁶ · Jonathan Marten²⁷ · Solomon K. Musani²⁸ · Ioanna Ntalla²⁹ · Tuomo Rankinen³⁰ · Melissa Richard ³¹ · Xueling Sim ³² · Albert V. Smith^{33,34} · Salman M. Tajuddin³⁵ · Bamidele O. Tayo³⁶ · Dina Vojinovic³⁷ · Helen R. Warren^{29,38} · Deng Xuan³⁹ · Maris Alver^{40,41} · Mathilde Boissel ⁴² · Jin-Fang Chai³² · Xu Chen ⁴³ · Kaare Christensen ⁴⁴ · Jasmin Divers⁴⁵ · Evangelos Evangelou^{46,47} · Chuan Gao⁴⁸ · Giorgia Grotto^{49,50} · Sarah E. Harris ⁵¹ · Meian He⁵² · Fang-Chi Hsu⁴⁵ · Brigitte Kühnel^{53,54} · Federica Laguzzi⁵⁵ · Xiaoyin Li^{56,57} · Leo-Pekka Lyytikäinen^{58,59} · Ilja M. Nolte ⁶⁰ · Alaitz Poveda⁶¹ · Rainer Rauramaa⁶² · Muhammad Riaz⁶³ · Rico Rueedi^{64,65} · Xiao-ou Shu⁶⁶ · Harold Snieder ⁶⁰ · Tamar Sofer⁶⁷ · Fumihiko Takeuchi ⁶⁸ · Niek Verweij⁶⁹ · Erin B. Ware ⁷⁰ · Stefan Weiss ^{71,72} · Lisa R. Yanek ⁷³ · Najaf Amin³⁷ · Dan E. Arking⁷⁴ · Donna K. Arnett⁷⁵ · Sven Bergmann ^{64,65} · Eric Boerwinkle^{7,76} · Jennifer A. Brody ⁷⁷ · Ulrich Broeckel⁷⁸ · Marco Brumat⁴⁹ · Gregory Burke⁷⁹ · Claudia P. Cabrera^{29,38} · Mickaël Canouil ⁴² · Miao Li Chee⁸⁰ · Yii-Der Ida Chen⁸ · Massimiliano Cocca⁵⁰ · John Connell⁸¹ · H. Janaka de Silva⁸² · Paul S. de Vries⁷ · Gudny Eiriksdottir³⁴ · Jessica D. Faul⁷⁰ · Virginia Fisher³⁹ · Terrence Forrester⁸³ · Ervin F. Fox⁸⁴ · Yechiel Friedlander⁸⁵ · He Gao^{46,86} · Bruna Gigante ^{87,88} · Franco Giulianini⁸⁹ · Chi Charles Gu² · Dongfeng Gu⁹⁰ · Tamara B. Harris⁹¹ · Jiang He^{92,93} · Sami Heikkinen ^{94,95} · Chew-Kiat Heng^{96,97} · Steven Hunt^{98,99} · M. Arfan Ikram ^{37,100} · Marguerite R. Irvin¹⁰¹ · Mika Kähönen^{102,103} · Maryam Kavousi³⁷ · Chiea Chuen Khor^{20,104} · Tuomas O. Kilpeläinen^{105,106} · Woon-Puay Koh^{32,107} · Pirjo Komulainen⁶² · Aldi T. Kraja⁵ · J. E. Krieger²³ · Carl D. Langefeld⁴⁵ · Yize Li ² · Jingjing Liang⁵⁶ · David C. M. Liewald ⁵¹ · Ching-Ti Liu ³⁹ · Jianjun Liu ^{20,32} · Kurt K. Lohman¹⁰⁸ · Reedik Mägi⁴⁰ · Colin A. McKenzie⁸³ · Thomas Meitinger^{109,110} · Andres Metspalu^{40,41} · Yuri Milaneschi¹¹¹ · Lili Milani ⁴⁰ · Dennis O. Mook-Kanamori^{25,112} · Mike A. Nalls^{113,114} · Christopher P. Nelson ^{115,116} · Jill M. Norris¹¹⁷ · Jeff O'Connell^{118,119} · Adesola Ogunniyi¹²⁰ · Sandosh Padmanabhan¹²¹ · Nicholette D. Palmer ¹²² · Nancy L. Pedersen⁴³ · Thomas Perls¹²³ · Annette Peters^{54,124} · Astrid Petersmann¹²⁵ · Patricia A. Peyser ¹⁶ · Ozren Polasek^{126,127,128} · David J. Porteous^{17,51}

Leslie J. Raffel¹²⁹ · Treva K. Rice² · Jerome I. Rotter⁸ · Igor Rudan¹³⁰ · Oscar-Leonel Rueda-Ochoa³⁷ · Charumathi Sabanayagam^{18,19} · Babatunde L. Salako¹²⁰ · Pamela J. Schreiner¹³¹ · James M. Shikany¹³² · Stephen S. Sidney¹³³ · Mario Sims²⁸ · Colleen M. Sitlani⁷⁷ · Jennifer A. Smith^{16,70} · John M. Starr^{134,135} · Konstantin Strauch^{136,137} · Morris A. Swertz¹³⁸ · Alexander Teumer^{72,139} · Yih Chung Tham¹⁸ · André G. Uitterlinden^{37,140} · Dhananjay Vaidya⁷³ · M. Yldau van der Ende⁶⁹ · Melanie Waldenberger^{53,54,124} · Lihua Wang⁵ · Ya-Xing Wang¹⁴¹ · Wen-Bin Wei¹⁴² · David R. Weir⁷⁰ · Wanqing Wen⁶⁶ · Jie Yao⁸ · Bing Yu⁷ · Caizheng Yu⁵² · Jian-Min Yuan^{143,144} · Wei Zhao¹⁶ · Alan B. Zonderman¹⁴⁵ · Diane M. Becker⁷³ · Donald W. Bowden¹²² · Ian J. Deary⁵¹ · Marcus Dörr^{72,146} · Tõnu Esko^{40,147} · Barry I. Freedman¹⁴⁸ · Philippe Froguel^{42,149} · Paolo Gasparini^{49,50} · Christian Gieger^{53,150} · Jost Bruno Jonas^{151,152} · Candace M. Kammerer¹⁵³ · Norihiro Kato⁶⁸ · Timo A. Lakka^{62,94,154} · Karin Leander⁵⁵ · Terho Lehtimäki^{58,59} · Lifelines Cohort Study¹⁵⁵ · Patrik K. E. Magnusson⁴³ · Pedro Marques-Vidal¹⁵⁶ · Brenda W. J. H. Penninx¹¹¹ · Nilesh J. Samani^{115,116} · Pim van der Harst^{69,157} · Lynne E. Wagenknecht⁷⁹ · Tangchun Wu⁵² · Wei Zheng⁶⁶ · Xiaofeng Zhu⁵⁶ · Claude Bouchard³⁰ · Richard S. Cooper³⁶ · Adolfo Correa²⁸ · Michele K. Evans³⁵ · Vilmundur Gudnason^{34,158} · Caroline Hayward²⁷ · Bernardo L. Horta²¹ · Tanika N. Kelly⁹² · Stephen B. Kritchevsky¹⁵⁹ · Daniel Levy^{160,161} · Walter R. Palmas¹⁶² · A. C. Pereira²³ · Michael M. Province⁵ · Bruce M. Psaty^{163,164} · Paul M. Ridker^{12,89} · Charles N. Rotimi⁶ · E. Shyong Tai^{32,107,165} · Rob M. van Dam^{32,165} · Cornelia M. van Duijn^{37,166} · Tien Yin Wong^{18,19} · Kenneth Rice¹⁶⁷ · W. James Gauderman¹⁶⁸ · Alanna C. Morrison⁷ · Kari E. North¹⁶⁹ · Sharon L. R. Kardia¹⁶ · Mark J. Caulfield^{29,38} · Paul Elliott^{46,86} · Patricia B. Munroe^{29,38} · Paul W. Franks^{61,170} · Dabeeru C. Rao² · Myriam Fornage³¹

¹ Cardiovascular Division, Department of Medicine, Washington University, St. Louis, MO 63110, USA

² Division of Biostatistics, Washington University School of Medicine, St. Louis, MO 63110, USA

³ Section of Gerontology and Geriatrics, Department of Internal Medicine, Leiden University Medical Center, Leiden 2333ZA, The Netherlands

⁴ Department of Genetic Epidemiology, University of Regensburg, 93051 Regensburg, Germany

⁵ Division of Statistical Genomics, Department of Genetics, Washington University School of Medicine, St. Louis, MO 63108, USA

⁶ Center for Research on Genomics and Global Health, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD 20892, USA

⁷ Human Genetics Center, Department of Epidemiology, Human Genetics, and Environmental Sciences, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX 77030, USA

⁸ The Institute for Translational Genomics and Population Sciences, Division of Genomic Outcomes, Department of Pediatrics, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, CA 90502, USA

⁹ Clinical and Translational Epidemiology Unit, Massachusetts General Hospital, Boston, MA 02114, USA

¹⁰ Department of Medicine, Harvard Medical School, Boston, MA 02115, USA

¹¹ Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA 02215, USA

¹² Harvard Medical School, Boston, MA 02115, USA

¹³ Department of Epidemiology, Harvard School of Public Health, Boston, MA 02115, USA

¹⁴ Centre de Bioinformatique, Biostatistique et Biologie Intégrative (C3BI), Institut Pasteur, Paris 75724, France

¹⁵ Cardiovascular Health Research Unit, Biostatistics and Medicine, University of Washington, Seattle, WA 98101, USA

¹⁶ Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, MI 48109, USA

¹⁷ Centre for Genomic & Experimental Medicine, Institute of Genetics & Molecular Medicine, University of Edinburgh, Edinburgh EH4 2XU, UK

¹⁸ Ocular Epidemiology, Singapore Eye Research Institute, Singapore National Eye Centre, Singapore 169856, Singapore

¹⁹ Ophthalmology & Visual Sciences Academic Clinical Program (Eye ACP), Duke-NUS Medical School, Singapore 169857, Singapore

²⁰ Genome Institute of Singapore, Agency for Science Technology and Research, Singapore 138672, Singapore

²¹ Postgraduate Programme in Epidemiology, Federal University of Pelotas, Pelotas, RS 96020-220, Brazil

²² Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol BS8 2BN, UK

²³ Laboratory of Genetics and Molecular Cardiology, Heart Institute (InCor), University of São Paulo Medical School, São Paulo, SP 5403000, Brazil

²⁴ Epidemiology and Biostatistics, University of Georgia at Athens College of Public Health, Athens, GA 30602, USA

²⁵ Department of Clinical Epidemiology, Leiden University Medical Center, Leiden 2333ZA, Netherlands

²⁶ Public Health Sciences, Epidemiology and Prevention, Wake Forest University Health Sciences, Winston-Salem, NC 27157, USA

- 27 Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh EH4 2XU, UK
- 28 Jackson Heart Study, Department of Medicine, University of Mississippi Medical Center, Jackson, MS 39213, USA
- 29 Clinical Pharmacology, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK
- 30 Human Genomics Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA, USA
- 31 Institute of Molecular Medicine, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX 77030, USA
- 32 Saw Swee Hock School of Public Health, National University Health System and National University of Singapore, Singapore 117549, Singapore
- 33 Department of Biostatistics, University of Michigan, Ann Arbor, MI 48109, USA
- 34 Icelandic Heart Association, Kopavogur 201, Iceland
- 35 Health Disparities Research Section, Laboratory of Epidemiology and Population Sciences, National Institute on Aging, National Institutes of Health, Baltimore, MD 21224, USA
- 36 Department of Public Health Sciences, Loyola University Chicago, Maywood, IL 60153, USA
- 37 Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands
- 38 NIHR Barts Cardiovascular Biomedical Research Unit, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, London EC1M 6BQ, UK
- 39 Biostatistics, Boston University School of Public Health, Boston, MA 02118, USA
- 40 Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu 51010, Estonia
- 41 Department of Biotechnology, Institute of Molecular and Cell Biology, University of Tartu, Tartu 51010, Estonia
- 42 CNRS UMR 8199, European Genomic Institute for Diabetes (EGID), Institut Pasteur de Lille, University of Lille, Lille 59000, France
- 43 Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Stockholm 17177, Sweden
- 44 Unit of Epidemiology, Biostatistics and Biodemography, Department of Public Health, Southern Denmark University, Odense 5000, Denmark
- 45 Biostatistical Sciences, Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA
- 46 Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London W2 1PG, UK
- 47 Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina 45110, Greece
- 48 Molecular Genetics and Genomics Program, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA
- 49 Medical Genetics, Department of Medicine, Surgery and Health Sciences, University of Trieste, Trieste 34100, Italy
- 50 Institute for Maternal and Child Health–IRCCS “Burlo Garofolo”, Trieste 34100, Italy
- 51 Department of Psychology, Centre for Cognitive Ageing and Cognitive Epidemiology, The University of Edinburgh, Edinburgh EH8 9JZ, UK
- 52 Department of Occupational and Environmental Health and State Key Laboratory of Environmental Health for Incubating, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China
- 53 Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany
- 54 Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany
- 55 Unit of Cardiovascular and Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm 17177, Sweden
- 56 Department of Population and Quantitative Health Sciences, Case Western Reserve University, Cleveland, OH 44106, USA
- 57 Department of Mathematics and Statistics, University of Minnesota, Duluth, MN 55812, USA
- 58 Department of Clinical Chemistry, Fimlab Laboratories, Tampere 33520, Finland
- 59 Department of Clinical Chemistry, Finnish Cardiovascular Research Center - Tampere, Faculty of Medicine and Health Technology, University of Tampere, Tampere 33014, Finland
- 60 University of Groningen, University Medical Center Groningen, Department of Epidemiology, Groningen 9700RB, The Netherlands
- 61 Department of Clinical Sciences, Genetic and Molecular Epidemiology Unit, Lund University Diabetes Centre, Skåne University Hospital, Malmö, Skåne 205 02, Sweden
- 62 Foundation for Research in Health Exercise and Nutrition, Kuopio Research Institute of Exercise Medicine, Kuopio 70100, Finland
- 63 College of Medicine, Biological Sciences and Psychology, Health Sciences, The Infant Mortality and Morbidity Studies (TIMMS), Leicester LE1 7RH, UK
- 64 Department of Computational Biology, University of Lausanne, Lausanne 1011, Switzerland
- 65 Swiss Institute of Bioinformatics, Lausanne 1015, Switzerland
- 66 Division of Epidemiology, Department of Medicine, Vanderbilt University School of Medicine, Nashville, TN 37203, USA
- 67 Division of Sleep and Circadian Disorders, Brigham and Women’s Hospital, Boston, MA 02115, USA

- 68 Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine, Tokyo 1628655, Japan
- 69 University of Groningen, University Medical Center Groningen, Department of Cardiology, Groningen 9700, The Netherlands
- 70 Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, MI 48104, USA
- 71 Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, 9713GZ Greifswald, Germany
- 72 DZHK (German Centre for Cardiovascular Health), Partner Site Greifswald, 17475 Greifswald, Germany
- 73 Division of General Internal Medicine, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21287, USA
- 74 McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA
- 75 Dean's Office, University of Kentucky College of Public Health, Lexington, KY 40536, USA
- 76 Human Genome Sequencing Center, Baylor College of Medicine, Houston, TX 77030, USA
- 77 Cardiovascular Health Research Unit, Medicine, University of Washington, Seattle, WA 98101, USA
- 78 Section of Genomic Pediatrics, Department of Pediatrics, Medicine and Physiology, Medical College of Wisconsin, Milwaukee, WI 53226, USA
- 79 Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC 27109, USA
- 80 Statistics Unit, Singapore Eye Research Institute, Singapore National Eye Centre, Singapore 169856, Singapore
- 81 Ninewells Hospital & Medical School, University of Dundee, Dundee, Scotland DD1 9SY, UK
- 82 Department of Medicine, Faculty of Medicine, University of Kelaniya, Ragama, Sri Lanka
- 83 Tropical Metabolism Research Unit, Tropical Medicine Research Institute, University of the West Indies, Mona JMAAW15, Jamaica
- 84 Cardiology, Medicine, University of Mississippi Medical Center, Jackson, MS 39216, USA
- 85 Braun School of Public Health, Hebrew University-Hadassah Medical Center, Jerusalem 91120, Israel
- 86 MRC-PHE Centre for Environment and Health, Imperial College London, London W2 1PG, UK
- 87 Cardiovascular Unit, Bioclinicum, Department of Medicine, Karolinska Hospital, Stockholm 17164, Sweden
- 88 Division of Cardiovascular Medicine, Department of Clinical Sciences, Danderyd University Hospital, Stockholm 18288, Sweden
- 89 Brigham and Women's Hospital, Boston, MA 02215, USA
- 90 Department of Epidemiology, State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China
- 91 Laboratory of Epidemiology and Population Sciences, National Institute on Aging, National Institutes of Health, Bethesda, MD 20892, USA
- 92 Epidemiology, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA 70112, USA
- 93 Medicine, Tulane University School of Medicine, New Orleans, LA 70112, USA
- 94 Institute of Biomedicine, School of Medicine, University of Eastern Finland, Kuopio Campus, Kuopio 70211, Finland
- 95 Institute of Clinical Medicine, Internal Medicine, University of Eastern Finland, Kuopio 70211, Finland
- 96 Department of Paediatrics, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119228, Singapore
- 97 Khoo Teck Puat – National University Children's Medical Institute, National University Health System, Singapore 119228, Singapore
- 98 Cardiovascular Genetics, Department of Internal Medicine, University of Utah, Salt Lake City, UT 84108, USA
- 99 Weill Cornell Medicine in Qatar, Doha, Qatar
- 100 Department of Radiology and Nuclear Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands
- 101 Department of Biostatistics, University of Alabama at Birmingham, Birmingham, AL 35294, USA
- 102 Department of Clinical Physiology, Tampere University Hospital, Tampere 33521, Finland
- 103 Department of Clinical Physiology, Finnish Cardiovascular Research Center–Tampere, Faculty of Medicine and Health Technology, University of Tampere, Tampere 33014, Finland
- 104 Department of Biochemistry, National University of Singapore, Singapore 117596, Singapore
- 105 Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen DK-2200, Denmark
- 106 Department of Environmental Medicine and Public Health, The Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
- 107 Health Services and Systems Research, Duke-NUS Medical School, Singapore 169857, Singapore
- 108 Public Health Sciences, Biostatistics and Data Science, Wake Forest University Health Sciences, Winston-Salem, NC 27157, USA
- 109 Institute of Human Genetics, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany
- 110 Institute of Human Genetics, Technische Universität München, 80333 Munich, Germany

- 111 Department of Psychiatry, Amsterdam Neuroscience and Amsterdam Public Health Research Institute, VU University Medical Center, Amsterdam 1081 BT, The Netherlands
- 112 Department of Public Health and Primary Care, Leiden University Medical Center, Leiden 2333ZA, Netherlands
- 113 Molecular Genetics Section, Laboratory of Neurogenetics, National Institute on Aging, Bethesda, MD 20895, USA
- 114 Data Tecnica International, Glen Echo, MD 20812, USA
- 115 Department of Cardiovascular Sciences, University of Leicester, Leicester LE3 9QP, UK
- 116 NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester LE3 9QP, UK
- 117 Department of Epidemiology, University of Colorado Denver, Aurora, CO 80045, USA
- 118 Division of Endocrinology, Diabetes, and Nutrition, University of Maryland School of Medicine, Baltimore, MD, USA
- 119 Program for Personalized and Genomic Medicine, University of Maryland School of Medicine, Baltimore, MD, USA
- 120 Department of Medicine, University of Ibadan, Ibadan, Oyo, Nigeria
- 121 British Heart Foundation Glasgow Cardiovascular Research Centre, Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8TA, UK
- 122 Biochemistry, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA
- 123 Department of Medicine, Geriatrics Section, Boston Medical Center, Boston University School of Medicine, Boston, MA 02118, USA
- 124 DZHK (German Centre for Cardiovascular Research), Partner Site Munich Heart Alliance, 85764 Neuherberg, Germany
- 125 Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, 17475 Greifswald, Germany
- 126 University of Split School of Medicine, Split, Croatia
- 127 University Hospital Split, Split, Croatia
- 128 Psychiatric Hospital "Sveti Ivan", Zagreb, Croatia
- 129 Division of Genetic and Genomic Medicine, Department of Pediatrics, University of California, Irvine, CA 92868, USA
- 130 Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh EH8 9AG, UK
- 131 Division of Epidemiology & Community Health, School of Public Health, University of Minnesota, Minneapolis, MN 55454, USA
- 132 Division of Preventive Medicine, School of Medicine, University of Alabama at Birmingham, Birmingham, AL 25249, USA
- 133 Division of Research, Kaiser Permanente of Northern California, Oakland, CA, USA
- 134 Alzheimer Scotland Dementia Research Centre, The University of Edinburgh, Edinburgh EH8 9AZ, UK
- 135 Centre for Cognitive Ageing and Cognitive Epidemiology, The University of Edinburgh, Edinburgh, UK
- 136 Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany
- 137 Institute of Medical Informatics Biometry and Epidemiology, Ludwig-Maximilians-Universität München, 80539 Munich, Germany
- 138 University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen 9700RB, The Netherlands
- 139 Institute for Community Medicine, University Medicine Greifswald, 17475 Greifswald, Germany
- 140 Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands
- 141 Beijing Institute of Ophthalmology, Beijing Tongren Eye Center, Beijing Ophthalmology and Visual Science Key Lab, Beijing Tongren Hospital, Capital Medical University, 100730 Beijing, China
- 142 Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, 100730 Beijing, China
- 143 Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA 15261, USA
- 144 Division of Cancer Control and Population Sciences, UPMC Hillman Cancer, University of Pittsburgh, Pittsburgh, PA 15232, USA
- 145 Behavioral Epidemiology Section, Laboratory of Epidemiology and Population Sciences, National Institute on Aging, National Institutes of Health, Baltimore, MD 21224, USA
- 146 Department of Internal Medicine B, University Medicine Greifswald, 17475 Greifswald, Germany
- 147 Broad Institute of the Massachusetts Institute of Technology and Harvard University, Boston, MA 02142, USA
- 148 Section on Nephrology, Department of Internal Medicine, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA
- 149 Department of Genomics of Common Disease, Imperial College London, London W12 0NN, UK
- 150 German Center for Diabetes Research (DZD e.V.), 85764 Neuherberg, Germany
- 151 Department of Ophthalmology, Medical Faculty Mannheim, University Heidelberg, 68167 Mannheim, Germany
- 152 Beijing Institute of Ophthalmology, Beijing Ophthalmology and Visual Science Key Lab, Beijing Tongren Eye Center, Capital Medical University, 100730 Beijing, China
- 153 Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA 15261, USA
- 154 Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, Kuopio 70211, Finland
- 155 Lifelines Cohort, Groningen 9700RB, The Netherlands

- ¹⁵⁶ Department of Medicine, Internal Medicine, Lausanne University Hospital and University of Lausanne, Lausanne 1011, Switzerland
- ¹⁵⁷ Durrer Center for Cardiogenetic Research, ICIN-Netherlands Heart Institute, Utrecht, The Netherlands
- ¹⁵⁸ Faculty of Medicine, University of Iceland, Reykjavik 101, Iceland
- ¹⁵⁹ Sticht Center for Health Aging and Alzheimer's Prevention, Department of Internal Medicine, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA
- ¹⁶⁰ NHLBI Framingham Heart Study, Framingham, MA 01702, USA
- ¹⁶¹ Population Sciences Branch, Division of Intramural Research, National Heart, Lung, and Blood Institute, NIH, Bethesda, MD 20892, USA
- ¹⁶² Division of General Medicine, Department of Medicine, Columbia University Medical Center, New York, NY 10032, USA
- ¹⁶³ Cardiovascular Health Research Unit, Epidemiology, Medicine and Health Services, University of Washington, Seattle, WA 98101, USA
- ¹⁶⁴ Kaiser Permanente Washington Health Research Institute, Seattle, WA 98101, USA
- ¹⁶⁵ Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119228, Singapore
- ¹⁶⁶ Nuffield Department of Population Health, University of Oxford, Oxford, UK
- ¹⁶⁷ Department of Biostatistics, University of Washington, Seattle, WA 98195, USA
- ¹⁶⁸ Biostatistics, Preventive Medicine, University of Southern California, Los Angeles, CA 90032, USA
- ¹⁶⁹ Epidemiology, University of North Carolina Gilling School of Global Public Health, Chapel Hill, NC 27514, USA
- ¹⁷⁰ Department of Public Health & Clinical Medicine, Umeå University, Umeå, Västerbotten 901 85, Sweden