



Universiteit  
Leiden

The Netherlands

**Evaluation and improvement of integrated cardiac care:  
efficient and effective care in patients with chronic  
coronary artery disease and chronic heart failure**

Vester, M.P.M.

**Citation**

Vester, M. P. M. (2022, December 13). *Evaluation and improvement of integrated cardiac care: efficient and effective care in patients with chronic coronary artery disease and chronic heart failure*. Retrieved from <https://hdl.handle.net/1887/3497444>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3497444>

**Note:** To cite this publication please use the final published version (if applicable).

# CHAPTER 4

## UTILIZATION OF DIAGNOSTIC RESOURCES AND COSTS IN PATIENTS WITH SUSPECTED CARDIAC CHEST PAIN

M.P.M. Vester<sup>1</sup>, D.C. Eindhoven<sup>1</sup>, T.N. Bonten<sup>2</sup>, H. Wagenaar<sup>3</sup>, H.J. Holthuis<sup>3</sup>, M.  
J. Schalij<sup>1</sup>, G.J. de Grooth<sup>1</sup>, P.R.M. van Dijkman<sup>1</sup>

<sup>1</sup>Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands

<sup>2</sup>Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands

<sup>3</sup>Performation-HOTflo, Bilthoven, The Netherlands

## ABSTRACT

**Objective:** Non-acute chest pain is a common complaint and can be caused by various conditions. With the rising healthcare expenditures of today, it is necessary to use our healthcare resources effectively. This study aims to give insight into the diagnostic effort and costs for patients with non-acute chest pain.

**Design and methods:** Financial data of patients without a cardiac history from 4 hospitals (January 2012–October 2018), who were registered with the national diagnostic code 'No cardiac pathology' (ICD-10 Z13.6), 'Chest wall syndrome' (ICD-10 R07.4) or 'stable angina pectoris' (ICD-10 I20.9) were extracted.

**Results:** In total, 74,091 patients were included for analysis and divided into the following final diagnosis groups: No cardiac pathology: N= 19,688 (age 53±18), 46% male), Chest wall syndrome: N= 40,858 (age 56±15), 45% male), and stable angina pectoris: N= 13,545 (age 67±11), 61% male). A total of approximately €142.7 million was spent during diagnostic work-up. The total expenditure during diagnostic effort was €1.97, €8.13, and €10.7 million respectively for no cardiac pathology, chest wall syndrome, and stable AP per year. After 8 years follow up ≥95% of the patients diagnosed with no cardiac pathology or chest wall syndrome had an (cardiac) ischemic free survival.

**Conclusion:** The diagnostic expenditure and clinical effort to ascertain non-cardiac chest pain is high. We should define what we as society find acceptable as 'assurance costs' with an increasing pressure on the healthcare system and costs.

## INTRODUCTION

Chest pain may be a symptom of ischemic heart disease.(1) Every year 0.7-2.7% of the general population consult the general practitioner for having chest pain.(2-5) Multiple underlying causes have been described varying from a musculoskeletal origin, gastro-oesophageal reflux disease to potential life-threatening events such as coronary artery disease (CAD). Fortunately, only a minority of patients have chest pain symptoms due to ischemic heart disease.(6)

Because of limited diagnostic resources in a primary care setting, patients suspected of a cardiac cause of chest pain are often referred to a cardiologist for additional diagnostic testing. However, of the referred patients with chest pain, no cardiac cause has been found in 55% to 90%.(5, 7) As the health expenditures are increasing, so does the need to spend the available resources sparingly.(8) Ideally, healthcare providers aim to offer the highest quality of care and use the minimal required additional diagnostic procedures to make the correct diagnosis. On the other hand, in the diagnostic trajectory of a patient with chest pain, a wide variety of diagnostic tests are available and it appears that besides the use of guidelines, the choice of diagnostics depends on the opinion of the consultant doctor and the location of the consultation.(9-11)

Numerous studies have investigated the final diagnosis of patients with chest pain in primary care.(2, 3, 5, 12) However, to our knowledge, studies examining the healthcare trajectory of patients with non-acute, or chronic, chest pain in a hospital setting are scarce. A recent study verified that financial databases are a valid source of information in the evaluation of a patient's care chain.(13) Given the high prevalence of chest pain and high healthcare expenditures on CAD, investigating this care chain is of relevance for the everyday cardiology practice. The current study aims to gain insight into the amount of referred patients, utilization of diagnostic resources and costs, and clinical outcome in patients with non-acute chest pain in a hospital setting using financial data.

## METHODS

### Patient population and data sources

All Dutch citizens are covered by a basic mandatory insurance. Treatments and diagnoses supplied by health services are coded according to a national financial coding system (DOT: Diagnose Behandelings Combinatie (DBC) Op weg naar Transparantie) combined with the World Health Organization (WHO) International Classification of Diseases (ICD). Declaration data from unique patients of 18 years and older with a new onset of suspected cardiac chest pain seen in the outpatient clinic were extracted from the financial database of Performance-HOTflo from January 2012 until October 2018.

Performance-HOTflo (Bilthoven, the Netherlands) is a healthcare consultancy company that provides patient-level costing and benchmarking products for different healthcare services across Europe.<sup>(14)</sup> Hospital selection was based on the region of South-Holland (Zuid-Holland) and the availability of cost-price information by Performance-HOTflo. The relevant hospitals were requested to give consent for using their data for this study. Four hospitals, of which one academic hospital and three regional hospitals, gave consent.

Suspected cardiac chest pain was defined by the following three diagnosis codes 'No cardiac pathology' (DOT code 0320.101 similar to ICD-10 Z13.6), 'Chest wall syndrome' (DOT code 0320.201 similar to ICD-10 R07.4) or 'stable angina pectoris' (DOT code 0320.202 similar to ICD-10 I20.9). The diagnosis code was recorded after the first diagnosis. Subsequently, patients with an ischemic cardiac history or with another cardiac history were excluded from the analysis. The remaining patients were divided into three different groups based on diagnosis code:

- Group I: No cardiac pathology. Also defined as chest pain of no cardiac origin.
- Group II: Chest wall syndrome. Also defined as chest pain of no cardiac origin.
- Group III: Stable Angina Pectoris. Also defined as cardiac chest pain.

In a combination of two or more previously mentioned diagnosis codes within the same patient, the code Stable Angina Pectoris had priority above Chest wall syndrome and No Cardiac Pathology. The code Chest wall syndrome had priority above No Cardiac Pathology.

### Activities

The following characteristics were retrieved after first diagnosis registration: age, gender and all healthcare utilization with the associated admission. To gain more insight into the care process and expenditures, the used resources were divided into the following categories: 'Cardiac Invasive Diagnostics or Treatment', 'Cardiac non-

invasive diagnostics', 'Emergency Department', 'Inpatient care', 'Outpatient care' and 'Other'. Other included i.e. the use of materials and administrative costs. (Supplemental Table 1).

### Cost analysis

By using Time-Driven Activity-Based Costing (TD-ABD) methodology costs were calculated at patient level resource utilization.<sup>(15)</sup> TD-ABD is a micro-costing method, and calculates two parameters per activity: the costs per time unit to perform each activity and the overall time units spent performing the activity. As cost price calculations are standardized by Performance-HOTflo, it was possible to compare participating hospitals. The database contained information about the period of treatment, the differently registered diagnostics and the registered interventions. All available data from January 2012 until October 2018 were obtained and the most recent cost price model was used for calculations. Therefore differences in the cost price calculations due to inflation were avoided.

### Statistical analyses

Continuous data are presented as mean and 95% CI when normally distributed. Categorical data are presented as numbers and percentages. A chi-square test was used for comparing the baseline characteristics of the different patient groups. P-values <0.05 were considered statistically significant. The duration of follow-up was calculated from the date when the first diagnosis code was registered, the inclusion of the patient until the last date activity for one of the above-mentioned diagnosis or resource code was registered. The time to an ischemic event after the inclusion of a patient is presented in a Kaplan Meier plot. An ischemic event was based on diagnosis codes and resource use after the first registration:

- Unstable angina pectoris or Infarction: patients who developed unstable angina pectoris (DOT code 0320.11.203 similar to ICD-10 I20.0), a Non-ST-Segment Elevation Myocardial Infarction (NSTEMI) (DOT code 0320.11.205 similar to ICD-10 I21.4), a ST-Segment Elevation Myocardial Infarction (STEMI) (DOT code 0320.11.204 similar to ICD-10 I21.9) or who were followed up after acute coronary syndrome (DOT code 0320.11.801 similar to ICD-10 Z86.7).
- Percutaneous Coronary Intervention (PCI) or Coronary artery bypass grafting (CABG): patient who underwent a PTCA or a CABG (DOT code 0320.11.802 or 0320.11.810 similar to ICD-10 Z09.0) and the cardiothoracic declaration codes from the Performance Hotflo database (Cardiothoracic (CTC) codes for a CABG coded as 2320, 2400, 2415, 2425 or 2470 similar to ICD-10-PCS codes based on procedure).

Statistical analyses were performed using SPSS (version 23.0, IBM SPSS Statistics).


### **Ethical considerations**

Performation-Hotflo is ISO 27001 and NEN 7510 certified, meaning that all patient data are used in strict confidence. All members of the research team signed a statement of confidentiality for processing the data. The participating hospitals formally consented to participate. The local medical ethics committee of the Leiden University Medical Center, in its capacity as the coordinating center of this retrospective study, approved the study design and waived the need for individual patient informed consent.

## **RESULTS**

### **Population**

In total, 90.436 unique patients in four hospitals between January 2012 until October 2018 were included. Patients with a history of ischemic cardiac disease (N=7.805, 9%) or another cardiac history (N=8.540, 9%) were excluded from analysis. Subsequently, a total of 74.091 (mean age  $57 \pm 16$ , 48% men) with no cardiac history and new onset of cardiac chest pain remained. Of these patients, 19.688 (27%) had no cardiac pathology, 40.858 (55%) had chest wall syndrome and 13.545 (18%) had stable angina pectoris as the final diagnosis (Fig. 1). The stable AP patients (age  $67 \pm 11$ , men 61%) were older and consisted of more men in comparison to both other patient groups (non-cardiac chest pain, no cardiac pathology (age  $53 \pm 18$ , 46% men) and chest wall syndrome (age  $56 \pm 15$ , 45% men). (Table 1)



**Figure 1.** Flow chart of included patients. Patients were subdivided into groups, based on the different diagnostic codes: No cardiac pathology\* (coded as 0320101 similar to ICD-10 Z13.6), Idiopathic thoracic complaints (coded as 0320.201 similar to ICD-10 R07.4) and stable angina pectoris (AP) (coded as 0320.202 similar ICD-10 I20.9).

\*Ischemic Cardiac History: see Table S1.

\*\*Other Cardiac History: see Table S2


**Table 1.** Baseline characteristics

|                    | No Cardiac Pathology<br>N = 19,688 (27%) | Chest Wall Syndrome<br>N = 40,858 (55%) | Stable AP<br>N = 13,545 (18%) | Total<br>N = 74,092 | P-value |
|--------------------|------------------------------------------|-----------------------------------------|-------------------------------|---------------------|---------|
| Age (mean±SD )     | 53<br>±18                                | 56<br>±15                               | 67<br>±11                     | 57<br>±16           | P<0.001 |
| Gender (n. % male) | 8,971<br>46%                             | 18,477<br>45%                           | 8,203<br>61%                  | 35,651<br>48%       | P<0.001 |

Stable AP = stable angina pectoris

### Activities and costs


A total of €142,702,110 was spent during the diagnostic work-up period. The group with no cardiac pathology covered 9% (€13,485,941) of the total costs, the group with chest wall syndrome 39% (€55,557,410). The majority of the expenditures were covered by the group with stable AP (52%, €73,658,759). Fig. 2a shows the overall costs per year per diagnosis: €1,973,552 in the no cardiac pathology group, €8,130,352 chest wall syndrome group and €10,779,330 in the stable AP group. This corresponds with a mean expenditure of €685, €1,360 and € 5,483 per patient with no cardiac pathology, chest wall syndrome and stable AP, respectively. (Fig. 2b) As is shown in Table 2, most money was spent on 'Inpatient Care' (€ 50,428,212, 35%) followed by 'Cardiac non-invasive diagnostics' (€ 39,924,124, 28%), 'Cardiac invasive diagnostics or treatment' (€ 28,627,656, 20%), 'Outpatient Care' (€ 13,535,585, 10%), 'Emergency Department' (€ 8,275,256, 6%) and 'Other' €1,911,276, 1%). The no cardiac pathology group and the chest wall syndrome group spent more money in the Emergency department, compared with the stable AP group. A possible explanation could be that because of more uncertainty and concern these patients are referred to the emergency department to rule out an acute coronary syndrome. Table 3 shows the percentage of patients per group who underwent non-invasive diagnostic tests. In all groups, almost every patient underwent a non-invasive diagnostic test. The most applied non-invasive diagnostic test was a ECG in no cardiac pathology, chest wall syndrome, and stable AP, 80%, 87%, 79%, respectively. Followed by laboratory tests in the group with chest wall syndrome (76%) and stable AP (76%) and cardiac ultrasound in the group with no cardiac pathology (54%). The proportion of applied diagnostics in the groups with chest wall syndrome and stable AP were comparable with each other.



**Figure 2.** A total of €142,702,110 (2011-2018) was spent between January 2012 until October 2018. Panel A shows the total costs in the group with no cardiac pathology (N=19,688), chest wall syndrome (N=40,858) and stable AP (13,545) in millions (€) per year. Panel B shows the mean costs (€) per patient per group. Stable AP: Stable angina pectoris.

### Follow up

Fig. 3 shows a Kaplan Meier plot of the patient with an ischemic free survival. After one year 99% of the patient in the group with no cardiac pathology and chest wall syndrome, had an ischemic free survival. The one-year ischemic free survival of the group with stable AP was 77%. The percentage of patients with an ischemic free survival at 5 and 8 years follow-up included 99% and 98% in the group with no cardiac pathology, respectively. In the group with chest wall syndrome, the ischemic free survival was 96% at 5 years follow-up and 95% at 8 years follow-up. In the group with stable AP the ischemic free survival was lower, with 66% at 5 years and 63% at 8 years. There seemed to be a slightly lower chance of developing cardiac symptoms for the 'No cardiac pathology' when compared to the 'Chest wall syndrome' group. This is surprising, since both groups are considered to not have any cardiac issues after discharge. It might be that cardiologists are more likely to give a 'No cardiac pathology' diagnosis when there is more clarity on the cause of the chest pain.



**Figure 3.** Kaplan Meier ischemic free survival curve for eight years follow-up

**Table 2.** Overview of healthcare expenditures per group

| Category (N,%)*      | No Cardiac Pathology<br>N = 19,688 (27%) | Chest Wall Syndrome<br>N = 40,858 (55%) | Stable AP<br>N = 13,545 (18%) | Total                |
|----------------------|------------------------------------------|-----------------------------------------|-------------------------------|----------------------|
| Cardiac Invasive     | € 1.112.551<br>8                         | € 4.054.831<br>7                        | € 23.460.274<br>32            | € 28.627.656<br>20   |
| Cardiac Non Invasive | € 4.945.849<br>37                        | € 19.902.132<br>36                      | € 15.076.143<br>21            | € 39.924.124<br>28   |
| Emergency Department | € 1.682.815<br>13                        | € 4.881.403<br>9                        | € 1.711.038<br>2              | € 8.275.256<br>6     |
| Inpatient Care       | € 3.498.977<br>26                        | € 20.832.042<br>38                      | € 26.097.193<br>35            | € 50.428.212<br>35   |
| Outpatient Care      | € 2.142.629<br>16                        | € 5.743.350<br>10                       | € 5.649.606<br>8              | € 13.535.585<br>10   |
| Other                | € 103.119<br>1                           | € 143.653<br>0                          | € 1.664.504<br>2              | € 1.911.276<br>1     |
| Total                | € 13.485.941<br>9                        | € 55.557.410<br>39                      | € 73.658.759<br>51            | € 142.702.110<br>100 |

Stable AP = Stable angina pectoris

\*Distinction of the different categories can be seen in Supplemental Table 1.

**Table 3.** The amount of patients (in percentages) that underwent a non-invasive diagnostic test per group

|                                 | No Cardiac Pathology<br>N = 19.688 (27%) |    | Chest Wall Syndrome<br>N = 40.858 (55%) |    | Stable AP<br>N = 13.545 (18%) |    |
|---------------------------------|------------------------------------------|----|-----------------------------------------|----|-------------------------------|----|
| Non Invasive Diagnostics (N, %) | 18.783                                   | 90 | 40.461                                  | 99 | 12.545                        | 93 |
| ECG                             | 15.682                                   | 80 | 35.646                                  | 87 | 10.734                        | 79 |
| Cardiac ultrasound              | 10.678                                   | 54 | 16.216                                  | 40 | 8.672                         | 64 |
| Laboratory                      | 7.762                                    | 39 | 30.857                                  | 76 | 10.232                        | 76 |
| Exercise test                   | 3.780                                    | 19 | 19.638                                  | 48 | 7.217                         | 53 |
| X-Ray                           | 2.562                                    | 13 | 14.986                                  | 37 | 5.628                         | 42 |
| Holter monitor                  | 2.440                                    | 12 | 3.907                                   | 10 | 1.731                         | 13 |
| Cardiac CT scan                 | 1.233                                    | 6  | 9.845                                   | 24 | 3.256                         | 24 |
| Cardiac Nuclear/SPECT scan      | 510                                      | 3  | 3.585                                   | 9  | 2.787                         | 21 |
| Rhythm monitoring               | 469                                      | 2  | 4.154                                   | 10 | 2.732                         | 20 |
| Cardiac MRI                     | 268                                      | 1  | 1.793                                   | 4  | 1.331                         | 10 |
| ABPM                            | 135                                      | 1  | 62                                      | 2  | 447                           | 3  |

Stable AP = Stable angina pectoris; ECG = electrocardiogram; Cardiac CT scan = Cardiac Computed Tomography scan; SPECT = Single Photon Emission Computed Tomography; MRI = Magnetic Resonance imaging; ABPM = Ambulatory Blood Pressure Monitoring

## DISCUSSION

This study was conducted to gain insight into the diagnostic effort for patients with non-acute chest pain in the hospital. The findings of this study can be summarized as follows: (i) 82% of the patients referred to the hospital with suspicion of cardiac chest pain had a non-cardiac origin; (ii) a total of € 142.7.- million has been spent on 74.091 patients, of which €67 million was spent on non-cardiac patients; (iii) after 8 years follow up ≥95% of the patient diagnosed with no cardiac pathology or chest wall syndrome had an ischemic free survival. Data from one tertiary and three general hospitals was analysed. Two of the four hospitals were percutaneous coronary intervention-centers. A total of 180.000 patients without a cardiac history are observed with chest pain per year in the Netherlands.(9) In this study, 74.091 patients were included over a period of 5,7 years. This represents approximately 13.000 patients per year amounting to 7% of the national population. Patients with a cardiac history (18%, Fig. 1) were excluded, this will lead to an underestimation of the total incidence of non-acute cardiac chest pain. These patients were excluded to facilitate interpretation of the data obtained, as patients have the same cardiac history (i.e. none). Compared with previous studies of chest pain in a non-acute setting baseline characteristics were comparable. (5, 16)

Since the cohort can be considered representable for the national population, we believe that the findings in this study are representable too.

### **The accuracy of patient referral**

The current study showed a high incidence of non-cardiac chest pain in the referred patients (82%): 27% of the patients were diagnosed with no cardiac pathology and 55% of the patients were diagnosed with chest wall syndrome. As the patients were referred to the hospital with the suspicion of a cardiac cause, the 'hit rate' of an actual cardiac cause, for non-acute cases, seems low.

A study from Dumville et al. studied the long-term outcome of patients with chest pain who were referred from primary care towards a Rapid Access Chest Pain clinic in a non-acute setting.<sup>(17)</sup> A total of 52% had non-cardiac chest pain after a follow-up of 6 months. Similar results were found in a study by Byrne et al., where 633 patients were referred to a rapid access chest pain clinic.<sup>(16)</sup> An incidence of low risk or non-cardiac chest pain in 51% of the patients was found after 8 months. In the current study, an incidence of 82% was found. The large difference can have several causes. For instance, the current study included patients based only on the initial diagnosis. In contrast, both other studies allow for a change in diagnosis during follow-up (i.e. from non-cardiac to cardiac). Furthermore, both studies by Dumville et al. and Byrne et al. were performed in the UK. It is possible that the country-specific circumstances with regard to referral to hospitals in the case of non-acute chest pain are different, resulting in a seemingly higher hit-rate of the British GPs in the described situation.

### **Costs and healthcare utilization.**

The total cost to obtain the diagnoses 'no cardiac pathology' was €13.5 million (€685 per patient). For 'chest wall syndrome' the total diagnostic expenditure was €55.5 million (€1360 per patient). A study from Mourad et al. analyzed the extent of costs in secondary care, incurred by patients (N=199) with non-cardiac chest pain and compared this with the costs of patients with acute myocardial infarction and AP.<sup>(18)</sup> The annual cost per patient with non-cardiac chest pain was €10.068, which included also costs in primary care, indirect costs on productivity, loss due to sick-leave and medical costs. In the current study, only the direct costs (TD-ABC) were calculated per patient and could explain the lower costs found in the current study.

To extrapolate these costs to the total Dutch population, data from Dutch National Healthcare Institute is used.<sup>(9)</sup> In this report, the annual amount of new patients with diagnostic codes 201 and 202 were reported to be n = 180.000. In this report, the diagnostic codes 201 (55%), 202 (18%), and 101 (27%) were included. To accurately compare the data, the mentioned 180.000 only represents 73% of the hypothetical national cohort as considered in this study. The total cohort would then be n = 247.000. The annual diagnostic cost for 'no cardiac pathology' is then €45.6 million, and the

annual diagnostic cost for 'chest wall syndrome' is €184,4 million. The total annual expenditure to ascertain the absence of a cardiac cause for chest pain is then €230 million. We could call these 'assurance costs'.

In the current study, referred patients were subjected to a wide variety of diagnostic tests. Furthermore the utilization rate of non-invasive diagnostics was high in all groups:  $\geq 90\%$ . Hoornweg et al. investigated the utilization of diagnostic tests in an observational study, by including patients (N=281) with acute and non-acute chest pain in primary care.<sup>(5)</sup> A total of 44% of the patients underwent diagnostic testing. The higher utilization rate of the current study might be explained by the secondary care setting and the focus on non-acute chest pain: the clinical presentation might not be as clear as in an acute setting and therefore requires more diagnostic tests. In addition, the high utilization rate emphasizes the need to increase the compliance with the current guidelines. The ESC-guidelines recommend no non-invasive testing in patients with a pre-test probability (PTP) score of  $<15\%$ .<sup>(19, 20)</sup> The guidelines do leave room for testing below 15% in particular when symptoms are limiting, but this should only apply to a small part of the cohort.<sup>(20)</sup> The present study showed a diagnosis of cardiac chest pain in 18% of the total cohort. Under the assumption that the total incidence of cardiac chest pain of 18% implies an average PTP score of 18%, it is highly unlikely that more than 90% of the cohort had a PTP of more than 15%. Over investigation is a well-known problem and paralleled with high healthcare expenditures, hence an important factor in reducing healthcare costs.<sup>(21)</sup>

### **Clinical outcome and follow up**

The current study showed that  $\geq 95\%$  of the patients in the group with no cardiac pathology and the group with chest wall syndrome had an ischemic free survival after 8 years. A total of 68% of the patient with stable angina pectoris had an ischemic free survival after 8 years. It can be concluded that the current system is very effective in distinguishing those that have an underlying cardiac cause for their chest pain and those that have not.

### **Limitations**

Some limitations should be acknowledged. First, this is an analysis based on a large-scale financial database. Previous studies showed a correlation between medical charts and financial data.<sup>(13)</sup> However, in this study, this was not verified and might result in over- or under-diagnosis. For similar data, the Dutch National Healthcare Institute used a 15% extrapolation factor. Since TD-ABD costing is used in this study, there is no available data to come to a similar extrapolation factor. Since we do not use an extrapolation factor, we are confident our costing data is conservative. Secondly, there is currently no universal definition for non-acute chest pain. Depending on the doctor, a patient with non-cardiac chest pain can be diagnosed with 'no cardiac pathology' or with 'chest wall syndrome'. For this reason, we defined non-acute

suspected cardiac chest pain by including 'no cardiac pathology', 'chest wall syndrome' or 'stable angina pectoris'. The diagnosis 'no cardiac pathology' can also be given to patients that are not referred due to chest pain. This might also be reflected in Table 3, where a different amount of diagnostic resources are used among the group with 'no cardiac pathology' compared with the groups with 'chest wall syndrome' and 'stable AP'. The focus of this study was to investigate the diagnostic course of non-acute cardiac sounding chest pain, in the outpatient clinic. By careful selection of included diagnostic codes and exclusion of emergency department patients, we are confident that the vast majority of the cohort is from the outpatient clinic. Despite this focus, it is possible that patients were included who came with chest pain in an acute setting. Fourth, information about medication was not available in the database and has not been included in our analysis. This probably leads to an underestimation of the real costs. Fifth, this research analyzed anonymous data from four different hospitals, in case a patient has been treated in multiple hospitals that patient could be included two or more times in the database. Finally, it is worth noting that the incidence of ECGs was not nearing 100%, which one would expect based on diagnostic guidelines (20, 22, 23) It is likely that not all ECGs were accounted for in the declaration database. In fact it can be expected that an ECG was made in all patients.

### Future implications

The Dutch National Healthcare Institute showed that the current guidelines were not always followed in obtaining a diagnosis. This might be a reason for the large variety of applied diagnostics. This is also shown in the current study. The variety of applied diagnostics was large and most notably, the amount of performed exercise ECGs was large, even though the sensitivity of this test is quite low at 58% (95% CI 46-69%).(19) The ESC and NICE guidelines clearly underline the importance of a careful history ie. a clear distinction between typical, atypical and non-specific chest pain in the diagnosis of cardiac sounding chest pain.(20, 23) The outcomes in the study by Sekhri et al. supported the distinctions (atypical, typical and non-specific) in these guidelines. (24) It is worthwhile to investigate why guidelines are frequently not followed, as the goal of these guidelines is to deliver effective and efficient healthcare.

The mentioned 'assurance costs' are currently considered a necessary expenditure. As discussed earlier, to stay ahead of the curve with regard to the rising healthcare costs, it follows that we should search for ways to drive the assurance costs down. One such way could be to switch to an integrated care model, which has shown promise in earlier research. (25) Integrated care may play a large role in value based-healthcare; measuring costs in combination with health outcomes and patient experiences. (26) Several previous studies correlated patient-level costs and outcome analyses to improvement in healthcare quality.(27, 28) Nation-wide quality measurement and benchmarking feedback are already conducted in surgery.(28) In future perspective and in the pursuit of improving cardiac care, extending the analysis of this study by

benchmarking the hospitals on a nation-wide level, or benchmarking the performances of healthcare systems in countries can provide useful data. Similar studies such as this study can be performed for other diagnostic courses where there is a high expenditure, and a desire to find out where the effort is spent. Mourad et al. showed that patients with no cardiac chest pain use a significant amount of healthcare resources and costs society a substantially amount of money. (18) To gain insight in the entire diagnostic course of non-acute chest pain and its corresponding costs, it would be interesting to investigate also the behaviour and potential recurrent referrals of patients who were diagnosed with non-cardiac chest pain.

## **Conclusion**

The prevalence of patients with non-acute chest pain is high and a significant amount of €142.7 million is spent on this particular patient group of 74.091 patients. This included a mean expenditure of €685, €1.360 and € 5.483 per patient with no cardiac pathology (27% of cohort), chest wall syndrome (55%) or stable AP (18%), respectively. In the majority (82%) no cardiac cause of chest pain was found. Furthermore, ≥95% of the patients in the group with no cardiac pathology and chest wall syndrome did not develop an ischemic event after 8 years follow-up.

In conclusion, it is found that the Dutch healthcare system is very effective in determining underlying cardiac cause for non-acute chest pain. Extrapolating the data, it is also found that the diagnostic expenditure to ascertain non-cardiac chest pain is €230 million per year in the Netherlands. Not only is the monetary expenditure high, the time expenditure by the healthcare system is also high. In a time where the healthcare workers are under a permanently high workload, and where recent events have shown that the healthcare system needs extra capacity to deal with crises, we should define what we as society find acceptable as 'assurance costs'.

## **Acknowledgements**

The data underlying this article were provided by Performance-HOTflo after approval of the concerned hospitals. Data will be shared on request to the corresponding author with permission of Performance-HOTflo. Conflict of Interest: none declared

## REFERENCES

- Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;3(11).
- Ruigomez A, Rodriguez LA, Wallander MA, Johansson S, Jones R. Chest pain in general practice: incidence, comorbidity and mortality. *Fam Pract*. 2006;23(2):167-74.
- Bosner S, Haasenritter J, Becker A, Karatolios K, Vaucher P, Gencer B, et al. Ruling out coronary artery disease in primary care: development and validation of a simple prediction rule. *CMAJ*. 2010;182(12):1295-300.
- Bosner S, Becker A, Haasenritter J, Abu Hani M, Keller H, Sonnichsen AC, et al. Chest pain in primary care: epidemiology and pre-work-up probabilities. *Eur J Gen Pract*. 2009;15(3):141-6.
- Hoorweg BBN, Willemssen RTA, Cleef LE, Boogaerts T, Buntinx F, Glats JFC, et al. Frequency of chest pain in primary care, diagnostic test performed and final diagnoses. *Heart*. 2017;103:1727-32.
- Haasenritter J, Biroga T, Keunecke C, Becker A, Donner-Banzhoff N, Dornieden K, et al. Causes of chest pain in primary care – a systematic review and meta-analysis. *Croatian Medical Journal*. 2015;56(5):422-30.
- Rathod KS, Ward H, Farooqi F. Chest pain symptom scoring can improve the quality of referrals to Rapid Access Chest Pain Clinic. *BMJ Qual Improv Rep*. 2014;3(1).
- Network EH. European Cardiovascular Disease Statistics. 2017.
- Zorginstituut Nederland. Zinnige Zorg: Verbetersignalement Pijn op de borst (verdenking) Stabiele Angina Pectoris. 2017.
- Poldervaart JM, Reitsma JB, Backus BE, Koffijberg H, Veldkamp RF, Ten Haaf ME, et al. Effect of Using the HEART Score in Patients With Chest Pain in the Emergency Department: A Stepped-Wedge, Cluster Randomized Trial. *Ann Intern Med*. 2017;166(10):689-97.
- Nieuwets A, Poldervaart JM, Reitsma JB, Buitendijk S, Six AJ, Backus BE, et al. Medical consumption compared for TIMI and HEART score in chest pain patients at the emergency department: a retrospective cost analysis. *BMJ Open*. 2016;6(6):e010694.
- Nilsson S, Scheike M, Engblom D, Karlsson LG, Mölstad S, Åkerlind I, et al. Chest pain and ischaemic heart disease in primary care. *British Journal of General Practice*. 2003;53:378-82.
- Eindhoven DC, van Staveren LN, van Erkelens JA, Ikkersheim DE, Cannegieter SC, Umans V, et al. Nationwide claims data validated for quality assessments in acute myocardial infarction in the Netherlands. *Neth Heart J*. 2018;26(1):13-20.
- Perfromation-HOTflo. 2018 [11-01-2019]. Available from: <https://www.perfromation.com/>.
- Kaplan R, Anderson S. Time-Driven Activity-Based Costing. *Harvard Business Review* 2004;82(11):131-8.
- Byrne J, Murdoch D, Morrison C, McMurray J. An audit of activity and outcome from a daily and a weekly "one stop" rapid assessment chest pain clinic. *Postgrad Med J*. 2002;78:43-6.
- Dumville JC, MacPherson H, Griffith K, Miles JN, Lewin RJ. Non-cardiac chest pain: a retrospective cohort study of patients who attended a Rapid Access Chest Pain Clinic. *Fam Pract*. 2007;24(2):152-7.
- Mourad G, Alwin J, Strömberg A, Jaarsma T. Societal costs of non-cardiac chest pain compared with ischemic heart disease - a longitudinal study. *BMC Health Services Research*. 2013;13(403).
- Knuuti J, Ballo H, Juarez-Orozco LE, Saraste A, Kolh P, Rutjes AWS, et al. The performance of non-invasive tests to rule-in and rule-out significant coronary artery stenosis in patients with stable angina: a meta-analysis focused on post-test disease probability. *Eur Heart J*. 2018;39(35):3322-30.
- Knuuti J, Wijns W, Saraste A, Capodanno D, Barbato E, Funck-Brentano C, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes. *Eur Heart J*. 2019.

21. OECD. Tackling Wasteful Spending on Health 2017.
22. Task Force M, Montalescot G, Sechtem U, Achenbach S, Andreotti F, Arden C, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2013;34(38):2949-3003.
23. National Institute for Health and Care Excellence (NICE). Chest pain of recent onset: assessment and diagnosis 2010 [Available from: <https://www.nice.org.uk/guidance/cg95>].
24. Sekhri N, Feder GS, Junghans C, Hemingway H, Timmis AD. How effective are rapid access chest pain clinics? Prognosis of incident angina and non-cardiac chest pain in 8762 consecutive patients. *Heart*. 2007;93(4):458-63.
25. Vester MPM, de Grooth GJ, Bonten TN, van der Hoeven BL, de Doelder MS, Eindhoven DC, et al. Implementation of an integrated care model between general practitioner and cardiologist. *Journal of Integrated Care*. 2019;27(4):305-15.
26. Porter ME. What is value in health care? *N Engl J Med*. 2010;363(26):2477-81.
27. Lee VS, Kawamoto K, Hess R, Park C, Young J, Hunter C, et al. Implementation of a Value-Driven Outcomes Program to Identify High Variability in Clinical Costs and Outcomes and Association With Reduced Cost and Improved Quality. *JAMA*. 2016;316(10):1061-72.
28. Govaert JA. Value-Based healthcare in Colorectal Cancer Surgery. Improving quality and reducing costs. Leiden: Leiden University; 2017.



