



Universiteit
Leiden
The Netherlands

Lipoprotein(a) lowering by alirocumab reduces the total burden of cardiovascular events independent of low-density lipoprotein cholesterol lowering: ODYSSEY OUTCOMES trial

Szarek, M.; Bittner, V.A.; Aylward, P.; Baccara-Dinet, M.; Bhatt, D.L.; Diaz, R.; ... ;
ODYSSEY OUTCOMES Investigators

Citation

Szarek, M., Bittner, V. A., Aylward, P., Baccara-Dinet, M., Bhatt, D. L., Diaz, R., ...
Schwartz, G. G. (2020). Lipoprotein(a) lowering by alirocumab reduces the total burden of
cardiovascular events independent of low-density lipoprotein cholesterol lowering:
ODYSSEY OUTCOMES trial. *European Heart Journal*, 41(44), 4245-4255.
doi:10.1093/eurheartj/ehaa842

Version: Publisher's Version
License: [Creative Commons CC BY-NC 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3232713>

Note: To cite this publication please use the final published version (if applicable).

Lipoprotein(a) lowering by alirocumab reduces the total burden of cardiovascular events independent of low-density lipoprotein cholesterol lowering: ODYSSEY OUTCOMES trial

Michael Szarek ^{1*}, Vera A. Bittner ², Philip Aylward ³, Marie Baccara-Dinet ⁴, Deepak L. Bhatt ⁵, Rafael Diaz ⁶, Zlatko Fras ^{7,8}, Shaun G. Goodman ^{9,10}, Sigrun Halvorsen ¹¹, Robert A. Harrington ¹², J. Wouter Jukema ^{13,14}, Patrick M. Moriarty ¹⁵, Robert Pordy ¹⁶, Kausik K. Ray ¹⁷, Peter Sinnaeve ¹⁸, Sotirios Tsimikas ¹⁹, Robert Vogel ²⁰, Harvey D. White ²¹, Doron Zahger ²², Andreas M. Zeiher ²³, Ph. Gabriel Steg ^{24,25}, and Gregory G. Schwartz ²⁶; for the ODYSSEY OUTCOMES Investigators[†]

¹Department of Epidemiology and Biostatistics, State University of New York, Downstate School of Public Health, 450 Clarkson Avenue, MS 43, Brooklyn, NY 11203, USA;

²Division of Cardiovascular Disease, University of Alabama at Birmingham, 701 19th Street South - LHRB 310, Birmingham, AL 35294, USA; ³Department of Cardiology, South Australian Health and Medical Research Institute, Flinders University and Medical Centre, South Australia 5042, Australia; ⁴Sanofi R&D, Global Development, 371 Rue du Professeur Blayac, 34080 Montpellier, France; ⁵Department of Medicine, Brigham and Women's Hospital Heart and Vascular Center, Harvard Medical School, 75 Francis Street, Boston, MA 02115, USA; ⁶Estudios Clínicos Latinoamérica, Instituto Cardiovascular de Rosario, Paraguay 160, Rosario, Santa Fe, Rosario 2000, Argentina; ⁷Preventive Cardiology Unit, Department of Vascular Medicine, Division of Medicine, University Medical Centre Ljubljana, Zaloška cesta 7, SI-1525 Ljubljana, Slovenia; ⁸Faculty of Medicine, University of Ljubljana, Vrazov trg 2, SI-1000 Ljubljana, Slovenia; ⁹Canadian VIGOUR Centre, University of Alberta, 87 Ave NW, Edmonton, Alberta T6G 2E1, Canada; ¹⁰Division of Cardiology, St. Michael's Hospital, Room 6-034 Donnelly Wing, Toronto, Ontario M5B 1W8, Canada; ¹¹Department of Cardiology, Oslo Universitetssykehus HF Ullevål, and University of Oslo, Problemveien 7, 0315 Oslo, Norway; ¹²Stanford Center for Clinical Research, Department of Medicine, Stanford University, 291 Campus Drive, Li Ka Shing Building, Stanford, CA 94305-5101, USA; ¹³Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands; ¹⁴Netherlands Heart Institute, Utrecht, the Netherlands; ¹⁵Clinical Pharmacology—University of Kansas Medical Center, 3901 Rainbow Blvd, Kansas City, KS 66160, USA; ¹⁶Regeneron Pharmaceuticals, 777 Old Saw Mill River Rd., Tarrytown, NY 10591, USA; ¹⁷Imperial Centre for Cardiovascular Disease Prevention, Department of Primary Care and Public Health, Imperial College London, Reynolds Building, St Dunstons Road, London W6 8RP, UK; ¹⁸Department of Cardiovascular Medicine, University Hospitals Leuven, Herestraat 49, Leuven 3000, Belgium; ¹⁹Sulpizio Cardiovascular Center, Division of Cardiovascular Medicine, University of California San Diego, 9434 Medical Center Dr, La Jolla, CA 92037, USA; ²⁰Department of Medicine, University of Colorado Denver, 13001 E 17th Pl, Aurora, CO 80045, USA; ²¹Green Lane Cardiovascular Services, Auckland City Hospital, 5 Park Road, Grafton, Auckland 1142, New Zealand; ²²Department of Cardiology, Soroka University Medical Center, Faculty of Health Sciences, Ben Gurion University of the Negev, PO Box 151 Beer Sheva 8410, Israel; ²³Department of Medicine III, Goethe University, Theodor-Stern-Kai 7, Frankfurt am Main 60590, Germany; ²⁴Université de Paris, Assistance Publique-Hôpitaux de Paris, Hôpital Bichat, Université de Paris, FACT (French Alliance for Cardiovascular Trials), INSERM U1148, 46 Rue Henri Huchard, Paris 75018, France; ²⁵National Heart and Lung Institute, Imperial College, Royal Brompton Hospital, Sydney St, Chelsea, London SW3 6NP, UK; and ²⁶Division of Cardiology, University of Colorado School of Medicine, Box B130, Aurora, CO 80045, USA

Received 11 May 2020; revised 24 June 2020; editorial decision 21 July 2020; accepted 24 July 2020; online publish-ahead-of-print 14 October 2020

See page 4256 for the editorial comment on this article (doi: 10.1093/eurheartj/ehaa842)

Aims

Lipoprotein(a) concentration is associated with first cardiovascular events in clinical trials. It is unknown if this relationship holds for total (first and subsequent) events. In the ODYSSEY OUTCOMES trial in patients with recent

* Corresponding author. Tel: +1 646 696 8447, Fax: +1 718 270 2533, Email: michael.szarek@downstate.edu

[†] A complete list of the ODYSSEY OUTCOMES investigators is provided in the [Supplementary material online](#).

These results were reported in part at the American College of Cardiology—2019 Scientific Sessions.

© The Author(s) 2020. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

acute coronary syndrome (ACS), the proprotein convertase subtilisin/kexin type 9 inhibitor alirocumab reduced lipoprotein(a), low-density lipoprotein cholesterol (LDL-C), and cardiovascular events compared with placebo. This *post hoc* analysis determined whether baseline levels and alirocumab-induced changes in lipoprotein(a) and LDL-C [corrected for lipoprotein(a) cholesterol] independently predicted total cardiovascular events.


Methods and results

Cardiovascular events included cardiovascular death, non-fatal myocardial infarction, stroke, hospitalization for unstable angina or heart failure, ischaemia-driven coronary revascularization, peripheral artery disease events, and venous thromboembolism. Proportional hazards models estimated relationships between baseline lipoprotein(a) and total cardiovascular events in the placebo group, effects of alirocumab treatment on total cardiovascular events by baseline lipoprotein(a), and relationships between lipoprotein(a) reduction with alirocumab and subsequent risk of total cardiovascular events. Baseline lipoprotein(a) predicted total cardiovascular events with placebo, while higher baseline lipoprotein(a) levels were associated with greater reduction in total cardiovascular events with alirocumab (hazard ratio $P_{trend} = 0.045$). Alirocumab-induced reductions in lipoprotein(a) (median -5.0 [-13.6, 0] mg/dL) and corrected LDL-C (median -51.3 [-67.1, -34.0] mg/dL) independently predicted lower risk of total cardiovascular events. Each 5-mg/dL reduction in lipoprotein(a) predicted a 2.5% relative reduction in cardiovascular events.

Conclusion

Baseline lipoprotein(a) predicted the risk of total cardiovascular events and risk reduction by alirocumab. Lipoprotein(a) lowering contributed independently to cardiovascular event reduction, supporting the concept of lipoprotein(a) as a treatment target after ACS.

Graphical Abstract



Keywords

Lipoprotein • Alirocumab • LDL • Acute coronary syndrome

Introduction

Elevated baseline levels of lipoprotein(a), a genetically determined low-density lipoprotein particle, have been associated with increased risk of cardiovascular events in epidemiological studies¹ and Mendelian randomization analyses.^{2,3} In keeping with these observations, recent updates to the European Society of Cardiology (ESC) and European Atherosclerosis Society (EAS) lipid guidelines⁴ have

highlighted the role of lipoprotein(a) as a risk enhancer in primary prevention and recommend its measurement at least once in each adult's lifetime and in those at intermediate risk for cardiovascular disease events, with a family history of premature cardiovascular disease, or a family history of elevated lipoprotein(a)⁴ to optimize lipid modification therapies for the level of absolute risk.

Until recently, no intervention had demonstrated that modification in lipoprotein(a) levels would translate to a predicted risk reduction

Table 1 Categories of total events

Event	Alirocumab (n = 9462)	Placebo (n = 9462)	Total (n = 18 924)
Cardiovascular	2658	3111	5769
Coronary heart disease death	205	222	427
Non-fatal myocardial infarction ^a	877	1008	1885
Ischaemia-driven coronary revascularization	875	1003	1878
Non-fatal heart failure requiring hospitalization	287	277	564
Unstable angina requiring hospitalization	39	64	103
Death related to underlying coronary heart disease ^b	6	14	20
Death related to peripheral disease	10	11	21
Ischaemic stroke	130	185	315
Fatal	10	16	26
Non-fatal	120	169	289
Haemorrhagic stroke	23	25	48
Fatal	9	8	17
Non-fatal	14	17	31
Peripheral artery disease ^c	157	240	397
Limb revascularization ^c	109	165	274
Critical limb ischaemia ^c	48	75	123
Venous thromboembolism event ^c	49	62	111
Deep venous thrombosis ^c	29	33	62
Pulmonary embolism ^c	20	29	49
Non-cardiovascular death	94	121	215

Values are n.

^aTwenty-six events in the alicumab group and 43 events in the placebo group subsequently led to coronary heart disease death.

^bHeart failure, cardiogenic shock, cardiovascular procedure, cardiovascular haemorrhage.

^cInvestigator reported; not subject to adjudication by independent committee.

in subsequent cardiovascular events. However, two large placebo-controlled trials of inhibitors of proprotein convertase subtilisin/kexin type 9 (PCSK9) have provided such evidence. Reductions in lipoprotein(a) by alicumab were associated with subsequent reduced risk of first major cardiovascular events⁵ and, separately, first major peripheral artery disease (PAD) events or venous thromboembolism (VTE).⁶ Importantly, these relationships were independent of the simultaneous alicumab-induced reduction in low-density lipoprotein cholesterol (LDL-C) corrected for lipoprotein(a) cholesterol (LDL-C_{corr}). In addition, greater reduction in lipoprotein(a) concentration with the PCSK9 inhibitor evolucumab was associated with greater reduction in the risk of first major coronary events.⁷

From the perspective of patients and of healthcare delivery systems, the total burden of cardiovascular disease is more closely tied to total rather than first events. Accordingly, this *post hoc* analysis of the ODYSSEY OUTCOMES trial tested the hypothesis that baseline lipoprotein(a) predicts event burden and modification of this burden by alicumab, quantified by total cardiovascular events following an index acute coronary syndrome in patients receiving intensive statin therapy. In addition, to expand on previous observations from the study that reductions in lipoprotein(a) are associated with reduced risk of first cardiovascular event,^{5,6} we evaluated whether the decrease in lipoprotein(a) concentration under treatment with alicumab is associated with decreased risk of subsequent total cardiovascular events, independent of the concurrent reduction in LDL-C.

Methods

Patients and treatments

The design,⁸ primary results,⁹ and total events results¹⁰ of the ODYSSEY OUTCOMES trial have been published. The trial was performed at 1315 sites in 57 countries; the institutional review board at each site approved the protocol. All participants provided informed consent. Randomization in a 1:1 ratio to treatment with alicumab 75 mg or matching placebo, stratified by country, was performed in 18 924 patients meeting study entry criteria that included age ≥ 40 years, hospitalization with an acute coronary syndrome (myocardial infarction or unstable angina) 1–12 months before randomization, and LDL-C ≥ 70 mg/dL (1.81 mmol/L), non-high-density lipoprotein cholesterol ≥ 100 mg/dL (2.59 mmol/L), or apolipoprotein B ≥ 80 mg/dL assessed during stable treatment with atorvastatin 40–80 mg daily, rosuvastatin 20–40 mg daily, or the maximum-tolerated dose of either statin. All doses of study medication were given by subcutaneous injection every 2 weeks.

Outcomes

The primary efficacy outcome was time to first occurrence of death from coronary heart disease, non-fatal myocardial infarction, fatal or non-fatal ischaemic stroke, or unstable angina requiring hospitalization. The trial continued until 1613 patients experienced a primary outcome event or all surviving patients (other than those enrolled in China) were eligible to be followed for at least 2 years. Ultimately, the latter condition determined the end date of the trial. For the present analysis, we examined all cardiovascular outcomes collected in the trial including cardiovascular death and total (first and subsequent) non-fatal cardiovascular events.

The latter category included non-fatal primary outcome events, haemorrhagic stroke, hospitalization for heart failure, ischaemia-driven coronary revascularization, major PAD events (critical limb ischaemia, lower extremity revascularization procedures, and amputation for ischaemia), and VTE (deep vein thrombosis and pulmonary embolism). Non-cardiovascular deaths were also recorded during follow-up; subcategories of fatal and non-fatal events are listed in *Table 1*. All events included in the analyses except PAD events and VTE were adjudicated by an independent committee blinded to treatment assignment. PAD events and VTE were reported by investigators blinded to treatment assignment on a specific case-report form.

Measurement of lipoproteins

Details of lipoprotein measurements have been described.⁵ Lipoprotein(a) mass was measured at COVANCE Central Laboratories (Los Angeles, California, USA) on a Siemens BNII nephelometric analyser using an immunoturbidometric assay with rabbit polyclonal anti-lipoprotein(a) detection antibody and interassay coefficient of variation 3.1–4.8% depending on lipoprotein(a) concentration (Siemens, Healthcare Diagnostics, Malvern, PA, USA). To account for the fact that measured LDL-C includes cholesterol contained in lipoprotein(a) particles [$\sim 30\%$ of lipoprotein(a) mass is cholesterol], corrected LDL-C was calculated using the formula¹¹:

$$\text{LDL-C}_{\text{corr}} = \text{LDL-C} - 0.3 \times \text{lipoprotein(a) mass}$$

Additional analyses varied the percentage from 30% to determine if results were sensitive to the selection of this correction factor.

Statistical analysis

Lipoprotein(a) and LDL-C_{corr} distributions were assessed for the overall population and by the treatment group at baseline and at Month 4 (± 4 weeks) after randomization. If a patient had multiple values within the time window, the first value was included in the analyses. Missing values were imputed by prespecified methods.

We applied a marginal proportional hazards model that allows for the possibility of a given patient having multiple cardiovascular events while treating non-cardiovascular death as a competing event.¹² Relative effects on first and total cardiovascular events are summarized by hazard ratios (HRs), corresponding 95% confidence intervals (CIs), and *P*-values. Rates of events are expressed as the number of events per 100 patient-years of follow-up.

Relationships between baseline lipoprotein(a) and total cardiovascular events in the placebo group were determined by models using baseline lipoprotein(a) quartile as the predictor variable, adjusted for demographic and clinical variables (age, sex, race, geographic region, body mass index, smoking history, diabetes, time from index acute coronary syndrome to randomization), and baseline LDL-C_{corr}. A *P*-value was computed for linear trend in the estimated log HRs across baseline lipoprotein(a) quartiles. Cumulative incidence functions were used to estimate, by baseline lipoprotein(a) quartile, rates of total cardiovascular events through 4 years of follow-up in the placebo group in the presence of competing non-cardiovascular death.

Heterogeneity in the relative and absolute effects of alirocumab treatment on total cardiovascular events was assessed according to baseline lipoprotein(a) quartile. To assess heterogeneity in relative treatment effects, we constructed a Cox proportional hazards model with baseline lipoprotein(a) quartile, treatment, and their interaction as predictors and computed a *P*-value for linear trend across the estimated log HRs. To assess heterogeneity in absolute treatment effects, we constructed absolute risk reductions with alirocumab treatment, quantified as differences

in the event rates per 100 person-years of follow-up, along with associated 95% CIs.¹³

To determine the association between modification in lipoprotein(a) levels by alirocumab treatment and total events, the relationships between the change in lipoprotein(a) from baseline to Month 4 and the risk of cardiovascular events after Month 4 were described using models with patients in the alirocumab group. The following models were developed: a model without covariates (Model 1), a model adjusted for baseline lipoprotein(a) (Model 2); a model additionally adjusted for baseline LDL-C_{corr} and change from baseline to Month 4 in LDL-C_{corr} (Model 3); and a model adjusted for all variables in Model 3 as well as the demographic and clinical variables indicated above (Model 4). A comparison of Models 2 and 3 indicates whether the relationship between the change in lipoprotein(a) and total events is modified by adjustment for the simultaneous change in LDL-C_{corr}, while a comparison of Models 3 and 4 indicates if this relationship is confounded by other baseline characteristics of the patients. Sensitivity analyses excluded patients who had protocol-specified blinded substitution of placebo for alirocumab in response to consecutive LDL-C measurements < 15 mg/dL. Effects are summarized by the observed median reduction in lipoprotein(a) (all models) and LDL-C_{corr} (Models 3 and 4) at Month 4. The predicted risk reduction attributable to reduction in lipoprotein(a) as a percentage of the joint predicted risk reduction by decreases in lipoprotein(a) and LDL-C_{corr} was determined from the estimated log HRs from Model 3.

Continuous variables are expressed as median (Quartile 1, Quartile 3) whereas categorical variables are expressed as counts and percentages. All analyses were conducted according to intention-to-treat, including all patients and events from randomization to the common study end date (11 November 2017). Analyses were conducted by an independent academic statistical team at the State University of New York Downstate School of Public Health using SAS version 9.4.

Results

Categories of total vascular events

The median follow-up was 2.8 (2.3, 3.4) years. The types and counts of total cardiovascular events and non-cardiovascular deaths after randomization by treatment assignment are presented in *Table 1*. The absolute reduction in total events by alirocumab was primarily due to effects on non-fatal myocardial infarction, ischaemia-driven coronary revascularization, and major PAD events. Competing non-cardiovascular deaths were relatively infrequent, with fewer in the alirocumab group than in the placebo group.

Table 2 summarizes the distributions of cardiovascular events and non-cardiovascular deaths by event number. The total number of cardiovascular events, 5769, was 1.85 times the number of first events, 3114. As illustrated in *Supplementary material online, Figure S1*, there were 192 fewer first and 453 fewer total cardiovascular events with alirocumab than with placebo (2658 vs. 3111). Although the majority of patients did not experience a cardiovascular event during the study, a meaningful subset—48% of the patients with a non-fatal first cardiovascular event (1382 of 2860), corresponding to 7% of the total study population—experienced more than one cardiovascular event. In addition, cardiovascular death occurred in a relatively small proportion (2.7%) of the study population, and only 1.3% of patients had a cardiovascular death without a preceding non-fatal cardiovascular event, so that nearly all patients were at risk for multiple events.

Table 2 Distribution of first and subsequent cardiovascular events and non-cardiovascular death

Event number and type	Alirocumab (n = 9462)	Placebo (n = 9462)	Total (n = 18 924)
First			
Cardiovascular	1461	1653	3114
Fatal	125	129	254
Non-fatal	1336	1524	2860
Non-cardiovascular death	62	84	146
Second			
Cardiovascular	633	749	1382
Fatal	62	61	123
Non-fatal	571	688	1259
Non-cardiovascular death	16	17	33
Third			
Cardiovascular	253	321	574
Fatal	30	35	65
Non-fatal	223	286	509
Non-cardiovascular death	6	10	16
Fourth and subsequent			
Cardiovascular	311	388	699
Fatal	23	46	69
Non-fatal	288	342	630
Non-cardiovascular death	10	10	20
Total			
Cardiovascular	2658	3111	5769
Fatal	240	271	511
Non-fatal	2418	2840	5258
Non-cardiovascular death	94	121	215

Values are n.

Baseline lipoprotein(a) and total cardiovascular events in the placebo group

The cumulative incidence functions for total cardiovascular events over time by baseline lipoprotein(a) quartile are shown for the placebo group in Figure 1. The relationship between baseline lipoprotein(a) quartile and the number of events per 100 patient-years of follow-up in the placebo group is presented in Figure 2. There was a monotonic gradient of risk in relation to baseline lipoprotein(a) quartile, with the largest number of total cardiovascular events in the highest baseline lipoprotein(a) quartile (HR $P_{\text{trend}} < 0.0001$). The magnitude of the gradient was substantial: over 4 years, the estimated number of cardiovascular events per 100 patients was 53.7 among patients in the highest quartile (≥ 59.6 mg/dL) vs. 37.5 among patients in the lowest quartile (< 6.7 mg/dL), corresponding to an adjusted 60% higher risk ($P < 0.0001$).

Effect of alirocumab on total cardiovascular events by baseline lipoprotein(a) quartile

Relative and absolute treatment effects on total cardiovascular events stratified by baseline lipoprotein(a) quartile are shown in Figure 2. Overall, the alirocumab: placebo HR for first

cardiovascular event was 0.88 (95% CI, 0.82–0.94; $P = 0.0002$). For total cardiovascular events, the HR was 0.85 (95% CI, 0.78–0.93; $P = 0.0004$) with an absolute risk reduction of 1.7 events per 100 patient-years of follow-up. There was a significant linear trend in the HR for total cardiovascular events across baseline lipoprotein(a) quartiles ($P = 0.045$), decreasing from 0.95 in Quartile 1 to 0.75 in Quartile 4. Coupled with the relationship between baseline lipoprotein(a) and risk in the placebo group, the absolute risk reduction increased more than seven-fold, from 0.5 events per 100 patient-years of follow-up in Quartile 1 to 3.7 events per 100 patient-years in Quartile 4.

Effect of alirocumab-induced changes in lipoprotein(a) and LDL-C_{corr} on total cardiovascular events

This analysis included 9129 of the 9462 (96.5%) patients randomized to alirocumab with lipoprotein measurements at baseline and Month 4, who experienced a total of 2223 vascular events after their Month 4 assessment. Among these patients, the median absolute changes from baseline to Month 4 were -5.0 (-13.6, 0) mg/dL for lipoprotein(a) and -51.3 (-67.1, -34.0) mg/dL for LDL-C_{corr}. The median changes in lipoprotein(a) by baseline quartile were 0 (-1.4, 0), -5.1 (-7.9, -2.3), -9.9 (-16.3, -3.2), and -20.1 (-34.0, -8.0) mg/dL for Quartiles 1, 2, 3, and 4, respectively. Corresponding median changes

in LDL-C_{corr} by baseline lipoprotein(a) quartile were -53.3 (-68.8, -35.5), -52.7 (-69.2, -36.1), -51.2 (-66.5, -34.0), and -47.2 (-63.5, -31.4) mg/dL. The randomized treatment HR after the Month 4 assessments




Figure 1 Cumulative incidence functions for total cardiovascular events by quartile of baseline lipoprotein(a) (<6.7, 6.7 to <21.2, 21.2 to <59.6, and ≥59.6 mg/dL) in the placebo group. The estimated number of events per 100 patients at 4 years was 37.5, 43.5, 44.7, and 53.7 for Quartiles 1, 2, 3, and 4, respectively. The hazard ratios (95% confidence intervals) for Quartile 2, Quartile 3, and Quartile 4, relative to Quartile 1, are 1.05 (0.88–1.25), 1.17 (0.99–1.38), and 1.60 (1.34–1.89), respectively ($P_{\text{trend}} < 0.0001$). Hazard ratios and P -value reflect adjustment for age, sex, race, geographic region, body mass index, smoking status, diabetes status, time from index acute coronary syndrome to randomization, and baseline low-density lipoprotein cholesterol corrected for lipoprotein(a) cholesterol.

(HR, 0.85; 95% CI, 0.77–0.93) was nearly identical to the overall treatment HR (0.85; 95% CI, 0.78–0.93).

Table 3 summarizes the results of sequential marginal proportional hazards models relating the observed median change in lipoprotein(a) on alirocumab treatment to the risk of total vascular events. In an unadjusted model, a relationship between change in lipoprotein(a) and the risk of vascular events was not evident (Model 1). Larger changes in lipoprotein(a) under alirocumab treatment were associated with higher baseline lipoprotein(a) levels, and therefore higher risk. A significant relationship between reduction in lipoprotein(a) under alirocumab treatment and reduction in risk of total cardiovascular events was therefore revealed after adjustment for baseline lipoprotein(a) (Model 2). Additional adjustment for baseline and change in LDL-C_{corr} (Model 3) and further adjustment for demographic and clinical variables (Model 4) had a minor influence on this relationship. In all cases, reductions in both lipoprotein(a) and LDL-C_{corr} were associated with lower risk of cardiovascular events. In the model with lipoprotein(a) and LDL-C_{corr} (Model 3), the predicted relative risk reductions associated with median changes were 2.5% (HR, 0.975) for lipoprotein(a) and 15.8% (HR, 0.842) for LDL-C_{corr}; 13% of the predicted joint risk reduction was attributable to lipoprotein(a). Results were similar after excluding 715 patients with per-protocol blinded substitution of placebo for alirocumab (Supplementary material online, Table S1), and when the lipoprotein(a) correction factor for the calculation was specified to be 15% (Supplementary material online, Table S2) or 55% (Supplementary material online, Table S3).

Given that as baseline lipoprotein(a) increases, the alirocumab-induced change in lipoprotein(a) also increases while the alirocumab-induced change in LDL-C_{corr} remains relatively constant, the predicted risk reductions associated with reductions in lipoprotein(a) and LDL-C_{corr} were investigated by baseline lipoprotein(a) quartile; the results are summarized in the Take home figure with details in Table 4. Lipoprotein(a) did not contribute to the joint predicted risk





Figure 2 Relative and absolute treatment effects on total cardiovascular events, overall and by quartile of baseline lipoprotein(a). The rates of total vascular events per 100 patient-years of follow-up are shown for the alirocumab and placebo groups stratified by baseline quartile of lipoprotein(a) and for the overall population. The forest plots depict relative and absolute risk reduction with alirocumab compared with placebo. For relative risk reduction, there was a significant linear trend in the log hazard ratios across baseline quartiles, with point estimates progressively further below 1.00 for higher quartiles. CI, confidence interval; HR, hazard ratio.

Table 3 Relationship between reductions in lipoprotein(a) and low-density lipoprotein cholesterol from baseline to Month 4 and total cardiovascular events after Month 4 in the alirocumab group

Model	Model adjustments	Change parameter	HR (95% CI) for observed median reduction	P-value
1	None	Lipoprotein(a)	0.990 (0.964–1.017)	0.46
2	Baseline lipoprotein(a)	Lipoprotein(a)	0.972 (0.950–0.995)	0.016
3	Baseline lipoprotein(a), baseline LDL-C _{corr} , change from baseline to Month 4 in LDL-C _{corr}	Lipoprotein(a)	0.975 (0.953–0.997)	0.029
		LDL-C _{corr}	0.842 (0.755–0.940)	0.002
4	Baseline lipoprotein(a), baseline LDL-C _{corr} , change from baseline to Month 4 in LDL-C _{corr} , demographic and clinical characteristics	Lipoprotein(a)	0.978 (0.958–0.999)	0.036
		LDL-C _{corr}	0.820 (0.734–0.916)	0.001

Observed median reductions in lipoprotein(a) and LDL-C_{corr} were 5.0 and 51.3 mg/dL, respectively. The randomized treatment HR after Month 4 lipoprotein(a) and LDL-C measurements was 0.85 (95% CI 0.77–0.93).

CI, confidence interval; HR, hazard ratio; LDL-C_{corr}, low-density lipoprotein cholesterol corrected for lipoprotein(a) cholesterol.



reduction in Quartile 1, but contributed 39% of the joint reduction in Quartile 4. Consequently, the relative contribution of lipoprotein(a) change to the reduction in the risk of total cardiovascular events becomes greater as baseline lipoprotein(a) level increases.

Discussion

Among patients with recent acute coronary syndrome receiving intensive or maximally tolerated statin treatment, it was previously reported from the ODYSSEY OUTCOMES trial that baseline lipoprotein(a) level predicted risk of a first primary outcome event (death from coronary heart disease, non-fatal myocardial infarction,

ischaemic stroke, or hospitalization for unstable angina) among patients treated with placebo. In patients treated with alirocumab, the absolute reduction in lipoprotein(a) predicted the absolute reduction in risk of a first event, independent of the concurrent reduction in LDL-C_{corr}.⁵ Similar findings were reported for first major PAD or VTE events.⁶

The present report extends those findings and provides additional clinical context by including a substantially larger number of events (2223) than the previous first primary outcome (710) or first PAD or VTE event (116) analyses, and spanning a broader range of cardiovascular event categories, including ischaemia-driven coronary revascularization and hospitalization for heart failure. The disease burden of cardiovascular events was substantial; among patients who had a

Table 4 Relationship between reductions in lipoprotein(a) and low-density lipoprotein cholesterol corrected for lipoprotein(a) cholesterol from baseline to Month 4 and total cardiovascular events after Month 4 in the alirocumab group by quartile of baseline lipoprotein(a)^a

Baseline Lp(a) quartile	Change parameter	HR (95% CI) for observed median reduction within quartile	Joint predicted risk reduction attributable to lipoprotein(a) change (%) ^b
Quartile 1 (<6.7 mg/dL)	Lipoprotein(a)	1.000	0
	LDL-C _{corr}	0.836 (0.746–0.937)	
Quartile 2 (6.7 to <21.2 mg/dL)	Lipoprotein(a)	0.975 (0.952–0.997)	13
	LDL-C _{corr}	0.838 (0.749–0.938)	
Quartile 3 (21.2 to <59.6 mg/dL)	Lipoprotein(a)	0.951 (0.909–0.995)	23
	LDL-C _{corr}	0.842 (0.755–0.940)	
Quartile 4 (≥59.6 mg/dL)	Lipoprotein(a)	0.903 (0.824–0.990)	39
	LDL-C _{corr}	0.854 (0.772–0.944)	

CI, confidence interval; HR, hazard ratio; LDL-C_{corr}, low-density lipoprotein cholesterol corrected for lipoprotein(a) cholesterol.
^aObserved median reductions in lipoprotein(a) by baseline quartile were 0, 5.1, 9.9, and 20.1 mg/dL for Quartiles 1, 2, 3, and 4, respectively. Corresponding median reductions in LDL-C_{corr} by baseline lipoprotein(a) quartile were 53.3, 52.7, 51.2, and 47.2 mg/dL. Models reflect adjustments corresponding to Model 3 in Table 3.
^bLog(HR) for lipoprotein(a)/[log(HR) for lipoprotein(a) + log(HR) for LDL-C_{corr}].

non-fatal cardiovascular event, 48% experienced multiple events during the study. In the current analysis, baseline lipoprotein(a) predicted the burden of total cardiovascular events. The magnitude of lipoprotein(a) reduction by alirocumab, which was greater for higher levels of baseline lipoprotein(a), was associated with the benefit of treatment on those events, with greater relative and absolute risk reduction across successively higher quartiles of baseline lipoprotein(a). Furthermore, the association of alirocumab-induced reduction in lipoprotein(a) with reduced risk of total cardiovascular events was independent of the concurrent alirocumab-induced reduction in LDL-C_{corr}. Overall, each 5 mg/dL reduction in lipoprotein(a) with alirocumab predicted a 2.5% relative reduction in total cardiovascular events, which was similar in magnitude to that for first primary events.⁵ Although most of the reduction was attributable to LDL-C_{corr}, the contribution of lipoprotein(a) change to the reduction in risk of total cardiovascular events was substantial for higher levels of baseline lipoprotein(a).

In a *post hoc* regression-based analysis of the FOURIER (Further Cardiovascular Outcomes Research with PCSK9 Inhibition in Subjects with Elevated Risk) trial, which included patients in both the placebo and evolocumab groups, lipoprotein(a) concentration at week 12 of assigned treatment was associated with the subsequent risk of first major coronary event.⁷ Although these results are supportive of the present study, the analysis of FOURIER did not examine the relationship between change in lipoprotein(a) from baseline under evolocumab treatment and the risk of a subsequent event, or the relationship of achieved lipoprotein(a) levels to total event burden.

The findings from the present study are consistent with the 2019 ESC/EAS 2019 guidelines,⁴ which have highlighted the emerging importance of lipoprotein(a) in cardiovascular risk assessment and further extend the evidence base for lipoprotein(a) to a contemporary population with recent acute coronary syndrome. ODYSSEY OUTCOMES recruited patients on the basis of an LDL-C >1.8 mmol/L (70 mg/dL), and those in the highest quartile of lipoprotein(a) had as much as a 60% increased risk of total cardiovascular

events, corresponding to 16 extra events per 100 patients over 4 years, vs. those in the lowest quartile. Under the alirocumab treatment, this risk was reduced by 3.7 total events per 100 patient-years in the highest quartile, with 39% of the benefit attributable to lipoprotein(a) lowering. Likewise, if a 10% relative reduction in the risk of cardiovascular events is considered to be the minimum clinically meaningful effect, the present results indicate that with alirocumab given over the duration of the trial, patients with lipoprotein(a) ≥60 mg/dL would be expected to have their risk reduced by at least 10% consequent to reduction in lipoprotein(a), independent of any benefits from LDL-C reduction. Therefore, in patients with recent acute coronary syndrome, measurement of lipoprotein(a) levels might be a useful tool to identify patients at particularly high risk of recurrent events and who may derive substantial absolute and relative benefit from alirocumab treatment. It remains to be determined whether the findings from the present analysis are scalable to the larger reductions in lipoprotein(a) expected with antisense oligonucleotide and small-interfering RNA agents targeting the expression of apolipoprotein(a).¹⁴

A limitation of the analyses is that the majority of events were cardiac, restricting the ability to investigate if relationships with lipoprotein(a) potentially differ by vascular territory (e.g. cerebrovascular, peripheral). In addition, lipoprotein(a) was assessed by a mass concentration assay, and measurement by a molar concentration assay might have affected the observed relationships between alirocumab-induced changes in lipoprotein(a) with subsequent cardiovascular events. Furthermore, the applied LDL-C corrected formula used 30% lipoprotein(a) mass to estimate the cholesterol content of lipoprotein(a), and while results were consistent with higher or lower correction factors, the actual content may vary over a range around these nominal values.¹⁵ Finally, the current analysis does not define the mechanisms by which PCSK9 inhibition lowers lipoprotein(a) concentration. Kinetic studies indicate that in statin-treated patients with elevated lipoprotein(a) levels, alirocumab reduces those levels by increasing clearance.¹⁶

Conclusions

Recent results from ODYSSEY OUTCOMES have demonstrated that, among patients receiving intensive statin therapy following a recent acute coronary syndrome, patient-level reductions in lipoprotein(a) by alirocumab independently contributed to reduced risk of a first primary outcome event and first PAD or VTE event. The present analyses demonstrate that baseline lipoprotein(a) predicted total cardiovascular event risk in the placebo group and relative and absolute risk reduction in the alirocumab group. This risk modification was partially attributable to changes in lipoprotein(a), where reduction in lipoprotein(a) by alirocumab was associated with fewer total cardiovascular events independent of LDL-C_{corr} reduction. These findings support the concept of lipoprotein(a) as a treatment target among high-risk patients with a recent acute coronary syndrome.

Supplementary material

Supplementary material is available at *European Heart Journal* online.

Acknowledgements

The authors thank the patients, study coordinators, and investigators who participated in this trial. Sophie K. Rushton-Smith (MedLink Healthcare Communications, London) provided editorial assistance in the preparation of the manuscript (limited to editing for style and referencing, as well as figure and table editing) and was funded by Sanofi, Paris, France. Kausik K. Ray acknowledges support from the NIHR Imperial Biomedical Research Centre.

Data Sharing

Individual participant data are not available.

Funding

ODYSSEY OUTCOMES (ClinicalTrials.gov: NCT01663402) was funded by Sanofi and Regeneron Pharmaceuticals.

Conflict of interest: M.S. reports serving as a consultant or on advisory boards (or both) for CiVi, Resverlogix, Baxter, Esperion, Sanofi, and Regeneron Pharmaceuticals, Inc. V.A.B reports grant support from Sanofi, AstraZeneca, DalCor, Esperion, Bayer, The Medicines Company, and Amgen, all paid directly to her institution, and personal fees from Sanofi. P.A. reports research grants from Sanofi, CSL, AstraZeneca, Da Cor, and advisory boards and speaker fees from Sanofi, Amgen, CSL, AstraZeneca, Bayer, Novartis, and Boehringer Ingelheim. D.L.B reports advisory board fees from Cardax, Cereno Scientific, Elsevier Practice Update Cardiology, Medscape Cardiology, PhaseBio, and Regado Biosciences; Board of Directors membership for Boston VA Research Institute, Society of Cardiovascular Patient Care, and TobeSoft; position of Chair for the American Heart Association Quality Oversight Committee; membership of Data Monitoring Committees for the Baim Institute for Clinical Research (formerly Harvard Clinical Research Institute, for the PORTICO trial, funded by St. Jude Medical, now Abbott), Cleveland Clinic (including for the ExCEED trial, funded by Edwards), Duke Clinical Research Institute, Mayo Clinic, Mount Sinai School of Medicine (for the ENVISAGE trial, funded by Daiichi-Sankyo), Population Health Research Institute; honoraria from the American College of Cardiology (Senior Associate Editor, Clinical Trials and News, ACC.org; Vice-Chair, ACC

Accreditation Committee), Baim Institute for Clinical Research (formerly Harvard Clinical Research Institute; RE-DUAL PCI clinical trial steering committee funded by Boehringer Ingelheim; AEGIS-II executive committee funded by CSL Behring), Belvoir Publications (Editor in Chief, Harvard Heart Letter), Duke Clinical Research Institute (clinical trial steering committees, including for the PRONOUNCE trial, funded by Ferring Pharmaceuticals), HMP Global (Editor in Chief, Journal of Invasive Cardiology), Journal of the American College of Cardiology (Guest Editor; Associate Editor), Medtelligence/ReachMD (CME steering committees), Population Health Research Institute (for the COMPASS operations committee, publications committee, steering committee, and USA national co-leader, funded by Bayer), Slack Publications (Chief Medical Editor, Cardiology Today's Intervention), Society of Cardiovascular Patient Care (Secretary/Treasurer), and WebMD (CME steering committees); other relationships with *Clinical Cardiology* (Deputy Editor), NCDR-ACTION Registry Steering Committee (Chair), and VA CART Research and Publications Committee (Chair); research funding from Abbott, Afimmune, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Chiesi, CSL Behring, Eisai, Ethicon, Ferring Pharmaceuticals, Forest Laboratories, Fractyl, Idorsia, Ironwood, Ischaemix, Lilly, Medtronic, PhaseBio, Pfizer, PLx Pharma, Regeneron, Roche, Sanofi-Aventis, Synaptic, and The Medicines Company; royalties from Elsevier (Editor, Cardiovascular Intervention: A Companion to Braunwald's Heart Disease); position as a site co-investigator for Biotronik, Boston Scientific, CSI, St. Jude Medical (now Abbott), and Svelte; position as a trustee for the American College of Cardiology; and unfunded research for FlowCo, Merck, Novo Nordisk, and Takeda. R.D. reports research grants from Sanofi, DalCor Pharmaceuticals, Population Health Research Institute, Duke Clinical Research Institute, the TIMI group, Amgen, Cirius, Montreal Health Innovations Coordinating Center, and Lepetit and personal fees, as a member of the Executive Steering Committee, from Amgen and Cirius. Z.F. reports serving as a National Coordinator for the ODYSSEY OUTCOMES trial (Duke Research Institute/Sanofi-Regeneron) and has received research support and speaker's/consulting fees from Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Krka, Novo Nordisk, Pfizer, Sanofi, and Servier. S.G.G. reports research grants from Daiichi-Sankyo, Luitpold Pharmaceuticals, Merck, Novartis, Servier, Regeneron Pharmaceuticals, Inc., Sanofi, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, CSL Behring, Eli Lilly, Pfizer, and Tenax Therapeutics; honoraria from Bristol-Myers Squibb, Eli Lilly, Esperion, Fenix Group International, Ferring Pharmaceuticals, Merck, Novartis, Pfizer, Servier, Regeneron Pharmaceuticals, Inc., Sanofi, Amgen, AstraZeneca, Bayer, and Boehringer Ingelheim; and serving as a consultant or on advisory boards (or both) for AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly, HLS Therapeutics, Pfizer, Servier, Tenax Therapeutics, Sanofi, Amgen, and Bayer. S.H. reports personal fees from Sanofi and Amgen. R.A.H. reports research grants (all DSMB related) from AstraZeneca, Janssen, and Bristol-Myers Squibb, serving on advisory boards for Gilead (uncompensated) and WebMD, and serving on the boards of directors (unpaid) for the American Heart Association and Stanford HealthCare. J.W.J. reports research grants from the Netherlands Heart Foundation, the Interuniversity Cardiology Institute of the Netherlands, and the European Commission Seventh Framework Programme and research support from Amgen, Astellas, AstraZeneca, Daiichi-Sankyo, Lilly, Merck-Schering-Plough, Pfizer, Roche, and Sanofi. P.M.M. reports personal fees from Regeneron Pharmaceuticals, Sanofi, Academic CME, Amarin, Ambry Genetics, National Lipid Association, Esperion, Lilly, and BioPharma; research grants from Akcea, Familial Hypercholesterolemia Foundation, GB Life Sciences, Ionis, Kowa, Novartis, RegenX, Amgen, Kaneka, and RegenXBio; and grants and personal fees from Stage II Innovations/Renew. R.P. is an employee of Regeneron Pharmaceuticals, Inc. K.K.R.

reports personal fees from Abbvie, AstraZeneca, Medco, Resverlogix, Akcea, Boehringer Ingelheim, Novo Nordisk, Takeda, Kowa, Algorithm, Cipla, Cerenis, Dr Reddys, Lilly, and Zuellig Pharma; Novartis, and Silence Therapeutics; and grants and personal fees from Amgen, Sanofi, Regeneron Pharmaceuticals, MSD, Daiichi-Sankyo, and Pfizer. P.S. reports personal fees from Sanofi, Abbott, Boehringer Ingelheim, BMS, Pfizer, and Amgen; personal fees and non-financial support from AstraZeneca and Merck Sharp Dohme; and grants and personal fees from Daiichi-Sankyo and Bayer. S.T. is an employee of Ionis Pharmaceuticals and reports personal fees (as a co-founder) from Oxitope, Inc., and Kleanthi Diagnostics, LLC and also has the following patents: methods of identifying a subject having or at risk of having or developing coronary artery disease; methods for assessing atherogenesis by determining oxidized phospholipid to apolipoprotein B ratios; methods for assessing atherogenesis by determining oxidized phospholipid to apolipoprotein B ratios; and oxidative biomarkers in predicting the risk of stroke, transient ischaemic attack, and peripheral arterial disease (royalties for all three are paid to Kleanthi Diagnostics). R.V. reports research grants and speaker fees from Sanofi and Regeneron Pharmaceuticals. H.D.W. reports receiving grant support paid to the institution and fees for serving on a steering committee for the ODYSSEY OUTCOMES trial (Evaluation of Cardiovascular Outcomes After an Acute Coronary Syndrome During Treatment With Alirocumab) from Sanofi-Aventis and Regeneron Pharmaceuticals, for the ACCELERATE study (A Study of Evacetrapib in High-Risk Vascular Disease) from Eli Lilly, for the STRENGTH trial (Outcomes Study to Assess Statin Residual Risk Reduction With EpaNova in High CV Risk Patients With Hypertriglyceridaemia) from Omthera Pharmaceuticals, for the SPIRE trial (The Evaluation of Bococizumab [PF-04950615; RN 316] in Reducing the Occurrence of Major Cardiovascular Events in High Risk Subjects) from Pfizer USA, for the HEART-FID study (Randomized Placebo-Controlled Trial of FCM as Treatment for Heart Failure With Iron Deficiency) from American Regent, for the CAMELLIA-TIMI study [A Study to Evaluate the Effect of Long-term Treatment With BELVIQ (Lorcaserin HC) on the Incidence of Major Adverse Cardiovascular Events and Conversion to Type 2 Diabetes Mellitus in Obese and Overweight Subjects With Cardiovascular Disease or Multiple Cardiovascular Risk Factors] from Eisai, Inc., for the dal-GenE study (Effect of Dalcetrapib vs. Placebo on CV Risk in a Genetically Defined Population With a Recent ACS) from DalCor Pharma UK, Inc., for the AEGIS-II study from CSL Behring, for the SCORED trial (Effect of Sotagliflozin on Cardiovascular and Renal Events in Patients With Type 2 Diabetes and Moderate Renal Impairment Who Are at Cardiovascular Risk) and the SOLOIST-WHF trial (Effect of Sotagliflozin on Cardiovascular Events in Patients With Type2 Diabetes Post Worsening Heart Failure) from Sanofi-Aventis Australia Pty Ltd, and for the CLEAR Outcomes Study [Evaluation of Major Cardiovascular Events in Patients with, or at High Risk for, Cardiovascular Disease Who Are Statin Intolerant Treated with Bempedoic Acid (ETC-1002) or Placebo] from Esperion Therapeutics, Inc., was on the Advisory Boards for Acetelion, Sirtex, and Genentech, Inc. (an affiliate of F. Hoffmann-La Roche Ltd, 'Roche'; Lytics Post-PCI Advisory Board at European Society of Cardiology), and received lecture fees from AstraZeneca. D.Z. reports personal fees from Sanofi, Bayer, Pfizer, Novonordisk, and Astra Zeneka. A.M.Z. reports receiving fees for serving on a steering committee for the ODYSSEY OUTCOMES trial from Sanofi and advisory board and speaker fees from Sanofi, Amgen, Boehringer Ingelheim, Bayer, Novartis, Pfizer, AstraZeneca, and Vifor. P.G.S. reports grants, personal fees, and non-financial support from Sanofi; grants and personal fees from Amarin, Servier and Bayer; and personal fees from Amgen, AstraZeneca, BMS, Boehringer Ingelheim, Idorsia, Pfizer, and Novartis and also has a patent

use of alirocumab to reduce risk after ACS (royalties to Sanofi) pending. G.G.S. reports research grants to the University of Colorado from Resverlogix, Sanofi, The Medicines Company, and Roche and is a coinventor of pending US patent 62/806 313 (Methods for Reducing Cardiovascular Risk) assigned in full to the University of Colorado.

References

- Nordestgaard BG, Langsted A. Lipoprotein (a) as a cause of cardiovascular disease: insights from epidemiology, genetics, and biology. *J Lipid Res* 2016;**57**: 1953–1975.
- Burgess S, Ference BA, Staley JR, Freitag DF, Mason AM, Nielsen SF, Willeit P, Young R, Surendran P, Karthikeyan S, Bolton TR, Peters JE, Kamstrup PR, Tybjaerg-Hansen A, Benn M, Langsted A, Schnohr P, Vedel-Krogh S, Kobylecki CJ, Ford I, Packard C, Trompet S, Jukema JW, Sattar N, Di Angelantonio E, Saleheen D, Howson JMM, Nordestgaard BG, Butterworth AS, Danesh J; for the European Prospective Investigation Into Cancer and Nutrition–Cardiovascular Disease (EPIC-CVD) Consortium. Association of LPA variants with risk of coronary disease and the implications for lipoprotein(a)-lowering therapies: a Mendelian randomization analysis. *JAMA Cardiol* 2018;**3**:619–627.
- Lamina C, Kronenberg F, Lp GC; for the Lp(a)-GWAS-Consortium. Estimation of the required lipoprotein(a)-lowering therapeutic effect size for reduction in coronary heart disease outcomes: a Mendelian randomization analysis. *JAMA Cardiol* 2019;**4**:575–579.
- Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L, Chapman MJ, De Backer GG, Delgado V, Ference BA, Graham IM, Halliday A, Landmesser U, Mihaylova B, Pedersen TR, Riccardi G, Richter DJ, Sabatine MS, Taskinen M-R, Tokgozoglul L, Wiklund O, Mueller C, Drexler H, Aboyans V, Corsini A, Doehner W, Farnier M, Gigante B, Kayikcioglu M, Krstacic G, Lambrinou E, Lewis BS, Masip J, Moulin P, Petersen S, Petronio AS, Piepoli MF, Pinto X, Raber L, Ray KK, Reiner Z, Riesen WF, Roffi M, Schmid J-P, Shlyakhto E, Simpson IA, Stroes E, Sudano I, Tselepis AD, Viigimaa M, Vindis C, Vonbank A, Vrablik M, Vrsalovic M, Zamorano JL, Collet J-P, Koskinas KC, Casula M, Badimon L, John Chapman M, De Backer GG, Delgado V, Ference BA, Graham IM, Halliday A, Landmesser U, Mihaylova B, Pedersen TR, Riccardi G, Richter DJ, Sabatine MS, Taskinen M-R, Tokgozoglul L, Wiklund O, Windecker S, Aboyans V, Baigent C, Collet J-P, Dean V, Delgado V, Fitzsimons D, Gale CP, Grobbee D, Halvorsen S, Hindricks G, Iung B, Juni P, Katus HA, Landmesser U, Leclercq C, Lettino M, Lewis BS, Merkely B, Mueller C, Petersen S, Petronio AS, Richter DJ, Roffi M, Shlyakhto E, Simpson IA, Sousa-Uva M, Touyz RM, Nibouche D, Zelveian PH, Siostrzonek P, Najafav R, van de Borne P, Pojskic B, Postadzhiyan A, Kypris L, Špinar J, Larsen ML, Eldin HS, Viigimaa M, Strandberg TE, Ferrières J, Agladze R, Laufs U, Rallidis L, Bajnok L, Gudjonsson T, Maher V, Henkin Y, Gulizia MM, Mussagaliyeva A, Bajraktari G, Kerimkulova A, Latkovskis G, Hamoui O, Slapikas R, Visser L, Dingli P, Ivanov V, Boskovic A, Nazzi M, Visseren F, Mitevska I, Retterstøl K, Jankowski P, Fontes-Carvalho R, Gaita D, Ezhov M, Foscoli M, Giga V, Pella D, Fras Z, de Isla LP, Hagström E, Lehmann R, Abid L, Ozdogan O, Mitchenko O, Patel RS; Group ESCSD. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020; **41**:111–188.
- Bittner VA, Szarek M, Aylward PE, Bhatt DL, Diaz R, Edelberg JM, Fras Z, Goodman SG, Halvorsen S, Hanotin C, Harrington RA, Jukema JW, Loizeau V, Moriarty PM, Moryusef A, Pordy R, Roe MT, Sinnaeve P, Tsimikas S, Vogel R, White HD, Zahger D, Zeiher AM, Steg PG, Schwartz GG; ODYSSEY OUTCOMES Investigators. Effect of alirocumab on lipoprotein(a) and cardiovascular risk after acute coronary syndrome. *J Am Coll Cardiol* 2020; **75**:133–144.
- Schwartz GG, Steg PG, Szarek M, Bittner VA, Diaz R, Goodman SG, Kim YU, Jukema JW, Pordy R, Roe MT, White HD, Bhatt DL; for the ODYSSEY OUTCOMES Committees and Investigators. Peripheral artery disease and venous thromboembolic events after acute coronary syndrome: role of lipoprotein(a) and modification by alirocumab: prespecified analysis of a randomized clinical trial. *Circulation* 2020;**141**:1608–1617.
- O'Donoghue ML, Fazio S, Giugliano RP, Stroes ESG, Kanevsky E, Gouni-Berthold I, Im KAH, Lira Pineda A, Wasserman SM, Česka R, Ezhov MV, Jukema JW, Jensen HK, Tokgözoğlu SL, Mach F, Huber K, Sever PS, Keech AC, Pedersen TR, Sabatine MS. Lipoprotein(a), PCSK9 inhibition, and cardiovascular risk. *Circulation* 2019; **139**:1483–1492.
- Schwartz GG, Bessac L, Berdan LG, Bhatt DL, Bittner V, Diaz R, Goodman SG, Hanotin C, Harrington RA, Jukema JW, Mahaffey KW, Moryusef A, Pordy R, Roe MT, Rorick T, Sasiela WJ, Shirodaria C, Szarek M, Tamby JF, Tricoci P, White H, Zeiher A, Steg PG. Effect of alirocumab, a monoclonal antibody to PCSK9, on long-term cardiovascular outcomes following acute coronary syndromes: rationale and design of the ODYSSEY outcomes trial. *Am Heart J* 2014;**168**:682–689.

9. Schwartz GG, Steg PG, Szarek M, Bhatt DL, Bittner VA, Diaz R, Edelberg JM, Goodman SG, Hanotin C, Harrington RA, Jukema JW, Lecorps G, Mahaffey KW, Moryusef A, Pordy R, Quintero K, Roe MT, Sasiela WJ, Tamby JF, Tricoci P, White HD, Zeiher AM; for the ODYSSEY OUTCOMES Committees and Investigators. Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med* 2018;**379**:2097–2107.
10. Szarek M, White HD, Schwartz GG, Alings M, Bhatt DL, Bittner VA, Chiang CE, Diaz R, Edelberg JM, Goodman SG, Hanotin C, Harrington RA, Jukema JW, Kimura T, Kiss RG, Lecorps G, Mahaffey KW, Moryusef A, Pordy R, Roe MT, Tricoci P, Xavier D, Zeiher AM, Steg PG; ODYSSEY OUTCOMES Committees and Investigators. Alirocumab reduces total nonfatal cardiovascular and fatal events: the ODYSSEY OUTCOMES trial. *J Am Coll Cardiol* 2019;**73**:387–396.
11. Yeang C, Witztum JL, Tsimikas S. 'LDL-C' = LDL-C + Lp(a)-C: implications of achieved ultra-low LDL-C levels in the proprotein convertase subtilisin/kexin type 9 era of potent LDL-C lowering. *Curr Opin Lipidol* 2015;**26**:169–178.
12. Andersen PK, Angst J, Ravn H. Modeling marginal features in studies of recurrent events in the presence of a terminal event. *Lifetime Data Anal* 2019;**25**: 681–695.
13. Stukel TA, Glynn RJ, Fisher ES, Sharp SM, Lu-Yao G, Wennberg JE. Standardized rates of recurrent outcomes. *Stat Med* 1994;**13**:1781–1791.
14. Landmesser U, Poller W, Tsimikas S, Most P, Paneni F, Lüscher TF. From traditional pharmacological towards nucleic acid-based therapies for cardiovascular diseases. *Eur Heart J* 2020;doi:10.1093/eurheartj/ehaa229.
15. Kinpara K, Okada H, Yoneyama A, Okubo M, Murase T. Lipoprotein(a)-cholesterol: a significant component of serum cholesterol. *Clin Chim Acta* 2011;**412**: 1783–1787.
16. Watts GF, Chan DC, Pang J, Ma L, Ying Q, Aggarwal S, Marcovina SM, Barrett PHR. PCSK9 Inhibition with alirocumab increases the catabolism of lipoprotein(a) particles in statin-treated patients with elevated lipoprotein(a). *Metabolism* 2020;**107**:154221.

Corrigendum

doi:10.1093/eurheartj/ehz826

Online publish-ahead-of-print 14 November 2019

Corrigendum to: 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk [*Eur Heart J* (2019); doi:10.1093/eurheartj/ehz455].

The following post-publication corrections have been made to these Guidelines:

In row 4 of Table 3, 'should' has been corrected to 'may';

in 4.2.1, paragraph 3, 'should' has been corrected to 'may' to read 'Overall, CAC score assessment with CT may be considered in individuals. . .'. This has also been corrected in the second row of 'Recommendations for cardiovascular imaging for risk assessment of atherosclerotic cardiovascular disease,' and the Class has been corrected to 'Ib';

in the second paragraph of 7.5.2, '5–10 mg of monacolin K' has been corrected to '2.5–10 mg';

and in the Key messages section, number 4, 'ApoB may be a better measure of an individual's exposure to atherosclerotic lipoproteins' has been corrected to 'ApoB may be a better measure of an individual's exposure to pro atherogenic lipoproteins'.

© The European Society of Cardiology and the European Atherosclerosis Association 2019. All rights reserved. For permissions please email: journals.permissions@oup.com.