



Universiteit
Leiden
The Netherlands

Immune-based therapies for hematological malignancies: an update by the EHA SWG on immunotherapy of hematological malignancies

Einsele, H.; Briones, J.; Ciceri, F.; Garcia-Cadenas, I.; Falkenburg, F.; Bolanos, N.; ... ; Loosdrecht, A.A. van de

Citation

Einsele, H., Briones, J., Ciceri, F., Garcia-Cadenas, I., Falkenburg, F., Bolanos, N., ... Loosdrecht, A. A. van de. (2020). Immune-based therapies for hematological malignancies: an update by the EHA SWG on immunotherapy of hematological malignancies. *Hemasphere*, 4(4). doi:10.1097/HS9.0000000000000423

Version: Publisher's Version
License: [Creative Commons CC BY-NC 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3279622>

Note: To cite this publication please use the final published version (if applicable).

Immune-based Therapies for Hematological Malignancies: An Update by the EHA SWG on Immunotherapy of Hematological Malignancies

Hermann Einsele¹, Javier Briones², Fabio Ciceri³, Irene García-Cadenas², Fred Falkenburg⁴, Natacha Bolaños⁵, H.M. Mirjam Heemskerk⁴, Roch Houot^{6,7}, Michael Hudecek¹, Franco Locatelli⁸, Kate Morgan⁹, C. Emma Morris¹⁰, Michael O'Dwyer¹¹, Jordi Sierra Gil^{2,12}, Marcel van den Brink¹³, Arjan A. van de Loosdrecht¹⁴

Correspondence: Hermann Einsele (e-mail: Einsele_h@ukw.de).

Immunotherapy is revolutionizing the treatment of certain hematological malignancies and providing cure to patients even with some advanced diseases (eg, B-ALL, lymphoma). However, immunotherapy, especially cellular immunotherapy, is an emerging field and on-going research will have to address several issues to improve safety and efficacy of this novel treatment modality. There is also the need to extend the application to additional hematological malignancies. The EHA scientific working group for immunotherapy has defined major research questions for the future development of immunotherapy for patients with hematological malignancies.

The incidence of hematological malignancies continues to rise, while the underlying biological mechanisms of tumorigenesis remain unknown. The treatment landscape for hematological malignancies is diverse and immunotherapies are clearly entering the arena. Immune-based therapies for hematological malignancies aim at generating new agents, such as monoclonal antibodies,

immunotoxins, engineered monoclonal antibodies called bispecific T-cell engagers, cell therapies involving cells of the innate and adoptive immune system, adoptive cell transfer therapy with T cells engineered to express chimeric antigen receptors or T-cell receptors, immune cell redirection strategies, gene modified immune cells, vaccines and checkpoint inhibitors (i.e. PD-1, CTLA-4, and IDO), which are less toxic and potentially more effective when compared to conventional chemo- and radiotherapy or even allogeneic HSCT. These various forms of immune-based therapies have shown significant promise, leading to the change in improving patient outcomes.

Monoclonal antibodies

Monoclonal antibodies are effective in a number of hematological malignancies.¹⁻³ Most of the current identified targets for monoclonal antibodies are also expressed on non-malignant cells.⁴⁻⁶ However, in contrast with either gene-modified T cells (eg, CAR T-cells) or bispecific antibodies, the on-target toxicity of monoclonal antibodies on non-malignant cells is mostly tolerable.^{7,8} New strategies to improve therapy with monoclonal antibodies include genetically engineering the structure and function of these antibodies, an approach shown to significantly improve their effectiveness.¹⁻³

Vaccination strategies

Vaccines activating the autologous immune system for prevention and treatment of infections and other diseases might also have a major impact on human healthcare. Compared to other immunotherapies, such as checkpoint blockade or adoptive T cell therapy, most cancer vaccines to date have failed to demonstrate notable clinical efficacy.^{9,10} One of the key obstacles to the development of an effective cancer vaccine is the difficulty in antigen selection and the requirement to overcome tolerance to self. In the past, most of the cancer vaccines were targeting tumor-associated antigens (TAAs) which are overexpressed in many cancers and were seen as universal targets for the treatment of patients with hematological malignancies. Unfortunately, TAAs are also expressed on normal tissues and, thus, central and peripheral tolerance can interfere with the efficacy of vaccination or alternatively it can induce auto-immunity/auto-reactivity against normal tissues. In contrast to these non-mutated self-

The authors have indicated they have no potential conflicts of interest to disclose.

¹Universitätsklinikum Würzburg, Medizinische Klinik und Poliklinik II, Würzburg, Germany

²Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

³Vita-Salute San Raffaele University and Division of Transplantation, Immunology and Transplantation Diseases, IRCCS San Raffaele Scientific Institute, Milan, Italy

⁴Department of Hematology, Leiden University Medical Center, Leiden, the Netherlands

⁵Lymphoma Coalition Europe, Madrid, Spain

⁶CHU de Rennes, Université de Rennes, INSERM U1236, Rennes, France

⁷Dana-Farber Cancer Institute, Boston, Massachusetts, USA

⁸IRCCS, Bambino Gesù Children's Hospital, Sapienza University of Rome, Rome, Italy

⁹Myeloma UK, Edinburgh, UK

¹⁰UCL Institute of Immunity and Transplantation Royal Free Hospital, London, UK

¹¹Department of Haematology, University Hospital Galway and National University of Ireland, Galway

¹²Sant Pau Biomedical Research Institute and Jose Carreras Leukemia Research Institute, Barcelona, Spain

¹³Memorial Sloan Kettering Cancer Center, New York, New York, USA

¹⁴Amsterdam UMC, VU Medical Center, Amsterdam, the Netherlands.

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the European Hematology Association. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and build upon the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

HemaSphere (2020) 4:4(e423). <http://dx.doi.org/10.1097/HS9.0000000000000423>.

Received: 14 May 2020 / Accepted: 25 May 2020

antigens, neoantigens which are derived from random somatic mutations in tumor cells and are not present in normal cells, can be recognized as non-self by the host immune system and thus be used as attractive targets for immunotherapies with potentially increased specificity, efficacy and safety.^{11,12} Preclinical and clinical studies have demonstrated neoantigen-specific T cells represent the most potent tumor-reactive immune cell subpopulation. However, the inefficient presentation of neoantigens and the low clonal frequency, unfortunately, lead to a mostly very small population of neoantigen-specific cytotoxic T cells in cancer patients. Thus, vaccination with neoantigens may be required to elicit and expand the T-cell population specifically targeting neoantigens. The most frequently used adjuvants for personalized immunotherapy based on vaccines are autologous dendritic cells.¹³ In some cancer patients, dendritic cell/peptide vaccines have demonstrated clinical benefit in phase III trials.

A clear shortcoming of cancer vaccines is the difficult standardization of the antigenic material and the reduced efficacy of adjuvants. Therefore, most investigators favor mixing relevant immunogenic peptides to be used directly as peptide vaccines or loaded onto dendritic cells as professional antigen-presenting cells.¹³ The development of more effective adjuvants will probably also improve the efficacy of cancer vaccines, especially based on neoantigens. Alternatively, off-the-shelf dendritic cell-based vaccine strategies are developed to overcome some of these shortcomings by a fully standardized production of tumor vaccines.¹⁴

The post-transplant setting, especially the period following allogeneic stem cell transplantation is a potentially promising platform for vaccination due to cytoreduction and relative depletion of inhibitory accessory cells fostering greater immune responsiveness. Another source of non-self antigens especially in the setting of allogeneic stem cell transplantation are the minor histocompatibility antigens (MiHA), which are derived from single nucleotide differences between patient and donor.^{15,16} MiHAs are recognized as non-self by the donor immune system and thus can be used as attractive targets for immunotherapy after allogeneic stem cell transplantation in form of vaccination.^{17–20} In addition, currently clinical studies are ongoing with the HA-1 TCR in high risk hematological malignancies (Leiden, Seattle).

Donor lymphocyte infusions (DLI) and antigen-specific DLI

Donor lymphocyte infusions have been used for more than 40 years now to treat overt relapse, mixed chimerism or residual tumor cells following allogeneic stem cell transplantation.^{21,22} In addition, the adoptive transfer of antigen-specific cytotoxic T lymphocytes and CD4⁺ T-helper cells has been widely used to treat viral infections (especially CMV, EBV, adenovirus infections and more recently BK and JC viruses).^{23–26} Currently early phase trials are in progress to study the role of donor lymphocyte infusions in the management of refractory, invasive fungal infections. One of the greatest challenges of the next few years will be that of transitioning the initial promising results, mainly obtained in proof-of-concept studies, to a wider application of these therapies using standardized methodological approaches.

Checkpoint inhibitors

Checkpoint inhibitor blockade releases the brakes on tumor-specific T cells, allowing them to persist and expand to attack malignant cells. Cancers can grow, at least in part, as a

consequence of cancer-induced immunosuppression. In many individuals, immunosuppression is mediated by CTLA4 and PD-1, 2 immunomodulatory receptors expressed on T cells. Monoclonal antibody-based therapies targeting CTLA4 or PD-1 have shown significant clinical effects in patients with hematological malignancies, especially patients with Hodgkin's lymphoma, patients with primary mediastinal B-cell lymphoma,^{27–30} and are currently being explored in several other hematological malignancies and after allogeneic transplantation despite the expected risk of increasing the rate and severity of graft versus host disease reactions.^{31–33} Synergistic efficacy has been shown for CTLA4 and PD-1 blocking antibodies especially in the setting of solid tumors (eg, metastatic melanoma). Furthermore, antibodies against additional checkpoint molecules like TIM3, TIGIT, LAG-3, etc. have been developed and are being tested in early clinical trials in patients with hematological malignancies.^{34,35}

Innate immune cells

Next to dendritic cells, other innate immune cells, especially natural killer (NK) cells play a key role in anti-tumor immunity.³⁶ NK cell function is finely tuned by both inhibitory and activating signals. NK cells do not require tumor-antigen recognition or clonal expansion before killing cancer cells. The most important inhibitory signals for NK cells are mediated via their killer immune-globulin-like receptors (KIRs), which recognize major histocompatibility complex/HLA class I antigens on autologous cells. Activating receptors bind to ligands expressed on tumor cells and induce activation of NK cells.

Recently $\gamma\delta$ T cells, a subset of T cells expressing $\gamma\delta$ TCRs rather than the conventional $\alpha\beta$ TCR, which can exert a direct anti-tumor effect have been used for immunotherapy of hematological malignancies.^{37,38} Donor-derived $\gamma\delta$ T cells^{38,39} selected after depletion of $\alpha\beta$ TCR T cells and infused following lymphodepletion were found to induce partial or even complete responses in patients with acute myeloid leukemia and multiple myeloma. In addition, infusions of NKT cells have been used in the treatment of hematological malignancies.^{40,41}

Bispecific antibodies: T Cell redirecting antibodies

Bispecific antibodies that recruit and redirect T cells to attack tumor cells have tremendous potential for the treatment of hematological malignancies. These antibody constructs promote tumor cell killing by crosslinking a CD3 component of the T cell receptor complex with the tumor-associated antigen on the surface of the tumor cell. Importantly, this mode of action does not rely on a cognate interaction between the T cell receptor and a peptide/HLA complex. Thus, this strategy is not dependent on HLA restriction or on HLA expression (this representing a significant advantage since in tumors HLA class I and II molecules may be downregulated). Therefore, bispecific antibodies may find a key role in hematological malignancies with low neoantigen content and a low inflammation and these novel immunotherapeutics may productively be combined with checkpoint blockade.

Extensive optimization and process development have progressed a large number of bispecific/trispecific antibodies into clinical trials for a wide range of indications, with promising signs of therapeutic activity. As an example, Blinatumomab has already been approved for treatment of refractory, relapsed, BCP-ALL and also for patients with molecularly resistant disease

following intensive chemotherapy.^{42–44} Very promising efficacy data for bispecific antibodies were demonstrated⁴⁵ in the treatment of AML and, especially, advanced multiple myeloma.

Gene-modified T cells

T cells recognize antigens through a unique antigen-specific T-cell receptor (TCR) promoting the elimination of a given target and amplifying the attack through the recruitment of other components of the immune response. T cells can target peptides derived from both intracellular and extracellular proteins, including those encoded by genes mutated in cancer cells. T cells can actively distribute within tissues and in the tumor environment, and have the potential for *in vivo* expansion and self-maintenance, as they can establish a memory compartment.

The last decade has witnessed technological advances, which have allowed genetic modifications of T cells providing personalized cellular therapies that target specific tumor-associated antigens. Gene transfer into human T cells can be accomplished by several means. Long-term culture of the genetically modified T cells is often required to reach meaningful clinical doses, with the functional impact of prolonged *ex vivo* expansion potentially adversely affecting subsequent long-term persistence *in vivo*. Gene delivery (or transfer) is mainly achieved through the use of viral (retroviral or lentiviral) vectors. These vectors can be manufactured to clinical grade on a large-scale producing stable integration into the genome of the T cell and its progeny. Adverse consequences due to insertional mutagenesis in T cells have not yet been reported and are unexpected given the mature differentiation status of the T cell at the time of exogenous gene integration.

However, lentiviral vectors are particularly attractive when less differentiated T cell subsets are targeted for modification, as they have the unique ability to infect T cells even upon minimal activation, a property lacking in retroviral vectors. Novel non-viral systems (eg, Sleeping Beauty and PiggyBac) allow larger fragments of DNA to be inserted than viral vectors permit.⁴⁶ These novel strategies of non-viral gene transduction are likely to reduce the cost of genetic modification of immune cells in the future by avoiding the need for large scale manufacture of clinical grade viral vectors. When combined with CRISPR/Cas (or other) gene-editing technologies, site directed insertion of the gene encoding the TCR or the chimeric antigen receptor (CAR) is achievable.^{47–50} A major advantage of gene-modified T cells is that they are a living drug; thus, which can expand and proliferate in the patient. Persistence over time has been demonstrated in responding patients.

CAR T-cells

T cells redirected to specific surface antigens on malignant cells by engineered CARs are emerging as powerful therapies for hematological malignancies. In contrast to bispecific antibodies, which link the activated T cells to the tumor cells by a small molecule with binding domains for both CD3 and a surface antigen expressed on the tumor cell, here T cells are engineered to express a new antigen recognition receptor, which targets the antigen on the tumor cell surface.

An increasing number of clinical trials using CAR T-cells for the treatment of hematological malignancies have been reported, especially targeting B-cell lymphoma, BCP-ALL and multiple myeloma. Two cell products have been approved in 2017/2018 for clinical use: Kymriah (Tisagenlecleucel) and Yescarta

(axicabtagene ciloleucel), which received FDA and European marketing authorization for the treatment of relapsed or refractory CD19+ BCP-ALL and for the treatment of diffuse large B-cell lymphoma (DLBCL). These approvals were based on impressive responses observed in patients with BCP-ALL including high risk patients.⁵¹ The overall response rate and the role of complete remission in B cell lymphoma is lower than in ALL, but long-term remissions without further therapy have been reported for up to 40% of patients, enough to encourage the approval and speed up the ongoing research.^{52,53} Promising initial results have been also reported in the treatment of patients with multiple myeloma, where CAR T-cells target the B-cell membrane antigen (BCMA).⁵⁴

TCR gene-modified T cells

T cell receptor (TCR) modified T cells are a novel alternative of adoptive cell therapy designed to treat hematological malignancies and non-hematological solid tumors. Typically the genes encoding the alpha and beta chains of the TCR are cloned into retroviral or lentiviral vectors for gene transfer into autologous T cells. Novel non-viral transduction technologies, which are increasingly developed and established, are also being optimized for clinical grade TCR transfer. TCR modified T cells can mediate anti-tumor efficacy and have been used to target several antigens like NYESO-1, MAGE-A3 and PRAME, MAGE-A10 and WT1. Clinical trials have been performed to treat patients with multiple myeloma and AML.⁵⁵

European research contributions

Europe has played a leading role in the development of bispecific antibodies. Indeed, bispecific antibodies were approved for clinical use by both the FDA and EMA after having been developed in Europe from the first pilot clinical trials until the larger phase II studies (also performed in Europe). In 2018 the FDA granted accelerated approval for the treatment of adult and pediatric patients with B-cell precursor ALL in first or second complete remission with minimal residual disease greater than or equal to 0.1%. The whole development program of Blinatumomab from the pilot trial until the final approval by the FDA and the EMA was performed under the leadership of European scientists. Currently most of the new indications for treatment of Bite/bispecific antibodies are clinically tested in Europe. New formats of bispecific antibodies are being developed by European biotech companies. Promising data on the use of bispecific antibodies also in multiple myeloma and partially in AML have been generated in European centers.⁴⁵

Technologies of gene modification of T cells including CAR and TCR gene-transfer have been intensively developed in Europe. Patients in the UK and other EU countries have been some of the first in the world to receive these therapies outside the US. European scientists and clinicians are leading the development of gene-modified neo-antigen specific T cells, which are predicted to have a major impact in the management of solid hematological and non-hematological tumors. Other major contributions have been the development and validation of novel strategies for immune cell selection.

Despite these major contributions, Europe is currently lagging behind in the initiation of CAR T cell therapy trials, which are mainly conducted in the US and China. Currently, European patients are queuing up to receive treatment with CAR T-cells for hematological malignancies in US and Chinese centers due to the

lack of open clinical trials in Europe. Of more than 400 clinical trials listed as investigating CAR T-cell therapy in the treatment of hematological or solid cancer around the world about 87% are performed in the USA and China and only a minority in Europe. The much larger pharma market, the more rapid approval for novel agents and the better support for early clinical trials explains this development.

Conclusions

Immunotherapy is revolutionizing the treatment of certain hematological malignancies and provides cure to patients even with advanced disease. Research ongoing in Europe is focusing on improving the safety and efficacy of this novel treatment modality. The goal of the future development in immunotherapy is to make it even more successful and safer as well as to extend its use to additional hematological malignancies.

Disclosures

Hermann Einsele: Consultation for BMS, Janssen, Novartis, Amgen, Takeda; Grants from BMS, Amgen, Janssen; Javier Briones: Honoraria from Roche, Takeda, Celgene, Novartis, Gilead; Consultation or advisory role for Takeda, Janssen, Celgene, Gilead/Kyte; Travel/accommodations/expenses from Roche, Takeda, Celgene, Janssen, Gilead; Fabio Ciceri: Employment/consultation for MolMed; Irene García Cadenas: Nothing to disclose; Fred Falkenburg: Financial affiliation with the Dutch Cancer Society; Natacha Bolaños: Lymphoma Coalition (LC) received grants (funds directly to LC) from Takeda, Janssen, Novartis, Amgen, Kyowa Kirin, Roche, Celgene, BMS; H.M. Mirjam Heemskerk: Grants from KWF, EU: Enacting, LUMC, Health Holland; Roch Houot: Royalties of payments for lectures/manuscript preparations from Bristol-Myers Squibb, MSD, Gilead, Kite, Roche, Novartis, Janssen, Celgene; Michael Hudecek: Royalties on patents; Franco Locatelli: Employment/consultation for Novartis, Amgen, Bellicum Pharmaceuticals, Miltenyi, Jazz, Gilead, Medac, Takeda, Janssen; Kate Morgan: Myeloma Patients Europe (MPE) receives unrestricted, core funding from Janssen, Celgene, Novartis, Takeda, Amgen, Oncopeptides, Mundipharma, Pfizer, Roche, AstraZeneca, GSK, Sanofi; MPE receives funding from the European Commission through Horizon2020 and the Innovative Medicines Initiative; C. Emma Morris: Employment/consultation for Quell Therapeutics Limited; Grants/pending grants for NIHR, MRC, Wellcome Trust; Royalties for lectures/manuscript preparations for Gilead, Shire; Scientific founder (shares) of Quell Therapeutics Ltd; Michael O'Dwyer: Consultation with Janssen, Celgene, AbbVie, Adaptive Biotechnology; Equity in Onk Therapeutics, Carrick Therapeutics; Research support from BMS, Janssen, Onk Therapeutics; Jordi Sierra Gil: Employment/consultation for Pfizer, Amgen, Daiichi Sankyo, Astellas, Novartis, CSL Behring; Grants/pending grants from Novartis; Royalties for lectures/manuscript preparations from Astellas; Marcel van den Brink: Employment/consultation for: DKMS Chairman of the Board, WindMIL Therapeutics, Memorial Sloan Kettering Cancer Center and Sloan Kettering Institute; Grants/pending grants from R01-CA228358, R01-CA228308, R01 HL147584, MSKCC Cancer Center, NHLBI, NIA, NIAID, Tri-Institutional Stem Cell Initiative, The Lymphoma Foundation, The Susan and Peter Solomon Divisional Genomics Program, Parker Institute for Cancer Immunotherapy at Memorial Sloan Kettering Cancer Center, Seres Therapeutics (M.V.D.B); Research support from Seres Therapeutics; Consulted/received honorarium from or participated in advisory boards for Seres Therapeutics, Flagship Ventures, Novartis, Evelo, Jazz Pharmaceuticals, Therakos, Amgen, Magenta Therapeutics, WindMIL Therapeutics, Merck & Co, Inc., Acute Leukemia Forum (ALF), DKMS Medical Council (Board); Has IP Licensing

with Seres Therapeutics, and Juno Therapeutics; Arjan A van de Loosdrecht: Nothing to disclose.

References

- Weiner GJ. Building better monoclonal antibody-based therapeutics. *Nat Rev Cancer*. 2015;15:361–370.
- Hutchings CJ, Koglin M, Olson WC, et al. Opportunities for therapeutic antibodies directed at G-protein-coupled receptors. *Nat Rev Drug Discov*. 2017;16:787–810.
- Kennedy PJ, Oliveira C, Granja PL, et al. Monoclonal antibodies: technologies for early discovery and engineering. *Crit Rev Biotechnol*. 2018;38:394–408.
- van de Donk NW, Janmaat ML, Mutis T, et al. Monoclonal antibodies targeting CD38 in hematological malignancies and beyond. *Immunol Rev*. 2016;270:95–112.
- Salles G, Barrett M, Foà R, et al. Rituximab in B-Cell hematologic malignancies: a review of 20 years of clinical experience. *Adv Ther*. 2017;34:2232–2273.
- Tazi I, Nafil H, Mahmal L. Monoclonal antibodies in hematological malignancies: past, present and future. *J Cancer Res Ther*. 2011;7:399–407.
- Hansel TT, Kropshofer H, Singer T, et al. The safety and side effects of monoclonal antibodies. *Nat Rev Drug Discov*. 2010;9:325–338.
- Weiner LM, Surana R, Wang S. Monoclonal antibodies: versatile platforms for cancer immunotherapy. *Nat Rev Immunol*. 2010;10:317–327.
- Melero I, Gaudernack G, Gerritsen W, et al. Therapeutic vaccines for cancer: an overview of clinical trials. *Nat Rev Clin Oncol*. 2014;11:509–524.
- Finn OJ. The dawn of vaccines for cancer prevention. *Nat Rev Immunol*. 2018;18:183–194.
- Hu Z, Ott PA, Wu CJ. Towards personalized, tumour-specific, therapeutic vaccines for cancer. *Nat Rev Immunol*. 2018;18:168–182.
- Mullard A. Economists propose a US\$30 billion boost to biomedical R&D. *Nat Rev Drug Discov*. 2012;11:735–737.
- Palucka K, Banchereau J. Cancer immunotherapy via dendritic cells. *Nat Rev Cancer*. 2012;12:265–277.
- van de Loosdrecht AA, van Wetering S, Santegoets SJAM, et al. A novel allogeneic off-the-shelf dendritic cell vaccine for post-remission treatment of elderly patients with acute myeloid leukemia. *Cancer Immunol Immunother*. 2018;67:1505–1518.
- Griffioen M, van Bergen CA, Falkenburg JH. Autosomal minor histocompatibility antigens: how genetic variants create diversity in immune targets. *Front Immunol*. 2016;7:100.
- Goulmy E, Schipper R, Pool J, et al. Mismatches of minor histocompatibility antigens between HLA-identical donors and recipients and the development of graft-versus-host disease after bone marrow transplantation. *N Engl J Med*. 1996;334:281–285.
- Griffioen M, Honders MW, van der Meijden ED, et al. Identification of 4 novel HLA-B*40:01 restricted minor histocompatibility antigens and their potential as targets for graft-versus-leukemia reactivity. *Haematologica*. 2012;97:1196–1204.
- van Bergen CA, van Luxemburg-Heijs SA, de Wreede LC, et al. Selective graft-versus-leukemia depends on magnitude and diversity of the alloreactive T cell response. *J Clin Invest*. 2017;127:517–529.
- Bleakley M, Riddell SR. Exploiting T cells specific for human minor histocompatibility antigens for therapy of leukemia. *Immunol Cell Biol*. 2011;89:396–407.
- Warren EH, Greenberg PD, Riddell SR. Cytotoxic T-lymphocyte-defined human minor histocompatibility antigens with a restricted tissue distribution. *Blood*. 1998;91:2197–2207.
- Kolb HJ, Mittermüller J, Clemm C, et al. Donor leukocyte transfusions for treatment of recurrent chronic myelogenous leukemia in marrow transplant patients. *Blood*. 1990;76:2462–2465.
- Falkenburg F, Ruggiero E, Bonini C, et al. Prevention and treatment of relapse after stem cell transplantation by cellular therapies. *Bone Marrow Transplant*. 2019;54:26–34.
- Riddell SR, Watanabe KS, Goodrich JM, et al. Restoration of viral immunity in immunodeficient humans by the adoptive transfer of T cell clones. *Science*. 1992;257:238–241.
- Einsele H, Roosnek E, Rufer N, et al. Infusion of cytomegalovirus (CMV)-specific T cells for the treatment of CMV infection not responding to antiviral chemotherapy. *Blood*. 2002;99:3916–3922.
- Feuchtinger T, Opherck K, Bethge WA, et al. Adoptive transfer of pp65-specific T cells for the treatment of chemorefractory cytomegalovirus

- disease or reactivation after haploidentical and matched unrelated stem cell transplantation. *Blood*. 2010;116:4360–4367.
26. Neuenhahn M, Albrecht J, Odendahl M, et al. Transfer of minimally manipulated CMV-specific T cells from stem cell or third-party donors to treat CMV infection after allo-HSCT. *Leukemia*. 2017;31:2161–2171.
 27. von Keudell G, Younes A. Novel therapeutic agents for relapsed classical Hodgkin lymphoma. *Br J Haematol*. 2019;184:105–112.
 28. Goodman A, Patel SP, Kurzrock R. PD-1-PD-L1 immune-checkpoint blockade in B-cell lymphomas. *Nat Rev Clin Oncol*. 2017;14:203–220.
 29. Bair RM, Strelec LE, Feldman TA, et al. Outcomes and toxicities of programmed death-1 (PD-1) inhibitors in hodgkin lymphoma patients in the United States: a real-world, multicenter retrospective analysis. *Oncologist*. 2019;24:955–962.
 30. Moy RH, Younes A. Immune checkpoint inhibition in Hodgkin lymphoma. *HemaSphere*. 2018;2:e20p1-p8.
 31. Meryman RW, Kim HT, Zinzani PL, et al. Safety and efficacy of allogeneic hematopoietic stem cell transplant after PD-1 blockade in relapsed/refractory lymphoma. *Blood*. 2017;129:1380–1388.
 32. Haverkos BM, Abbott D, Hamadani M, et al. PD-1 blockade for relapsed lymphoma post-allogeneic hematopoietic cell transplant: high response rate but frequent GVHD. *Blood*. 2017;130:221–228.
 33. Soiffer RJ. Checkpoint inhibition to prevent or treat relapse in allogeneic hematopoietic cell transplantation. *Bone Marrow Transplant*. 2019;54 (Suppl 2):798–802.
 34. Hattori N, Kawaguchi Y, Sasaki Y, et al. Monitoring TIGIT/DNAM-1 and PVRL2 immune checkpoint expression levels in allogeneic stem cell transplantation for acute myeloid leukemia. *Biol Blood Marrow Transplant*. 2019;25:861–867.
 35. Ok CY, Young KH. Checkpoint inhibitors in hematological malignancies. *J Hematol Oncol*. 2017;10:103, p1-p16.
 36. Dolstra H, Roeven MWH, Spanholtz J, et al. Successful Transfer of Umbilical Cord Blood CD34+ Hematopoietic Stem and Progenitor-derived NK Cells in Older Acute Myeloid Leukemia Patients. *Clin Cancer Res*. 2017;23:4107–4118.
 37. de Witte MA, Kuball J, Miller JS. NK Cells and $\gamma\delta$ T cells for relapse protection after allogeneic hematopoietic cell transplantation (HCT). *Curr Stem Cell Rep*. 2017;3:301–311.
 38. de Witte MA, Sarhan D, Davis Z, et al. Early reconstitution of NK and $\gamma\delta$ T cells and its implication for the design of post-transplant immunotherapy. *Biol Blood Marrow Transplant*. 2018;24:1152–1162.
 39. Kunzmann V, Smetak M, Kimmel B, et al. Tumor-promoting versus tumor-antagonizing roles of $\gamma\delta$ T cells in cancer immunotherapy: results from a prospective phase I/II trial. *J Immunother*. 2012;35:205–213.
 40. Zhu Y, Smith DJ, Zhou Y, et al. Development of hematopoietic stem cell-engineered invariant natural killer T cell therapy for cancer. *Cell Stem Cell*. 2019;25:542–557.
 41. Kobayashi T, Doff BL, Rearden RC, et al. NKT cell-targeted vaccination plus anti-4-1BB antibody generates persistent CD8 T cell immunity against B cell lymphoma. *Oncimmunology*. 2015;4:e990793, p1-p11.
 42. Topp MS, Gökbuğet N, Zugmaier G, et al. Phase II trial of the anti-CD19 bispecific T cell-engager linatumomab shows hematologic and molecular remissions in patients with relapsed or refractory B-precursor acute lymphoblastic leukemia. *J Clin Oncol*. 2014;32: 4134–4140.
 43. Topp MS, Gökbuğet N, Stein AS, et al. Safety and activity of blinatumomab for adult patients with relapsed or refractory B-precursor acute lymphoblastic leukaemia: a multicentre, single-arm, phase 2 study. *Lancet Oncol*. 2015;16:57–66.
 44. Kantarjian H, Stein A, Gökbuğet N, et al. Blinatumomab versus chemotherapy for advanced acute lymphoblastic leukemia. *N Engl J Med*. 2017;376:836–847.
 45. Topp MS, Duell J, Zugmaier G, et al. Anti-B-Cell maturation antigen BiTE molecule AMG 420 induces responses in multiple myeloma. *J Clin Oncol*. 2020;38:775–783.
 46. Querques I, Mades A, Zuliani C, et al. A highly soluble Sleeping Beauty transposase improves control of gene insertion. *Nat Biotechnol*. 2019;37:1502–1512.
 47. Schober K, Müller TR, Gökmen F, et al. Orthotopic replacement of T-cell receptor α - and β -chains with preservation of near-physiological T-cell function. *Nat Biomed Eng*. 2019;3:974–984.
 48. Eyquem J, Mansilla-Soto J, Giavridis T, et al. Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances tumour rejection. *Nature*. 2017;543:113–117.
 49. Liu X, Zhang Y, Cheng C, et al. CRISPR-Cas9-mediated multiplex gene editing in CAR-T cells. *Cell Res*. 2017;27:154–157.
 50. Jensen TI, Axelgaard E, Bak RO. Therapeutic gene editing in haematological disorders with CRISPR/Cas9. *Br J Haematol*. 2019; 185:821–835.
 51. Authorisation of tisagenlecleucel in the EU. European Medicines Agency (EMA). <https://www.ema.europa.eu/en/medicines/human/EPAR/kymriah>. Accessed June 22, 2020.
 52. Authorisation of axicabtagene ciloleucel in the EU. European Medicines Agency (EMA). <https://www.ema.europa.eu/en/medicines/human/EPAR/ycarta>. Accessed June 22, 2020.
 53. Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in adult relapsed or refractory diffuse large B-Cell lymphoma. *N Engl J Med*. 2019;380:45–56.
 54. Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-cell therapy bb2121 in relapsed or refractory multiple myeloma. *N Engl J Med*. 2019;380:1726–1737.
 55. Mastaglio S, Genovese P, Magnani Z, et al. NY-ESO-1 TCR single edited stem and central memory T cells to treat multiple myeloma without graft-versus-host disease. *Blood*. 2017;130:606–618.