

# 2-AG-mediated control of GABAergic signaling is impaired in a model of epilepsy

Colangeli, R.; Morena, M.; Werner, A.; Thompson, R.J.; Stelt, M. van der; Pittman, Q.J.; ...; Teskey, G.C.

# Citation

Colangeli, R., Morena, M., Werner, A., Thompson, R. J., Stelt, M. van der, Pittman, Q. J., ... Teskey, G. C. (2022). 2-AG-mediated control of GABAergic signaling is impaired in a model of epilepsy. *Journal Of Neuroscience*. doi:10.1523/JNEUROSCI.0541-22.2022

Version:Accepted ManuscriptLicense:Leiden University Non-exclusive licenseDownloaded from:https://hdl.handle.net/1887/3505028

Note: To cite this publication please use the final published version (if applicable).



Research Articles: Systems/Circuits

# 2-AG-mediated Control of GABAergic Signaling is Impaired in a Model of Epilepsy

https://doi.org/10.1523/JNEUROSCI.0541-22.2022

Cite as: J. Neurosci 2022; 10.1523/JNEUROSCI.0541-22.2022

Received: 17 March 2022 Revised: 21 October 2022 Accepted: 25 November 2022

This Early Release article has been peer-reviewed and accepted, but has not been through the composition and copyediting processes. The final version may differ slightly in style or formatting and will contain links to any extended data.

Alerts: Sign up at www.jneurosci.org/alerts to receive customized email alerts when the fully formatted version of this article is published.

Copyright © 2022 the authors

# 2-AG-mediated Control of GABAergic Signaling is Impaired in a Model of Epilepsy

Roberto Colangeli<sup>1,2\*</sup>, Maria Morena<sup>1,3,4</sup>, Allison Werner<sup>1</sup>, Roger J. Thompson<sup>1</sup>, Mario van der Stelt<sup>5</sup>, Quentin J. Pittman<sup>6</sup>, Matthew N. Hill<sup>1</sup>, G. Campbell Teskey<sup>1,6</sup>.

<sup>1</sup>Department of Cell Biology and Anatomy, and Hotchkiss Brain Institute, University of Calgary, Calgary, AB, T2N4N1, Canada.

<sup>2</sup>Department Experimental and Clinical Medicine, Università Politecnica delle Marche, 60126 Ancona, Italy.

<sup>3</sup>Department of Physiology and Pharmacology, Sapienza University of Rome, 00185 Rome, Italy.

<sup>4</sup>Neuropsychopharmacology Unit, IRCSS Fondazione Santa Lucia, 00143 Rome, Italy.

<sup>5</sup>Department of Molecular Physiology, Leiden Institute of Chemistry, Leiden University, 2333 CC Leiden, The Netherlands.

<sup>6</sup>Department of Physiology and Pharmacology, and Hotchkiss Brain Institute, University of Calgary, Calgary, AB, T2N4N1, Canada.

\*Corresponding Author

Roberto Colangeli, PhD

Department Experimental and Clinical Medicine,

Università Politecnica delle Marche, Ancona, Italy.

Phone # +390712206050

E-mail: r.colangeli@univpm.it

Number of pages: 26

Number of figures: 8

Abstract: 237 words

Introduction: 465 words

Discussion: 1945 words

# **Conflict of interest**

The authors declare no competing financial interests.

# Acknowledgements

This work was supported by Canadian Institutes for Health Research (CIHR) MOP-130495 to G.C.T. and M.N.H. and CIHR PJT-159454 to Q.J.P. R.C. received salary support from Eyes High postdoctoral fellowship. M.M. received salary support from Alberta Innovates and CIHR postdoctoral fellowships.

# **Running title**

Enduring 2-AG signaling maladaptation in epilepsy

# Significant statement

The plastic reorganization of endocannabinoid (eCB) signaling after seizures and during epileptogenesis may contribute to the negative neurobiological consequences associated with seizure activity. Therefore, a deeper understanding of the molecular basis underlying the pathologic long-term eCB signaling remodelling following seizure activity will be crucial to develop novel therapies for epilepsy which not only target seizure activity, but, most importantly, the epileptogenesis and the comorbid conditions associated with epilepsy.

Repeated seizures result in a persistent maladaptation of endocannabinoid signaling, mediated in part by anandamide signaling deficiency in the basolateral amygdala (BLA) that manifests as aberrant synaptic function and altered emotional behaviour. Here, we determined the effect of repeated seizures (kindling) on 2-AG signaling on GABA transmission by directly measuring tonic and phasic eCB-mediated retrograde signaling in an in vitro BLA slice preparation from male rats. We report that both activity-dependent and muscarinic receptor (mAChR)-mediated depression of GABA synaptic transmission was reduced following repeated seizure activity. These effects were recapitulated in sham rats by preincubating slices with the 2-AG synthesising enzyme inhibitor DO34. Conversely, preincubating slices with the 2-AG degrading enzyme inhibitor KML29 rescued activitydependent 2-AG signaling, but not mAChR-mediated synaptic depression, over GABA transmission in kindled rats. These effects were not attributable to a change in CB1 receptor sensitivity or altered 2-AG tonic signaling since the application of the highly selective CB1 receptor agonist CP55,940 provoked a similar reduction in GABA synaptic activity in both sham and kindled rats, while no effect of either DO34 or of the CB1 inverse agonist AM251 was observed on frequency and amplitude of spontaneous inhibitory postsynaptic currents (sIPSCs) in either sham or kindled rats. Collectively, these data provide evidence that repeated amygdala seizures persistently alter phasic 2-AG mediated-retrograde signaling at BLA GABAergic synapses, probably by impairing stimulus-dependent 2-AG synthesis/release, which contributes to the enduring aberrant synaptic plasticity associated with seizure activity.

# Introduction

Seizure activity results in aberrant neuronal plasticity (Colangeli et al., 2017; Farrell et al., 2020) which manifests as learning and memory impairments (Farrell et al., 2017). However, the pathophysiological underpinnings of the aberrant rewiring of brain circuits following seizures remain elusive. A large body of evidence supports the view that the endocannabinoid (eCB) system plays an important role in the pathophysiology of epilepsy (Soltesz et al., 2015; Sugaya and Kano, 2018; Colangeli et al., 2021; Sugaya and Kano, 2021). As a general rule, activation of the cannabinoid type 1 (CB1) receptor typically protects against seizure activity by producing a net reduction in overall neuronal excitability (Marsicano et al., 2003; Sugaya et al., 2016; Colangeli et al., 2017). Conversely, both acute (Farrell et al., 2021) and persistent (Chen et al., 2003; Chen et al., 2007; Colangeli et al., 2020) maladaptation of eCB signaling following seizure activity may account for some of the negative neurobiological consequences of seizures.

eCBs are lipid molecules that modulate synaptic function to regulate homeostatic excitatory/inhibitory balance and thereby finely tune neuronal excitability (Kano et al., 2009; Castillo et al., 2012). eCB signaling subserves both tonic and phasic control of synaptic activity in several brain regions (Kano et al., 2009), including the basolateral amygdala (BLA) (Azad et al., 2004; Yoshida et al., 2011; Colangeli et al., 2020; Yasmin et al., 2020). The BLA is a common epileptogenic temporal lobe structure, principally involved in seizure generation and expression (Engel, 2001) and, for this reason, is often targeted in kindling studies (Teskey, 2020).

In the BLA, phasic eCB signaling encompasses transient and long-lasting changes in presynaptic transmission (Marsicano et al., 2002; Yoshida et al., 2011). eCB-mediated transient changes in neurotransmitter release occur following postsynaptic depolarization with a subsequent rise in Ca<sup>2+</sup>, which triggers eCB production (Kano et al., 2009). eCB mobilization can also occur upon activation of  $G_q$ -coupled GPCRs to induce inhibitory long-term depression (Wettschureck et al., 2006; Heifets and Castillo, 2009). In addition to the phasic regulation of synaptic activity in the BLA, tonic eCB signaling also contributes to the homeostatic control of neurotransmitter release (Hill and Tasker, 2012). We have previously reported that repeated seizures in the amygdala cause a persistent maladaptive downregulation of anandamide (AEA) signaling in the BLA, which drives aberrant excitatory synaptic function and altered emotional behaviour (Colangeli et al., 2020). Here, we

determined the effect of repeated amygdala seizures on 2-AG signaling on GABA transmission in the BLA. We provide evidence that repeated seizure activity in the amygdala is associated with enduring alterations of both transient and long-term GABAergic synaptic plasticity in the BLA; these synaptic alterations were mimicked by the disruption of 2-AG signaling in sham rats by inhibiting 2-AG synthesis. Inhibition of the 2-AG degrading enzyme MAGL completely rescued transient but not long-term GABAergic plasticity following kindling.

### Methods

### Animals

108 male Hooded Long-Evans rats (300-350g) at the start of electrophysiological experiments (Charles River, Canada) were housed in a temperature-controlled  $(23 \pm 1^{\circ}C)$  vivarium room and maintained under specified pathogen-free conditions in a 12 h/12 h light/dark cycle (7:00 A.M. to 7:00 P.M. lights on). Food and water were available ad libitum. All experimental procedures complied with protocols approved by the University of Calgary Animal Care Committee and guidelines from the Canadian Council on Animal Care. All efforts were made to minimize animal suffering and to reduce the number of animals used.

# Surgery

Rats were anaesthetized with a 5% isoflurane in  $O_2$ , maintained between 1% and 2% and then positioned in a stereotaxic frame (David Kopf Instruments). One Teflon-coated stainless steel bipolar electrode (A-M Systems, Sequim, WA) was chronically implanted into the right BLA (coordinates: anteroposterior, - 2.8 mm, and mediolateral, - 4.8 mm from Bregma; dorsoventral, - 8.0 mm from dura) and anchored to the skull using dental cement and three stainless steel screws. One of the three screws served as a ground electrode. Rats were given buprenorphine, housed individually, and allowed to recover from surgery for one week before the start of kindling procedures.

# **Kindling procedure**

The electrical kindling procedure was performed as previously reported (Colangeli et al., 2020). Briefly, twenty, twice daily, electrical stimulations of the right amygdala were elicited

using standard kindling stimulation (1 s train of 1 ms biphasic square wave pulses at 60 Hz) through a Grass S88 stimulator (Natus Neurology, Warwick, RI). Seizure duration and stage were recorded, and seizure score was classified as follows: Class 1: immobility, Class 2: orofacial automatisms with head nodding, Class 3: unilateral forelimb clonus, Class 4: rearing with bilateral forelimb clonus, Class 5: rearing with bilateral forelimb clonus followed by falling. Using this classification system, rats are considered "kindled" after three consecutive Class 5 convulsions (Teskey, 2020). Sham rats were treated identically but no current was delivered.

### Electrophysiology

One week after the last evoked sham or kindled seizure, rats were decapitated under deep anaesthesia with isoflurane and transcardially perfused with ice cold slicing solution containing (in mM): 87 NaCl, 2.5 KCl, 25 NaHCO<sub>3</sub>, 0.5 CaCl<sub>2</sub>, 7 MgCl<sub>2</sub>, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 25 D-glucose, and 75 sucrose. Coronal brain slices of the BLA (300 µm) were prepared with Leica VT1200S slicer in ice-cold slicing solution. Slices were incubated in a holding chamber with oxygenated aCSF containing in mM (126 NaCl, 2.5 KCl, 2.5 CaCl<sub>2</sub>, 1.5 MgCl<sub>2</sub>, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 26 NaHCO<sub>3</sub>, and 10 D-glucose at pH 7.4) for 30 min at 32 °C. Slices were then incubated for at least 45 min in regular aCSF or vehicle (DMSO), DO34 (1 µM) or KML29 (1µM) containing aCSF, depending on the experimental design. We applied drugs directly to the slices, rather than perform i.p. systemic injection, to narrow the potential extra-amygdala effects of eCB signaling manipulation in other brain regions, such as hippocampus (Segev et al., 2018) and frontal cortex (Morena et al., 2014; Vogel et al., 2016; Marcus et al., 2020), which may, in turn, affect eCB-mediated control of synaptic transmission and plasticity in the BLA. After incubation, slices were placed in the recording chamber and continuously superfused with either regular aCSF, or vehicle, DO34 or KML29 containing aCSF at a flow rate of 1.5 ml/min. All experiments were performed at room temperature (22-25°C) (Azad et al., 2004; Chevaleyre et al., 2007). Whole-cell patch-clamp recordings were performed on BLA principal neurons. GABA-A receptor-mediated inhibitory postsynaptic currents (IPSCs) were recorded by using glass electrodes  $(3-5 \text{ M}\Omega)$  filled with a solution containing in mM: 90 CsCH<sub>3</sub>SO<sub>3</sub>, 50 CsCl, 1 EGTA, 10 HEPES, 4.6 MgCl<sub>2</sub>, 0.1 CaCl<sub>2</sub>, 5 QX314, and 0.3 Na-GTP, 4 Mg-ATP, pH 7.3 adjusted with CsOH. IPSCs were isolated by adding to the bath 40 μM d(-)-2-amino-5-phosphono-pentanoic acid (d-AP-5) and 20 µM 6,7-Dinitroquinoxaline-2,3dione (DNQX) to block glutamatergic transmission. To elicit evoked IPSCs (eIPSCs), putative GABAergic fibers were stimulated using monopolar aCSF-filled patch-type pipettes

(1 M $\Omega$ ) placed within the BLA. For depolarization-induced suppression of inhibition (DSI) studies, stimuli were applied at 0.2 Hz. The magnitude of DSI was determined as the reduction from the mean amplitude of 12 successive eIPSCs evoked just before a depolarization (5 sec at 0 mV) of pyramidal cell (baseline, BL), compared to the mean amplitude of 3 successive eIPSCs taken just after the pulse (DSI) and the mean amplitude of the last 5 eIPSCs (90-120 s; recovery). Whenever possible, 2-4 DSI trials were averaged to obtain mean DSI in each condition. For mAChR-induced eCB mobilization experiments and CB1 sensitivity, evoked simulations were elicited every 15 s with 4 stimulations being averaged to obtain one data point per minute. mAChR-driven synaptic suppression was assessed by 10 min bath application of Carbachol (CCh, 10 µM) after 5 min BL recording, and the magnitude of LTD was calculated for 10 min (from 30 to 40 min after the end of CCh application) as a percentage of BL responses. Two neurons insensitive to CCh effect during its application were excluded from the LTD analysis. CB1 sensitivity was assessed by bath application of the CB1 receptor agonist CP55,940 (3µM) and calculated as the average of CB1-induced eIPSC amplitude depression 10-15 min from agonist application. eIPSCs were analysed with Clampfit 10.7 software. The amplitudes and frequencies of spontaneous IPSCs (sIPSCs) were detected (and confirmed by visual inspection) by continuous recording over 300 sec by using Mini Analysis 6.0.7 software (Synaptosoft) using thresholds of 5 times RMS noise levels. Cell capacitance and access resistance (initial value < 30 M $\Omega$ ) were monitored during experiments, and recordings were accepted for analysis if either variable did not change by more than 20%. Signals were collected via an Axopatch 200B amplifier (Molecular Devices). The pCLAMP 9 software (Molecular Devices) was used for data acquisition.

# Enzyme activity assays.

Brain structures were excised on ice and samples were then immediately snap frozen and stored at -80 °C. Brain tissues were homogenized, and membrane fractions were isolated as described previously (Vecchiarelli et al., 2021). The monoacylglycerol lipase (MAGL) activity was measured as the conversion of [3 H]-2- oleoylglycerol (2-OG) to [3 H]-glycerol. The maximal hydrolytic activity (Vmax) of MAGL and the binding affinity (Km) of 2-AG for MAGL were determined by fitting the data to the Michaelis-Menten equation.

### Statistical analyses

Data were analysed using Prism 8 (GraphPad, San Diego, CA, USA, RRID:SCR\_002798). Differences between groups were analysed using Student's t-test for unpaired or paired data, two-way or repeated measures (RM) ANOVA, followed by Tukey's post hoc test for multiple comparisons. Significance was accepted at p < 0.05. Statistical design for each experiment is detailed in the results section.

# Results

# Enduring loss of activity-dependent 2-AG retrograde signaling at GABA synapses after repeatedly evoked seizures.

To determine if repeated seizure activity persistently alters 2-AG actions at BLA GABAergic synapses one week after the last evoked seizure, we used the most well-established form of 2-AG-mediated retrograde eCB signaling expressed widely throughout the CNS; the depolarization-induced suppression of inhibition (DSI). DSI has been reliably reported in the BLA (Zhu and Lovinger, 2005; Patel et al., 2009; Yoshida et al., 2011) and, consistent with those data, we found that depolarization of BLA pyramidal neuron from -70 to 0 mV for 5 sec resulted in a transient depression of the amplitude of the evoked GABA-A receptormediated inhibitory postsynaptic currents (eIPSCs) which was followed by a complete recovery of inhibitory activity (RM ANOVA;  $F_{2,32} = 6.304$ , p = 0.017; DSI, 82.07% + 4.20%, of BL, p = 0.004; recovery: 99.17%  $\pm$  5.63%, of BL; p > 0.05, n = 11 neurons, 9 rats, Fig. 1A,B). In kindled rats, depolarization of pyramidal neurons did not induce a significant decrease of eIPSC magnitude (RM ANOVA; F<sub>2,35</sub> =0.260, p > 0.05; DSI; 96.44% + 4.152%, of BL, p > 0.05, recovery; 98.38%  $\pm$  5.526%, of BL, p > 0.05, n = 12 neurons, 9 rats, Fig. 1A,C). In the brain, DSI has been shown to be exclusively dependent on 2-AG retrograde signaling as it is completely absent in DAGL KO mice that are unable to synthesize 2-AG (Tanimura et al., 2010) and abolished by the DAGL inhibitor DO34 (Ogasawara et al., 2016). Here, we preincubated brain slices from sham rats with either vehicle or DO34 (1µM) (Ogasawara et al., 2016) and then tested DSI. We found that DO34 completely abolished DSI, thus mimicking what was observed in kindled rats in the absence of DO34. Indeed, while in vehicle-treated slices of sham rats, depolarization of pyramidal neurons induced a significant transient suppression of eIPSC (RM ANOVA; F<sub>2,23</sub> = 12.26, p = 0.002; DSI:  $81.79\% \pm 4.309\%$ , of BL; p = 0.009, n = 8 neurons, 5 rats) which eventually recovered to BL values (recovery:  $99.49\% \pm 4.894\%$ , p > 0.05 of BL, Fig. 1D,E), the transient suppression of eIPSCs after the depolarization step was absent in slices treated with DO34 (RM ANOVA;  $F_{2,20} = 0.219$ , p > 0.05; DSI: 97.37%  $\pm$  4.903%, of BL; recovery: 100.1%  $\pm$  5.104%, n = 7 neurons, 6 rats, **Fig. 1D,F**).

### 2-AG restoration normalizes DSI in kindled rats.

We then determined whether blocking 2-AG deactivation and thereby amplifying its actions at CB1 receptors could rescue the DSI impairment observed in kindled rats. We first assessed the effect of KML29 (1 $\mu$ M), an inhibitor of the 2-AG degrading enzyme, MAGL, on DSI in slices obtained from sham rats. As shown in Fig. 2, depolarization of post-synaptic neurons caused a similar transient suppression of eIPSC amplitude in slices treated with either vehicle (RM ANOVA;  $F_{2,23} = 6.64$ , p = 0.015; DSI:  $86.94\% \pm 4.29\%$ , of BL; p = 0.040, n = 8neurons, 7 rats, Fig. 2A,B), or KML29 (RM ANOVA;  $F_{2,41} = 17.36$ , p < 0.001; DSI: 88.26%  $\pm$  2.17%, of BL; p <0.001, n = 14 neurons, 5 rats, Fig. 2A,C), indicating that occlusion did not occur at the drug concentration used in these experiments. In addition, KML29 significantly increased the late component of DSI (recovery:  $84.91\% \pm 3.17\%$ , of BL; p = 0.01, Fig. 2C), thereby significantly prolonging the suppression of IPSC amplitude following postsynaptic depolarization. In the kindled group, preincubating slices with KML29 rescued DSI. Specifically, the transient suppression of eIPSC amplitude following depolarization step could be induced in KML29 treated slices (RM ANOVA;  $F_{2.35} = 22.45$ , p < 0.001; DSI:  $84.24\% \pm 3.09\%$ , of BL; p = 0.001, n = 12 neurons, 6 rats, Fig. 2D,F) but not in vehicle treated slices (RM ANOVA;  $F_{2,29} = 1.03$ , p = 0.36; DSI: 93.36%  $\pm$  5.40%, of BL, n = 10 neurons, 7 rats, Fig. 2D,E) of kindled rats. Moreover, as observed in sham rats, KML29 significantly prolonged DSI in kindled rats (recovery: 84.43% + 2.58%, of BL; p > 0.001, Fig. 2D,F).

# Repeated seizure activity attenuated mAChR-dependent depression on inhibitory transmission in the BLA.

Besides strong depolarization of neurons, eCB release can be triggered by activation of Gqcoupled receptors such as group I metabotropic glutamate receptors (mGluR1) and mAChRs (Maejima et al., 2001; Fukudome et al., 2004) through the PLC-DAGL cascade (Hashimotodani et al., 2005; Maejima et al., 2005). In the BLA, mGluR1 appears to stimulate AEA production, rather than 2-AG, independently of the PLC–DAGL activity (Azad et al., 2004). On the other hand, mAChR activation triggers PLC-dependent eCB release in many brain regions (Kim et al., 2002; Ohno-Shosaku et al., 2003; Neu et al., 2007) but has not yet been demonstrated in the BLA. Therefore, we first asked whether prolonged mAChR activation would also result in a long-term depression (LTD) in the BLA and whether this form of plasticity is dependent on the CB1 receptor. To activate mAChRs, after 5 min BL period, carbachol (CCh, 10 µM) was bath applied for 10 min and eIPSCs were recorded for 40 minutes in BLA pyramidal neurons from control non-operated rats (Fig. 3A). We found that CCh induced a transient depression followed by a long-lasting depression that remained upon washout (n = 9 neurons, 7 rats; paired t-test.  $t_8 = 4.36$ , p < 0.02 vs BL, Fig. **3B**). The long lasting depression after CCh washout was still present when the mAChR were blocked by scopolamine (n = 4 neurons, 4 rats; RM two-way ANOVA: SCOP X CCh interaction:  $F_{(1,11)} = 0.46$ , p > 0.05; CCh effect:  $F_{1,11} = 22.33$ , p < 0.001; SCOP effect:  $F_{1,11} = 0.46$ , p > 0.05, Fig. 3B), while it was reversed when slices were continuously superfused with the CB1 receptor inverse agonist AM251 (n= 7 neurons, 7 rats; RM two-way ANOVA: AM251 X CCh interaction:  $F_{1,14} = 5.18$ , p = 0.039; CCh effect:  $F_{1,14} = 21.93$ , p < 0.001; AM251 effect:  $F_{1,14} = 5.18$ , p = 0.039, Fig. 3C,D). Collectively these data provide evidence that BLA pyramidal neurons undergo CB1-dependent, mAChR-induced LTD. We then determined whether this form of plasticity is altered by seizure activity. Similarly to control non-operated rats, both sham and kindled rats displayed CCh-induced LTD (Fig. 4A) however, RM twoway ANOVA followed by Tukey's post-hoc test revealed that the reduction in GABA synaptic efficacy after CCh wash out was significantly more pronounced in sham rats (n = 8neurons, 7 rats) compared with kindled rats (n = 6 neurons, 4 rats); (RM two-way ANOVA: seizures X CCh interaction: F<sub>1,12</sub> = 12.87, p = 0.004; CCh effect: F<sub>1,12</sub> = 111.22, p < 0.001; seizure effect:  $F_{1,12} = 12.87$ , p = 0.004, Fig. 4B). Having established that inhibitory synapses in the BLA can undergo CB1-mediated, mAChR-dependent LTD and this form of plasticity is attenuated in kindled rats, we next asked if it requires 2-AG signaling. To this aim we preincubated slices from sham rats with the 2-AG synthesis inhibitor, DO34 (n = 4 neurons, 4 rats) and then bath applied CCh (Fig. 4C). In these conditions the long-lasting depression recorded after CCh wash out was significantly attenuated in the presence of DO34 (RM twoway ANOVA: DO34 X CCh interaction effect:  $F_{1,10} = 18.00$ , p = 0.002; CCh effect:  $F_{1,10} =$ 112.55, p < 0.001; DO34 effect:  $F_{1,10} = 18.00$ , p = 0.002, Fig. 4D), thus resembling the mAChR-LTD obtained in slices from kindled rats superfused with aCSF only. These findings support the possibility that 2-AG signaling, at least in part, subserves mAChR-LTD in the BLA and repeated seizures negatively impacted 2-AG-mediated plasticity in the BLA.

# Inhibition of the 2-AG degrading enzyme does not restore mAChR-dependent depression on inhibitory transmission in the BLA.

We then asked whether inhibiting 2-AG degradation might normalize mAChR-LTD in kindled rats by amplifying 2-AG activity at GABA terminals. To test this possibility, brain slices were treated with the MAGL inhibitor KML29 (1µM) before bath application of CCh. We first tested KML29 in slices from sham rats (n = 5 neurons, 4 rats; Fig. 5A). In contrast with what we expected, KML29 reduced, rather than enhanced mAChR-LTD in slices from sham rats (Fig. 5B). Indeed, the reduction in eIPSCs amplitude recorded in KML29 treated slices during CCh wash out was significantly less pronounced compared to eIPSCs amplitude recorded in vehicle treated slices (RM two-way ANOVA: KML29 X CCh interaction:  $F_{1,11} =$ 11.99, p = 0.005; CCh effect:  $F_{1,11}$  = 176.86, p < 0.001; KML29 effect:  $F_{1,11}$  = 11.99, p = 0.005, Fig. 5B). Moreover, despite a tendency to rescue pharmacologically induced LTD in slices prepared from kindled rats, KML29 was not sufficient to fully restore GABA synaptic depression during CCh wash out (Fig. 5C). Indeed, no significant effect was observed when we compared the reduction of eIPSCs amplitude respect to the BL during CCh wash out in vehicle (n = 6 neurons, 4 rats) and KML29 (n= 6 neurons, 3 rats) treated slices from kindled rats (RM two-way ANOVA: KML29 X CCh interaction:  $F_{1,10} = 3.09$ , p = 0.110; CCh effect:  $F_{1,10} = 49.68$ , p < 0.001; KML29 effect:  $F_{1,10} = 3.09$ , p = 0.110, Fig. 5D).

# Kindling-induced alteration of phasic eCB signaling is not mediated by altered 2-AG tone over GABA transmission.

We previously reported that tissue levels of AEA, but not 2-AG, were reduced in kindled rats one week after the last evoked seizure, which mainly impacted the tonic eCB control over glutamate transmission (Colangeli et al., 2020). However, alteration of the 2-AG tone over GABA transmission may occur at the synaptic scale and thereby not be revealed by analysing the whole amygdala. Since prolonged tonic signaling can cause desensitization and functional antagonism of CB1 receptors, the possibility that the alteration of synaptic plasticity was caused by changes in the 2-AG tone existed. To address this point, we examined basal GABAergic synaptic input onto BLA pyramidal neurons of sham or kindled rats in the presence of either vehicle or DO34 (**Fig. 6A,B**). If 2-AG signaling is tonically present under basal conditions to suppress GABAergic transmission, then blockade of its synthesis by DO34 should unmask a facilitation of inhibitory transmission. DO34 treatment did not change spontaneous IPSCs (sIPSC) frequency and amplitude in either sham rats (unpaired ttest for frequency:  $t_{21} = 0.143$ , p = 0.887; amplitude:  $t_{21} = 0.643$ , p = 0.527, aCSF, n = 12 neurons, 9 rats, DO34; 11 neurons, 9 rats; **Fig. 6A**) or kindled rats (unpaired t-test for frequency:  $t_{19} = 0.635$ , p = 0.533; amplitude:  $t_{19} = 0.0735$ , p = 0.942, aCSF; n = 10 neurons, 8 rats, DO34; n = 11 neurons, 4 rats, **Fig. 6B**). Moreover, to test whether there is a tonic CB1 receptor-mediated control over GABA transmission, presumably mediated by AEA, we performed a time course analysis of sIPSC before and after bath application of the CB1 receptor inverse agonist AM251. Again, we found no difference in either frequency or amplitude of sIPSC in both sham (paired t-test for frequency:  $t_{(6)} = 0.77$ , p > 0.05; amplitude:  $t_{(6)} = 1.76$ , p > 0.05, n = 7 neurons, 7 rats, **Fig. 6C**) and kindled rats (paired t-test for frequency:  $t_{(6)} = 1.85$ , p > 0.05; amplitude:  $t_{(6)} = 1.64$ , p > 0.05, n = 7 neurons, 6 rats, **Fig. 6D**). These data confirm previous evidence showing that GABAergic inputs in the BLA are not under a tonic control of eCB signaling (Colangeli et al., 2020; Yasmin et al., 2020) and providing evidence that altered tonic 2-AG signaling does not account for the disrupted activity-dependent 2-AG action over GABA transmission.

# Kindling-induced alteration of phasic eCB signaling is not mediated by changes in CB1 receptor sensitivity.

Repeated seizures could cause functional tolerance of CB1 receptor and thereby lead to an apparent abolition of eCB-mediated plasticity. To test whether repeated seizure activity may directly impair CB1 receptor functionality, we bath-applied the cannabinoid agonist CP55,940 ( $3\mu$ M) and recorded eIPSCs (**Fig. 7A**). In both groups, CP55,940 significantly inhibited synaptic transmission similarly (RM two-way ANOVA; CP55,940 X seizures interaction effect: F<sub>1,9</sub> = 0.3170, p = 0.587; CP55,940 effect: F<sub>1,9</sub> = 44.42, p < 0.001; seizures effect: F<sub>1,9</sub> = 0.3170 p = 0.587, sham; n = 5 neurons, 3 rats, kindled; n = 6 neurons, 6 rats; **Fig. 7B**). This finding rules out the possibility that altered CB1 receptor sensitivity contributes to the loss of 2-AG signaling in kindled rats.

# Enhanced MAGL enzymatic breakdown activity does not account for the seizureinduced 2-AG-signaling alteration.

After having assessed that neither basal 2-AG signaling nor CB1 receptor sensitivity are altered by seizure activity, the loss of activity-dependent 2-AG signaling over GABA transmission after repetitive seizure activity may be attributable to impairments of 2-AG synthesis/release from the postsynaptic cell or enhanced enzymatic breakdown activity. Our data so far indicate that the seizure-induced alteration in 2-AG signaling may be due to a

dysfunction in 2-AG synthesis/release machinery since DO34, a synthesis inhibitor, is able to recapitulate these changes; nonetheless, enhanced 2-AG degradation in BLA may lead to a reduction in 2-AG levels thus contributing to the loss of stimulus-dependent 2-AG signaling in BLA. Thereby, in both sham and kindled rats, we a) assessed the effects of KML29, an inhibitor of 2-AG degradation by MAGL, on GABA transmission and 2) tested the enzymatic activity of MAGL in amygdala tissue. If there is an enhanced enzymatic MAGL activity following repeated seizure activity, KML29 treatment should affect sIPSCs in kindled rats more profoundly than sham rats. However, in brain slices prepared from sham rats sIPSC frequency, but not the amplitude, was significantly reduced by KML29 treatment (unpaired ttest for frequency:  $t_{23} = 3.06$ , p = 0.006; amplitude:  $t_{23} = 0.886$ , p > 0.05, aCSF; n = 11neurons, 5 rats, KML29; n = 14 neurons, 5 rats; Fig. 8A) while this effect was absent in brain slices prepared from kindled rats (unpaired t-test for frequency:  $t_{23} = 0.67$ , p > 0.05; amplitude: t<sub>23</sub> = 0.20, p > 0.05, aCSF; n = 12 neurons, 5 rats, KML29; n = 13 neurons, 6 rats; Fig. 8B). Moreover, when MAGL enzymatic activity was assessed in the amygdala, the  $V_{\text{max}}$ of MAGL was unchanged after kindling (unpaired t-test:  $t_9 = 0.692$ , p = 0.506), but  $K_m$  for MAGL was found increased compared with sham rats (unpaired t-test:  $t_9 = 2.406$ , p = 0.040; Fig. 8C). This latter finding clearly indicates a decrease in the affinity of MAGL for its substrate, thus a less efficient capability of the enzyme to catabolize 2-AG. Collectively, these results rule out the possibility that enhanced 2-AG degradation accounts for the altered 2-AG mediated synaptic plasticity alterations over GABA transmission induced by seizure activity.

### Discussion

The main findings of the present work are that repeated amygdala seizure activity impairs BLA GABAergic plasticity and that these effects are recapitulated in non-kindled rats by pharmacological disruption of 2-AG synthesis. Moreover, augmentation of 2-AG signaling restores activity-dependent, but not mAChR-induced synaptic plasticity of GABA transmission. Neither CB1 receptor sensitivity nor 2-AG degradation was altered by seizures, thus, providing evidence that the most likely mechanism underlying aberrant 2-AG signaling over GABA transmission following repeated seizure activity is dysfunctional 2-AG synthesis/release machinery. We previously reported that both tonic AEA control over glutamatergic transmission and phasic AEA signaling over GABA transmission are impaired

by repeated seizure activity (Colangeli et al., 2020). We expand these findings by demonstrating that activity-dependent 2-AG signaling dysfunction also contributes to the persistent aberrant GABA plasticity in the BLA following repeated seizures, providing evidence that amygdala kindling causes an enduring alteration of both AEA and 2-AG signaling within the BLA.

Persistent alteration of neuroplasticity is an important hallmark of the pathophysiology of epilepsy and contributes to the negative neurobiological consequences associated with seizure activity (Farrell et al., 2017; Farrell et al., 2020). It is well established that facilitation of eCB signaling acutely dampens seizure expression and provides neuroprotection during seizure activity, whereas eCB signaling blockade often does the opposite (Marsicano et al., 2003; Katona and Freund, 2008; Colangeli et al., 2017; Sugaya and Kano, 2018; Colangeli et al., 2019; Di Maio et al., 2019). Conversely, evidence of pathologic long-term remodelling of the eCB signaling following seizure activity is less consistent across studies and the relative contributions of AEA and 2-AG are still a matter of debate (Soltesz et al., 2015; Sugaya and Kano, 2018).

DSI is a widespread, 2-AG-mediated phenomenon crucially involved in proper brain plasticity, and alteration of DSI is associated with a wide range of neurological diseases (Patel et al., 2009; Wamsteeker et al., 2010; Wang et al., 2010; Jenniches et al., 2016; Folkes et al., 2020). Earlier studies have demonstrated hippocampal and cortical long-term dynamic changes of eCB signaling in animal models of chronic epilepsy, with the most consistent findings being an upregulation of CB1 receptor function at GABAergic terminals following status epilepticus (Falenski et al., 2007; Falenski et al., 2009; Maglòczky et al., 2010) and a persistent potentiation of the activity-dependent 2-AG signaling selectively at inhibitory (DSI) synapses following febrile seizures (Chen et al., 2003). As a consequence of the enduring upregulation of the eCB system over GABA transmission, suppression, rather than enhancement, of eCB signaling seemed to provide protection during epileptogenesis (Chen et al., 2007). In contrast to these findings, here we demonstrate that activity-dependent eCB retrograde signaling at BLA GABA synapses is absent in amygdala kindled rats, and this alteration is recapitulated in slices of sham rats treated with the 2-AG synthesis inhibitor DO34. Importantly, pharmacological normalization of 2-AG levels restores DSI in kindled rats, thus providing evidence that persistent 2-AG signaling deficiency, rather than enhancement, occurs following amygdala kindling. The apparent controversy between our findings and previous works might be attributable to the model used to induce seizures and an epileptic state. Indeed, while the aforementioned studies used an initial insult to trigger epileptogenesis (i.e. febrile seizure and chemically-induced status epilepticus), studies that used kindling consistently indicate that deficiency of the 2-AG signaling occurs, since augmentation of 2-AG signaling retarded epileptogenesis (Griebel et al., 2015; von Ruden et al., 2015) while disruption of 2-AG synthesis increased seizure susceptibility (Sugaya et al., 2016). Thus, the temporal evolution of the pathological plasticity of the eCB signaling system may differ depending on the epilepsy model employed.

It is also important to note that, when pathological alterations are found at the presynaptic level, the electrophysiological experimental paradigms used to investigate the synaptic activity may account for discrepancies among different findings. For example, extracellular  $Ca^{2+}$  concentration used in the recording solution might alter release probability (Mintz et al., 1995), synaptic plasticity (Inglebert et al., 2020), and eCB signaling (Lenz and Alger, 1999; Varma et al., 2002). The extracellular solution in these experiments (2.5 mM  $Ca^{2+}$  and 1.5 mM  $Mg^{2+}$ ) included concentrations of  $Ca^{2+}$  in the upper value of physiological levels in rat brain (1–3 mM) (Rusakov and Fine, 2003) as it is widely used to test eCB control of synaptic transmission (Báldi et al., 2016).

The eCB system reorganization following seizures may also depend on the brain region involved in seizure activity. In the BLA, eCB signaling controls both glutamatergic and GABAergic transmission, thus we speculate that repeated electrically evoked seizures may cause a persistent and unselective (glutamate vs GABA) dysregulation of eCB signaling in the BLA when it is required for retrograde suppression of synaptic transmission. The consequences of eCB signaling breakdown are a lack of control over seizure generation when impacting glutamatergic terminals (mainly AEA) and aberrant synaptic plasticity when impacting GABAergic transmission (mainly 2-AG). In this context, augmentation of eCB signaling provides neuroprotection against epileptiform discharges and restores proper GABA plasticity, crucially involved in amygdala-mediated emotional behaviour (Ehrlich et al., 2009).

Besides the 2-AG-mediated short-term plasticity, the production of eCBs is also tied to the activity of the metabotropic receptors such as postsynaptic mGluR and mAChR (Kano et al., 2009), which are critical for eCB-mediated long-term depression (LTD) at central synapses (Chevaleyre and Castillo, 2003; Katona and Freund, 2012). Several studies have demonstrated that, in the BLA, low frequency stimulation can mobilise eCBs, and

particularly AEA, via activation of postsynaptic mGluR1 (Marsicano et al., 2002; Azad et al., 2004; Chevaleyre et al., 2007) and consistently, seizure-induced alteration of LTD of inhibitory transmission is completely rescued by AEA signaling augmentation (Colangeli et al., 2020). On the other hand, "on demand" eCB synthesis is not linked to special metabotropic receptors (Heifets and Castillo, 2009) and, independently of the mGluR system, mAChR activation mediates LTD through 2-AG signaling in several brain regions (Tanimura et al., 2010). Since BLA pyramidal neurons richly express mAChR, which are closely positioned to DAGL clusters (Yoshida et al., 2011), we tested whether, besides AEA-mGluRdriven LTD, mAChR activation also triggers eCB-LTD at BLA GABAergic synapses and if it requires 2-AG/CB1 signaling. Our results further confirmed our previous report that seizures severely impact LTD at GABA transmission within the BLA (Colangeli et al., 2020), as mAChR-driven LTD was consistently induced in sham rats while significantly attenuated in kindled rats. However, our results do not fully elucidate whether this seizure-induced alteration of LTD might be ascribed to differences in 2-AG signaling. Indeed, while DSI was completely abolished with DO34, we were unable to completely abolish mAChR-induced synaptic depression by inhibiting the DAGL enzyme in the sham group. It is important to note that different sensitivities to DAGL inhibitors between metabotropic receptor-driven 2-AG mobilisation and stimulus-dependent 2-AG mobilisation have been previously reported. For example, pharmacological DAGL inhibition fully blocked DSI but only partially prevented mGluR-driven 2-AG mobilization in the CA1 region of the hippocampus (Zhang et al., 2011). This could explain why mAChR-induced LTD might be less sensitive to DO34 than DSI.

Another possible explanation is that, unlike other brain regions, 2-AG activity plays only a partial role in the expression of mAChR-driven LTD at GABA transmission in the BLA. In support of this hypothesis, we found that AM251 treatment completely blocked mAChR-induced LTD in naïve rats; thus, the residual synaptic depression observed after CCh washout in DO34 treated rats is likely to be ascribed to AEA signaling released by the activation of mAChR receptors (van der Stelt et al., 2005). This latter hypothesis, together with our previous findings showing an altered AEA signaling in the amygdala of kindled rats, could explain why boosting the 2-AG signal, which fully restored DSI in kindled rats, did not completely rescue kindling-induced LTD impairments, as AEA signaling dysfunction may contribute to LTD dysfunction. The possible contribution of AEA signaling on CCh-induced LTD may also explain why KML29 treatment impaired, rather than enhanced, CCh-induced

LTD in sham rats. Indeed, if both AEA and 2-AG are involved in this form of plasticity, boosting 2-AG signaling might occlude LTD in sham rats by interfering with AEA signaling.

A further possibility is that other neuromodulators might influence eCB-mediated LTD and may account for the attenuated plasticity seen in kindled rats. For example, in some conditions, nitric oxide appears to be necessary for the induction of eCB-mediated LTD (Crosby et al., 2011) and eCB-mediated short-term regulation of GABAergic synaptic transmission (Makara et al., 2007). Intriguingly, in this latter finding, nitric oxide is involved in DSI expression only in the presence of cholinergic receptor agonists (Makara et al., 2007). Given the contribution of nitric oxide signaling in seizure expression (Gupta and Dettbarn, 2003; Kato et al., 2005; Kovács et al., 2009), a potential involvement of nitric oxide signaling of seizure-induced disruption of eCB-mediated plasticity is an interesting mechanism that it is worth further investigation.

We have previously reported (Colangeli et al., 2020) and confirmed in the current work (data not shown) that sIPSC frequency is significantly reduced in the kindled group. Since the eCB signaling subserves tonic constraint of basal synaptic activity in many brain regions, it is possible that enhanced 2-AG tone over GABA transmission in kindled rats occurred. This would lead to a reduction of the basal GABA synaptic activity and may also result in occlusion of activity-dependent 2-AG depression of inhibitory transmission. However, we found that neither CB1 receptor blockade nor interference with either 2-AG synthesis or degradation altered the sIPSC frequency reduction found in kindled rats. Together with the lack of effect of AEA degradation inhibitor on sIPSC reduction found in kindled rats in our previous paper (Colangeli et al., 2020), we ruled out any possible occlusion of phasic 2-AG signaling after repeated seizure activity performed by an enhanced tonic eCB signaling over GABA transmission. Conversely, we found that inhibiting the degradation of 2-AG in sham rats treated with MAGL inhibitor might be ascribed to an occlusion effect.

A further possible explanation for the seizure-induced loss of 2-AG signaling in kindled rats is an altered CB1 receptor functionality. It has been recently shown that seizure activity induces intense and prolonged activation of CB1 receptors by 2-AG during ictal events, which quickly returns to baseline soon after seizure cessation (Farrell et al., 2021). Thus, it is possible that seizure-evoked transient but repeated 2-AG signaling activation of CB1 receptors can cause a persistent compensatory CB1 receptor desensitization and, in turn, an enduring deficiency in the 2-AG-mediated responses over GABA transmission. However, we excluded this possibility since our results provide evidence that CB1 receptor sensitivity is not altered after twenty evoked amygdala seizures in BLA.

We finally asked whether the most likely mechanism subserving phasic 2-AG signaling collapse following repeated seizures is a reduction in the stimulus-dependent 2-AG mobilization or an enhanced 2-AG degradation. 2-AG is generated postsynaptically, but its synaptic effect is regulated strictly by presynaptic MAGL activity. Here we found that while MAGL activity in the amygdala is not altered following seizure activity the binding affinity of 2-AG for MAGL enzyme was found to be reduced. This reduction in binding affinity indicates a less efficient capability of the enzyme to catabolize 2-AG, which would actually lead to increased 2-AG signaling. This counterintuitive finding is likely explained by the reduction of MAGL binding affinity for 2-AG as a compensatory mechanism to counteract the impaired capability of postsynaptic neurons to mobilise 2-AG when needed.

In conclusion, we show that persistent 2-AG signaling maladaptation occurs in the BLA following repeated seizures, leading to aberrant plasticity in a brain region crucially involved in emotional behaviour. Thereby enhancing 2-AG-CB1 signaling via MAGL inhibition could represent an attractive therapeutic approach for the treatment of epilepsy-related emotional disorders.

# References

- Azad SC, Monory K, Marsicano G, Cravatt BF, Lutz B, Zieglgansberger W, Rammes G (2004) Circuitry for associative plasticity in the amygdala involves endocannabinoid signaling. J Neurosci 24:9953-9961.
- Báldi R, Ghosh D, Grueter BA, Patel S (2016) Electrophysiological Measurement of Cannabinoid-Mediated Synaptic Modulation in Acute Mouse Brain Slices. Current protocols in neuroscience 75:6.29.21-26.29.19.
- Castillo PE, Younts TJ, Chàvez AE, Hashimotodani Y (2012) Endocannabinoid signaling and synaptic function. Neuron 76:70-81.
- Chen K, Neu A, Howard AL, Foldy C, Echegoyen J, Hilgenberg L, Smith M, Mackie K, Soltesz I (2007) Prevention of plasticity of endocannabinoid signaling inhibits persistent limbic hyperexcitability caused by developmental seizures. J Neurosci 27:46-58.
- Chen K, Ratzliff A, Hilgenberg L, Gulyas A, Freund TF, Smith M, Dinh TP, Piomelli D, Mackie K, Soltesz I (2003) Long-term plasticity of endocannabinoid signaling induced by developmental febrile seizures. Neuron 39:599-611.

- Chevaleyre V, Castillo PE (2003) Heterosynaptic LTD of hippocampal GABAergic synapses: a novel role of endocannabinoids in regulating excitability. Neuron 38:461-472.
- Chevaleyre V, Heifets BD, Kaeser PS, Sudhof TC, Castillo PE (2007) Endocannabinoidmediated long-term plasticity requires cAMP/PKA signaling and RIM1alpha. Neuron 54:801-812.
- Colangeli R, Teskey GC, Di Giovanni G (2021) Endocannabinoid-serotonin systems interaction in health and disease. Prog Brain Res 259:83-134.
- Colangeli R, Morena M, Pittman QJ, Hill MN, Teskey GC (2020) Anandamide Signaling Augmentation Rescues Amygdala Synaptic Function and Comorbid Emotional Alterations in a Model of Epilepsy. J Neurosci 40:6068-6081.
- Colangeli R, Pierucci M, Benigno A, Campiani G, Butini S, Di Giovanni G (2017) The FAAH inhibitor URB597 suppresses hippocampal maximal dentate afterdischarges and restores seizure-induced impairment of short and long-term synaptic plasticity. Sci Rep 7:11152.
- Colangeli R, Di Maio R, Pierucci M, Deidda G, Casarrubea M, Di Giovanni G (2019) Synergistic action of CB1 and 5-HT2B receptors in preventing pilocarpine-induced status epilepticus in rats. Neurobiol Dis 125:135-145.
- Crosby KM, Inoue W, Pittman QJ, Bains JS (2011) Endocannabinoids gate state-dependent plasticity of synaptic inhibition in feeding circuits. Neuron 71:529-541.
- Di Maio R, Colangeli R, Di Giovanni G (2019) WIN 55,212-2 Reverted Pilocarpine-Induced Status Epilepticus Early Changes of the Interaction among 5-HT2C/NMDA/CB1 Receptors in the Rat Hippocampus. ACS Chem Neurosci 10:3296-3306.
- Ehrlich I, Humeau Y, Grenier F, Ciocchi S, Herry C, Luthi A (2009) Amygdala inhibitory circuits and the control of fear memory. Neuron 62:757-771.
- Engel J, Jr. (2001) Mesial temporal lobe epilepsy: what have we learned? The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry 7:340-352.
- Falenski KW, Blair RE, Sim-Selley LJ, Martin BR, DeLorenzo RJ (2007) Status epilepticus causes a long-lasting redistribution of hippocampal cannabinoid type 1 receptor expression and function in the rat pilocarpine model of acquired epilepsy. Neuroscience 146:1232-1244.
- Falenski KW, Carter DS, Harrison AJ, Martin BR, Blair RE, DeLorenzo RJ (2009) Temporal characterization of changes in hippocampal cannabinoid CB(1) receptor expression following pilocarpine-induced status epilepticus. Brain Res 1262:64-72.
- Farrell JS, Colangeli R, Wolff MD, Wall AK, Phillips TJ, George A, Federico P, Teskey GC (2017) Postictal hypoperfusion/hypoxia provides the foundation for a unified theory of seizure-induced brain abnormalities and behavioral dysfunction. Epilepsia 58:1493-1501.
- Farrell JS, Colangeli R, Dudok B, Wolff MD, Nguyen SL, Jackson J, Dickson CT, Soltesz I, Teskey GC (2020) In vivo assessment of mechanisms underlying the neurovascular basis of postictal amnesia. Sci Rep 10:14992.
- Farrell JS, Colangeli R, Dong A, George AG, Addo-Osafo K, Kingsley PJ, Morena M, Wolff MD, Dudok B, He K, Patrick TA, Sharkey KA, Patel S, Marnett LJ, Hill MN, Li Y, Teskey GC, Soltesz I (2021) In vivo endocannabinoid dynamics at the timescale of physiological and pathological neural activity. Neuron 109:2398-2403.e2394.
- Folkes OM, Báldi R, Kondev V, Marcus DJ, Hartley ND, Turner BD, Ayers JK, Baechle JJ, Misra MP, Altemus M, Grueter CA, Grueter BA, Patel S (2020) An endocannabinoidregulated basolateral amygdala-nucleus accumbens circuit modulates sociability. J Clin Invest 130:1728-1742.

- Fukudome Y, Ohno-Shosaku T, Matsui M, Omori Y, Fukaya M, Tsubokawa H, Taketo MM, Watanabe M, Manabe T, Kano M (2004) Two distinct classes of muscarinic action on hippocampal inhibitory synapses: M2-mediated direct suppression and M1/M3mediated indirect suppression through endocannabinoid signalling. Eur J Neurosci 19:2682-2692.
- Griebel G et al. (2015) Selective blockade of the hydrolysis of the endocannabinoid 2arachidonoylglycerol impairs learning and memory performance while producing antinociceptive activity in rodents. Sci Rep 5:7642.
- Gupta RC, Dettbarn WD (2003) Prevention of kainic acid seizures-induced changes in levels of nitric oxide and high-energy phosphates by 7-nitroindazole in rat brain regions. Brain Res 981:184-192.
- Hashimotodani Y, Ohno-Shosaku T, Tsubokawa H, Ogata H, Emoto K, Maejima T, Araishi K, Shin HS, Kano M (2005) Phospholipase Cbeta serves as a coincidence detector through its Ca2+ dependency for triggering retrograde endocannabinoid signal. Neuron 45:257-268.
- Heifets BD, Castillo PE (2009) Endocannabinoid signaling and long-term synaptic plasticity. Annu Rev Physiol 71:283-306.
- Hill MN, Tasker JG (2012) Endocannabinoid signaling, glucocorticoid-mediated negative feedback, and regulation of the hypothalamic-pituitary-adrenal axis. Neuroscience 204:5-16.
- Inglebert Y, Aljadeff J, Brunel N, Debanne D (2020) Synaptic plasticity rules with physiological calcium levels. Proc Natl Acad Sci U S A 117:33639-33648.
- Jenniches I, Ternes S, Albayram O, Otte DM, Bach K, Bindila L, Michel K, Lutz B, Bilkei-Gorzo A, Zimmer A (2016) Anxiety, Stress, and Fear Response in Mice With Reduced Endocannabinoid Levels. Biol Psychiatry 79:858-868.
- Kano M, Ohno-Shosaku T, Hashimotodani Y, Uchigashima M, Watanabe M (2009) Endocannabinoid-mediated control of synaptic transmission. Physiol Rev 89:309-380.
- Kato N, Sato S, Yokoyama H, Kayama T, Yoshimura T (2005) Sequential changes of nitric oxide levels in the temporal lobes of kainic acid-treated mice following application of nitric oxide synthase inhibitors and phenobarbital. Epilepsy Res 65:81-91.
- Katona I, Freund TF (2008) Endocannabinoid signaling as a synaptic circuit breaker in neurological disease. Nat Med 14:923-930.
- Katona I, Freund TF (2012) Multiple functions of endocannabinoid signaling in the brain. Annu Rev Neurosci 35:529-558.
- Kim J, Isokawa M, Ledent C, Alger BE (2002) Activation of muscarinic acetylcholine receptors enhances the release of endogenous cannabinoids in the hippocampus. J Neurosci 22:10182-10191.
- Kovács R, Rabanus A, Otáhal J, Patzak A, Kardos J, Albus K, Heinemann U, Kann O (2009) Endogenous nitric oxide is a key promoting factor for initiation of seizure-like events in hippocampal and entorhinal cortex slices. J Neurosci 29:8565-8577.
- Lenz RA, Alger BE (1999) Calcium dependence of depolarization-induced suppression of inhibition in rat hippocampal CA1 pyramidal neurons. J Physiol 521 Pt 1:147-157.
- Maejima T, Hashimoto K, Yoshida T, Aiba A, Kano M (2001) Presynaptic inhibition caused by retrograde signal from metabotropic glutamate to cannabinoid receptors. Neuron 31:463-475.
- Maejima T, Oka S, Hashimotodani Y, Ohno-Shosaku T, Aiba A, Wu D, Waku K, Sugiura T, Kano M (2005) Synaptically driven endocannabinoid release requires Ca2+-assisted metabotropic glutamate receptor subtype 1 to phospholipase Cbeta4 signaling cascade in the cerebellum. J Neurosci 25:6826-6835.

- Maglòczky Z, Tóth K, Karlòcai R, Nagy S, Eross L, Czirjàk S, Vajda J, Ràsonyi G, Kelemen A, Juhos V, Halàsz P, Mackie K, Freund TF (2010) Dynamic changes of CB1receptor expression in hippocampi of epileptic mice and humans. Epilepsia 51 Suppl 3:115-120.
- Makara JK, Katona I, Nyíri G, Németh B, Ledent C, Watanabe M, de Vente J, Freund TF, Hájos N (2007) Involvement of nitric oxide in depolarization-induced suppression of inhibition in hippocampal pyramidal cells during activation of cholinergic receptors. J Neurosci 27:10211-10222.
- Marcus DJ, Bedse G, Gaulden AD, Ryan JD, Kondev V, Winters ND, Rosas-Vidal LE, Altemus M, Mackie K, Lee FS, Delpire E, Patel S (2020) Endocannabinoid Signaling Collapse Mediates Stress-Induced Amygdalo-Cortical Strengthening. Neuron 105:1062-1076 e1066.
- Marsicano G, Wotjak CT, Azad SC, Bisogno T, Rammes G, Cascio MG, Hermann H, Tang J, Hofmann C, Zieglgansberger W, Di Marzo V, Lutz B (2002) The endogenous cannabinoid system controls extinction of aversive memories. Nature 418:530-534.
- Marsicano G, Goodenough S, Monory K, Hermann H, Eder M, Cannich A, Azad SC, Cascio MG, Gutierrez SO, van der Stelt M, Lopez-Rodriguez ML, Casanova E, Schutz G, Zieglgansberger W, Di Marzo V, Behl C, Lutz B (2003) CB1 cannabinoid receptors and on-demand defense against excitotoxicity. Science 302:84-88.
- Mintz IM, Sabatini BL, Regehr WG (1995) Calcium control of transmitter release at a cerebellar synapse. Neuron 15:675-688.
- Morena M, Roozendaal B, Trezza V, Ratano P, Peloso A, Hauer D, Atsak P, Trabace L, Cuomo V, McGaugh JL, Schelling G, Campolongo P (2014) Endogenous cannabinoid release within prefrontal-limbic pathways affects memory consolidation of emotional training. Proc Natl Acad Sci U S A 111:18333-18338.
- Neu A, Foldy C, Soltesz I (2007) Postsynaptic origin of CB1-dependent tonic inhibition of GABA release at cholecystokinin-positive basket cell to pyramidal cell synapses in the CA1 region of the rat hippocampus. J Physiol 578:233-247.
- Ogasawara D, Deng H, Viader A, Baggelaar MP, Breman A, den Dulk H, van den Nieuwendijk AM, Soethoudt M, van der Wel T, Zhou J, Overkleeft HS, Sanchez-Alavez M, Mori S, Nguyen W, Conti B, Liu X, Chen Y, Liu QS, Cravatt BF, van der Stelt M (2016) Rapid and profound rewiring of brain lipid signaling networks by acute diacylglycerol lipase inhibition. Proc Natl Acad Sci U S A 113:26-33.
- Ohno-Shosaku T, Matsui M, Fukudome Y, Shosaku J, Tsubokawa H, Taketo MM, Manabe T, Kano M (2003) Postsynaptic M1 and M3 receptors are responsible for the muscarinic enhancement of retrograde endocannabinoid signalling in the hippocampus. Eur J Neurosci 18:109-116.
- Patel S, Kingsley PJ, Mackie K, Marnett LJ, Winder DG (2009) Repeated homotypic stress elevates 2-arachidonoylglycerol levels and enhances short-term endocannabinoid signaling at inhibitory synapses in basolateral amygdala. Neuropsychopharmacology 34:2699-2709.
- Rusakov DA, Fine A (2003) Extracellular Ca2+ depletion contributes to fast activitydependent modulation of synaptic transmission in the brain. Neuron 37:287-297.
- Segev A, Korem N, Mizrachi Zer-Aviv T, Abush H, Lange R, Sauber G, Hillard CJ, Akirav I (2018) Role of endocannabinoids in the hippocampus and amygdala in emotional memory and plasticity. Neuropsychopharmacology 43:2017-2027.
- Soltesz I, Alger BE, Kano M, Lee SH, Lovinger DM, Ohno-Shosaku T, Watanabe M (2015) Weeding out bad waves: towards selective cannabinoid circuit control in epilepsy. Nat Rev Neurosci 16:264-277.

- Sugaya Y, Kano M (2018) Control of excessive neural circuit excitability and prevention of epileptic seizures by endocannabinoid signaling. Cell Mol Life Sci 75:2793-2811.
- Sugaya Y, Kano M (2021) Endocannabinoid-Mediated Control of Neural Circuit Excitability and Epileptic Seizures. Front Neural Circuits 15:781113.
- Sugaya Y, Yamazaki M, Uchigashima M, Kobayashi K, Watanabe M, Sakimura K, Kano M (2016) Crucial Roles of the Endocannabinoid 2-Arachidonoylglycerol in the Suppression of Epileptic Seizures. Cell Rep 16:1405-1415.
- Tanimura A, Yamazaki M, Hashimotodani Y, Uchigashima M, Kawata S, Abe M, Kita Y, Hashimoto K, Shimizu T, Watanabe M, Sakimura K, Kano M (2010) The endocannabinoid 2-arachidonoylglycerol produced by diacylglycerol lipase alpha mediates retrograde suppression of synaptic transmission. Neuron 65:320-327.
- Teskey GC (2020) Kindling. In: Biological Foundations of Psychology, Cognitive Psychology/Neuroscience, Methods and Approaches in Psychology (Psychology. OREo, ed). Oxford Oxford University
- van der Stelt M, Trevisani M, Vellani V, De Petrocellis L, Schiano Moriello A, Campi B, McNaughton P, Geppetti P, Di Marzo V (2005) Anandamide acts as an intracellular messenger amplifying Ca2+ influx via TRPV1 channels. The EMBO journal 24:3026-3037.
- Varma N, Brager D, Morishita W, Lenz RA, London B, Alger B (2002) Presynaptic factors in the regulation of DSI expression in hippocampus. Neuropharmacology 43:550-562.
- Vecchiarelli HA, Morena M, Keenan CM, Chiang V, Tan K, Qiao M, Leitl K, Santori A, Pittman QJ, Sharkey KA, Hill MN (2021) Comorbid anxiety-like behavior in a rat model of colitis is mediated by an upregulation of corticolimbic fatty acid amide hydrolase. Neuropsychopharmacology 46:992-1003.
- Vogel E, Krabbe S, Grundemann J, Wamsteeker Cusulin JI, Luthi A (2016) Projection-Specific Dynamic Regulation of Inhibition in Amygdala Micro-Circuits. Neuron 91:644-651.
- von Ruden EL, Bogdanovic RM, Wotjak CT, Potschka H (2015) Inhibition of monoacylglycerol lipase mediates a cannabinoid 1-receptor dependent delay of kindling progression in mice. Neurobiol Dis 77:238-245.
- Wamsteeker JI, Kuzmiski JB, Bains JS (2010) Repeated stress impairs endocannabinoid signaling in the paraventricular nucleus of the hypothalamus. J Neurosci 30:11188-11196.
- Wang W, Sun D, Pan B, Roberts CJ, Sun X, Hillard CJ, Liu QS (2010) Deficiency in endocannabinoid signaling in the nucleus accumbens induced by chronic unpredictable stress. Neuropsychopharmacology 35:2249-2261.
- Wettschureck N, van der Stelt M, Tsubokawa H, Krestel H, Moers A, Petrosino S, Schütz G, Di Marzo V, Offermanns S (2006) Forebrain-specific inactivation of Gq/G11 family G proteins results in age-dependent epilepsy and impaired endocannabinoid formation. Molecular and cellular biology 26:5888-5894.
- Yasmin F, Colangeli R, Morena M, Filipski S, van der Stelt M, Pittman QJ, Hillard CJ, Teskey GC, McEwen BS, Hill MN, Chattarji S (2020) Stress-induced modulation of endocannabinoid signaling leads to delayed strengthening of synaptic connectivity in the amygdala. Proc Natl Acad Sci U S A 117:650-655.
- Yoshida T, Uchigashima M, Yamasaki M, Katona I, Yamazaki M, Sakimura K, Kano M, Yoshioka M, Watanabe M (2011) Unique inhibitory synapse with particularly rich endocannabinoid signaling machinery on pyramidal neurons in basal amygdaloid nucleus. Proc Natl Acad Sci U S A 108:3059-3064.

- Zhang L, Wang M, Bisogno T, Di Marzo V, Alger BE (2011) Endocannabinoids generated by Ca2+ or by metabotropic glutamate receptors appear to arise from different pools of diacylglycerol lipase. PLoS One 6:e16305.
- Zhu PJ, Lovinger DM (2005) Retrograde endocannabinoid signaling in a postsynaptic neuron/synaptic bouton preparation from basolateral amygdala. J Neurosci 25:6199-6207.

**Figure legends** 

# Fig.1. DSI is impaired in kindled rats, an effect mimicked by inhibition of 2-AG synthesis.

(A) Time course data showing a transient inhibition of eIPSC amplitude following postsynaptic depolarization (vertical dashed line) in sham and kindled rats. (B) Top, Baseline (BL, 1) trace represents the average of 12 stimulations before depolarization, "DSI" (2) trace represents the average of three stimulations immediately after the depolarization and "recovery" (3) trace represents the average of the last five stimulations (95-115 s) corresponding to the indicated regions in (A). (B) Bottom, summary data corresponding to shaded regions in (A) showing that DSI is induced in sham (B) but not in kindled rats (C). Significance is reported with respect to the preceding shaded BL region. Preincubating and superfusing slices from sham rats with the DAGL inhibitor DO34 1 $\mu$ M recapitulated the effect of seizures on DSI as shown in the time course graph (D) and in the summary data

graph (E) and (F), respectively. Scale bars, 200 pA, 40 ms. Post hoc versus BL p-values are shown as follows: \*\*p < 0.01. Data are shown as mean  $\pm$  SEM.

# Fig.2. Inhibition of 2-AG degradation with KML29 prolongs DSI in sham rats and restores DSI in kindled rats.

(A) Summary time course of eIPSC amplitude normalized to the BL period obtained from slices of sham rats treated with either vehicle or the MAGL inhibitor KML29 (1µM). Representative traces and summary data corresponding to shaded regions in (A) in slices from sham rats treated with either vehicle (B) or the MAGL inhibitor KML29 (C), showing that KML29 did not affect max-DSI but significantly prolonged the duration of DSI. Significance is reported with respect to the preceding shaded BL region. (D) Time course data showing the effect of vehicle and KML29 containing aCSF on DSI in kindled rats. Representative traces and summary data corresponding to shaded regions in (D) in slices from kindled rats treated with either vehicle (E) or KML29 (F) showing that DSI is restored by KML29 treatment in kindled rats. Significance is reported with respect to the preceding shaded BL region. and summary data corresponding to shaded regions in (D) in slices from kindled rats treated with either vehicle (E) or KML29 (F) showing that DSI is restored by KML29 treatment in kindled rats. Significance is reported with respect to the preceding shaded BL region. Scale bars, 200 pA, 40 ms. Post hoc versus BL p-values are shown as follows: \*\*\*p < 0.001, \*p < 0.01, \*p < 0.05. Data are shown as mean ± SEM.

# Fig.3 BLA pyramidal neurons display mAChR-dependent long-term synaptic depression on inhibitory transmission which is mediated by CB1 receptor activation.

(A) Time course graph showing the effect of mAChR activation on eIPSC amplitude on naive rats (CTRL). eIPSCs were evoked every 15 s and then averaged to obtain a value per minute. CCh (10 $\mu$ M) was bath applied for 10 min and then washed out to assess long-term suppression of eIPSCs. In a subset of cells, scopolamine (SCOP, 1 $\mu$ M) was bath applied during CCh washout to demonstrate that the synaptic depression is not attributable to a lasting effect of CCh that has not been completely washed out of slices. (B) Bar graphs quantifying the percentage of control of eIPSC amplitude by CCh 10-20 min post-application (2) with respect to the BL period (1) in aCSF and SCOP treated slices. (C) AM251 treatment (2 $\mu$ M) in neurons of CTRL rats completely blocked LTD after CCh washout. (D) Bar graphs quantifying the percentage of control eIPSC amplitude by CCh recorded 10-20 min post-application (2) respect to the BL period (1) in aCSF and AM251 treated slices from CTRL rats. Scale bars, 200 pA, 40 ms. Post hoc CTRL aCSF versus CTRL AM251 p-values: \*\*p <

0.01. Data are shown as mean  $\pm$  SEM. CTRL aCSF data are repeated in the graphs as they belong to the same experimental design.

# Fig.4. mAChR-dependent synaptic depression on inhibitory transmission in the BLA is attenuated in the kindled rats, and this effect is mimicked by application of the 2-AG synthesis inhibitor DO34.

(A) Time course graph showing the effect of acute mAChR activation on eIPSC amplitude on sham and kindled rats. eIPSCs were evoked every 15 s and then averaged to obtain a dot per minute. CCh (10 $\mu$ M) was bath applied for 10 min and then washed out to assess long term suppression of eIPSCs. (B) Bar graphs quantifying the percentage inhibition of eIPSC amplitude by CCh 10-20 min post-application (2) respect to the BL period (1) in sham and kindled rats. Kindled rats display a significant attenuation of LTD after CCh washout. (C) Time course graph showing the effect of mAChR activation on eIPSC amplitude on slices from sham rats, treated with vehicle or DO34. (D) Bar graphs quantifying the percentage inhibition of eIPSC amplitude by CCh recorded 10-20 min post-application (2) respect to the BL period (1) in sham and DO34 treated slices. DO34 did not affect acute suppression of eIPSC amplitude during CCh application but attenuated LTD after CCh washout, thus mimicking the impaired mAChR-induced LTD observed in kindled rats. Scale bars, 200 pA, 40 ms. Post hoc sham versus kindled p-values: \*\*\*p < 0.001; Post hoc sham versus DO34 p-values: \*\*\*p < 0.001. Data are shown as mean  $\pm$  SEM. Sham aCSF data are repeated in the graphs as they belong to the same experimental design.

Fig.5. Inhibition of 2-AG degradation with KML29 reduces mAChR-dependent depression on inhibitory transmission in the BLA of sham rats and it only partially reverts seizure-induced impairment of mAChR-dependent depression on inhibitory transmission in the BLA.

(A) Time course graph showing the effect of mAChR activation on eIPSC amplitude on slices from sham rats, treated with vehicle or KML29. (B) Bar graphs quantifying the percentage inhibition of evoked IPSC amplitude by CCh (10-20 min post-application) showed that CCh produces a long-term suppression of evoked IPSC amplitude, which is significantly attenuated in the presence of KML29. (C) KML29 treatment is not sufficient to

fully restore mAChR-LTD in neurons recorded from kindled rats. (**D**) Bar graph quantifying the percentage inhibition of evoked IPSC amplitude by CCh showed no significant difference in mACh-induced LTD in slices from kindled rats treated with either vehicle or KML29. Scale bars, 200 pA, 40 ms. Post hoc sham aCSF versus sham KML29 p-values: \*\*\*p < 0.001. Data are shown as mean  $\pm$  SEM. Sham aCSF data are repeated in the graphs as they belong to the same experimental design.

### Fig.6. 2-AG synthesis inhibition does not alter GABA transmission in the BLA.

The effect of DO34 on sIPSC frequency and amplitude on slices from sham (A) and kindled rats (B) is shown. DO34 did not change either the frequency or the amplitude of sIPSCs, ruling out the possibility that 2-AG signaling tonically constrains GABA transmission in the BLA. Effect of AM251 ( $2\mu$ M) on sIPSC frequency and amplitude in sham (C) and kindled (D) rats. AM251 was bath applied after 5 min baseline recording and sIPSCs were assessed 10-15 min after AM251 application. Scale bars: 50 pA, 1 s.

# Fig.7. No effects of repeated seizures on CB1 receptor sensitivity.

(A) CP55,940 (CP55;  $3\mu$ M), a specific CB1 receptor agonist, induced a similar depression of eIPSCs in the BLA in both sham and kindled rats. (B) The effect of CB1 receptor activation on inhibitory transmission was assessed by measuring the average eIPSC amplitude normalized to the BL obtained 10-15 min after CP55,940 application in both sham and kindled rats. Scale bars, 200 pA, 40 ms. Post hoc versus BL p-values are shown as follows: \*\*\*p < 0.001, \*\*p < 0.01. Data are shown as mean ± SEM.

# Fig.8. Enhanced 2-AG degradation does not account for eCB signaling alteration in kindled rats.

The effect of KML29, an inhibitor of 2-AG degradation, on sIPSC frequency and amplitude on slices from sham (A) and kindled (B) rats is shown. KML29 reduced sIPSC frequency but not amplitude in sham rats, while it had no effect on either the frequency or the amplitude of sIPSCs in kindled rats. Scale bars: 50 pA, 1 s. sham aCSF versus sham KML29; \*\*p < 0.001.

Data are shown as mean  $\pm$  SEM. (C) Seizures did not change V<sub>max</sub> but significantly increased K<sub>m</sub> of MAGL enzyme in the amygdala, which is indicative of reduced binding affinity without alteration of hydrolytic activity of the enzyme for its substrate. sham versus kindled; \*p < 0.05. Data are expressed as mean  $\pm$  SEM (n = 5–6 per group).







50

0

BL-

-ISO

recovery





# **JNeurosci Accepted Manuscript**





0

BL-

-ISO

recovery-











Α

O CTRL aCSF O CTRL AM251





В

D



elPSCs amp (%) 22 20 50 52 25-BL CCh (10µM) post CCh Time (min)

○ sham aCSF ○ sham DO34



















В

# **JNeurosci Accepted Manuscript**







