



Universiteit  
Leiden  
The Netherlands

## Identifying and appraising patient-reported outcome measures on treatment satisfaction in acne: a systematic review

Zuuren, E.J. van; Arents, B.W.M.; Miklas, M.; Schoones, J.W.; Tan, J.

### Citation

Zuuren, E. J. van, Arents, B. W. M., Miklas, M., Schoones, J. W., & Tan, J. (2020). Identifying and appraising patient-reported outcome measures on treatment satisfaction in acne: a systematic review. *British Journal Of Dermatology*, 185(1), 36-51.  
doi:10.1111/bjd.19675

Version: Publisher's Version  
License: [Creative Commons CC BY-NC 4.0 license](#)  
Downloaded from: <https://hdl.handle.net/1887/3280086>

**Note:** To cite this publication please use the final published version (if applicable).

# Identifying and appraising patient-reported outcome measures on treatment satisfaction in acne: a systematic review\*

E.J. van Zuuren <sup>1</sup> B.W.M. Arents <sup>2</sup> M. Miklas,<sup>3</sup> J.W. Schoones<sup>4</sup> and J. Tan <sup>3,5</sup>

<sup>1</sup>Dermatology Department and <sup>4</sup>Walaes Library, Leiden University Medical Centre, Leiden, the Netherlands

<sup>2</sup>Skin Patients Netherlands (Huidpatiënten Nederland), Nieuwegein, the Netherlands

<sup>3</sup>Windsor Clinical Research Inc., Windsor, ON, Canada

<sup>5</sup>Western University, London, ON, Canada

**Linked Comment:** J. Kottner and J. Schmitt. *Br J Dermatol* 2021; **185**:3–4.

## Abstract

### Correspondence

Esther J. van Zuuren.

Email: e.j.van\_zuuren@lumc.nl

### Accepted for publication

9 November 2020

### Funding sources

None.

### Conflicts of interest

J.T. has been an advisor, consultant, speaker and/or investigator for Almirall, Bausch Health, Boots/Walgreens, Botanix, Cipher, Galderma, L'Oreal and Sun Pharma. E.J.v.Z., B.W.M.A., M.M. and J.W.S. declare they have no conflicts of interest.

E.J.v.Z. and B.W.M.A. contributed equally to this work.

\*Plain language summary available online

DOI 10.1111/bjd.19675

**Background** After dermatitis, acne is the next skin disease to contribute most to the burden of skin diseases worldwide. Recently, seven core outcome domains have been identified, which together form an Acne Core Outcome Set (ACORN). One of these was satisfaction with acne treatment.

**Objectives** To identify studies that described the development of patient-reported outcome measures (PROMS), evaluated one or more measurement properties of a PROM, or evaluated the interpretability of a PROM in patients with acne regarding treatment satisfaction.

**Methods** The Consensus-based Standards for the selection of health Measurement INstruments (COSMIN) search strategy for identifying PROMS on acne treatment satisfaction was used. We searched PubMed, MEDLINE, Embase, LILACS, Web of Science, Cochrane Library, Emcare, PsycINFO and Academic Search premier (June 2020). Study selection, data extraction and assessment of methodological quality according to COSMIN guidance were carried out independently by two authors.

**Results** Only one study could be included, describing the development of a treatment satisfaction measure in patients with acne. The development was assessed as inadequate and data on measurement properties were lacking. Additionally, we found 188 studies reporting treatment satisfaction solely as an outcome, using a wide variety of methods, none of them standardized or validated.

**Conclusions** We could not find a PROM on treatment satisfaction to recommend for a core outcome set in acne. There is an unmet need for a PROM on treatment satisfaction in acne that is robustly developed, designed and validated.

### What is already known about this topic?

- Core outcome sets are consensus-based minimum outcome measures that should be reported in clinical trials of a specific disease or target condition.
- The Acne Core Outcomes Research Network identified the following domains important for acne: satisfaction with appearance; extent of dark marks and scars; long-term acne control; signs and symptoms; satisfaction with treatment; health-related quality of life; and adverse events.

**What does this study add?**

- We could not find a PROM on treatment satisfaction that can be recommended for a core outcome set in acne.
- Many studies reported treatment satisfaction as an outcome, with a wide variety of methods, none of them standardized or validated.
- There is an unmet need for a PROM measuring treatment satisfaction in acne that is robustly developed and validated according to COSMIN standards.

Core outcome sets are consensus-based minimum outcome measures that should be reported in clinical trials of a specific disease or target condition.<sup>1</sup> Outcomes can be conceived as constructs or domains, credibly established by patients and other relevant stakeholders, reflecting what is to be measured in trial participants to evaluate the effect of an intervention. Outcome measures are instruments or tools to measure the quality or quantity of the intervention on each of the domains. These measures can range from clinical examinations to patient responses to questionnaires to laboratory findings and to imaging studies.<sup>2</sup> Outcome measures to be included in a core outcome set are based on a systematic search, assessment of quality and recommendations to a voting panel for inclusion of only one outcome measurement for each domain where possible. In the absence of an adequate measure, a need for development would be established.<sup>3</sup>

In clinical trials investigating acne treatments, multiple outcome measures have been used with no established standards.<sup>4,5</sup> This multiplicity, heterogeneity and lack of quality have been problematic in data synthesis by impeding comparative outcomes research and contributing to resource wastage. An international consensus on a core outcome set for clinical trials in acne could address this unmet need by standardizing and harmonizing existing outcome measures, and identifying those that might be lacking.

Development of an acne core outcome set for acne clinical trials was initiated with the Acne Core Outcomes Research Network (ACORN), initially funded by a US National Institutes of Health/National Institute of Arthritis and Musculoskeletal and Skin Diseases grant (1U01AR065109-01). In a landmark study involving stakeholders worldwide, including 307 patients or their parents, 218 healthcare professionals, 45 nonclinical researchers, 17 industry employees and nine journal editors, the most important domains for an acne core outcome set were identified.<sup>6</sup> These included satisfaction with appearance, extent of dark marks and scars, long-term acne control, signs and symptoms, satisfaction with treatment, health-related quality of life and adverse events (AEs).<sup>6</sup>

A hierarchy of treatment satisfaction can extend from procedures, therapies, activity limitations and dietary restrictions, along with included medications. A more circumscribed concept of satisfaction with intervention or medication impacting on disease signs/symptoms and potential side-effects is relevant to clinical trials.<sup>7</sup>

In this study, we focused on addressing treatment satisfaction in acne clinical trials, more specifically on satisfaction

with any intervention or with the (general) care received. Our aims were to identify patient-reported outcome measures (PROMs) of satisfaction with acne treatments and to evaluate their quality with Consensus-based Standards for the selection of health Measurement INstruments (COSMIN) methodology.<sup>3</sup>

**Materials and methods**

This systematic review conformed to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.<sup>8</sup> A prespecified protocol was submitted on 29 October 2019 to the International Prospective Register of Systematic Reviews (PROSPERO) and was registered on 18 May 2020 (CRD42020156473). Owing to the delay with PROSPERO, the protocol was also published at the Open Science Framework (www.osf.io) on 2 February 2020. We followed the COSMIN guideline for systematic reviews of PROMs.<sup>9</sup>

**Eligible studies**

Any study of people with acne vulgaris that described the development of a PROM, evaluated one or more measurement properties of a PROM, or evaluated the interpretability of a PROM regarding treatment satisfaction was eligible. Treatment satisfaction was interpreted as patients' satisfaction with any intervention or with the (general) care received. Studies that solely used a PROM on treatment satisfaction as part of, for example, assessing the outcomes of an intervention would be excluded.<sup>9</sup> However, the latter were tabulated separately to create an overview of instruments used to measure treatment satisfaction. Studies on acne conglobata, rosacea and hidradenitis suppurativa were excluded.

**Literature search**

A systematic literature search was conducted on 1 November 2019. The following bibliographic databases were searched from their inception for reports of outcome measures assessing satisfaction with treatment of acne: PubMed, MEDLINE (OVID), Embase (OVID), LILACS, Web of Science, COCHRANE Library, Emcare (OVID), PsycINFO (EbscoHOST) and Academic Search Premier (EbscoHOST). There were no language restrictions. The COSMIN search strategy for identifying all PROMs on satisfaction with acne treatment was used (Appendix S1; see Supporting Information), and contained the

following: (i) construct – comprehensive search terms regarding treatment satisfaction; and (ii) population – comprehensive search terms regarding people with acne; and (iii) type of instrument – COSMIN PROM filter NOT COSMIN exclusion filter, as described by Terwee *et al.*<sup>10</sup>

A rerun of the search was conducted on 10 June 2020. Search results were uploaded into RAYYAN to facilitate selection of potentially eligible studies (<http://rayyan.qcri.org/>). Two authors (E.J.v.Z. and B.W.M.A.) independently assessed the eligibility of the studies based on title, abstract and keywords. Records were only excluded when there was sufficient information to support exclusion. From all the other records, also those lacking data, full-text copies were obtained. These full-text papers were independently assessed for eligibility by two authors, and the references of eligible studies were independently checked for additional studies. Any disagreement was resolved by discussion and consensus. As recommended by Prinsen *et al.*,<sup>9</sup> studies that only used the PROM as an outcome measurement instrument were excluded based on the full text. These were as prespecified in the protocol, tabulated by collecting study details and outcomes using a predetermined form designed for this purpose and serving as supportive information for the ACORN project.

### Data extraction and methodological quality of included studies

For data extraction of the included studies on PROM development, validation and evaluation, predefined COSMIN forms were used, as per the methodology.<sup>3</sup> The authors only included data if there was an independently attained consensus. The methodological quality of the included studies on PROM development and validation was assessed using the COSMIN Risk of Bias checklist.<sup>11</sup> The COSMIN Risk of Bias checklist consists of 10 boxes, each with multiple items that can be scored as very good, adequate, doubtful, inadequate or not applicable.<sup>9</sup> The lowest rating of any standard determines the overall quality of the PROM. Box 1 addresses PROM development, while the other nine address measurement properties.<sup>9,11</sup> The overall assessment of included PROMs was performed according to the COSMIN guideline for systematic reviews of PROMs.<sup>3,9</sup>

## Results

### Search results


The search identified 705 records (after removal of duplicates) for which abstracts were screened. A total of 451 references were excluded. Of the remaining 254, the full texts were obtained. Screening of the reference lists of these papers resulted in one additional study, making the total 255. Of these 254 were excluded.<sup>12–265</sup> Reasons for exclusion were that treatment satisfaction was not, in fact, an outcome ( $n = 56$ ),<sup>12–67</sup> only using a treatment satisfaction instrument for acne treatments ( $n = 91$ )<sup>68–158</sup> or for acne scar treatment ( $n = 97$ ),<sup>159–255</sup> or a report on the same study group ( $n = 10$ ).<sup>256–265</sup> Therefore, just one study met the inclusion criteria (see Figure 1).<sup>266</sup>

### Description of the included study

The study of Alomar *et al.* was published in 2004 and described the development of a treatment satisfaction questionnaire in patients with acne.<sup>266</sup> The authors initiated the process by reviewing the literature in MEDLINE (1980–2002), with the objective of deriving a preliminary list of aspects considered important regarding patient satisfaction with treatment in general and specifically in dermatology. Subsequently, an exploratory method with a moderator-led focus group was used to discuss aspects from the literature (semi-structured script), such as degree of knowledge of acne and acne treatments; psychological impact of acne; effects on physical appearance; relationships; information and expectations of treatments; comfort and ease of application or intake; associated AEs; and overall satisfaction with treatment. The focus group consisted of just six women, of whom four were middle-aged and two were adolescents; all were using isotretinoin or had used isotretinoin. After the 2-h focus group session, all gathered information was used for the development of an 11-item questionnaire. More details on this study are provided in Tables S1 and S2 (see Supporting Information). The items referred to satisfaction of various aspects related to treatment such as improvement of symptoms, satisfaction with provided information, satisfaction with treatment, mood, social life, AEs, route of administration, daily activities and the treatment in general. Responses were presented as 4- or 5-point Likert scales. The overall scoring system was not clearly explained. The study authors concluded by stating that a validation study to evaluate the measurement properties still had to be completed. However, the authors confirmed that such a study had not been completed nor published. Furthermore, we found no evidence that this questionnaire was validated or used in other studies.

The methodological quality of this study was assessed with the COSMIN Risk of Bias checklist,<sup>11</sup> and independently conducted by two authors (E.J.v.Z. and B.W.M.A.). As the study of Alomar *et al.* only described development of the questionnaire we were only able to complete the first part of box 1 of that checklist,<sup>266</sup> which addresses general design requirements and concept elicitation. The second part assesses the comprehensiveness and comprehensibility of the questionnaire; however, this was not reported in the study (see Table 1).

A key criterion is whether a PROM is developed in a sample of patients representing the target population for which the PROM is intended.<sup>267</sup> The intention of Alomar *et al.* was to develop a treatment satisfaction questionnaire for patients with acne, which implies patients of both sexes, all ages, all treatment modalities and encompassing those with a spectrum of acne severity.<sup>266</sup> However, only six women were included, with apparently more severe acne because they used or had used isotretinoin. Therefore, this sample was not representative of the target population of people with acne. According to COSMIN standards, this PROM development was considered to be inadequate (see Table 1). As the study of Alomar *et al.* contained no data on any further validation, we were unable to complete boxes 2–10 of the COSMIN Risk of Bias



**Figure 1** Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram. n.a., not available; PROM, patient-reported outcome measure.

checklist.<sup>266</sup> This shortcoming could not be supplemented with data from other validation studies regarding this PROM.

### Evaluation of the PROM

Following COSMIN guidance, we reviewed this PROM against the 10 criteria for good content validity and applied the modified GRADE (Grading of Recommendations, Assessment, Development and Evaluation) approach regarding the quality of the supporting evidence (high, moderate, low or very low).<sup>9</sup> Based on that, we assessed the content validity of this PROM as insufficient [very low quality evidence; Table S3 (see Supporting Information) and Table 1]. For the other measurement

properties, we would have provided summary of findings tables per measurement property if more studies were available. As this was not the case, we opted to summarize all the ratings and the quality of the available evidence in one table, based on the spreadsheet provided by COSMIN (see Table 2). Owing to the lack of studies and data we were unable to describe the interpretability and feasibility of the PROM.

The overall results of our findings were that, per the COSMIN definitions, this PROM was categorized as ‘category B’: there is no sufficient evidence for content validity, but high-quality evidence for an insufficient measurement property is not available either. This means that this PROM may have a potential to be recommended, but further validation studies are needed.

**Table 1** Quality of the patient-reported outcome measure (PROM) development

PROM	PROM design			Concept elicitation <sup>b</sup>		Total PROM design	CI study <sup>a</sup>	Comprehensibility	Comprehensiveness	Total CI study	Total PROM development
Treatment satisfaction in acne	General design requirements						General design requirements				
	Clear construct	Clear origin of construct	Clear target population for which the PROM was developed	Clear context of use	PROM developed in sample representing the target population		CI study performed in sample representing the target population				
Alomar <i>et al.</i> (2004) <sup>266</sup>	A	D	A	A	I					I	I

CI, cognitive interview; A, adequate; D, doubtful; I, inadequate. <sup>a</sup>Empty cells indicate that a CI study (or part of it) was not performed. <sup>b</sup>When the PROM was not developed in a sample representing the target population, the concept elicitation was not further rated.

## Studies that only reported treatment satisfaction as an outcome

Studies that only used the PROM as an outcome measurement instrument were excluded, but study details were tabulated, per protocol. We identified 188 studies that reported treatment satisfaction as an outcome: 91 on treatment of acne<sup>68–158</sup> and 97 on treating acne scars (a possible sequela of acne).<sup>159–255</sup> These numbers show that clinicians and researchers do consider treatment satisfaction as an important outcome. However, the methods used clearly demonstrate the current diversity (Figures 2 and 3). A summary of the studies is supplied in Table 3 and characteristics of the studies in Tables S4 and S5 (see Supporting Information).

## Discussion

In dermatology, various collaborative groups have been established to facilitate the development of core outcome sets, including the International Dermatology Outcome Measures group (IDEOM)<sup>268</sup> and the Cochrane Skin – Core Outcome Set Initiative (CS-COUSIN).<sup>269</sup> Disease-specific efforts have been initiated in atopic dermatitis, psoriasis and hidradenitis suppurativa.<sup>270–272</sup> That such an elaborate worldwide undertaking has an effect was shown with the Harmonising Outcome Measures for Eczema (HOME) initiative for atopic dermatitis, as more trials adhered to its recommendations.<sup>273</sup>

For acne this has been the remit of ACORN.<sup>274</sup> As part of that initiative, this systematic review was undertaken to identify and evaluate the quality of outcome measures relevant to one of the seven previously identified core domains<sup>6</sup> – treatment satisfaction. The methodology we followed was established by COSMIN.<sup>3,9,11</sup> Briefly, the steps involved were a search for existing instruments (from systematic reviews, literature searches and other sources) and quality assessment of the instruments found. This would then lead to recommendations on selection and a consensus procedure for final agreement on an outcome measure – in this case treatment satisfaction.

Treatment satisfaction has previously been defined as a patient-assessed domain addressing attributes of the process and outcome of the treatment experience.<sup>275</sup> This would be inclusive of benefits in relief of signs and symptoms and risk of AEs, inconvenience and cost. In clinical trials, cost of treatment would be excluded as medications are typically provided to patients, therefore not reflecting real-world experience. Furthermore, in preapproval trials, cost of treatment may not have yet been established by the manufacturer. Regulatory authorities are increasingly recognizing the importance of these patient-reported measures on aspects of treatment and disease impact that are not readily accessible or evaluable by other methodologies.

From our literature search, numerous studies were found to have treatment satisfaction of acne and of acne scars as a patient-reported outcome. These findings underline the importance of treatment satisfaction in research and support the




Figure 2 Types of tools to measure satisfaction with acne treatment. NRS, numerical rating scale; VAS, visual analogue scale.




Figure 3 Types of tools to measure satisfaction with acne scar treatment. NRS, numerical rating scale; VAS, visual analogue scale.



**Table 2** Quality of the evidence for measurement properties of the patient-reported outcome measure (treatment satisfaction in acne)

	Overall rating	Quality of evidence High, moderate, low, very low
PROM is not named	+/-/ ?	low
Content validity	–	Very low
Relevance	–	Very low
Comprehensiveness	–	Very low
Comprehensibility	–	Very low
Structural validity		
Internal consistency		
Cross-cultural validity		
Measurement invariance		
Reliability		
Measurement error		
Criterion validity		
Construct validity		
Responsiveness		
Empty cells indicate that there was no evidence available to assess. (+), sufficient; (?), Indeterminate; (–), insufficient.		

previous consensus involving patients with acne and other relevant stakeholders in identifying it as an important core outcome domain.<sup>6</sup> This was also reflected by the increase in studies evaluating treatment satisfaction over the years (Figure 4). However, while investigators of these studies considered treatment satisfaction important and reported data for this outcome, none used a standardized, validated instrument – underlining the need for harmonization. Furthermore, we found instances where the degree of improvement was interpreted as a degree of treatment satisfaction, despite these constructs not being identical.

We found only one study that addressed the development of a PROM for treatment satisfaction in acne, but it was assessed as not yet to be recommended by COSMIN standards (category B).<sup>266</sup> This means that there is still an unmet need for a well-developed and validated PROM for treatment satisfaction in acne. PROM development and validation is not a trivial endeavour. Guidance from academia and regulatory authorities are available to facilitate the development of such a PROM.<sup>276–278</sup> Development and validation should meet COSMIN standards, especially as it is intended to be part of a core outcome set to be used in clinical trials. In its absence, we observed that researchers are left to develop ad hoc, nonstandardized, unvalidated measures for evaluation of a domain they consider important in their clinical trials. Although it reflects the researchers' involvement and creativity, it was this diversity and variety of measures that impeded comparison of outcomes on treatment satisfaction in acne between studies or the ability to conduct a systematic review with quantitative meta-analyses on this important outcome.

The limitations are that our search strategy would not have detected studies that do not contain the construct treatment

**Table 3** Characteristics of studies measuring patient-reported treatment satisfaction in acne

	Treatment of acne (n = 91)	Treatment of acne of scars (n = 97)	Total (n = 188)
<b>Participants</b>			
Total	32 811	3791	36 602
Men	10 917 (33.3)	903 (23.8)	11 820 (32.3)
Women	15 664 (47.7)	1774 (46.8)	17 438 (47.6)
Sex unknown/not reported	6230 (19.0)	1114 (29.4)	7344 (20.1)
Mean age (years)	22.7	33.0	
Minimum in a study (n)	10	3	
Maximum in a study (n)	5131	352	
Mean per study (n)	364.6	39.1	
Median per study (n)	62	25	
<b>Location</b>			
Asia	20 (22)	27 (28)	47 (25.0)
Australia	–	2 (2)	2 (1.1)
Europe	25 (27)	15 (15)	40 (21.3)
Latin America	4 (4)	7 (7)	11 (5.9)
Middle East	12 (13)	17 (17)	29 (15.4)
Multiple countries	9 (10)	6 (6)	15 (8.0)
North America	20 (22)	23 (24)	43 (22.9)
South Africa	1 (1)	–	1 (0.5)
<b>Study design (%)</b>			
Cross-sectional	11 (12)	1 (1)	12 (6.4)
Open label	29 (32)	56 (58)	85 (45.2)
RCT	43 (47)	24 (25)	67 (35.6)
Other	8 (9)	16 (16)	24 (12.8)
<b>Study duration (weeks)</b>			
Minimum (n)	2	4	
Maximum (n)	156	208	
Mean (n)	18.6	33.5	
Median (n)	12	26	
<b>Measurement methods</b>			
One question	46 (51)	58 (60)	104 (55.3)
Questionnaire	35 (39)	27 (28)	62 (33.0)
Unclear/not reported	10 (11)	12 (12)	22 (11.7)
<b>Answer options</b>			
NRS/VAS	19 (21)	18 (19)	37 (19.7)

(continued)



Table 3 (continued)

	Treatment of acne (n = 91)	Treatment of acne of scars (n = 97)	Total (n = 188)
3-point Likert	1 (1)	5 (5)	6 (3.2)
4-point Likert	16 (18)	22 (23)	38 (20.2)
5-point Likert	21 (23)	27 (28)	48 (25.5)
6-point Likert	2 (2)	1 (1)	3 (1.6)
7-point Likert	1 (1)	–	1 (0.5)
Other	4 (4)	1 (1)	5 (2.7)
Unclear/not reported	27 (30)	23 (24)	50 (26.6)

Data are n (%) unless otherwise indicated. RCT, randomized controlled trial; NRS, numeric rating scale; VAS, visual analogue scale.

satisfaction and associated terms in the meta-data of the consulted databases. This could have underestimated the number of trials evaluating treatment satisfaction in acne. The strengths of our review were the comprehensive search in multiple databases for potentially eligible studies without language restriction, ensuring no risk of language bias. Furthermore, although only one study could be included (in Spanish), we compiled a comprehensive list of measures of patient treatment satisfaction in acne research. As such, this review could serve as a resource for developing a PROM on treatment satisfaction in acne.

This study was undertaken to identify and assess the quality of existing measures corresponding to a core domain in a core outcome set for acne: treatment satisfaction. However, in our search and subsequent assessments, we were unable to find a single measure that fulfilled the necessary criteria. We did find a plethora of ad hoc scales addressing treatment satisfaction in acne and acne scarring, none standardized or validated. This represents an unmet need for a PROM measuring treatment satisfaction in acne that is robustly developed and validated according to COSMIN standards.




Figure 4 Number of studies reporting treatment satisfaction over the years.

## References

- Williamson PR, Altman DG, Blazeby JM *et al.* Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012; **13**:132.
- COSMIN. I'm developing a core outcome set (COS). Available at: <https://www.cosmin.nl/finding-right-tool/developing-core-outcome-set/> (last accessed 30 June 2020).
- Mokkink LB, Prinsen CA, Patrick DL *et al.* COSMIN methodology for systematic reviews of patient-reported outcome measures (PROMs). Available at: [https://www.cosmin.nl/wp-content/uploads/COSMIN-syst-review-for-PROMs-manual\\_version-1\\_feb-2018.pdf](https://www.cosmin.nl/wp-content/uploads/COSMIN-syst-review-for-PROMs-manual_version-1_feb-2018.pdf) (last accessed 30 June 2020).
- Lehmann HP, Robinson KA, Andrews JS *et al.* Acne therapy: a methodologic review. *J Am Acad Dermatol* 2002; **47**:231–40.
- Zarchi K, Jemec GBE. Severity assessment and outcome measures in acne vulgaris. *Curr Derm Rep* 2012; **1**:131–6.
- Layton AM, Eady EA, Thiboutot DM, Tan J. Identifying what to measure in acne clinical trials: first steps towards development of a core outcome set. *J Invest Dermatol* 2017; **137**:1784–6.
- Shikhar R, Rentz AM. Satisfaction with medication: an overview of conceptual, methodologic, and regulatory issues. *Value Health* 2004; **7**:204–15.
- Liberati A, Altman DG, Tetzlaff J *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009; **339**:b2700.
- Prinsen CAC, Mokkink LB, Bouter LM *et al.* COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018; **27**:1147–57.
- Terwee CB, Jansma EP, Riphagen II, de Vet HC. Development of a methodological PubMed search filter for finding studies on measurement properties of measurement instruments. *Qual Life Res* 2009; **18**:1115–23.
- Mokkink LB, de Vet HCW, Prinsen CAC *et al.* COSMIN Risk of Bias checklist for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018; **27**:1171–9.
- Anonymous. Acne treatments come out a wash. *Consum Rep* 2012; **77**:9.
- Anonymous. Patient perspectives: treating acne with isotretinoin. *Pediatr Dermatol* 2015; **32**:e324–e325.
- Al-Talib H, Hameed A, Al-Khateeb A, Murugaiah C. Efficacy and safety of superficial chemical peeling in treatment of active acne vulgaris. *An Bras Dermatol* 2017; **92**:212–6.
- Amichae B. Long-term mini-doses of isotretinoin in the treatment of relapsing acne. *J Dermatol* 2003; **30**:572.
- Barolet D, Boucher A. Radiant near infrared light emitting diode exposure as skin preparation to enhance photodynamic therapy inflammatory type acne treatment outcome. *Lasers Surg Med* 2010; **42**:171–8.
- Berson DS, Chalker DK, Harper JC *et al.* Current concepts in the treatment of acne: report from a clinical roundtable. *Cutis* 2003; **72** (Suppl.):5–13.
- Blome C, Augustin M, Behechtnejad J, Rustenbach SJ. Dimensions of patient needs in dermatology: subscales of the patient benefit index. *Arch Dermatol Res* 2011; **303**:11–7.
- Boehncke W-H, Ochsendorf F, Paeslack I *et al.* Decorative cosmetics improve the quality of life in patients with disfiguring skin diseases. *Eur J Dermatol* 2002; **12**:577–80.
- Bogle MA, Dover JS, Arnd KA, Mordon S. Evaluation of the 1,540-nm erbium:glass laser in the treatment of inflammatory facial acne. *Dermatol Surg* 2007; **33**:810–7.
- Cho SB, Lee SJ, Kang JN *et al.* Combined fractional laser treatment with 1550-nm erbium glass and 10 600-nm carbon dioxide lasers. *J Dermatolog Treat* 2010; **21**:221–8.
- Cotellessa C, Manunta T, Ghersetich I *et al.* The use of pyruvic acid in the treatment of acne. *J Eur Acad Dermatol Venerol* 2004; **18**:275–8.
- Cunliff WJ, van de Kerkof PC, Caputo R *et al.* Roaccutane treatment guidelines: results of an international survey. *Dermatology* 1997; **194**:351–7.
- Draeos Z, Hornby S, Walters R *et al.* Hydrophobically modified polymers can minimize skin irritation potential caused by surfactant-based cleansers. *J Cosmet Dermatol* 2013; **12**:314–21.
- Draeos Z. Evaluating consumer dermatologic devices. *J Cosmet Dermatol* 2009; **8**:75–6.
- Dreno B. Assessing quality of life in patients with acne vulgaris – implications for treatment. *Am J Clin Dermatol* 2006; **7**:99–106.
- Dunlap FE, Mills OH, Tuley MR *et al.* Adapalene 0.1% gel for the treatment of acne vulgaris: its superiority compared to tretinoin 0.025% cream in skin tolerance and patient preference. *Br J Dermatol* 1998; **139** (Suppl.):17–22.
- Elman M, Lebzelter J. Light therapy in the treatment of acne vulgaris. *Dermatol Surg* 2004; **30**:139–46.
- Fabbrocini G, Donnarumma M, Russo G, Monfrecola G. Informative flashcards to increase therapy compliance in acne patients receiving adapalene 0.1%/benzoyl peroxide 2.5%. *Esper Dermatol* 2016; **18**:178–82.
- Fabbrocini G, Cacciapuoti S, Fardella N *et al.* CROSS technique: chemical reconstruction of skin scars method. *Dermatol Ther* 2008; **21** (Suppl.):S29–32.
- Fabbrocini G, Fardella N, Proietti I, Innoncenzi D. Acne scarring treatment using skin needling. *Clin Exp Dermatol* 2009; **34**:874–9.
- Fagundes DS, Fraser JM, Klauda HC. New therapy update – a unique combination formulation in the treatment of inflammatory acne. *Cutis* 2003; **72** (Suppl.):16–9.
- Feldman SR, Fried RG, Herndon JH *et al.* Digital videography assessment of patients experiences using adapalene-benzoyl peroxide gel in the treatment of acne vulgaris. *J Drugs Dermatol* 2012; **11**:919–25.
- Fenton C, Wellington K, Moen MD, Robinson DM. Drospirenone/ethinylestradiol 3mg/20 µg (24/4 day regimen) – a review of its use in contraception, premenstrual dysphoric disorder and moderate acne vulgaris. *Drugs* 2007; **67**:1749–65.
- Fulton JE Jr. Dermabrasion, chemabrasion, and laserabrasion. Historical perspectives, modern dermabrasion techniques, and future trends. *Dermatol Surg* 1996; **22**:619–28.
- Gold MH, Heath AD, Biron JA. Clinical evaluation of the Smart-Skin fractional laser for the treatment of photodamage and acne scars. *J Drugs Dermatol* 2009; **8** (Suppl.):S4–8.
- Gonzalez MJ, Sturgill WH, Ross EV, Uebelhoefer NS. Treatment of acne scars using the plasma skin regeneration (PSR) system. *Lasers Surg Med* 2008; **40**:124–7.
- Hayashi N, Kawashima M. Efficacy of oral antibiotics on acne vulgaris and their effects on quality of life: a multicenter randomized controlled trial using minocycline, roxithromycin and faropenem. *J Dermatol* 2011; **38**:111–9.
- Hesseler MJ, Shyam N. Platelet-rich plasma and its utility in the treatment of acne scars: a systematic review. *J Am Acad Dermatol* 2019; **80**:1730–45.
- Jordan R, Cummins C, Burls A. Laser resurfacing of the skin for the improvement of facial acne scarring: a systematic review of the evidence. *Br J Dermatol* 2000; **142**:413–23.
- Kainz JT, Berghammer G, Auer-Grumbach P *et al.* Azelaic acid 20 % cream: effects on quality of life and disease severity in adult female acne patients. *J Dtsch Dermatol Ges* 2016; **14**:1249–59.

- 42 Katz B. Efficacy of a new fractional CO<sub>2</sub> laser in the treatment of photodamage and acne scarring. *Dermatol Ther* 2010; **23**:403–6.
- 43 Khunger N, Bhardwaj D, Khunger M. Evaluation of CROSS technique with 100% TCA in the management of ice pick acne scars in darker skin types. *J Cosmet Dermatol* 2011; **10**:51–7.
- 44 Kilmer SL, Chotzen V, Zelickson BD *et al.* Full-face laser resurfacing using a supplemented topical anesthesia protocol. *Arch Dermatol* 2003; **139**:1279–83.
- 45 Kim K-H, Lee S-C, Park Y-B, Park Y-J. Cardiff acne disability index: cross-cultural translation in Korean and its relationship with clinical acne severity, pathological patterns, and general quality of life. *J Tradit Chin Med* 2017; **37**:702–9.
- 46 Klassen AF, Lipner S, O'Malley M *et al.* Development of a new patient-reported outcome measure to evaluate treatments for acne and acne scarring: the ACNE-Q. *Br J Dermatol* 2019; **181**:1121–2.
- 47 Korkut C, Piskin S. Benzoyl peroxide, adapalene, and their combination in the treatment of acne vulgaris. *Dermatol* 2005; **32**:169–73.
- 48 Layton AM. Acne scarring – reviewing the need for early treatment of acne. *J Dermatolog Treat* 2000; **11**:3–6.
- 49 Lowe NJ, Lowe PL, St Clair Roberts J. A phase IIa open-label dose-escalation pilot study using allogeneic human dermal fibroblasts for nasolabial folds. *Dermatol Surg* 2010; **36**:1578–85.
- 50 Magin PJ, Adams J, Heading GS *et al.* Complementary and alternative medicine therapies in acne, psoriasis, and atopic eczema: results of a qualitative study of patients' experiences and perceptions. *J Altern Complement Med* 2006; **12**:451–7.
- 51 Malhi HK, Tu J, Riley TV *et al.* Tea tree oil gel for mild to moderate acne; a 12 week uncontrolled, open-label phase II pilot study. *Australas J Dermatol* 2017; **58**:205–10.
- 52 Mariwalla K, Rohrer TE. Use of lasers and light-based therapies for treatment of acne vulgaris. *Lasers Surg Med* 2005; **37**:333–42.
- 53 McMullen E, Cox NH. The British Association of Dermatologists isotretinoin questionnaire for patients: a useful clinical tool. *Clin Exp Dermatol* 2006; **31**:713–4.
- 54 Miglani A, Manchanda RK. Observational study of Arctium lappa in the treatment of acne vulgaris. *Homeopathy* 2014; **103**:203–7.
- 55 Miyachi Y, Hayashi N, Furukawa F *et al.* Acne management in Japan: study of patient adherence. *Dermatology* 2011; **223**:174–81.
- 56 Newton JN, Malon E, Klassen A *et al.* The effectiveness of acne treatment: an assessment by patients of the outcome of therapy. *Br J Dermatol* 1997; **137**:563–7.
- 57 Ng QX, Koh SSH, Shin D *et al.* Use of polymethylmethacrylate (PMMA) microspheres collagen to treat atrophic acne scars. *Med Hypotheses* 2017; **108**:115–6.
- 58 Orringer JS, Kang S, Maier L *et al.* A randomized, controlled, split-face clinical trial of 1320-nm Nd:YAG laser therapy in the treatment of acne vulgaris. *J Am Acad Dermatol* 2007; **56**:432–8.
- 59 Petkova VB, Hussain S, Filinada D, Karamisheva V. Acne in Bulgaria and Greece: comparison study of patient adherence. *Acta Pol Pharm* 2019; **76**:913–6.
- 60 Raulin C, Grema H. Single-pass carbon dioxide laser skin resurfacing combined with cold-air cooling: efficacy and patient satisfaction of a prospective side-by-side study. *Arch Dermatol* 2004; **140**:1333–6.
- 61 Robertson KM. Acne vulgaris. *Facial Plast Surg Clin N Am* 2004; **12**:347–55.
- 62 Sadick NS. Handheld LED array device in the treatment of acne vulgaris. *J Drugs Dermatol* 2008; **7**:347–50.
- 63 Shahmoradi Z, Iraj F, Siadat AH, Ghorbaini A. Comparison of topical 5% nicotinamid gel versus 2% clindamycin gel in the treatment of the mild-moderate acne vulgaris: a double-blinded randomized clinical trial. *J Res Med Sci* 2013; **18**:115–7.
- 64 Snyder S, Crandell I, Davis SA, Feldman SR. Medical adherence to acne therapy: a systematic review. *Am J Clin Dermatol* 2014; **15**:87–94.
- 65 Tanghetti E, Mirkov M, Sierra RA. Delivery of light to the skin through ablated conduits. *Lasers Surg Med* 2017; **49**:69–77.
- 66 Wanitphakdeedecha R, Manuskiatti W, Siriphukpong S, Chen TM. Treatment of punched-out atrophic and rolling acne scars in skin phototypes III, IV, and V with variable square pulse erbium: yttrium-aluminum-garnet laser resurfacing. *Dermatol Surg* 2009; **35**:1376–83.
- 67 Xu Y, Deng Y. Ablative fractional CO<sub>2</sub> laser for facial atrophic acne scars. *Facial Plast Surg* 2018; **34**:205–19.
- 68 Abdel Hay R, Hegazy R, Abdel Hady M, Saleh N. Clinical and dermoscopic evaluation of combined (salicylic acid 20% and azelaic acid 20%) versus trichloroacetic acid 25% chemical peel in acne: an RCT. *J Dermatolog Treat* 2019; **30**:572–7.
- 69 Adalatkhah H, Pourfarzi F, Sadeghi-Bazargani H. Flutamide versus a cyproterone acetate-ethinyl estradiol combination in moderate acne: a pilot randomized clinical trial. *Clin Cosmet Investig Dermatol* 2011; **4**:117–21.
- 70 Alanazi TM, Alajroush W, Alharthi RM *et al.* Prevalence of acne vulgaris, its contributing factors, and treatment satisfaction among Saudi population in Riyadh, Saudi Arabia: a cross-sectional study. *J Dermatol Dermatol Surg* 2020; **24**:33–7.
- 71 Alsubeeh NA, Alsharafi AA, Ahamed SS, Alajlan A. Treatment adherence among patients with five dermatological diseases and four treatment types – a cross-sectional study. *Patient Prefer Adherence* 2019; **13**:2029–38.
- 72 Baker M, Tuley M, Busdiecker FL *et al.* Adapalene gel 0.1% is effective and well tolerated in acne patients in a dermatology practice setting. *Cutis* 2001; **68**:41–7.
- 73 Barakat MT, Mofteh NH, El Khayyat MAM, Abdelhakim ZA. Significant reduction of inflammation and sebaceous glands size in acne vulgaris lesions after intense pulsed light treatment. *Dermatol Ther* 2017; **30**:e12418.
- 74 Baugh WP, Kucaba WD. Nonablative phototherapy for acne vulgaris using the KTP 532 nm laser. *Dermatol Surg* 2005; **31**:1290–6.
- 75 Brodell RT, Schlosser BJ, Rafal E *et al.* A fixed-dose combination of adapalene 0.1%-BPO 2.5% allows an early and sustained improvement in quality of life and patient treatment satisfaction in severe acne. *J Dermatolog Treat* 2012; **23**:26–34.
- 76 Burger S, Truter I, Blignault SM, Venter DJL. Systemic isotretinoin in the management of acne – a patient questionnaire survey. *SA Fam Pract* 2009; **51**:427–33.
- 77 Campbell JL, Weiss JS. The results of the MORE trial: overview. *Cutis* 2006; **78**:5–11.
- 78 Cardinali C, Gimma A. Persistent acne in adult women: tolerability profile of clindamycin 1% in association with benzoyl peroxide 3% in comparison with tretinoin cream 0.05%. *Esper Dermatol* 2015; **17**:63–9.
- 79 Choi YS, Suh HS, Yoon MY *et al.* A study of the efficacy of cleansers for acne vulgaris. *J Dermatolog Treat* 2010; **21**:201–5.
- 80 International Clinical Trials Registry Platform. Effect of antihistamines in acne along with standard therapy. Available at <http://www.who.int/trialsearch/Trial2.aspx?TrialID=CTRI/2019/05/018951> (last accessed 31 March 2020).
- 81 Del Rosso JQ. A 6% benzoyl peroxide foaming cloth cleanser used in the treatment of acne vulgaris. *Clin Aesthet Dermatol* 2009; **2**:26–9.
- 82 Del Rosso JQ, Stein Gold L, Johnson SM *et al.* Efficacy and safety of adapalene 0.3%/benzoyl peroxide 2.5% gel plus oral doxycycline in subjects with severe inflammatory acne who are candidates for oral isotretinoin. *J Drugs Dermatol* 2018; **17**:264–73.

- 83 Dong Y, Zhou G, Chen J *et al.* A new LED device used for photodynamic therapy in treatment of moderate to severe acne vulgaris. *Photodiagnosis Photodyn Ther* 2016; **13**:188–95.
- 84 Donnarumma M, Fattore D, Greco V *et al.* How to increase adherence and compliance in acne treatment? a combined strategy of SMS and visual instruction leaflet. *Dermatology* 2019; **235**:1–8.
- 85 Dréno B, Thiboutot D, Gollnick H *et al.* Large-scale worldwide observational study of adherence with acne therapy. *Int J Dermatol* 2010; **49**:448–56.
- 86 Dréno B, Bissonnette R, Gagné-Henley A *et al.* Prevention and reduction of atrophic acne scars with adapalene 0.3%/benzoyl peroxide 2.5% gel in subjects with moderate or severe facial acne: results of a 6-month randomized, vehicle-controlled trial using intra-individual comparison. *Am J Clin Dermatol* 2018; **19**:275–86.
- 87 DuBois J, Ong GCW, Petkar G *et al.* Patient-reported outcomes in acne patients with skin of color using adapalene 0.3%-benzoyl peroxide 2.5%: a prospective real-world study. *J Drugs Dermatol* 2019; **18**:514–20.
- 88 Eichenfield LF, Sugarman JL, Guenin E *et al.* Novel tretinoin 0.05% lotion for the once-daily treatment of moderate-to-severe acne vulgaris in a preadolescent population. *Pediatr Dermatol* 2019; **36**:193–9.
- 89 Fabbrocini G, Staibano S, De Rosa G *et al.* Resveratrol-containing gel for the treatment of acne vulgaris: a single-blind, vehicle-controlled, pilot study. *Am J Clin Dermatol* 2011; **12**:133–41.
- 90 Franzke N, Zimmer L, Schäfer I *et al.* Quality of medical care of patients with acne vulgaris in Germany – nationwide survey of pharmacy clients. *J Dtsch Dermatol Ges* 2009; **7**:1060–3.
- 91 Fuchs CSK, Bay C, Adatto M *et al.* Acne treatment with light absorbing gold microparticles and optical pulses: an open-label European multi-centered study in moderate to moderately severe acne vulgaris patients. *Lasers Surg Med* 2019; **51**:686–93.
- 92 Gold MH, Biron J. Efficacy of a novel combination of pneumatic energy and broadband light for the treatment of acne. *J Drugs Dermatol* 2008; **7**:639–42.
- 93 Gold M. A New, once-daily, optimized, fixed combination of clindamycin phosphate 1.2% and low-concentration benzoyl peroxide 2.5% gel for the treatment of moderate-to-severe acne. *J Clin Aesthetic Dermatol* 2009; **2**:44–8.
- 94 Gold MH, Biron JA, Sensing W. Clinical and usability study to determine the safety and efficacy of the Silk'n Blue device for the treatment of mild to moderate inflammatory acne vulgaris. *J Cosmet Laser Ther* 2014; **16**:108–13.
- 95 Gollnick HPM, Friedrich M, Peschen M *et al.* Effect of adapalene 0.1%/benzoyl peroxide 2.5% topical gel on quality of life and treatment adherence during long-term application in patients with predominantly moderate acne with or without concomitant medication – additional results from the non-interventional cohort study ELANG. *J Eur Acad Dermatol Venereol* 2015; **29**:23–9.
- 96 Gonzalez P, Vela R, Cirigliano M. The tolerability profile of clindamycin 1%/benzoyl peroxide 5% gel vs. adapalene 0.1%/benzoyl peroxide 2.5% gel for facial acne: results of a randomized, single-blind, split-face study. *J Cosmet Dermatol* 2012; **11**:251–60.
- 97 Haedersdal M, Togsverd-Bo K, Wiegell SR, Wulf HC. Long-pulsed dye laser versus long-pulsed dye laser-assisted photodynamic therapy for acne vulgaris: a randomized controlled trial. *J Am Acad Dermatol* 2008; **58**:387–94.
- 98 Hermes B, Praetel C, Henz BM. Medium dose isotretinoin for the treatment of acne. *J Eur Acad Dermatol Venereol* 1998; **11**:117–21.
- 99 Ianosi S, Neagoe D, Calbureanu M, Ianosi G. Investigator-blind, placebo-controlled, randomized comparative study on combined vacuum and intense pulsed light versus intense pulsed light devices in both comedonal and papulopustular acne. *J Cosmet Laser Ther* 2013; **15**:248–54.
- 100 Iranian Registry of Clinical Trials. The effect of topical semisolid formulation of metformin on severity and number of skin lesions in acne patients (IRCT2017013132319N1). Available at: <https://en.irct.ir/trial/25214> (last accessed 31 March 2020).
- 101 Iranian Registry of Clinical Trials. Spironolactone for adult female acne (ISRCTN12892056). Available at: <http://www.isrctn.com/ISRCTN12892056> (last accessed 31 March 2020).
- 102 Jung HJ, Jang DH, Lee JI *et al.* [Study on the usage of acne patches in patients with acne]. *Korean J Dermatol* 2018; **56**:539–42 (in Korean).
- 103 Kellett N, West F, Finlay AY. Conjoint analysis: a novel, rigorous tool for determining patient preferences for topical antibiotic treatment for acne. A randomised controlled trial. *Br J Dermatol* 2006; **154**:524–32.
- 104 Khodaeiani E, Fouladi RF, Yousefi N *et al.* Efficacy of 2% metronidazole gel in moderate acne vulgaris. *Indian J Dermatol* 2012; **57**:279–81.
- 105 Kim MR, Kerrouche N. Combination of benzoyl peroxide 5% gel with liquid cleanser and moisturizer SPF 30 in acne treatment results in high levels of subject satisfaction, good adherence and favorable tolerability. *J Dermatolog Treat* 2018; **29**:49–54.
- 106 Kim S. The dual treatment of acne vulgaris using two kinds of ELOS (electro optical synergy) system: a simultaneous split-face trial. *J Cosmet Laser Ther* 2008; **10**:213–6.
- 107 Kwon HH, Choi SC, Jung JY *et al.* Comparison of novel dual mode vs conventional single pass of a 1450-nm diode laser in the treatment of acne vulgaris for Korean patients: a 20-week prospective, randomized, split-face study. *J Cosmet Dermatol* 2018; **17**:1063–8.
- 108 Kwon HH, Park HY, Choi SC *et al.* Novel device-based acne treatments: comparison of a 1450-nm diode laser and microneedling radiofrequency on mild-to-moderate acne vulgaris and seborrhoea in Korean patients through a 20-week prospective, randomized, split-face study. *J Eur Acad Dermatol Venereol* 2018; **32**:639–44.
- 109 Kwon HH, Choi SC, Jung JY *et al.* A novel combined light-based treatment of acne vulgaris with 1,450-nm diode laser and 450-nm blue light. *Dermatol Surg* 2019; **45**:1147–54.
- 110 Laquieze S, Czemielewski J, Rueda MJ. Beneficial effect of a moisturizing cream as adjunctive treatment to oral isotretinoin or topical tretinoin in the management of acne. *J Drugs Dermatol* 2006; **5**:985–90.
- 111 Lee JW, Kim BJ, Kim M *et al.* Selective sebaceous gland electrothermolysis as a treatment for acne: a prospective pilot study. *Int J Dermatol* 2012; **51**:339–44.
- 112 Lee KR, Lee EG, Lee HJ, Yoon MS. Assessment of treatment efficacy and sebosuppressive effect of fractional radiofrequency microneedle on acne vulgaris. *Lasers Surg Med* 2013; **45**:639–47.
- 113 Lekwuttikarn R, Tempark T, Chatproedprai S, Wanakul S. Randomized, controlled trial split-faced study of 595-nm pulsed dye laser in the treatment of acne vulgaris and acne erythema in adolescents and early adulthood. *Int J Dermatol* 2017; **56**:884–8.
- 114 Lubtikulthum P, Kamanamool N, Udompataikul M. A comparative study on the effectiveness of herbal extracts vs 2.5% benzoyl peroxide in the treatment of mild to moderate acne vulgaris. *J Cosmet Dermatol* 2019; **18**:1767–75.
- 115 Marron SE, Tomas-Aragones L, Boira S. Anxiety, depression, quality of life and patient satisfaction in acne patients treated with oral isotretinoin. *Acta Derm Venereol* 2013; **93**:701–6.
- 116 Martínez-Carpio PA, Alcolea-López JM, Vélez M. Efficacy of photodynamic therapy in the short and medium term in the

- treatment of actinic keratosis, basal cell carcinoma, acne vulgaris and photoaging: results from four clinical trials. *Laser Therapy* 2012; **21**:199–208.
- 117 Maruguchi Y, Maruguchi T. Treatment of inflammatory facial acne vulgaris: comparison of the 1450-nm diode laser and conventional physical treatment. *J Cosmet Laser Ther* 2006; **8**:167–9.
  - 118 Mazzetti A, Moro L, Gerloni M, Cartwright M. A phase 2b, randomized, double-blind vehicle controlled, dose escalation study evaluating clascoterone 0.1%, 0.5%, and 1% topical cream in subjects with facial acne. *J Drugs Dermatol* 2019; **18**:570–5.
  - 119 Min S, Park SY, Yoon JY, Suh DH. Comparison of fractional microneedling radiofrequency and bipolar radiofrequency on acne and acne scar and investigation of mechanism: comparative randomized controlled clinical trial. *Arch Dermatol Res* 2015; **307**:897–904.
  - 120 Min S, Park SY, Yoon JY *et al.* Fractional microneedling radiofrequency treatment for acne-related post-inflammatory erythema. *Acta Derm Venereol* 2016; **96**:87–91.
  - 121 Mofteh NH, Ibrahim SM, Wahba NH. Intense pulsed light versus photodynamic therapy using liposomal methylene blue gel for the treatment of truncal acne vulgaris: a comparative randomized split body study. *Arch Dermatol Res* 2016; **308**:263–8.
  - 122 Mohanan S, Parveen B, Malathy PA, Gomathi N. Use of intense pulse light for acne vulgaris in Indian skin – a case series. *Int J Dermatol* 2012; **51**:473–6.
  - 123 Mokhtari F, Faghihi G, Basiri A *et al.* Comparison effect of azithromycin gel 2% with clindamycin gel 1% in patients with acne. *Adv Biomed Res* 2016; **5**:72.
  - 124 Mokhtari F, Gholami M, Siadat AH *et al.* Efficacy of intense-pulsed light therapy with topical benzoyl peroxide 5% versus benzoyl peroxide 5% alone in mild-to-moderate acne vulgaris: a randomized controlled trial. *J Res Pharm Pract* 2017; **6**:199–205.
  - 125 Montefrecola G, Cacciapuoti S, Capasso C *et al.* Tolerability and camouflaging effect of corrective makeup for acne: results of a clinical study of a novel face compact cream. *Clin Cosmet Investig Dermatol* 2016; **9**:307–13.
  - 126 Myhill T, Coulson W, Nixon P. Use of supplementary patient education material increases treatment adherence and satisfaction among acne patients receiving adapalene 0.1%/benzoyl peroxide 2.5% gel in primary care clinics: a multicenter, randomized, controlled clinical study. *Dermatol Ther (Heidelb)* 2017; **7**:515–24.
  - 127 ClinicalTrials.gov. A randomized, controlled, evaluator-blinded pilot study to evaluate the effect of automated text message reminders on patient compliance with topical medications and its efficacy on skin disease control in adolescents and adults with mild to moderate acne (NCT01082523). Available at: <https://clinicaltrials.gov/ct2/show/NCT01082523> (last accessed 31 March 2020).
  - 128 Ohlson J, Dakovic R, Berg M. Observational study of clindamycin phosphate and tretinoin gel for the treatment of acne. *J Drugs Dermatol* 2019; **18**:328–34.
  - 129 Paradisi R, Fabbri R, Porcu E *et al.* Retrospective, observational study on the effects and tolerability of flutamide in a large population of patients with acne and seborrhea over a 15-year period. *Gynecol Endocrinol* 2011; **27**:823–9.
  - 130 Pariser DM, Rich P, Cook-Bolden FE, Korotzer A. An aqueous gel fixed combination of clindamycin phosphate 1.2% and benzoyl peroxide 3.75% for the once-daily treatment of moderate to severe acne vulgaris. *J Drugs Dermatol* 2014; **13**:1083–9.
  - 131 Picosse FR, Bonatto DC, Hassun KM *et al.* Treatment of moderate to severe acne vulgaris with an oral isotretinoin similar to the reference product. *Surg Cosmet Dermatol* 2016; **8**:121–7.
  - 132 Politi Y, Levi A, Enk CD, Lapidoth M. Integrated cooling-vacuum-assisted 1540-nm erbium:glass laser is effective in treating mild-to-moderate acne vulgaris. *Lasers Med Sci* 2015; **30**:2389–93.
  - 133 Prilepskaya VN, Serov VN, Zharov EV *et al.* Effects of a phasic oral contraceptive containing desogestrel on facial seborrhea and acne. *Contraception* 2003; **68**:239–45.
  - 134 Rapp DA, Brenes GA, Feldman SR *et al.* Anger and acne: implications for quality of life, patient satisfaction and clinical care. *Br J Dermatol* 2004; **151**:183–9.
  - 135 Rassai S, Mehri M, Yaghoobi R *et al.* Superior efficacy of azithromycin and levamisole vs. azithromycin in the treatment of inflammatory acne vulgaris: an investigator blind randomized clinical trial on 169 patients. *Int J Clin Pharmacol Ther* 2013; **51**:490–4.
  - 136 Rendon MI, Rodriguez DA, Kawata AK *et al.* Acne treatment patterns, expectations, and satisfaction among adult females of different races/ethnicities. *Clin Cosmet Investig Dermatol* 2015; **8**:231–8.
  - 137 See JA, Goh CL, Hayashi N, Suh DH. Optimizing the use of topical retinoids in Asian acne patients. *J Dermatol* 2018; **45**:522–8.
  - 138 Shamban AT, Enokibori M, Narurkar V, Wilson D. Photopneumatic technology for the treatment of acne vulgaris. *J Drugs Dermatol* 2008; **7**:139–45.
  - 139 Shin JU, Lee SH, Jung JY, Lee JH. A split-face comparison of a fractional microneedle radiofrequency device and fractional carbon dioxide laser therapy in acne patients. *J Cosmet Laser Ther* 2012; **14**:212–7.
  - 140 Smith SR, Kempers S. A study of 5.5% benzoyl peroxide microsphere cream versus 6% benzoyl peroxide gel in the treatment of acne vulgaris. *Cosmet Dermatol* 2006; **19**:537–42.
  - 141 Son BK, Yun Y, Choi IH. Efficacy of ah shi point acupuncture on acne vulgaris. *Acupunct Med* 2010; **28**:126–9.
  - 142 de Souza Sittart JA, da Cosa A, Mulinari-Brenner F *et al.* Multicenter study for efficacy and safety evaluation of a fixed dose combination gel with adapalene 0.1% and benzoyl peroxide 2.5% (Epiduo®) for the treatment of acne vulgaris in Brazilian population. *An Bras Dermatol* 2015; **90** (Suppl. 1):1–16.
  - 143 Stein Gold L, Weiss J, Rueda MJ *et al.* Moderate and severe inflammatory acne vulgaris effectively treated with single-agent therapy by a new fixed-dose combination adapalene 0.3%/benzoyl peroxide 2.5% gel: a randomized, double-blind, parallel-group, controlled study. *Am J Clin Dermatol* 2016; **17**:293–303.
  - 144 Suh DH, Shin JW, Min SU *et al.* Treatment-seeking behaviors and related epidemiological features in Korean acne patients. *J Korean Med Sci* 2008; **23**:969–74.
  - 145 Tan J, Stein Gold L, Schlessinger J *et al.* Short-term combination therapy and long-term relapse prevention in the treatment of severe acne vulgaris. *J Drugs Dermatol* 2012; **11**:174–80.
  - 146 Tan J, Bissonnette R, Gratton D *et al.* The safety and efficacy of four different fixed combination regimens of adapalene 0.1%/benzoyl peroxide 2.5% gel for the treatment of acne vulgaris: results from a randomised controlled study. *Eur J Dermatol* 2018; **28**:502–8.
  - 147 Tanghetti EA, Kircik LH, Green LJ *et al.* A phase 2, multicenter, double-blind, randomized, vehicle-controlled clinical study to compare the safety and efficacy of a novel tazarotene 0.045% lotion and tazarotene 0.1% cream in the treatment of moderate-to-severe acne vulgaris. *J Drugs Dermatol* 2019; **18**:542–8.
  - 148 Ting W. Randomized, observer-blind, split-face study to compare the irritation potential of 2 topical acne formulations over a 14-day treatment period. *Cutis* 2012; **90**:91–6.
  - 149 Tremblay JF, Sire DJ, Lowe NJ, Moy RL. Light-emitting diode 415 nm in the treatment of inflammatory acne: an open-label, multicentric, pilot investigation. *J Cosmet Laser Ther* 2006; **8**:31–3.



- 150 Troielli PA, Asis B, Bermejo A *et al.* Community study of fixed-combination adapalene 0.1% and benzoyl peroxide 2.5% in acne. *Skinmed* 2010; **8**:17–22.
- 151 Tucker RP, Maclure K, Paudyal V *et al.* An exploratory study of community pharmacist diagnosis and management of dermatitis and acne. *SelfCare* 2017; **8**:1–10.
- 152 Tying SK, Kircik LH, Pariser DM *et al.* Novel tretinoin 0.05% lotion for the once-daily treatment of moderate-to-severe acne vulgaris: assessment of efficacy and safety in patients aged 9 years and older. *J Drugs Dermatol* 2018; **17**:1084–91.
- 153 Veraldi S, Giovane GL, Guerriero C, Bettoli V. Efficacy and tolerability of topical 0.2% Myrtacine(R) and 4% vitamin PP for prevention and treatment of retinoid dermatitis in patients with mild to moderate acne. *G Ital Dermatol Venereol* 2012; **147**:491–7.
- 154 Veraldi S, Barbareschi M, Mauro M *et al.* Role of cleansers in the management of acne: results of an Italian survey in 786 patients. *J Dermatolog Treat* 2016; **27**:439–42.
- 155 Veraldi S, Micali G, Berardesca E *et al.* Results of a multicenter, randomized, controlled trial of a hydrogen peroxide-based kit versus a benzoyl peroxide-based kit in mild-to-moderate acne. *J Clin Aesthet Dermatol* 2016; **9**:50–4.
- 156 Weiss JS, Davis J, Matthew W. Improving patient satisfaction and acne severity in patients with mild to moderate acne: the BEST study. *Cutis* 2003; **71** (Suppl.):3–4.
- 157 Wheeland RG, Dhawan S. Evaluation of self-treatment of mild-to-moderate facial acne with a blue light treatment system. *J Drugs Dermatol* 2011; **10**:596–602.
- 158 Yu JNT, Huang P. Use of a TriPollar radio-frequency device for the treatment of acne vulgaris. *J Cosmet Laser Ther* 2011; **13**:50–3.
- 159 Abdel Kareem IM, Fouad MA, Ibrahim MK. Effectiveness of subcision using carboxytherapy plus fractional carbon dioxide laser resurfacing in the treatment of atrophic acne scars: comparative split face study. *J Dermatolog Treat* 2020; **31**:296–9.
- 160 Afra TP, Razmi TM, Narang T *et al.* Topical tazarotene gel, 0.1%, as a novel treatment approach for atrophic postacne scars: a randomized active-controlled clinical trial. *JAMA. Facial Plast Surg* 2019; **21**:125–32.
- 161 Alajlan AM, Alsuwaidan SN. Acne scars in ethnic skin treated with both non-ablative fractional 1,550 nm and ablative fractional CO<sub>2</sub> lasers: comparative retrospective analysis with recommended guidelines. *Lasers Surg Med* 2011; **43**:787–91.
- 162 Al-Dhalimi M, Jaber A. Treatment of atrophic facial acne scars with fractional Er: Yag laser. *J Cosmet Laser Ther* 2015; **17**:184–8.
- 163 Al-Dhalimi MA, Arnoos AA. Subcision for treatment of rolling acne scars in Iraqi patients: a clinical study. *J Cosmet Dermatol* 2012; **11**:144–50.
- 164 Asilian A, Faghihi G, Asemi Esfahani A *et al.* Comparison of two methods of subcision Nokor and blunt blade in acne scars treatment. *J Cosmet Dermatol* 2019; **18**:1788–93.
- 165 Alster TS, Tanzi EL, Lazarus M. The use of fractional laser photothermolysis for the treatment of atrophic scars. *Dermatol Surg* 2007; **33**:295–9.
- 166 Azzam OA, Atta AT, Sobhi RM, Mostafa PI. Fractional CO(2) laser treatment vs autologous fat transfer in the treatment of acne scars: a comparative study. *J Drugs Dermatol* 2013; **12**:e7–13.
- 167 Batra RS, Jacob CI, Hobbs L, Arndt KA, Dover JS. A prospective survey of patient experiences after laser skin resurfacing: results from 2 1/2 years of follow-up. *Arch Dermatol* 2003; **139**:1295–9.
- 168 Bernstein EF, Ferreira M, Anderson D. A pilot investigation to subjectively measure treatment effect and side-effect profile of non-ablative skin remodeling using a 532 nm, 2 ms pulse-duration laser. *J Cosmet Laser Ther* 2001; **3**:137–41.
- 169 Bernstein EF, Schomaker KT, Basilavacchio LD *et al.* Treatment of acne scarring with a novel fractionated, dual-wavelength, picosecond-domain laser incorporating a novel holographic beam-splitter. *Lasers Surg Med* 2017; **49**:796–802.
- 170 Bhatia AC, Dover JS, Arndt KA *et al.* Patient satisfaction and reported long-term therapeutic efficacy associated with 1,320 nm Nd:YAG laser treatment of acne scarring and photoaging. *Dermatol Surg* 2006; **32**:346–52.
- 171 Bjørn M, Stausbøl-Grøn B, Olesen AB, Hedelund L. Treatment of acne scars with fractional CO<sub>2</sub> laser at 1-month versus 3-month intervals: an intra-individual randomized controlled trial. *Lasers Surg Med* 2014; **46**:89–93.
- 172 Boss WK, Usal H, Fodor PB, Chernoff G. Autologous cultured fibroblasts: a protein repair system. *Ann Plast Surg* 2000; **44**:536–42.
- 173 Brunner E, Adamson PA, Harlock JN, Ellis DAF. Laser facial resurfacing: patient survey of recovery and results. *J Otolaryngol* 2000; **29**:377–81.
- 174 Cachafeiro T, Escobar G, Maldonado G *et al.* Comparison of non-ablative fractional erbium laser 1,340 nm and microneedling for the treatment of atrophic acne scars: a randomized clinical trial. *Dermatol Surg* 2016; **42**:232–41.
- 175 Casabona G. Combined use of microfocused ultrasound and a calcium hydroxylapatite dermal filler for treating atrophic acne scars: a pilot study. *J Cosmet Laser Ther* 2018; **20**:301–6.
- 176 Chan HHL, Lam LK, Wong DSY *et al.* Use of 1,320 nm Nd:YAG laser for wrinkle reduction and the treatment of atrophic acne scarring in Asians. *Lasers Surg Med* 2004; **34**:98–103.
- 177 Chan NPY, Ho SGY, Yeung CK *et al.* Fractional ablative carbon dioxide laser resurfacing for skin rejuvenation and acne scars in Asians. *Lasers Surg Med* 2010; **42**:615–23.
- 178 Chapas AM, Brightman L, Sukal S *et al.* Successful treatment of acneiform scarring with CO<sub>2</sub> ablative fractional resurfacing. *Lasers Surg Med* 2008; **40**:381–6.
- 179 Cho SI, Chung BY, Choi MG *et al.* Evaluation of the clinical efficacy of fractional radiofrequency microneedle treatment in acne scars and large facial pores. *Dermatol Surg* 2012; **38**:1017–24.
- 180 Chrastil B, Glaich AS, Goldberg LH, Friedman PM. Second-generation 1,550-nm fractional photothermolysis for the treatment of acne scars. *Dermatol Surg* 2008; **34**:1327–32.
- 181 Cole RP, Widdowson D, Moore JC. Outcome of erbium:yttrium aluminum garnet laser resurfacing treatments. *Lasers Med Sci* 2008; **23**:427–33.
- 182 Dai YX, Chuang YY, Chen PY, Chen CC. Efficacy and safety of ablative resurfacing with a high-energy 1,064 Nd-YAG picosecond-domain laser for the treatment of facial acne scars in Asians. *Lasers Surg Med* 2020; **52**:389–95.
- 183 Datz E, Schonberger C, Zeman F *et al.* Fractional carbon dioxide laser resurfacing of skin grafts: long-term results of a prospective, randomized, split-scar, evaluator-blinded study. *Lasers Surg Med* 2018; **50**:1010–6.
- 184 de Andrade LE. [Association of microneedling with phenol peeling: a new therapeutic approach for sagging, wrinkles and acne scars on the face]. *Surg Cosmet Dermatol* 2015; **7**:328–31 (in Portuguese).
- 185 de Andrade Lima EV. [Percutaneous induction of collagen with needles (IPCA) associated with pulsed radiofrequency with multi-needles (RPPM) in the management of depressed acne scars: treatment protocol]. *Surg Cosmet Dermatol* 2017; **9**:234–6 (in Portuguese).
- 186 Dierickx C, Larsson MK, Blomster S. Effectiveness and safety of acne scar treatment with nonanimal stabilized hyaluronic acid gel. *Dermatol Surg* 2018; **44** (Suppl. 1):S10–18.

- 187 Dierickx C. Using normal and high pulse coverage with picosecond laser treatment of wrinkles and acne scarring: long term clinical observations. *Lasers Surg Med* 2018; **50**:51–5.
- 188 Elawar A, Dahan S. Non-insulated fractional microneedle radiofrequency treatment with smooth motor insertion for reduction of depressed acne scars, pore size, and skin texture improvement: a preliminary study. *J Clin Aesthet Dermatol* 2018; **11**:41–4.
- 189 Epstein RE, Spencer JM. Correction of atrophic scars with Artefill: an open-label pilot study. *J Drugs Dermatol* 2010; **9**:1062–4.
- 190 Erol OO, Gulek A, Agaoglu G *et al.* Treatment of hypertrophic scars and keloids using intense pulsed light (IPL). *Aesth Plast Surg* 2008; **32**:902–9.
- 191 Faghihi G, Poostiyani N, Asilan A *et al.* Efficacy of fractionated microneedle radiofrequency with and without adding subcision for the treatment of atrophic facial acne scars: a randomized split-face clinical study. *J Cosmet Dermatol* 2017; **16**:223–9.
- 192 Firooz A, Rajabi-Estarabadi A, Nassiri-Kashani MH. Treatment of atrophic facial acne scars with fractional Er:YAG laser in skin phototype III–IV: a pilot study. *J Cosmet Laser Ther* 2016; **18**:204–7.
- 193 Galal O, Tawfik AA, Abdalla N, Soliman M. Fractional CO<sub>2</sub> laser versus combined platelet-rich plasma and fractional CO<sub>2</sub> laser in treatment of acne scars: image analysis system evaluation. *J Cosmet Dermatol* 2019; **18**:1665–71.
- 194 Gheisari M, Iranmanesh B, Saghi B. Blunt cannula subcision is more effective than Nokor needle subcision for acne scars treatment. *J Cosmet Dermatol* 2019; **18**:192–6.
- 195 Goel A, Gatne V. Use of nanofractional radiofrequency for the treatment of acne scars in Indian skin. *J Cosmet Dermatol* 2017; **16**:186–92.
- 196 Gold MH, Bron JA. Treatment of acne scars by fractional bipolar radiofrequency energy. *J Cosmet Laser Ther* 2012; **14**:172–8.
- 197 Goodman GJ. Carbon dioxide laser resurfacing: preliminary observations on short-term follow-up. A subjective study of 100 patients' attitudes and outcomes. *Dermatol Surg* 1998; **24**:665–72.
- 198 Gotkin RH, Sarnoff DS, Cannarozzo G *et al.* Ablative skin resurfacing with a novel microablative CO<sub>2</sub> laser. *J Drugs Dermatol* 2009; **8**:138–44.
- 199 Hedelund L, Haak CS, Togsverd-Bo K *et al.* Fractional CO<sub>2</sub> laser resurfacing for atrophic acne scars: a randomized controlled trial with blinded response evaluation. *Lasers Surg Med* 2012; **44**:447–52.
- 200 Hedelund L, Moreau KER, Beyer DM *et al.* Fractional nonablative 1,540-nm laser resurfacing of atrophic acne scars. A randomized controlled trial with blinded response evaluation. *Lasers Med Sci* 2010; **25**:749–54.
- 201 Hu S, Chen MC, Lee MC *et al.* Fractional resurfacing for the treatment of atrophic facial acne scars in Asian skin. *Dermatol Surg* 2009; **35**:826–32.
- 202 Hu S, Hsiao WC, Chen MC *et al.* Ablative fractional erbium-doped yttrium aluminum garnet laser with coagulation mode for the treatment of atrophic acne scars in Asian skin. *Dermatol Surg* 2011; **37**:939–44.
- 203 Ibrahim MK, Ibrahim SM, Salem AM. Skin microneedling plus platelet-rich plasma versus skin microneedling alone in the treatment of atrophic post acne scars: a split face comparative study. *J Dermatolog Treat* 2018; **29**:281–6.
- 204 Isarría MJ, Cornejo P, Muñoz E *et al.* Evaluation of clinical improvement in acne scars and active acne in patients treated with the 1540-nm non-ablative fractional laser. *J Drugs Dermatol* 2011; **10**:907–12.
- 205 Kang WH, Kim YJ, Pyo WS *et al.* Atrophic acne scar treatment using triple combination therapy: dot peeling, subcision and fractional laser. *J Cosmet Laser Ther* 2009; **11**:212–5.
- 206 Karnik J, Baumann L, Bruce S *et al.* A double-blind, randomized, multicenter, controlled trial of suspended polymethylmethacrylate microspheres for the correction of atrophic facial acne scars. *J Am Acad Dermatol* 2014; **71**:77–83.
- 207 Keller R, Belda Júnior W, Valente NY, Rodrigues CJ. Nonablative 1,064-nm Nd:YAG laser for treating atrophic facial acne scars: histologic and clinical analysis. *Dermatol Surg* 2007; **33**:1470–6.
- 208 Kim JE, Lee HW, Kim JK *et al.* Objective evaluation of the clinical efficacy of fractional radiofrequency treatment for acne scars and enlarged pores in Asian skin. *Dermatol Surg* 2014; **40**:988–95.
- 209 Kim H, Yo KH, Zheng Z, Cho SB. Pressure- and dose-controlled transcutaneous pneumatic injection of hypertonic glucose solution for the treatment of atrophic skin disorders. *J Cosmet Laser Ther* 2017; **19**:479–84.
- 210 Knight JM. Combined 400–600nm and 800–1200nm intense pulsed phototherapy of facial acne vulgaris. *J Drugs Dermatol* 2019; **18**:1116–22.
- 211 Koren A, Isman G, Cohen S *et al.* Efficacy of a combination of diluted calcium hydroxylapatite-based filler and an energy-based device for the treatment of facial atrophic acne scars. *Clin Exp Dermatol* 2019; **44**:e171–e176.
- 212 Lan T, Xiao Y, Tang L *et al.* Treatment of atrophic acne scarring with fractional micro-plasma radio-frequency in Chinese patients: a prospective study. *Lasers Surg Med* 2018; **50**:844–50.
- 213 Lee DH, Choi YS, Min SU *et al.* Comparison of a 585-nm pulsed dye laser and a 1064-nm Nd:YAG laser for the treatment of acne scars: a randomized split-face clinical study. *J Am Acad Dermatol* 2009; **60**:801–7.
- 214 Lee JW, Kim BJ, Kim MN, Lee CK. Treatment of acne scars using subdermal minimal surgery technology. *Dermatol Surg* 2010; **36**:1281–7.
- 215 Loss MJ, Leung S, Chien A *et al.* Adapalene 0.3% gel shows efficacy for the treatment of atrophic acne scars. *Dermatol Ther (Heidelb)* 2018; **8**:245–57.
- 216 Maas CS, Joseph JH. Safety and Effectiveness of microfocused ultrasound with visualization for the correction of moderate to severe atrophic acne scars. *J Drugs Dermatol* 2019; **18**:1109–14.
- 217 Mahmoud BH, Srivastava D, Janiga JJ *et al.* Safety and efficacy of erbium-doped yttrium aluminum garnet fractionated laser for treatment of acne scars in type IV to VI skin. *Dermatol Surg* 2010; **36**:602–9.
- 218 Majid I, Imran S. Fractional CO<sub>2</sub> laser resurfacing as monotherapy in the treatment of atrophic facial acne scars. *J Cutan Aesthet Surg* 2014; **7**:87–92.
- 219 Montesi G, Calvieri S, Balzani A, Gold MH. Bipolar radiofrequency in the treatment of dermatologic imperfections: clinicopathological and immunohistochemical aspects. *J Drugs Dermatol* 2007; **6**:890–6.
- 220 ClinicalTrials.gov. Comparison of treatments for atrophic acne scars (NCT02025088). Available at: <https://clinicaltrials.gov/show/NCT02025088> (last accessed 4 April 2020).
- 221 ClinicalTrials.gov. Restylane silk acne scar efficacy evaluation study (NCT02955381). Available at: <https://clinicaltrials.gov/show/NCT02955381> (last accessed 4 April 2020).
- 222 ClinicalTrials.gov. Comparison between microneedling daycare procedure and tazarotene 0.1% gel local application in acne scarring (NCT03170596). Available at: <https://clinicaltrials.gov/show/NCT03170596> (last accessed 4 April 2020).
- 223 Nirmal B, Pai SB, Sripathi H *et al.* Efficacy and safety of erbium-doped yttrium aluminium garnet fractional resurfacing laser for treatment of facial acne scars. *Indian J Dermatol Venereol Leprol* 2013; **79**:193–8.



- 224 Ortiz AE, Tremaine AM, Zachary CB. Long-term efficacy of a fractional resurfacing device. *Lasers Surg Med* 2010; **42**:168–70.
- 225 Panchaprateep R, Munavalli G. Low-fluence 585 nm Q-switched Nd:YAG laser: a novel laser treatment for post-acne erythema. *Lasers Surg Med* 2015; **47**:148–55.
- 226 Peterson JD, Palm MD, Kiripolsky MG *et al.* Evaluation of the effect of fractional laser with radiofrequency and fractionated radiofrequency on the improvement of acne scars. *Dermatol Surg* 2011; **37**:1260–7.
- 227 Petrov A. Efficiency of carbon dioxide fractional laser in skin resurfacing. *Maced J Med Sci* 2016; **4**:271–6.
- 228 Petrov A, Pljakovska V. Fractional carbon dioxide laser in treatment of acne scars. *Maced J Med Sci* 2016; **4**:38–42.
- 229 Politi Y, Levi A, Lapidoth M. Integrated cooling-vacuum-assisted non-fractional 1540 nm erbium:glass laser is effective in treating acne scars. *J Drugs Dermatol* 2016; **15**:1359–63.
- 230 Politi Y, Levi A, Snast I *et al.* Integrated cooling-vacuum-assisted non-fractional 1540-nm erbium:glass laser: a new modality for the simultaneous effective treatment of acne lesions and scars. *J Drugs Dermatol* 2018; **17**:1173–6.
- 231 Qin X, Li H, Jian X, Yu B. Evaluation of the efficacy and safety of fractional bipolar radiofrequency with high-energy strategy for treatment of acne scars in Chinese. *J Cosmet Laser Ther* 2015; **17**:237–45.
- 232 Rojas H, Vega NA, Lobos NM, Carvajal DA. Satisfacción de pacientes con cicatrices atróficas de acné tratados con láser ablativo y/o técnicas quirúrgicas. *Rev Chil Dermatol* 2015; **31**:365–8.
- 233 Saadawi AN, Esawy AM, Kandeel AH, El-Sayed W. Microneedling by dermapen and glycolic acid peel for the treatment of acne scars: comparative study. *J Cosmet Dermatol* 2018; **18**:107–14.
- 234 Samuthrsindh K, Helbig D, Paasch U *et al.* [Prospective, uncontrolled examination of ablative fractional photothermolysis on Asian and Caucasian skin]. *Med Laser Appl* 2011; **26**:3–9 (in German).
- 235 Sapra S, Stewart JA, Mraud K, Schupp R. A Canadian study of the use of poly-L-lactic acid dermal implant for the treatment of hill and valley acne scarring. *Dermatol Surg* 2015; **41**:587–94.
- 236 Tanzi EL, Alster TS. Comparison of a 1450-nm diode laser and a 1320-nm Nd:YAG laser in the treatment of atrophic facial scars: a prospective clinical and histologic study. *Dermatol Surg* 2004; **30**:152–7.
- 237 Tatliparmak A, Aksoy B, Shishehgharkhaneh LR *et al.* Use of combined fractional carbon dioxide laser and fractional microneedle radiofrequency for the treatment of acne scars: a retrospective analysis of 1-month treatment outcome on scar severity and patient satisfaction. *J Cosmet Dermatol* 2020; **19**:115–21.
- 238 Tay YK, Kwok C. Minimally ablative erbium:YAG laser resurfacing of facial atrophic acne scars in Asian skin: a pilot study. *Dermatol Surg* 2008; **34**:681–5.
- 239 Taylor MB, Zaleski-Larsen L, McGraw T. Single session treatment of rolling acne scars using tumescent anesthesia, 20% trichloroacetic acid extensive subcision, and fractional CO<sub>2</sub> laser. *Dermatol Surg* 2017; **43** (Suppl. 1):S70–S74.
- 240 Tenna S, Cogliandro A, Barone M *et al.* Comparative study using autologous fat grafts plus platelet-rich plasma with or without fractional CO<sub>2</sub> laser resurfacing in treatment of acne scars: analysis of outcomes and satisfaction with FACE-Q. *Aesthetic Plast Surg* 2017; **41**:661–6.
- 241 Thomas CL, Kim B, Lam J *et al.* Objective severity does not capture the impact of rosacea, acne scarring and photoaging in patients seeking laser therapy. *J Eur Acad Dermatol Venereol* 2017; **31**:361–6.
- 242 Trelles MA, Martínez-Carpio PA. Attenuation of acne scars using high power fractional ablative unipolar radiofrequency and ultrasound for transepidermal delivery of bioactive compounds through microchannels. *Lasers Surg Med* 2014; **46**:152–9.
- 243 Tzikas TI. Evaluation of the Radianse FN soft tissue filler for facial soft tissue augmentation. *Arch Facial Plast Surg* 2004; **6**:234–9.
- 244 Verner I. Clinical evaluation of the efficacy and safety of fractional bipolar radiofrequency for the treatment of moderate to severe acne scars. *Dermatol Ther* 2016; **29**:24–7.
- 245 Wang YS, Tay YK, Kwok C. Fractional ablative carbon dioxide laser in the treatment of atrophic acne scarring in Asian patients: a pilot study. *J Cosmet Laser Ther* 2010; **12**:61–4.
- 246 Weinstein A, Koren A, Sprecher E *et al.* The combined effect of tranilast 8% liposomal gel on the final cosmesis of acne scarring in patients concomitantly treated by isotretinoin: prospective, double-blind, split-face study. *Clin Exp Dermatol* 2020; **45**:41–7.
- 247 Yang Q, Huang W, Qian H *et al.* Efficacy and safety of 1550-nm fractional laser in the treatment of acne scars in Chinese patients: a split-face comparative study. *J Cosmet Laser Ther* 2016; **18**:312–6.
- 248 Yeung CK, Chan NPY, Shek SYN, Chan HHL. Evaluation of combined fractional radiofrequency and fractional laser treatment for acne scars in Asians. *Lasers Surg Med* 2012; **44**:622–30.
- 249 Yoo KH, Ahn JY, Kim JY *et al.* The use of 1540 nm fractional photothermolysis for the treatment of acne scars in Asian skin: a pilot study. *Photodermatol Photoimmunol Photomed* 2009; **25**:138–42.
- 250 Yu JN, Abat K. Safety and efficacy of hybrid energy and trifractional technologies in the treatment of acne scars: an open-label clinical trial. *J Cosmet Laser Ther* 2016; **18**:60–5.
- 251 Yu PX, Diao WQ, Qi ZL, Cai JL. Effect of dermabrasion and ReCell® on large superficial facial scars caused by burn, trauma and acnes. *Chin Med Sci J* 2016; **31**:173–9.
- 252 Yug A, Lane JE, Howard MS, Kent DE. Histologic study of depressed acne scars treated with serial high-concentration (95%) trichloroacetic acid. *Dermatol Surg* 2016; **32**:985–90.
- 253 Zhang M, Fang J, Wu Q, Lin T. Evaluation of the safety and efficacy of a picosecond alexandrite laser with DLA for acne scars in Chinese patients. *Lasers Surg Med* 2020; **52**:176–81.
- 254 Zhou BR, Zhang T, Bin Jameel AA *et al.* The efficacy of conditioned media of adipose-derived stem cells combined with ablative carbon dioxide fractional resurfacing for atrophic acne scars and skin rejuvenation. *J Cosmet Laser Ther* 2016; **18**:138–48.
- 255 Zhu JT, Xuan M, Zhang YN *et al.* The efficacy of autologous platelet-rich plasma combined with erbium fractional laser therapy for facial acne scars or acne. *Mol Med Rep* 2013; **8**:233–7.
- 256 Callender VD. Fitzpatrick skin types and clindamycin phosphate 1.2%/benzoyl peroxide gel: efficacy and tolerability of treatment in moderate to severe acne. *J Drugs Dermatol* 2012; **11**:643–8.
- 257 Cook-Bolden. Subject preferences for acne treatments containing adapalene gel 0.1%: results of the MORE trial. *Cutis* 2006; **78** (Suppl.):26–33.
- 258 Cook-Bolden FE. Treatment of moderate to severe acne vulgaris in a Hispanic population: a post-hoc analysis of efficacy and tolerability of clindamycin phosphate 1.2%/benzoyl peroxide 2.5% gel. *J Drugs Dermatol* 2012; **11**:455–9.
- 259 Fernandez-Obregon A, Davis MW. The BEST study: evaluating efficacy by selected demographic subsets. *Cutis* 2003; **71** (Suppl.):18–26.
- 260 Gold LS. Efficacy and tolerability of fixed-combination acne treatment in adolescents. *Cutis* 2013; **91**:152–9.
- 261 Rist T, Davis MW. Study design and selection criteria in the BEST study. *Cutis* 2003; **71** (Suppl.):5–9.

- 262 Rodriguez D, Davis MW. The BEST study: results according to prior treatment. *Cutis* 2003; **71** (Suppl.):27–34.
- 263 Shavin J, Weiss J. Implications of the BEST study. *Cutis* 2003; **71** (Suppl.):35–6.
- 264 Weiss J, Shavin J, Davis MW. Overall results of the BEST study following treatment of patients with mild to moderate acne. *Cutis* 2003; **71** (Suppl.):10–7.
- 265 Weiss J, Shavin J, Davis M. Preliminary results of a nonrandomized, multicenter, open-label study of patient satisfaction after treatment with combination benzoyl peroxide/clindamycin topical gel for mild to moderate acne. *Clin Ther* 2002; **24**:1706–17.
- 266 Alomar A, Guerra A, Perulero N *et al.* [Developing a questionnaire to assess treatment satisfaction in patients with acne]. *Actas Dermosifiliogr* 2004; **95**:491–5 (in Spanish).
- 267 Terwee CB, Prinsen CAC, Chiarotto A *et al.* COSMIN methodology for evaluating the content validity of patient-reported outcome measures: a Delphi study. *Qual Life Res* 2018; **27**:1159–70.
- 268 International Dermatology Outcome Measures (IDEOM). Available at: <http://dermoutcomes.org/index.html> (last accessed 30 June 2020).
- 269 Cochrane Skin. Core Outcome Set Initiative (COUSIN). Available at <http://cs-cousin.org/> (last accessed 30 June 2020).
- 270 Schmitt J, Apfelbacher C, Spuls PI *et al.* The Harmonizing Outcome Measures for Eczema (HOME) roadmap: a methodological framework to develop core sets of outcome measurements in dermatology. *J Invest Dermatol* 2015; **135**:24–30.
- 271 Callis Duffin K, Merola JF, Christensen R *et al.* Identifying a core domain set to assess psoriasis in clinical trials. *JAMA Dermatol* 2018; **154**:1137–44.
- 272 Ingram JR, Hadjieconomou S, Piguet V. Development of core outcome sets in hidradenitis suppurativa: systematic review of outcome measure instruments to inform the process. *Br J Dermatol* 2016; **175**:263–72.
- 273 Vincent R, Chalmers JR, McWilliams C *et al.* Assessing uptake of the Harmonising Outcome Measures for Eczema (HOME) Core Outcome Set and recommended instruments. *Br J Dermatol* 2020; **183**:566–8.
- 274 Acne Core Outcomes Research Network (ACORN). Available at: <https://sites.psu.edu/acnecoreoutcomes/> (last accessed 30 June 2020).
- 275 Revicki DA. Patient assessment of treatment satisfaction: methods and practical issues. *Gut* 2004; **53** (Suppl. 4):iv40–iv44.
- 276 Basch E, Spertus J, Dudley RA *et al.* Methods for developing patient-reported outcome-based performance measures (PRO-PMs). *Value Health* 2015; **18**:493–504.
- 277 Weldring T, Smith SM. Patient-reported outcomes (PROs) and patient-reported outcome measures (PROMs). *Health Serv Insights* 2013; **6**:61–8.
- 278 U.S. Department of Health and Human Services FDA Center for Drug Evaluation and Research; U.S. Department of Health and Human Services FDA Center for Biologics Evaluation and Research; U.S. Department of Health and Human Services FDA Center for Devices and Radiological Health. Guidance for industry: patient-reported outcome measures: use in medical product development to support labeling claims: draft guidance. *Health Qual Life Outcomes* 2006; **4**:79.

## Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website:

**Appendix S1** Search strategy.

**Table S1** Characteristics of the included patient-reported outcome measure.

**Table S2** Characteristics of the included study populations.

**Table S3** Content validity.

**Table S4** Study characteristics and results of studies evaluating satisfaction with acne treatment.

**Table S5** Study characteristics and results of studies evaluating satisfaction with acne scars treatment.

**Powerpoint S1** Journal Club Slide Set.

**Video S1** Author video.