



Universiteit
Leiden
The Netherlands

Internet-based treatment for eating disorders: bridging the treatment gap

Rohrbach, P.J.

Citation

Rohrbach, P. J. (2023, January 19). *Internet-based treatment for eating disorders: bridging the treatment gap*. Retrieved from <https://hdl.handle.net/1887/3512274>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3512274>

Note: To cite this publication please use the final published version (if applicable).

Chapter 7

Cost-effectiveness of three internet-based interventions for eating disorders: results from a randomized controlled trial

Published as: Rohrbach, P. J., Dingemans, A. E., Van Furth, E. F., Spinhoven, P., Van Ginkel, J. R., Bauer, S., & Van den Akker-Van Marle, M. E. (2022). Cost-effectiveness of three internet-based interventions for eating disorders: results from a randomized controlled trial. *International Journal of Eating Disorders*, 55, 1143-1155. <https://doi.org/10.1002/eat.23763>



Abstract

Objective: The primary aim was assessing the cost-effectiveness of an internet-based self-help program, expert-patient support, and the combination of both compared to a care-as-usual condition.

Method: An economic evaluation from a societal perspective was conducted alongside a randomized controlled trial. Participants aged 16 or older with at least mild eating disorder symptoms were randomly assigned to four conditions: (1) Featback, an online unguided self-help program, (2) chat or e-mail support from a recovered expert patient, (3) Featback with expert-patient support and (4) care-as-usual. After a baseline assessment and intervention period of 8 weeks, five online assessments were conducted over 12 months of follow-up. The main result constituted cost-utility acceptability curves with QALYs and societal costs over the entire study duration.

Results: No significant differences between the conditions were found regarding QALYs, health care costs and societal costs. Non-significant differences in QALYs were in favor of the Featback conditions and the lowest societal costs per participant were observed in the Featback only condition (€16,741) while the highest costs were seen in the care-as-usual condition (€28,479). The Featback only condition had the highest probability of being efficient compared to the alternatives for all acceptable willingness-to-pay values.

Conclusion: Featback, an internet-based unguided self-help intervention, was likely to be efficient compared to Featback with guidance from an expert patient, guidance alone and a care-as-usual condition. Results suggest that scalable interventions such as Featback may reduce health care costs and help individuals with eating disorders that are currently not reached by other forms of treatment.

Public significance statement

Internet-based interventions for eating disorders might reach individuals in society who currently do not receive appropriate treatment at low costs. Featback, an online automated self-help program for eating disorders, was found to improve quality of life slightly while reducing costs for society, compared to a do-nothing approach. Consequently, implementing internet-based interventions such as Featback likely benefits both individuals suffering from an eating disorder and society as a whole.

Introduction

Eating disorders are burdensome in terms of disability, quality of life and mortality (Arcelus et al., 2011; Smink et al., 2013) and also from an economic perspective (Erskine et al., 2016; Van Hoeken & Hoek, 2020). There exists a large treatment gap for eating disorders, meaning that many individuals with an eating disorder do not get help specifically for their eating disorder, despite having substantial symptoms (Austin et al., 2021). Nevertheless, they generally make more use of health care services compared to people without an eating disorder (Hart et al., 2011; Van Son et al., 2012; Weissman & Rosselli, 2017), which is reflected in higher health care costs (Ágh et al., 2016; Samnaliev et al., 2015). Eating disorder related costs may become larger still when also considering costs outside health care, such as productivity losses and caregiver costs (Deloitte Access Economics, 2020). These substantial costs warrant well-advised resource allocation decisions. In fact, investing in evidence-based treatment for eating disorders might ultimately result in cost savings (Bode et al., 2017). Apart from such policy changes, helping individuals with an eating disorder while reducing costs for society requires continued effort from researchers and clinicians to make treatments more effective, accessible and less expensive.

Cost-effectiveness research, where outcomes and costs of two different courses of action are compared, is necessary to distinguish interventions that are more efficient than others. Internet interventions, often coined as cost-effective alternative to other treatment options, have frequently been confirmed in their effectiveness (Linardon et al., 2020; Loucas et al., 2014; Melioli et al., 2016; Pittock et al., 2018), but cost-effectiveness research is scarce. Across mental disorders, evidence from systematic reviews cautiously suggests internet-based interventions might indeed be cost-effective, at least compared to do-nothing approaches (Ahern et al., 2018; Donker et al., 2015; Paganini et al., 2018). A few studies investigated the cost-effectiveness of internet interventions compared to face-to-face eating disorder treatment (Crow et al., 2009; König et al., 2018; Watson et al., 2018) and found internet interventions to be slightly less effective in reducing eating disorder symptoms, but also less costly. Consequently, such interventions might be especially efficient as a first step in a stepped-care treatment model, as they have the potential to reach individuals that currently do not get appropriate care for their eating disorder (Aardoom, Dingemans, & Van Furth, 2016). When researching these first step internet-based interventions for eating disorders, care as usual may be used as a reference, since it represents the, often inappropriate, care individuals with eating disorders in society receive. Unfortunately, there is a paucity of cost-effectiveness research comparing online interventions for eating disorders with care

as usual. A simulation study on US college students with eating disorders indicated that a stepped-care treatment model with online guided self-help was less costly and resulted in fewer individuals in need of additional treatment than usual care (Kass et al., 2017). Recently, Akers et al. (2021) showed an online version of the cognitive-dissonance based intervention 'the Body Project' to have health benefits compared to enhanced usual care, while health utilization was similar. Additionally, Aardoom, Dingemans, van Ginkel, et al. (2016) found that Featback, an online automatic monitoring and feedback system for people with an eating disorder, was cost-effective compared to a care-as-usual condition, regardless of whether the intervention was complemented with chat or e-mail support by a psychologist. Taken together, the limited evidence available suggests that online interventions for eating disorders may be cost-effective compared to care as usual, which is especially interesting considering that such interventions are scalable and easily accessible and can reach people at an early stage of eating disorder development. Recently, a second randomized controlled trial to replicate and extend the results on the effectiveness and cost-effectiveness of Featback compared to care as usual was conducted. In the first RCT (Aardoom, Dingemans, Spinhoven, et al., 2016), support by a psychologist did not add to the effectiveness of Featback. Possibly, support by expert patients (i.e., recovered individuals) is more fitting and effective for those reluctant to seek help (Rohrbach et al., 2019).

Aims

The primary aim of this study was to investigate the cost-effectiveness of the three conditions, 1) the fully automated internet intervention Featback, 2) chat or email support from expert patients and 3) the combination of both interventions compared to 4) a care-as-usual condition from a societal perspective. The three active online interventions were expected to be more efficient than care as usual.

Method

Design and randomization

This economic evaluation was part of a randomized controlled trial, pre-registered at the Dutch Trial Register (NL7065) and approved by an independent medical ethics committee (METC-LDD; NL64553.058.18). Detailed information on the interventions and methods can be found in the study protocol (Rohrbach et al., 2019). Results on the clinical effectiveness will be reported elsewhere. A two-by-two factorial design with the internet-based interventions Featback and expert-patient support was used, resulting in four conditions: (1) Featback, (2) Featback with expert-patient support, (3) expert-patient support and (4) care-as-usual condition. All conditions had a duration of eight weeks. Assessments on quality of life and costs were all online and completed by participants at post intervention and 3, 6, 9 and 12 month follow up. Participants were randomized and distributed across conditions in blocks of 40. For randomization, a computer-generated random numbers list was made by an independent researcher, concealing it from the principal investigator before and during the trial. The economic analysis maintains a societal perspective, meaning that both health

care costs and non-healthcare costs were included. Data concerning costs and utility covered a period of 14 months (i.e., 8 weeks intervention or waiting plus 12 months follow up).

Participants

Participants were recruited mainly via Proud2Bme, a Dutch online community for people with eating-related problems or eating disorders, from October 2018 to October 2019. After expressing interest to participate, they received a screening questionnaire. Eligible participants were 16 years or older, had internet access and reported at least mild eating disorder symptoms. Specifically, they scored 52 or higher on the Weight Concerns Scale (Killen et al., 1993) or reported a body mass index lower than or equal to 18.5, or one or more weekly binge eating episodes or compensatory behaviors in the past four weeks on the Short Evaluation of Eating Disorders (Bauer et al., 2005). Participants with severe eating disorder symptoms were advised to seek professional help, but were not excluded as they too may benefit from the offered interventions.

Interventions

Participants in all conditions were free to undergo any other type of intervention or treatment, representing individuals in society with varying levels of treatment. Consequently, the waiting list control condition can be seen as care as usual for individuals with eating disorder symptoms in (Dutch speaking) society.

Feedback

Participants could make weekly use of an automated monitoring and feedback system for eight weeks. Based on the answers of a short monitoring questionnaire, participants received a supportive feedback message with a summary of self-reported eating problems, psychoeducation, and guidance on how to counter eating disorder related symptoms. Current level of impairments as well as improvements or deteriorations in eating disorder related symptoms compared to the previous week were captured in the messages. Additionally, participants could access the Feedback website with psycho-educative material on eating disorders at their own convenience.

Expert-patient support

Five expert patients (sometimes referred to as peers or mentors) were recruited, who had a lived experience of an eating disorder and were fully recovered. They received a protocol and were trained on how to use their own experience to help others overcome their eating disorder via chat and e-mail. Monthly supervision from an experienced expert patient and clinical psychologist during the trial was included. Participants allocated to the conditions with expert-patient support were assigned to one of the expert patients for eight weeks and could schedule a 20-minute chat or e-mail session every week. Chat sessions closed automatically after 20 minutes. For e-mail sessions, participants sent an e-mail to their expert patient before the scheduled time slot and the expert patient responded during the appointment.

Feedback with expert-patient support

Participants in this condition were able to make use of both Feedback and weekly 20-minute chat or e-mail support from an expert patient.

Care-as-usual condition

Participants in this condition were placed on a waiting list for 14 months. After the waiting period, participants were offered eight weeks of Feedback with weekly expert-patient support.

Measures

Demographics

Assessed baseline variables were age, gender, nationality, education level, eating disorder treatment history, marital status, weight, height, eating disorder duration, internet usage, eating disorder symptoms assessed with the Eating Disorder Examination Questionnaire global scores (Fairburn & Beglin, 2008).

Quality of life

The primary outcome measure for the economic evaluation was quality-of-life adjusted life years (QALYs) as assessed with the EQ-5D-5L (EuroQol Group, 1990), which demonstrates adequate psychometric properties (Feng et al., 2021). The Dutch tariff (Versteegh, Vermeulen, et al., 2016) was used to translate EQ-5D-5L scores to utility values. Subsequently, QALYs were calculated over the 14 month follow-up period using the area-under-curve method.

Because generic health questionnaires like the EQ-5D-5L might be limited in their extent to detect changes in wellbeing for interventions aimed at mental health (Pietersma et al., 2013) the economic evaluation was also conducted using the ICECAP-A (Al-Janabi et al., 2012). Psychometric properties of the ICECAP-A have been found to be adequate (Afentou & Kinghorn, 2020; Rohrbach, Dingemans, Essers, et al., 2022). A capability value anchored at 0 (no capability) and 1 (full capability) was calculated for each participant using the ICECAP-A Dutch tariffs (Rohrbach et al., 2021) over the 14 month study period. Details on the used quality-of-life instruments and accompanying transformations can be found in Appendix F.1.

Costs

Health care costs included intervention costs and use of health care services. Intervention costs for Feedback included five minutes of technical support by a researcher (including setting up an account, redirecting participants to professional help in the case of severe symptom deterioration and responding to technical problems) multiplied by their hourly rate (€31.50). For expert-patient support, costs were calculated by multiplying their hourly rate (€22.31) with the time spent on support sessions (i.e., estimated at 30 minutes for each session, including preparation and administration). Additionally, supervision costs were calculated by

dividing the total time spent on supervision (i.e., 14 one-hour sessions attended by six expert patients with €22.31 hourly wage, one researcher with €31.50 hourly wage and one clinical psychologist with €106.17 hourly wage) by the amount of participants in the two conditions with the possibility of expert-patient support. All wages were determined based on the real wages during the conduct of the study.

Health care costs were measured with the TiC-P midi (Timman et al., 2015) at each assessment (i.e., over an 8-week period at post intervention and 3 months at all other follow-up assessments). The midi version was chosen over the full version as it reduced the time burden for participants while maintaining a reliable estimate of health service use (Timman et al., 2015). Finally, visits to the general practitioner, dietician, psychologist based in mental health institutions, the private section or hospitals, medical specialist, the emergency department, daycare in mental health institutions, and hospitalizations either in the hospital or a mental health institution were included as health care costs. After inspecting the data for errors and possible double counts, the amount of visits to each health care provider was multiplied with their cost prices as indicated by the Dutch guidelines for cost research in health care (Hakkaart-van Roijen et al., 2015; Kanters et al., 2017). All assessed health care services with their reference price are presented in Table 1.

Non-health care costs were measured with the Productivity Costs Questionnaire (PCQ) (Bouwman et al., 2015), including costs related to absence from work (absenteeism), reduced productivity at work because of health problems (presenteeism) and reduced productivity of unpaid work such as domestic chores because of health problems. Absenteeism costs were calculated by multiplying the recalled hours of missed work over the last 4 weeks extrapolated to 8 weeks (at post intervention) or 3 months (at follow-up measurements) by the average gross hourly wage of female working individuals in the Netherlands (Hakkaart-van Roijen et al., 2015). In cases of longer absence through illness the friction cost method was applied, meaning that no costs were incurred after being absent for 12 weeks, because initial production levels were expected to have been restored by that time. Presenteeism costs were calculated by multiplying the recalled hours with reduced productivity because of health problems over the last 4 weeks extrapolated to 8 weeks or 3 months by the average gross hourly wage of female working individuals in the Netherlands (Hakkaart-van Roijen et al., 2015). Lastly, costs related to reduced productivity of unpaid work was calculated by multiplying the recalled hours in which others had to perform domestic chores instead of the participant in the last 4 weeks extrapolated to 8 weeks or 3 months by the average gross hourly wage of a domestic worker (Hakkaart-van Roijen et al., 2015). Gross hourly wages are presented in Appendix 2.

All costs were indexed to the year 2021 using the Dutch consumer price index (OECD, 2021b). No discounting was applied to QALYs and costs, given that the time horizon was slightly more than 1 year.

Missing data

Baseline values of the EQ-5D-5L and ICECAP-A were not available. As these variables appear stable over a relatively short period (i.e., 8 weeks) of time, they were estimated to be equal to those at post intervention for the main analyses. This assumption was tested using sensitivity analyses.

Table 1. Price references

Category	Reference prize	CPI index 2014-2021	CPI index 2019-2021	Final cost price (2021)
Intervention costs^a				
Featback (5min researcher coordination per participant; hourly wage of €30.72)	€2.56		1.025	€2.62
Expert-patient support session (30min per session; hourly wage of €22.31)	€11.16		1.025	€11.44
Supervision costs per participant			€21.38	
Direct health care costs				
General practitioner	€33.00	1.095		€36.15
Dietician	€33.00	1.095		€36.15
Psychologist, psychotherapist or psychiatrist - mental health care	€98.00	1.095		€107.35
Psychologist, psychotherapist or psychiatrist - independent	€94.44	1.095		€103.45
Psychologist, psychotherapist or psychiatrist - hospital	€91.00	1.095		€99.68
Medical specialist	€91.00	1.095		€99.68
Emergency department	€259.00	1.095		€283.70
Day treatment - mental health care				€183.05
Hospitalization - mental health care	€302.36	1.095		€331.20
Hospitalization - hospital	€476.00	1.095		€521.40
Indirect costs				
Average gross hourly female wage	€31.60	1.095		€34.61
Average gross hourly domestic worker wage	€14.00	1.095		€15.34

CPI=Cost Price Index.

^a Wages of the research coordinator, expert patient and clinical psychologist (supervision) were based on the real wages during the conduct of the study.

Note. Dutch CPI indexes for 2021, 2019 and 2014 were €108.88, €106.20 and €99.40 respectively.

According to the intention-to-treat approach, all participants who completed baseline were included throughout the analyses. Missing data were multiply imputed (Rubin, 1986) using the software program R version 3.5.1. Details on the multiple imputation procedure can be found in Appendix F.2.

Statistical analyses

Costs, both health care and societal, and effects in terms of QALYs (EQ-5D-5L) and capabilities (ICECAP-A) over the 14 month period were compared between the four conditions using analyses of variance (ANOVA) pooled across imputations (Rubin, 1987; Van Ginkel & Kroonenberg, 2014). Multiple testing was corrected for using Holm's method (Holm, 1979). Cost-utility analyses were conducted with QALYs and societal costs over the 14 month follow-up period. Specifically, QALYs and costs were averaged over the 100 imputed datasets. Subsequently, a bootstrap procedure simulating 1000 samples drawn from the average imputation sample was conducted in Microsoft Excel to estimate the uncertainty regarding mean costs and QALYs. Mean costs and QALYs per study condition were used to calculate the incremental net benefit (INB) for each condition. To calculate the INB, first, society's willingness to pay (WTP) for one extra year lived in perfect health (i.e., 1 QALY) was multiplied with the QALY gain in a condition, which expresses the effect in monetary terms. Subtracting the costs for this condition resulted in its INB. The 1000 INBs for each condition were used to calculate the probability of a condition to be cost-effective compared to the other conditions for a range of WTP values. In the Netherlands the willingness to pay is assumed to vary between €20,000 per QALY for interventions in the context of 'low disease burden' to €80,000 per QALY in the context of severe diseases (Zwaap et al., 2015). To accommodate all relevant WTP values, the current study explored values ranging from €0 to €100,000. The results were presented in cost-utility acceptability curves for the four conditions separately.

Four sensitivity analyses were conducted to investigate the robustness of the results. Specifically, using the average imputation sample, cost-utility analyses were repeated with (1) capability values based on ICECAP-A scores resulting in cost-capability acceptability curves, (2) QALYs based on utility scores obtained from the visual analogue scale of the EQ-5D-5L (raw scores divided by 100) and (3) direct health care costs only instead of societal costs. Lastly, because baseline scores of the EQ-5D-5L were unavailable, a sensitivity analysis (4) was performed where baseline scores of the EQ-5D-5L were estimated using the 4-item Patient Health Questionnaire (PHQ-4). Equipercntile mapping was used to translate baseline PHQ-4 scores into EQ-5D-5L scores. These were then used to calculate adjusted QALYs for the cost-utility acceptability curves. Details on the mapping procedure can be found in Appendix F.3.

Results

Participants

In total, 355 participants completed informed consent and the baseline assessment and were included in the analyses. Retention of participants at baseline (T0), post intervention (8

weeks; T1) and 3, 6, 9 and 12 month follow-up (T2-T5) was 355 (100%), 280 (78.9%), 252 (71.0%), 244 (68.7%), 233 (65.6%) and 242 (68.2%) respectively. Study drop-out rates did not differ between conditions at post intervention, $\chi^2(3) = 3.99, p = .26$, or 12 month follow-up, $\chi^2(3) = 4.90, p = .18$. No differences in stopping with the intervention between the three active interventions were found, $\chi^2(2) = 1.24, p = .54$. Baseline characteristics of participants are presented in Table 2.

Table 2. Baseline characteristics of participants

Characteristics	Feedback (<i>N</i> = 88)	Feedback + Expert- patient support (<i>N</i> = 90)	Expert- patient support (<i>N</i> = 87)	Waiting list (<i>N</i> = 90)	Total sample (<i>N</i> = 355)
Gender					
Female (%)	89 (98.9)	82 (93.2)	84 (96.6)	88 (97.8)	343 (96.7)
Male (%)	1 (1.1)	5 (5.7)	1 (1.1)	2 (2.2)	9 (2.5)
Other (%)	0 (0.0)	1 (1.1)	2 (2.3)	0 (0.0)	3 (0.8)
Nationality					
Dutch (%)	80 (88.9)	78 (88.6)	80 (92.0)	81 (90.0)	319 (89.9)
Belgian (%)	9 (10.0)	9 (10.2)	6 (6.9)	8 (8.9)	32 (9.0)
Other (%)	1 (1.1)	1 (1.1)	1 (1.1)	1 (1.1)	4 (1.1)
Education					
Low (%)	12 (13.3)	5 (5.6)	12 (13.7)	18 (20.5)	47 (13.3)
Middle (%)	31 (34.4)	33 (37.5)	34 (39.0)	35 (39.3)	133 (37.6)
High (%)	47 (52.2)	50 (56.8)	41 (47.1)	36 (40.4)	174 (49.2)
Treatment history for ED					
Yes (%)	54 (54.0)	46 (52.3)	53 (60.9)	49 (54.4)	202 (56.9)
No (%)	36 (36.0)	42 (47.7)	34 (39.1)	41 (45.6)	153 (43.1)
Self-reported diagnosis status					
Officially diagnosed with ED	52 (59.1)	60 (66.7)	52 (59.8)	58 (64.4)	222 (62.5)
No diagnosis, but assumed to have ED	24 (27.3)	22 (24.4)	23 (26.4)	22 (24.4)	91 (25.6)
Eating problems, but likely no ED diagnosis	12 (13.6)	8 (8.9)	12 (13.7)	10 (11.1)	42 (11.8)
Marital status					
Married/living together (%)	22 (24.4)	20 (22.7)	26 (29.9)	30 (33.3)	98 (27.6)
Living alone (%)	66 (73.3)	68 (77.3)	58 (66.7)	58 (64.4)	250 (70.4)
Divorced (%)	1 (1.1)	0 (0.0)	3 (3.4)	2 (2.2)	6 (1.6)
Widow (%)	1 (1.1)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)
Age [Years]	28.0 (1.7)	28.3 (10.4)	26.8 (9.4)	28.1 (12.4)	27.8 (10.8)

Weight [kg]	64.0 (21.0)	62.2 (18.3)	63.6 (22.0)	64.7 (23.4)	63.6 (21.2)
Height [cm]	169.9 (7.2)	168.5 (6.9)	169.7 (7.1)	169.5 (6.9)	169.4 (7.0)
Years with ED	10.1 (9.1)	10.3 (8.8)	8.6 (8.2)	11.4 (12.0)	10.1 (9.7)
Internet usage [hours per day]	4.2 (2.6)	3.7 (2.2)	3.9 (2.3)	3.4 (2.8)	3.8 (2.5)
Eating disorder symptoms (EDE-Q)	3.9 (1.1)	4.1 (1.1)	4.3 (1.0)	4.3 (1.0)	4.1 (1.0)

Note. Data are presented as means (standard deviation) unless indicated otherwise.
ED = Eating Disorder; EDE-Q = Eating Disorder Examination Questionnaire.

Quality of life

EQ-5D-5L utility values and ICECAP-A index scores for all measurement points, as well as QALYs and capability values over the total study duration (12 months + 8 weeks) are presented in Table 3. Average QALYs were highest in the Feedback with expert-patient support condition and lowest for the care-as-usual condition. However, no significant differences in QALYs between the four conditions were found. Similarly, no differences in improvements on capabilities as derived from the ICECAP-A between the four conditions were found.

Costs

Intervention costs, health care costs and non-health care costs are presented in Table 4. Intervention costs were significantly higher in conditions with expert-patient support. Lowest health care costs were found in the Feedback only condition, while highest costs were found in the care-as-usual condition. The relatively low health care costs in the Feedback only condition could mostly be attributed to fewer participants being hospitalized in that condition. Average societal costs per participant over the study duration were again lowest in the Feedback only condition and highest in the care-as-usual condition. Although the omnibus test was significant, after a Holm correction for multiple testing, pooled ANOVA tests revealed no significant difference between the four conditions for health care costs and societal costs.

Table 3. Means (standard errors) of utilities, QALYs and capabilities

Category	Feedback (<i>N</i> = 88)	Feedback + Expert- patient support (<i>N</i> = 90)	Expert- patient support (<i>N</i> = 87)	Waiting list (<i>N</i> = 90)	Total sample (<i>N</i> = 355)	Pooled F-statistic
EQ-5D-5L utilities						
Post intervention (T1; 8 weeks)	0.68 (0.03)	0.68 (0.03)	0.61 (0.03)	0.58 (0.03)	0.64 (0.01)	<i>F</i> (3, 333) = 3.01, <i>p</i> = .03
3-month follow-up (T2)	0.62 (0.03)	0.68 (0.03)	0.60 (0.03)	0.58 (0.04)	0.62 (0.02)	<i>F</i> (3, 326) = 1.88, <i>p</i> = .13
6-month follow-up (T3)	0.69 (0.03)	0.69 (0.03)	0.62 (0.03)	0.60 (0.04)	0.65 (0.02)	<i>F</i> (3, 321) = 1.90, <i>p</i> = .13
9-month follow-up (T4)	0.61 (0.04)	0.65 (0.04)	0.63 (0.04)	0.62 (0.04)	0.63 (0.02)	<i>F</i> (3, 311) = 0.19, <i>p</i> = .91
12-month follow-up (T5)	0.66 (0.03)	0.71 (0.03)	0.66 (0.03)	0.64 (0.04)	0.67 (0.02)	<i>F</i> (3, 317) = 0.61, <i>p</i> = .61
EQ-5D-5L Visual Analogue Scale utilities						
Post intervention (T1; 8 weeks)	0.60 (0.02)	0.59 (0.02)	0.55 (0.02)	0.55 (0.02)	0.57 (0.01)	<i>F</i> (3, 331) = 1.05, <i>p</i> = .37
3-month follow-up (T2)	0.55 (0.02)	0.56 (0.02)	0.55 (0.02)	0.57 (0.02)	0.56 (0.01)	<i>F</i> (3, 333) = 0.22, <i>p</i> = .88
6-month follow-up (T3)	0.61 (0.02)	0.57 (0.02)	0.54 (0.02)	0.58 (0.02)	0.57 (0.01)	<i>F</i> (3, 325) = 1.77, <i>p</i> = .15
9-month follow-up (T4)	0.57 (0.02)	0.59 (0.02)	0.56 (0.02)	0.59 (0.02)	0.58 (0.01)	<i>F</i> (3, 318) = 0.43, <i>p</i> = .73
12-month follow-up (T5)	0.60 (0.02)	0.61 (0.02)	0.57 (0.02)	0.59 (0.02)	0.59 (0.01)	<i>F</i> (3, 326) = 0.52, <i>p</i> = .67
ICECAP-A capability values						
Post intervention (T1; 8 weeks)	0.69 (0.02)	0.68 (0.02)	0.63 (0.02)	0.65 (0.03)	0.66 (0.01)	<i>F</i> (3, 329) = 1.40, <i>p</i> = .24
3-month follow-up (T2)	0.68 (0.02)	0.69 (0.02)	0.62 (0.02)	0.66 (0.03)	0.66 (0.01)	<i>F</i> (3, 324) = 1.48, <i>p</i> = .22
6-month follow-up (T3)	0.70 (0.03)	0.69 (0.03)	0.64 (0.03)	0.65 (0.03)	0.67 (0.03)	<i>F</i> (3, 316) = 1.01, <i>p</i> = .39
9-month follow-up (T4)	0.67 (0.03)	0.68 (0.03)	0.65 (0.03)	0.69 (0.03)	0.67 (0.01)	<i>F</i> (3, 305) = 0.44, <i>p</i> = .72
12-month follow-up (T5)	0.72 (0.02)	0.72 (0.03)	0.64 (0.03)	0.72 (0.03)	0.70 (0.01)	<i>F</i> (3, 313) = 2.09, <i>p</i> = .10
Total QALYs	0.75	0.78	0.71	0.69	0.74	<i>F</i> (3, 337) =
EQ-5D-5L ^a	(0.03)	(0.03)	(0.03)	(0.03)	(0.02)	1.87, <i>p</i> = .14
Total QALYs	0.67	0.64	0.69	0.65	0.66	<i>F</i> (3, 339) =
EQ-5D Visual Analogue Scale ^a	(0.02)	(0.02)	(0.02)	(0.02)	(0.01)	0.74, <i>p</i> = .53
Total capability values ICECAP-A ^a	0.79 (0.02)	0.79 (0.02)	0.73 (0.02)	0.77 (0.03)	0.77 (0.01)	<i>F</i> (3, 328) = 1.31, <i>p</i> = .27

SE=Standard error.

^a Calculated over the entire 14-month study duration.

Table 4. Means (standard error) of costs per study condition over the course of 14 months in 2021 euros with the percentage of participants that incurred the costs

Category	Mean costs per participant (SE) [% of participants incurring costs]					Pooled F-statistic
	Feedback (N = 88)	Feedback + Support (N = 90)	Expert-patient support (N = 87)	Waiting list (N = 90)	Total sample (N = 355)	
Total intervention costs	3 (0)	65 (4)	72 (4)	0 (0)	35 (2)	$F(3, 351) = 226.45, p < .001$
Health care costs						
General practitioner	288 (33) [93%]	260 (39) [92%]	215 (28) [87%]	320 (36) [95%]	271 (17) [92%]	$F(3, 335) = 1.69, p = .17$
Dietician	122 (25) [52%]	217 (37) [61%]	133 (31) [53%]	202 (38) [58%]	169 (17) [56%]	$F(3, 334) = 2.12, p = .10$
Psych ^a – mental health care	1483 (298) [55%]	1745 (369) [58%]	2723 (448) [70%]	1810 (312) [66%]	1936 (184) [62%]	$F(3, 336) = 2.23, p = .08$
Psych ^a - independent	1005 (186) [57%]	869 (153) [59%]	1195 (222) [64%]	1052 (188) [63%]	1029 (95) [61%]	$F(3, 335) = 0.52, p = .67$
Psych ^a - hospital	617 (181) [40%]	241 (75) [32%]	714 (188) [43%]	549 (138) [43%]	528 (76) [39%]	$F(3, 340) = 1.83, p = .14$
Medical specialist	220 (50) [46%]	225 (47) [56%]	204 (43) [46%]	297 (53) [57%]	237 (24) [52%]	$F(3, 332) = 0.73, p = .53$
Emergency department	74 (68) [3%]	28 (27) [5%]	9 (11) [3%]	54 (26) [7%]	41 (20) [5%]	$F(3, 346) = 0.52, p = .67$
Day treatment - mental health care	827 (264) [25%]	642 (244) [24%]	1102 (338) [33%]	1542 (547) [28%]	1029 (186) [27%]	$F(3, 339) = 1.14, p = .33$
Hospitalization - mental health care	2557 (1105) [14%]	4385 (1758) [19%]	4417 (1908) [20%]	9158 (2524) [30%]	5150 (967) [21%]	$F(3, 338) = 2.24, p = .08$
Hospitalization – hospital	528 (282) [15%]	1537 (589) [19%]	1923 (900) [19%]	2836 (952) [27%]	1711 (370) [20%]	$F(3, 338) = 1.71, p = .16$
Total health care costs	7722 (1535)	10215 (2281)	12705 (2808)	17820 (3349)	12135 (1316)	$F(3, 340) = 2.81, p = .04$
Non-health care costs						
Absenteeism	1456 (388) [38%]	2602 (690) [49%]	1925 (514) [39%]	2117 (518) [43%]	2029 (271) [42%]	$F(3, 328) = 0.78, p = .51$
Presenteeism	3142 (630) [59%]	5122 (937) [77%]	1968 (437) [62%]	2588 (567) [68%]	3216 (342) [66%]	$F(3, 333) = 4.16, p = .01$
Substitution of unpaid work	4421 (870) [79%]	6042 (1183) [82%]	7022 (1489) [88%]	5954 (1139) [79%]	5858 (586) [82%]	$F(3, 337) = 0.80, p = .49$
Total societal costs	16741 (2023)	23980 (3277)	23620 (3365)	28479 (3736)	23238 (1612)	$F(3, 340) = 2.34, p = .07$


SE=Standard error.

^a Psychologist, psychotherapist or psychiatrist.

Cost-effectiveness

Cost-utility acceptability curves are presented in Figure 1. For values of the WTP for one additional QALY between €0 and €100,000, offering the Feedback only condition had the highest probability of being efficient for the four alternatives (66%-86%). In other words, between the four conditions, Feedback only had the highest probability of having the largest INB across the 1000 bootstrap samples, regardless of the WTP. At very high WTP values, the probability of the combination of Feedback with expert-patient support to be efficient compared to the alternatives increased, but still did not exceed the Feedback only condition. The care-as-usual condition had a probability of up to 13% of being optimal compared to the alternatives for all WTP values. This probability was around 1% across all WTP values for the expert-patient support only condition.


Figure 1. Cost-utility acceptability curves with EQ-5D QALYs for the four study conditions derived from 1000 bootstrap samples



Sensitivity analyses

As can be deduced from Figure 2, results were highly similar for cost-capability acceptability curves, where gains for a particular condition were measured as capability values as assessed with the ICECAP-A. Specifically, the Feedback only condition had the highest probability of being efficient compared to the other three conditions across all WTP values (72%-86%). A second sensitivity analysis using the visual analogue scale of the EQ-5D-5L to assess QALYs also showed results comparable to the main analysis. Third, cost-utility acceptability curves of the EQ-5D-5L with direct health care costs only showed the Feedback only condition to have the highest probability of being efficient (51%-78%) compared to the three alternatives for WTP values of €60,000 or less. For WTP values between €70,000 and €100,000, Feedback

Figure 2. Cost-capability acceptability curves with ICECAP-A capability values for the four study conditions derived from 1000 bootstrap samples



with expert-patient support had the highest probability of being efficient among the four conditions (52%-57%). Fourth, when baseline values of the EQ-5D-5L were estimated using equipercenile mapping with the PHQ-4, an almost identical pattern to the main analysis emerged with the Featback only condition having the highest probability to be efficient compared to the three other conditions across all WTP values (67%-95%). Cost-effectiveness acceptability curves of the last three sensitivity analyses can be found in Appendix F.4.

Discussion

In the current study, an economic evaluation based on a randomized controlled trial covering a period of 14 months was conducted comparing (1) a fully automated internet-based intervention 'Featback', (2) online support by expert patients via mail and chat and (3) the combination of these to (4) care as usual for people with eating disorders. Primarily, from a societal perspective, the Featback intervention had the highest probability of being efficient for a wide range of WTP values compared to the three other conditions. Secondly, expert-patient support alone and care as usual had very low probabilities of being efficient compared to the alternatives over the whole range of explored WTP values. Lastly, the combination of Featback and expert-patient support was more efficient than a care-as-usual condition for WTP values over €20,000, but less than Featback alone. The results suggest that, between the four investigated conditions, Featback is the intervention of choice from a (societal) economic perspective. Despite severe and long-lasting symptoms, 43% of participants in the sample never received treatment for their eating problems, demonstrating the potential of internet-based interventions to reach an underserved population. Notably, as Featback and

expert-patient support are brief interventions, it might be that their impact on quality of life or health care and societal costs is more distinct for people with less severe symptoms or at the beginning stages of their eating disorder, but it proved difficult to reach this group. Furthermore, while around 97% of the Dutch population older than 12 years has internet access, some individuals with eating disorder symptoms cannot be reached through internet or find it challenging to work with and require a different approach. However, implementing the unguided Featback intervention could help to reach individuals with an eating disorder who currently do not receive appropriate care and is likely to lead to similar or slightly better quality of life while reducing costs for society, compared to not implementing it.

The findings are in line with the two studies mentioned in the introduction, cautiously indicating online interventions for eating disorders to be cost-effective compared to care as usual (Akers et al., 2021; Kass et al., 2017). A previous trial concerning Featback indicated that Featback with or without guidance from a psychologist had higher probabilities of being efficient compared to a care-as-usual condition (Aardoom, Dingemans, Spinhoven, et al., 2016). In the current trial, the automated feedback messages were improved and more personalized towards users and support from a psychologist was replaced with expert-patient support, in an attempt to increase the effectiveness of and satisfaction with the interventions. Unexpectedly, conditions with expert-patient guidance were less efficient than Featback alone for all acceptable WTP values. The contrast with the previous trial, where Featback was equally efficient with and without therapist support, might partly be explained by the improvements to the automated monitoring system, possibly increasing its effectiveness. However, this is unlikely given that the combination condition also included the improved monitoring system, but was outperformed by Featback alone. Another explanation of the favorable probabilities of the Featback only condition in the cost-utility acceptability curves were the (non-significantly) lower costs for several categories, resulting in non-significantly lower total costs. Mainly, a low percentage of people in the Featback condition was hospitalized compared to the other conditions. The finding suggests that brief weekly monitoring and feedback messages can prevent hospitalization. Self-monitoring has been found to be important in preventing psychiatric hospitalization (Ådnanes et al., 2020). However, if the Featback monitoring system would have this effect, lower hospitalization rates in the combination condition would also be expected. Therefore, other explanations why the Featback condition had favorable results, such as chance, cannot be ruled out. Looking across mental disorders, adding guidance appears to increase the probability of being cost-effective compared to unguided alternatives (Donker et al., 2015). However, given the scarcity of evidence, more research directly comparing guided and unguided internet-based interventions is required for decisive conclusions.

More generally, an increasing number of studies is substantiating the claim that internet interventions are likely to be cost-effective compared to care as usual for a number of mental disorders (Ahern et al., 2018; Donker et al., 2015; Hedman et al., 2012; Paganini et al., 2018), but evidence is mixed (Kolovos et al., 2018). The systematic reviews and meta-analysis highlight the need to continue economic evaluation research, since heterogeneity in intervention content and application of guidance make it difficult to reach definitive conclusions. For eating disorder treatment, current evidence indicates that online self-monitoring is likely an efficient alternative to usual care, while more information is needed on whether and how to add guidance. Concordantly, internet-based interventions such as Featback have the

potential to help individuals currently not reached by traditional treatment options, while being worth the investment.

Strengths and limitations

The current study had several strengths, including a 12-month follow-up period with a societal perspective, a large sample size, several sensitivity analyses and multiple imputation procedures to handle missing data. Some limitations can also be noted. First, baseline values of quality of life and wellbeing questionnaires were unavailable. Although a sensitivity analysis with estimated baseline scores produced similar results, the missing values may have led to a slight underestimation of the QALYs and capability values in the active interventions. Relatedly, it could be worthwhile to study which generic preference-based measures are sensitive in eating disorder populations, as both the EQ-5D and ICECAP might be limited in this regard. A second limitation pertains to missing data due to dropout of the current sample. While missing data were handled adequately, imputing cost data was challenging since for some categories only a small percentage of people incurred relatively high costs while others incurred no costs. Thirdly, although many costs were accounted for, assessed direct and non-health care costs were not exhaustive. For example, medication costs and costs attributable to alleviating symptoms were not captured, which may have influenced results slightly. Lastly, cost data were based on self-reported health care visits and work productivity over a period of 4 weeks. This may have introduced recall bias.

Conclusion

A brief fully automated internet-based self-help program (Featback) for eating disorders was found to be efficient compared to care as usual. Results suggest that such interventions may be especially valuable as a first step in a stepped-care model of eating disorder treatment, as it is preferable over a do-nothing approach. Implementing highly scalable and low-threshold interventions such as Featback would not only benefit individuals suffering from an eating disorder, but society as a whole. Mental health professionals and researchers would profit from further investigating how to widely disseminate such interventions to optimize the potential benefits, both in terms of effects and costs. Furthermore, subsidy providers, policy makers and health insurances should consider wider funding to make installment of evidence-based internet interventions for eating disorders possible, as they appear to be worth the investment.

