



Universiteit  
Leiden  
The Netherlands

## **A revision of the CH<sub>4</sub> concentration in Jupiter's upper atmosphere from near-IR measurements**

López-Puertas, M.; Sanchez Lopez, A.; García-Comas, M.; Funke, B.; Fouchet, T.; Snellen, I.A.G.

### **Citation**

López-Puertas, M., Sanchez Lopez, A., García-Comas, M., Funke, B., Fouchet, T., & Snellen, I. A. G. (2022). A revision of the CH<sub>4</sub> concentration in Jupiter's upper atmosphere from near-IR measurements. *Epsc Abstracts*, (16), EPSC2022-975. doi:10.5194/epsc2022-975

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3561778>

**Note:** To cite this publication please use the final published version (if applicable).



## A revision of the CH<sub>4</sub> concentration in Jupiter's upper atmosphere from near-IR measurements

**Manuel López-Puertas**<sup>1</sup>, Alejandro Sánchez-López<sup>2</sup>, Maya García-Comas<sup>1</sup>, Bernd Funke<sup>1</sup>, Thierry Fouchet<sup>3</sup>, and Ignas Snellen<sup>2</sup>

<sup>1</sup>Instituto de Astrofísica de Andalucía, CSIC, Granada, Spain (puertas@iaa.es)

<sup>2</sup>Leiden Observatory, Leiden University, Leiden, The Netherlands (alexsl@strw.leidenuniv.nl)

<sup>3</sup>Observatoire de Paris-PSL, Paris, France (Thierry.Fouchet@obspm.fr)

Methane has a driving role in establishing the thermal structure of Jupiter's upper atmosphere. Hence, to know its spatial distribution, both in altitude and latitude, is essential for understanding that atmospheric region.

The abundance of CH<sub>4</sub> in Jupiter's upper atmosphere has been (or will soon be) derived using a number of different methods, e.g., solar occultation, the measurements of He and Ly-alpha airglow, and from the observations of near-IR emission near 3.3 μm (ISO/SWS, JUNO/JIRAM so far and, in the near future, from JWST/NIRSpec).

The retrieved values derived from these techniques show rather different values, particularly around the homopause. Even different studies of the same ISO/SWS spectral radiance yield very different CH<sub>4</sub> volume mixing ratio profiles.

Here, we will discuss the important role and the adequate use of the collisional relaxation of the CH<sub>4</sub> high-energy levels when deriving CH<sub>4</sub> abundances near the mesopause from NIR emission radiances. Further, we will also show some results on the important role of spectroscopic data of the CH<sub>4</sub> 3.3 μm hot bands for those analyses. Those studies will be presented for the particular case of the ISO/SWS measurements, although they are equally applicable to other measurement sets such as those of JUNO/JIRAM and JWST/NIRSpec.