



Universiteit
Leiden
The Netherlands

High-resolution transmission spectroscopic studies of hot and ultra-hot Jupiters

Stangret, M.B.; Palle Bago, E.; Orell-Miquel, J.; Esparza-Borges, E.; Casasayas Barris, N.

Citation

Stangret, M. B., Palle Bago, E., Orell-Miquel, J., Esparza-Borges, E., & Casasayas Barris, N. (2022). High-resolution transmission spectroscopic studies of hot and ultra-hot Jupiters. *Bulletin Of The American Astronomical Society*, (5). Retrieved from <https://hdl.handle.net/1887/3562720>

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3562720>

Note: To cite this publication please use the final published version (if applicable).

Bulletin of the AAS • Vol. 54, Issue 5

High-resolution transmission spectroscopic studies of hot and ultra-hot Jupiters.

**Monika B. Stangret¹ Enric Pallé Bago² Jaume Orell-Miquel²
Emma Esparza-Borges² Núria Casasayas-Barris³**

¹Instituto de Astrofísica de Canarias, ²Instituto de Astrofísica de Canarias, ³Leiden Observatory

Published on: Jun 20, 2022

URL: <https://baas.aas.org/pub/2022n5i102p304>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Hot and ultra-hot Jupiters, gas-giant planets with short orbital period, and hot, extended atmospheres, are most suitable objects in order to study the chemical composition of the atmospheres using emission and transmission spectroscopy. Their tidally-locked nature leads to big differences between night and day side temperatures, and in consequence differences in the atmospheric chemical composition on both terminators. Thanks to new generation of high-resolution ground based spectrographs such as HARPS-N, HARPS, CARMENES, and ESPRESSO, we are able to perform those studies.

During our talk we want to present a complex analysis of the chemical composition of a set of hot and ultra-hot Jupiters using high-resolution spectroscopic observations. Using cross-correlation method and transmission spectroscopy around single lines, we searched for a set of atoms and molecules in their atmospheres. Additionally, by analysing the Rossiter-McLaughlin effect we measured the obliquity of those systems.

Our work determines the temperature in which we differentiate between hot and ultra-hot Jupiters, and show the importance of complex studies of those objects using broad set of chemical species models.