



Universiteit
Leiden
The Netherlands

In between looking and seeing: recognition, referral and assessment of children and adolescents' mental health problems at the interface of primary care and secondary mental healthcare

Aydin, S.

Citation

Aydin, S. (2023, February 23). *In between looking and seeing: recognition, referral and assessment of children and adolescents' mental health problems at the interface of primary care and secondary mental healthcare*. Retrieved from <https://hdl.handle.net/1887/3564429>

Version: Publisher's Version


License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3564429>

Note: To cite this publication please use the final published version (if applicable).



We keep asking 'how to reduce the gap between science and practice' but
how about widening the common space between both?



Chapter 4

**The diagnostic process from primary care to child and adolescent mental healthcare services:
The incremental value of information conveyed through referral letters, screening questionnaires and structured multi-informant assessment**

Published as: Aydin, S., Siebelink, B.M., Crone, M.C., Van Ginkel, J., Numans, M.E., Vermeiren, R.R.J.M., & Westenberg, P.M. The diagnostic process from primary care to child and adolescent mental healthcare services: The incremental value of information conveyed through referral letters, screening questionnaires and structured multi-informant assessment. *British Journal of Psychiatry Open*, 2022, 8(3), E81, doi:10.1192/bjo.2022.47

The formulation of a clinical diagnosis is critical to the child and adolescent mental healthcare (CAMH).¹⁻³ The current approaches for the diagnostic process include the judgement of a clinician or the use of structured assessment instruments. Four decades of research support the use of structured instruments, which results in more consistent application of diagnostic criteria, a decrease in information variance and bias, and improved recognition of less obvious or secondary conditions.⁴⁻⁶ Clinical and evidence-based assessment (EBA) guidelines therefore recommend integration of both methods, to benefit from the nuance and parsimony associated with clinical judgement, combined with the accuracy and reliability intrinsic to structured assessment.⁷⁻⁸ As in clinical practice with stepped-care and matched-care approaches, assessment is conducted in sequential stages, with EBA the question is raised as to whether instruments meaningfully contribute to the diagnostic work-up and how far each additional information step overlaps. Although the various instruments have been studied for value as stand-alone measures,⁷ less is known about the incremental value of the various nodes of information. Given the tension between efficiency of information gathering and reliability in the diagnostic process,⁹ a better understanding is needed of the value of a validated diagnostic work-up; in this case, a work-up that captures the combined benefits of structured assessment and clinical judgement, suggesting potential for use at the interface between primary and secondary CAMH. Accordingly, the aim of the present study was to investigate the *incremental* value of routinely gathered successive assessments. We investigated the added value of referral letters, a screening questionnaire and a structured multi-informant assessment gathered during the registration procedure at an academic centre for child and adolescent psychiatry

The diagnostic procedure

In several countries it is standard practice for CAMH registration to take place via front-line practitioners such as paediatricians or general practitioners. If a decision is made, based on screening or clinical judgement, to refer to CAMH, a referral letter indicating the probable mental health diagnosis forms a bridge to CAMH. For many children and adolescents, referral letters represent the only form of information transfer from the referrer, and may contribute to the diagnostic and treatment process in CAMH. Although many professionals in the field believe that referral letters have no clinical value, in a recent study, we found that 42-93% of youth reasons for referral saw no change in later psychiatric diagnosis.¹⁰

Although these numbers are substantial, we also observed considerable variation between disorder groups, with internalizing problems in particular showing a relatively poor detection accuracy.

In EBA, a decision to refer should follow administration of a screening instrument. This procedure allows for the common false positives of screening instruments to be corrected by clinician judgement, and acknowledges that screening often helps improve detection of less obvious problems such as internalising disorders, thereby improving adequate referrals and access to treatment. Regrettably, the use of screening instruments is infrequent, a problem often attributed to the limited time available for patient consultation.¹¹ Many of the current short screening questionnaires were specifically developed to address this problem. Unintendedly, development of these questionnaires may have further limited their implementation, because understanding the pros and cons of the wide array of current screening instruments, together with interpretation of outcomes, has become more challenging.¹¹⁻¹⁴ A recent review of accessible CAMH instruments identified 672 questionnaires, of which only four broad screening instruments qualified as brief, free, accessible, and with excellent psychometric characteristics.¹⁴ One of these instruments is the strengths and difficulties questionnaire (SDQ), available in translation in over 70 languages.¹⁵ The SDQ was found to be as reliable and feasible as the much lengthier Achenbach scales (the Youth self report (YSR), Child behaviour checklist (CBCL) and Teacher report form (TRF)) that are frequently used in many European countries.¹⁶⁻¹⁸ The developers of the SDQ proposed using the instrument before a clinical appointment as a guide to decision-making.¹⁹ However, regarding recognition of emotional problems, studies suggest that the SDQ might be insufficient, a problem likely related to the limited number of questions in the scale, to differences in study samples and to general difficulties in detecting internalizing disorders.^{15 20}

The detection of mental health problems, including internalizing problems, often improves with the use of more extensive assessment methods. In EBA, more extensive assessment methods are in fact recommended in the case of individuals with high scores during screening. The Development and Well-Being Assessment (DAWBA) instrument combines the responses of various informants' (adolescents, parents and/or caregivers and teachers) to

closed-ended questions into so-called DAWBA band scores that indicate the likelihood of a child having any of 17 common mental health disorders.^{19 21} The DAWBA band scores were envisioned as a way to avoid the costly involvement of a clinician and to be a pragmatic solution for common issues at the point of care. Nonetheless, the value of DAWBA bands when accounted for the value of screening and clinical judgement in primary care is not yet investigated.

As part of the DAWBA, informants are also prompted to describe their problems and the context of their problems in their own words. These are then evaluated by a clinician who integrates the various factors to form a relatively nuanced image without the high cost of a full interview with a specialist clinician. DAWBA clinician ratings (CRs) were found to be conservative regarding the number of diagnoses made when compared with elaborate diagnostic interviews.²² Studies of clinician rated DAWBA found that it was useful in reducing unnecessary referral for externalizing disorders, and that they highlighted internalizing disorders that would not have been detected otherwise.^{23 24} Nevertheless, the exact extent to which clinician ratings supplements information from a primary care clinician , screening results and automatised DAWBA probability band scores remains an important but unanswered question.

Aims

In summary, the feasibility and psychometric properties of the DAWBA and SDQ have been individually well-researched in community, clinical and research settings in various European countries. However, less information is available regarding the predictive value of instruments when taking into account the usual overlap of information gained during successive steps in EBA. The aim of the present study was to determine both the unique and incremental predictive values for four sources of information in predicting a medical record consensus diagnosis: referral letters, a screening questionnaire (SDQ¹⁵), a more elaborate structured assessment (DAWBA band scores¹⁹) and the remote evaluation of structured and unstructured responses by a clinician (the clinician rated DAWBA). We hypothesized that each instrument would show incremental value in predicting the classification of five disorder groups commonly treated in CAMH: anxiety, depression, autism spectrum disorders (ASD), attention-deficit hyperactivity disorder (ADHD), and behavioural disorders.

Methods

Data source and procedure

The starting point for the sample consisted of children and adolescents who were referred to Leiden University Medical Centre Curium (LUMC Curium). LUMC Curium is an inpatient and outpatient mental health clinic delivering specialized care to young people aged 3 to 18 years.

About 70% of the yearly caseload at the institution consists of outpatient referrals that follow a routine procedure including referral letters, the SDQ and DAWBA. The remainder consists of inpatient referrals that follow a referral intake procedure adapted to cases in need of urgent evaluation, in which case questionnaires are not completed at registration. We included young people who I) registered between January 2015 and December 2017, II) followed the routine procedure including the SDQ and DAWBA, and III) had an accessible referral letter in the medical record. The procedures used to extract and code referral letters are described in detail in our recent publication on referral letters general practice.¹⁰ To briefly summarize, using an iterative process we created a manual to extract and code text in referral letters. The manual was then tested for inter-rater reliability by authors S.A., M.R.C., B.M.S. and P.M.W. ($\kappa=.77$ to $\kappa=.90$). We did not differentiate symptoms indicated in referral letters from suggested diagnoses. For instance, when a referral letter reported “treatment for anxiety disorders?” or “fearful”, both were coded as an indicator of the category anxiety disorders and related problems. Multiple indications were often found in referral letters and were thus coded. However, fewer than 20% of referral letters indicated more than four problems,¹⁰ which was also the case in the current sample.

The LUMC Medical Ethical committee waived a need for informed consent because of the retrospective nature of the study (approval number G18.080). Furthermore, the data management plan was approved by the scientific committee of the LUMC Departments of Public Health and Primary Care, LUMC Curium Child and Adolescent Psychiatry, and the Institute of Psychology of Leiden University.

Measures

All measures were extracted from medical records. We extracted referral letters as they were scanned and filed in individual patient medical records. The SDQ, structured DAWBA

data and classifications that are also outcome measure were extracted simultaneously from the medical record system.²⁵

In The Netherlands, only a healthcare professional can make a formal referral to youth and adolescent psychiatry, which then proceeds via either general practice, specialised healthcare (hospitals) or youth welfare offices (also called local youth teams). We did not include the type of professional as a covariate in the main analyses, as initial logistic regression analyses showed wide confidence intervals and no statistically significant differences between the various types of referrers.

Structured assessment: SDQ and DAWBA

During registration, families are provided with unique login codes for the online DAWBA package, which can be completed by up to two parents or caregivers, the young person themselves (if aged >11 years) and up to two teachers. The package always starts with the SDQ, and then moves on to the DAWBA instrument. Rules regarding skipping come into play when an informant shows low scores on a conceptually-related SDQ scale and provides negative answers to a gate-keeping question at the beginning of each DAWBA chapter.¹⁹ In the DAWBA package, SDQ scale scores and DAWBA probability band scores are generated for each informant individually and subsequently integrated into an overall SDQ score for each scale (0, 1, 2) and a DAWBA probability band score for each chapter (0 to 5). The cut-off scores and rules concerning integration of informant's scores can be found at www.sdqinfo.org and www.dawba.net. If not otherwise specified, we used integrated scores for all analyses. To analyse whether each assessment method indicated the presence of a disorder group, we dichotomized scores by separating the upper two scores from the lower score(s).^{21 24 26}

SDQ

The SDQ covers four problem areas (emotional, conduct, hyperactivity and peer problems scales) across 20 items, asks about children's strengths in five items (prosocial scale), and the impact and burden of problems in eight items. Informants rate items on a three-point Likert scale (0= not true, 1=somewhat true, 2=certainly true), with higher scores indicating more problems. Although the SDQ was not formally created to give indications of a probable autism spectrum disorder (ASD), in a later study Goodman et al²⁷ proposed use of a difference score by subtracting the total for the peer problems scale from the score for the

prosocial scale. We calculated this difference score solely based on parental scores, as the few studies available suggest that parents show the highest accuracy in detecting ASD.^{20 28 29}

DAWBA probability band scores

The DAWBA¹⁹ estimates the likelihood of the presence of 17 common mental health disorders. These so-called probability bands are automatically generated in the online DAWBA environment by integrating various informant responses to closed-ended questions.²¹ The questions are linked to the DSM criteria and result in probability band scores of 0, 1, 2, 3, 4 and 5, corresponding to prevalences found in the original British epidemiologic sample and approximating likelihoods <0.1% , 0.5%, 3%, 15%, 50% and >70%.²¹ Thus, a probability band score of 5 suggests that 70% or more of the cases with a similar response profile to the British reference sample were found to have that diagnostic outcome. When the DAWBA did not produce a score for a disorder group (e.g. behavioural disorders), we took the highest probability band score among the more specific disorders (i.e. the highest score among conduct and oppositional deviant disorder).²¹

Clinician rated DAWBA

Informants are also prompted to describe problems and their context in their own words. A senior clinical psychologist evaluated the open-ended questions, together with the SDQ and DAWBA probability band results, and scored the likelihood of a disorder on a three-point scale (absent, unsure, present). This final stage facilitates the incorporation of the diverse strands of information to develop a nuanced image without the accompanying cost of visiting a specialist clinician. The next step is to add a short report to a patient medical record in order to guide prioritization of appointments and to prevent tunnel vision during a face-to-face intake. In some study reports, clinician ratings are referred to as a DAWBA research diagnosis. In this paper, however, we use the term clinician rating to prevent confusion with the outcome classification.

Clinical Classification

The primary outcome measure was a patient's digital medical record classification according to the Longitudinal, Expert, and All Data (LEAD) procedure.³⁰ This is a product of all collected information and clinical judgement, including patient and family history, mental health treatment history, structured assessment and, if necessary, process diagnostics and additional assessment methods depending on suspected differential diagnoses.^{31 32} Based on

these insights, a case conceptualization is formed as a basis for treatment initiation and a classification selected and entered into the patient's medical record. Up to five different classifications could be recorded per case and all were extracted for this study.

Missing data

SDQ scale scores were available for all cases and DAWBA band scores were available for 97.7-98.9% of cases, depending on disorder group, but clinician-rated DAWBA data were available for only 52.1% of cases, as DAWBAs were not evaluated by a clinician during the first half of the study period. As this was a result of management decisions and unrelated to our research question, we could assume the data to be missing at random. To reliably estimate missing data, we applied multiple imputation (MI, with $m=100$) using the *mice* package in the R environment.³³⁻³⁷ Multiple imputation creates multiple sets with plausible values for missing cells by drawing values from the observed cases and by predicting from other associated variables in a dataset. Hence, it minimizes bias relative to complete case analysis. Generating multiple datasets enables estimation of the uncertainty in the imputation process as compared with, for example, simple mean imputation. In multiple imputation, it is necessary to balance the number of predictors and observed cases, as with regression analyses in general. Therefore, we limited the number of predictors during multiple imputation such that a minimum number of 15 cases had to be observed for each contributing predictor.

Statistical analysis

In the statistical analysis, we first computed diagnostic metrics such as sensitivity and specificity for each instrument. Next, we inspected youth diagnostic trajectories through the current sequence of four methods. To this end, we cross-tabulated frequencies of positive and negative indications in a four-layer table with each of the methods and the diagnostic outcome. To examine the effect of each added predictor on model fit, likelihood ratio tests³⁸ were performed with the *D3()* function in *mice*.³³ Multiple logistic regression analyses were performed, with each of the five diagnostic groups as the outcome and the assessment methods as the predictor, in order to quantify unique and corrected predictive values. Diagnostic odds ratios (ORs) of the instruments were extracted from the univariable and multivariable logistic regression models.

Results

The sample age ranged between 5 and 18 years (mean=11.08, s.d.=3.45) and 57.4% were boys (Table 1).

Univariable diagnostic metrics

The diagnostic metrics of the assessment methods as standalone measures are depicted in Table 2. The sensitivity and specificity of the successive assessment tools varied per mental health disorder. The value of referral letters in detecting patients with anxiety disorders was relatively low compared with the other disorder groups and to the other instruments: 46.9% of those eventually classified with an anxiety disorder had been indicated as such in referral letters. However, referral letters showed a relatively high specificity in excluding minors without the condition (85.9%). The highest sensitivity regarding anxiety disorders was found for the SDQ (95.1%), but was accompanied by a risk of being overinclusive (specificity 22.9%; false discovery rate 85.2%, supplementary material). The SDQ and referral letters showed the highest sensitivity or specificity, respectively, the DAWBA probability band and the clinician-rated DAWBA showed a more balanced profile.

We found that all instruments except the SDQ performed similarly in discriminating minors with or without depressive disorders (Table 2). In line with earlier studies, the SDQ frequently gave a positive indication in this clinical sample, yet often for the wrong persons (specificity 22.4%).

Upon inspecting the metrics for ASDs, the low number of positive indications by the DAWBA probability band was remarkable. Although the bands indicated ASDs infrequently, they did so for genuine cases, resulting in a high positive predictive value (78.3%, supplementary material) but low sensitivity (9.0%). The SDQ difference score (peer problems - prosocial score, see Methods) showed the highest sensitivity for ASDs as compared with other instruments. In contrast to high false positives for anxiety and depressive disorders, the SDQ showed a better specificity for ASDs (54.7%). Referral letters and clinician rated DAWBA scores showed a fairly even balance of sensitivity and specificity for ASD.

When considering ADHD, most instruments showed values similar to those for ASDs, with the DAWBA probability band showing the best performance in the detection of ADHD (sensitivity 59.3%).

Table 1. Sample Characteristics N=1259

		n (%)
Age, years	5-9	474 (37.6)
	10-14	508 (40.4)
	15-18	277 (22.0)
Gender	Male	723 (57.4)
	Female	536 (42.6)
CGAS	20-40	83 (6.6)
	41-50	503 (40.0)
	51-60	514 (40.8)
	>61	96 (7.6)
	Missing	63 (5.0)
Medical conditions	None classified	958 (76.1)
	Singular	99 (7.9)
	Complex	52 (4.1)
	Missing	150 (11.9)
Number of clinical classifications (comorbidity)	0	175 (13.9)
	1	544 (43.2)
	2	368 (29.2)
	3	125 (9.9)
	4	35 (2.8)
	5-6	12 (1.0)
Type of clinical classifications		
Neurodevelopmental disorders		727 (57.7)
Schizophrenia spectrum and other psychotic disorders		2 (0.2)
Depressive disorders		134 (10.6)
Anxiety disorders		174 (13.8)
Obsessive-compulsive and related disorders		13 (1.0)
Trauma and stressor-related disorders		68 (5.4)
Somatic symptom and related disorders		42 (3.3)
Feeding and eating disorders		54 (4.3)
Elimination disorders		19 (1.5)
Gender dysphoria		7 (0.6)
Disruptive, impulse-control, and conduct disorders		71 (5.6)
Substance-related and addictive disorders		2 (0.2)
Personality disorders		49 (3.9)

Distributions of the clinical classifications in the sample are depicted based on the higher order chapters of the DSM-5 (e.g. Neurodevelopmental disorders). The number of clinical classifications is depicted on the level of the specific disorders (e.g. ADHD and ASD). CGAS= Children's Global Assessment Scale score.

Behavioural disorders were frequently indicated by all instruments yet seldom classified.

This resulted in a very low predictive value. This frequent indication of behaviour problems resulted in relatively high sensitivity (86.4%).

After inspecting single descriptives, we explored frequencies of the instrument's successive positive and negative indications in order to gain insight into the potential of the sequence

Table 2. Two by two cross-tabulation of the instruments per disorder group

		Anxiety disorders		Depressive disorders		ASD		ADHD		Behavioural disorders	
		+	-	+	-	+	-	+	-	+	-
RL	+	38 (46.9)	81 (14.1)	39 (60.0)	73 (12.4)	108 (54.8)	89 (45.2)	114 (55.9)	99 (22.0)	26 (59.1)	156 (25.5)
	-	43 (53.1)	492 (85.9)	26 (40.0)	516 (87.6)	89 (45.2)	361 (80.2)	90 (44.1)	350 (78.0)	18 (40.9)	455 (74.5)
SDQ	+	77 (95.1)	442 (77.1)	62 (95.4)	457 (77.6)	140 (71.1)	204 (45.3)	181 (88.7)	230 (51.2)	38 (86.4)	328 (53.7)
	-	4 (4.9)	131 (22.9)	3 (4.6)	132 (22.4)	57 (28.9)	246 (54.7)	23 (11.3)	219 (48.8)	6 (13.6)	283 (46.3)
Band	+	57 (70.4)	185 (32.2)	45 (69.2)	94 (16.0)	18 (9.1)	5 (1.1)	121 (59.3)	78 (17.4)	16 (36.4)	225 (36.9)
	-	24 (29.6)	388 (67.7)	20 (30.8)	495 (84.0)	179 (90.9)	445 (98.9)	83 (40.7)	371 (82.6)	28 (63.6)	384 (63.1)
CR	+	62 (76.5)	194 (33.9)	49 (75.4)	104 (17.7)	151 (76.6)	154 (34.2)	170 (83.3)	158 (35.2)	26 (59.1)	200 (32.7)
	-	19 (23.5)	379 (66.1)	16 (24.6)	485 (82.3)	46 (23.4)	296 (65.8)	34 (16.7)	291 (64.8)	18 (40.9)	411 (67.3)

Frequency (%) of the positive and negative indications made per instrument and per disorder group, as a ratio of the total number of positive and negative cases. RL= referral letters, SDQ= strength and difficulties questionnaire, Band= DAWBA probability band score, CR= clinician rating in the DAWBA environment. Number of diagnoses and sample size were as follows: anxiety disorders n=81 and N=654; depressive disorder n=65 and N=654; autism spectrum disorders (ASD) n=197 and N=647; attention-deficit hyperactivity disorder (ADHD) n=204 and N=653; behavioural disorders n=44 and N=655. n= number of cases, N=total sample size.

for prognostic use. Of the youth with an anxiety disorder indicated by all four instruments, 48.8% were eventually classified with anxiety disorders (supplementary material). The classification rate was 54.9% for four successive indications of depressive disorders, 85.7% for ASDs, 70.0% for ADHDs and 10.7% for behavioural disorders.

When we considered the predictive value of successive negative indications, we found that 98.2% of those negative on all four instruments were not classified to anxiety disorders, 98.3% were not classified to depressive disorders, 90.5% to ASDs, 95.8% to ADHD and 99.1% were not classified to behavioural disorders.

Incremental and independent predictive values

When we examined the incremental value of the four assessment tools relative to each other, successive addition of a following instrument resulted in improvement in model fit for nearly all of the (4*5) models (Table 3). Only the fit for behavioural disorders did not improve with addition of the clinician-rated DAWBA scores to the model ($p=0.82$).

By controlling for the value of up to three other instruments, we explored independent associations of the four instruments with the outcome classifications (Figure 1). In these multivariable models most instruments showed predictive value. Only in the case of the SDQ did we see a failure to improve the prediction of depressive disorders and behavioural disorders (respectively OR=1.24, 95% CI 0.58-2.62; OR=1.85, 95% CI 0.82-4.16).

For most disorder groups and instruments, we found no differences in magnitude of the associations in the multivariable models as compared with the univariable prediction models. Similarly, no difference in patterns was observed when inspecting differences in the predictive value of the earlier instruments compared with the later instruments. The clinician rated DAWBA, for instance, did not show consistently higher predictive values as compared with the referral letters.

Table 3. Likelihood ratio test values comparing the effect of addition of instruments on model fit per disorder group

	RL	+SDQ	+DAWBA Band	+DAWBA CR
Anxiety disorders	92.74	33.47	41.81	15.1
Depressive disorders	136.81	8.28*	39.63	17.48
ASD	166.29	44.48	15.25	14.50
ADHD	203.53	79.52	42.23	39.58
Behavioural disorders	44.26	16.04	16.78	0.02**

Likelihood ratio test results depicting change in model fit by successive addition of the instruments, computed in the imputed dataset. All values are significant at the $p<.001$ level, except * $p=.004$ and ** $p=0.82$. ASD= Autism spectrum disorders. ADHD= attention-deficit hyperactivity disorders. PPV= positive predictive value. NPV= negative predictive value. Note the low frequency of four successive positive indications for ASDs and ADHD, as it was uncommon for these minors to have positive scores on all four instruments.




Figure 1. Univariable and multivariable odds ratios per instrument and per diagnostic outcome

Odds ratios per instrument and per disorder group for four models, computed in the imputed dataset. Each successive model contains one more instrument as predictor, presenting how the ORs change when controlling for overlap with more instruments. The vertical line presents OR=1.

Conclusion

To the best of our knowledge, this study is the first to compare the predictive value of referral letters, broad band screening, structured multi-informant assessment and a clinician's remote evaluation in predicting diagnostic outcome in a single population. We found that all four nodes of assessment generally showed a positive contribution to the prediction of common child adolescent mental health problems. Referral letters and SDQ scale scores showed either a high sensitivity or a high specificity, whereas DAWBA probability bands and clinician ratings were more balanced in terms of sensitivity and specificity. Referral letters performed especially well for depressive disorders, which might be related to an earlier observation made during the pilot phase of our previous study: professionals might focus on mood problems and associate it with risk of suicidal ideation.³⁹ For the other disorder groups, referral letters showed better performance in terms of specificity compared with sensitivity. The SDQ, by contrast, was overinclusive, particularly for emotional problems,²⁰ a finding in line with earlier conclusions that advised against complete reliance on the SDQ to guide referrals.¹⁸ To determine whether this might be a result of our categorization of the SDQ indication as positive from the upper two indications on, we reanalysed the data categorising only the upper category as positive. This resulted in a sensitivity decrease of 15 percentage points (to 80.5 for anxiety and 78.5 for depressive disorders), whereas specificity doubled to around 50% false positives. Nonetheless, in comparison with the other instruments, SDQ screening was still overinclusive, an issue inherent to a screening instrument's function (to detect problems), the clinic population, and, as underlined in the introduction, screening instruments should be accompanied by clinical judgement.

Although the SDQ does not officially have an ASD scale, we also included children and adolescents with ASD in the study to shed light on the issue of EBA in this clinically-widespread population. We used a difference score suggested by the SDQ developers²⁷ and found that youngsters with ASD were detected at a similar rate to other problem types on conceptually related SDQ scales. However, other studies have used other computational methods,^{20 29 40 41} and the respective methods have not yet been compared.

We also inspected frequencies of successive positive and negative indications as a first approach to the question of outcomes for young people who show successive positive or negative scores on a sequence of assessment instruments. In this explorative inspection we found that four successive indications of anxiety or depressive disorders resulted in only a one in two chance of being classified to these outcomes. By contrast, when all instruments indicated ASD or ADHD, cases were indeed clinically classified as such. Regarding behaviour problems, we found that even four successive positive indications were not predictive of a classification to behavioural disorders. When considering the opposite situation, those with four successive negative indications, we found that about 1% was classified to anxiety, depressive or behavioural disorders, whereas around 5% or 10% were still classified to ADHD or ASD, respectively. It is unsurprising that rates were highest for ASD, because if initial instruments fail to suggest this relatively difficult diagnosis further clinician based investigations subsequently detect ASD. These results underline the need for elaborate diagnostics, the inclusion of clinicians when aiming for specialised treatment and the importance of future studies with a diverse sample for better generalisability.

We found added benefits with each successive node of assessment, with only one exception for one outcome: the clinician ratings showed no improvement in the prediction of behavioural disorders relative to the three previous instruments combined. This might be because of the already marginal prediction of behavioural disorders and to the relatively conservative properties of the clinician rated DAWBA.²² With regards to the independent predictive value, we found that nearly all instruments remained individually associated with the outcome even when corrected for overlap with other instruments. Only the SDQ showed no independent value in predicting depressive and behavioural disorders when corrected for information provided by other nodes of assessment. In contrast to general literature suggesting that instruments applied later in a sequence might show stronger effects,⁴² we observed no increase in effect. While each instrument certainly contributed new information in our study, there was no indication that the most extensive assessment method should have been used first. Therefore, the study results give no support for use of the most elaborate instrument first and only, and support a stepwise approach to assessment.⁴³

Limitations

Although this study presented unique data on an important question, some limitations should be kept in mind. Firstly, people involved in classifying outcomes were not blinded to the instrument's results. To what extent results were viewed when formulating a diagnosis is not known. As regards the effect of the availability of DAWBA data, for instance, there are indications that it improves decision-making in the case of internalising problems but not in the case of externalizing problems.²⁴ In an effort to explore this type of potential effect, we split the sample between those with or without clinician ratings (see Methods section) but did not find differences in odds ratios between subsamples. Regardless, if disclosure had any effect it would likely result in the presented odds ratios overestimating associations. Looked at more positively, our research question concerned the relative predictive value of the instruments and, in principle, all instruments were accessible and have shown predictive value, also, in other studies with blinding.

Another limitation concerns discriminant ability of the instruments. If the aim is to predict the type and classification of a problem, insight into how scales relate to conceptually parallel classifications is not sufficient. Future studies could therefore focus on the discriminant ability of the tools and investigate cross relations between scales and types of problems. Furthermore, we focused only on the type of problems, whereas taking the staging and impact of symptoms into account could benefit clinical practice.⁴⁴

Implications

The questions addressed in this study are directly relevant to clinical practice. Referral letters are by definition available for many cases yet are seldom incorporated into the diagnostic process. In this study we found that referral letters add value, even when corrected for overlap with structured assessment instruments. Similarly, the DAWBA package has the potential to ease the assessment process by capturing the SDQ as a short yet sensitive screening instrument, the DAWBA structured questions as a broad assessment tool to “cast a wide net regarding the presenting problem of a client”,⁴⁵ and the clinician rated DAWBA to add some nuance regarding the fuller picture while not being overinclusive. When used within a sequential approach, the DAWBA package may help develop a shared language between primary care and specialised care professionals and parents, just as the DAWBA package also produces a report for parents when requested.⁴⁶ This in turn might stimulate

fruitful discussions within families and help ameliorate discrepancies between the problem perceptions of minors versus caregivers, the perceived focus of treatment and treatment outcomes.^{1 43 47 48} Moreover, a harmonised sequential diagnostic approach might facilitate real integration and joint working in the primary-secondary care interface, a challenge that has not been overcome despite decades of research and dissemination of the importance of EBA. The idea of working within and towards a complete and reliable work-up might be more palatable as compared with choosing from a list of measures purely based on one's own familiarity and time limits, without any insight regarding subsequent steps.^{6 48} Earlier studies found the DAWBA to be relatively conservative in terms of the number of diagnoses made and required administration time when compared with other elaborate diagnostic instruments.²² This suggests that it might hold potential for use at the primary-secondary care interface, as a second step for those with high scores on screening instruments in primary care and to prioritise referrals and registration in secondary mental healthcare.

Conclusion

In conclusion, our results suggest that integrating referral letters, screening questionnaires and information obtained from assessment is likely to facilitate diagnosis in clinical practice. Prospective studies could further quantify the clinical and economic value of this type of multi-tiered approach in relation to the facilitation of psychometrically sound and feasible decision-making, timely recognition of problems, determination of required care intensities and treatment outcomes.

Reference list Study 3

1. Jensen-Doss, A. and Weisz, J. R., *Diagnostic agreement predicts treatment process and outcomes in youth mental health clinics*. J Consult Clin Psychol, 2008. **76**(5): p. 711-22.
2. Youngstrom, E. A., Van Meter, A., Frazier, T. W., Hunsley, J., Prinstein, M. J., Ong, M. L., et al., *Evidence-based assessment as an integrative model for applying psychological science to guide the voyage of treatment*. Clinical Psychology: Science and Practice, 2017. **24**(4): p. 331-363.
3. Yates, B. T. and Taub, J., *Assessing the costs, benefits, cost-effectiveness, and cost-benefit of psychological assessment: we should, we can, and here's how*. Psychol Assess, 2003. **15**(4): p. 478-95.
4. Rettew, D. C., Lynch, A. D., Achenbach, T. M., Dumenci, L., and Ivanova, M. Y., *Meta-analyses of agreement between diagnoses made from clinical evaluations and standardized diagnostic interviews*. Int J Methods Psychiatr Res, 2009. **18**(3): p. 169-84.
5. Kuhn, C., Aebi, M., Jakobsen, H., Banaschewski, T., Poustka, L., Grimmer, Y., et al., *Effective Mental Health Screening in Adolescents: Should We Collect Data from Youth, Parents or Both?* Child Psychiatry Hum Dev, 2017. **48**(3): p. 385-392.
6. Jensen-Doss, A. and Hawley, K. M., *Understanding barriers to evidence-based assessment: clinician attitudes toward standardized assessment tools*. J Clin Child Adolesc Psychol, 2010. **39**(6): p. 885-96.
7. Youngstrom, E. A., Choukas-Bradley, S., Calhoun, C. D., and Jensen-Doss, A., *Clinical Guide to the Evidence-Based Assessment Approach to Diagnosis and Treatment*. Cognitive and Behavioral Practice, 2015. **22**(1): p. 20-35.
8. Johnston, C. and Murray, C., *Incremental validity in the psychological assessment of children and adolescents*. Psychol Assess, 2003. **15**(4): p. 496-507.
9. Sayal, K., Tischler, V., Coope, C., Robotham, S., Ashworth, M., Day, C., et al., *Parental help-seeking in primary care for child and adolescent mental health concerns: qualitative study*. Br J Psychiatry, 2010. **197**(6): p. 476-81.
10. Aydin, S., Crone, M. R., Siebelink, B. M., Ginkel, J. v., Numans, M. E., Vermeiren, R. R. J. M., et al., *Informative value of referral letters from general practice for child and adolescent mental healthcare*. European Child & Adolescent Psychiatry, 2021. **(forthcoming)**.
11. Beidas, R. S., Stewart, R. E., Walsh, L., Lucas, S., Downey, M. M., Jackson, K., et al., *Free, brief, and validated: Standardized instruments for low-resource mental health settings*. Cogn Behav Pract, 2015. **22**(1): p. 5-19.
12. Doss, A. J., *Evidence-based diagnosis: incorporating diagnostic instruments into clinical practice*. J Am Acad Child Adolesc Psychiatry, 2005. **44**(9): p. 947-52.
13. O'Brien, D., Harvey, K., Howse, J., Reardon, T., and Creswell, C., *Barriers to managing child and adolescent mental health problems: a systematic review of primary care practitioners' perceptions*. Br J Gen Pract, 2016. **66**(651): p. e693-707.
14. Becker-Haimes, E. M., Tabachnick, A. R., Last, B. S., Stewart, R. E., Hasan-Granier, A., and Beidas, R. S., *Evidence Base Update for Brief, Free, and Accessible Youth Mental Health Measures*. J Clin Child Adolesc Psychol, 2020. **49**(1): p. 1-17.

15. Goodman, R., *The Extended Version of the Strengths and Difficulties Questionnaire as a Guide to Child Psychiatric Caseness and Consequent Burden*. Journal of Child Psychology and Psychiatry, 1999. **40**(5): p. 791-799.
16. Becker, A., Woerner, W., Hasselhorn, M., Banaschewski, T., and Rothenberger, A., *Validation of the parent and teacher SDQ in a clinical sample*. Eur Child Adolesc Psychiatry, 2004. **13 Suppl 2**: p. II11-6.
17. van Widenfelt, B. M., Goedhart, A. W., Treffers, P. D., and Goodman, R., *Dutch version of the Strengths and Difficulties Questionnaire (SDQ)*. Eur Child Adolesc Psychiatry, 2003. **12**(6): p. 281-9.
18. Janssens, A. and Deboutte, D., *Screening for psychopathology in child welfare: the Strengths and Difficulties Questionnaire (SDQ) compared with the Achenbach System of Empirically Based Assessment (ASEBA)*. Eur Child Adolesc Psychiatry, 2009. **18**(11): p. 691-700.
19. Goodman, R., Ford, T., Richards, H., Gatward, R., and Meltzer, H., *The Development and Well-Being Assessment: Description and Initial Validation of an Integrated Assessment of Child and Adolescent Psychopathology*. Journal of Child Psychology and Psychiatry, 2000. **41**(5): p. 645-655.
20. Vugteveen, J., De Bildt, A., Hartman, C. A., and Timmerman, M. E., *Using the Dutch multi-informant Strengths and Difficulties Questionnaire (SDQ) to predict adolescent psychiatric diagnoses*. Eur Child Adolesc Psychiatry, 2018. **27**(10): p. 1347-1359.
21. Goodman, A., Heiervang, E., Collishaw, S., and Goodman, R., *The 'DAWBA bands' as an ordered-categorical measure of child mental health: description and validation in British and Norwegian samples*. Soc Psychiatry Psychiatr Epidemiol, 2011. **46**(6): p. 521-32.
22. Angold, A., Erkanli, A., Copeland, W., Goodman, R., Fisher, P. W., and Costello, E. J., *Psychiatric diagnostic interviews for children and adolescents: a comparative study*. J Am Acad Child Adolesc Psychiatry, 2012. **51**(5): p. 506-17.
23. Citroen, A., Siebelink, B., and van Lang, N., *DAWBA: de nieuwe poortwachter van de jeugd-ggz?* Tijdschrift voor gezondheidswetenschappen, 2017. **95**(6): p. 246-249.
24. Aebi, M., Kuhn, C., Metzke, C. W., Stringaris, A., Goodman, R., and Steinhausen, H. C., *The use of the development and well-being assessment (DAWBA) in clinical practice: a randomized trial*. Eur Child Adolesc Psychiatry, 2012. **21**(10): p. 559-67.
25. Nesvag, R., Jonsson, E. G., Bakken, I. J., Knudsen, G. P., Bjella, T. D., Reichborn-Kjennerud, T., et al., *The quality of severe mental disorder diagnoses in a national health registry as compared to research diagnoses based on structured interview*. BMC Psychiatry, 2017. **17**(1): p. 93.
26. Crone, M. R., Zeijl, E., and Reijneveld, S. A., *When do parents and child health professionals agree on child's psychosocial problems? Cross-sectional study on parent-child health professional dyads*. BMC Psychiatry, 2016. **16**: p. 151.
27. Goodman, A., Lamping, D. L., and Ploubidis, G. B., *When to use broader internalising and externalising subscales instead of the hypothesised five subscales on the Strengths and Difficulties Questionnaire (SDQ): data from British parents, teachers and children*. J Abnorm Child Psychol, 2010. **38**(8): p. 1179-91.
28. Iizuka, C., Yamashita, Y., Nagamitsu, S., Yamashita, T., Araki, Y., Ohya, T., et al., *Comparison of the strengths and difficulties questionnaire (SDQ) scores between children with high-functioning autism spectrum disorder (HFASD) and attention-deficit/hyperactivity disorder (AD/HD)*. Brain Dev, 2010. **32**(8): p. 609-12.

29. Vugteveen, J., de Bildt, A., Theunissen, M., Reijneveld, S. A., and Timmerman, M., *Validity Aspects of the Strengths and Difficulties Questionnaire (SDQ) Adolescent Self-Report and Parent-Report Versions Among Dutch Adolescents*. Assessment, 2021. **28**(2): p. 601-616.
30. Spitzer, R. L., *Psychiatric diagnosis: Are clinicians still necessary?* Compr Psychiatry, 1983. **24**: p. 399-411.
31. Jensen-Doss, A., Youngstrom, E. A., Youngstrom, J. K., Feeny, N. C., and Findling, R. L., *Predictors and moderators of agreement between clinical and research diagnoses for children and adolescents*. J Consult Clin Psychol, 2014. **82**(6): p. 1151-62.
32. Jensen, A. L. and Weisz, J. R., *Assessing match and mismatch between practitioner-generated and standardized interview-generated diagnoses for clinic-referred children and adolescents*. Journal of Consulting and Clinical Psychology, 2002. **70**(1): p. 158-168.
33. van Buuren, S. and Groothuis-Oudshoorn, K., *Multivariate Imputation by Chained Equations in R*. Journal of Statistical Software, 2011. **45**: p. 1-67.
34. van Ginkel, J. R., Linting, M., Rippe, R. C. A., and van der Voort, A., *Rebutting Existing Misconceptions About Multiple Imputation as a Method for Handling Missing Data*. J Pers Assess, 2020. **102**(3): p. 297-308.
35. Rubin, D. B., *Multiple Imputation for Nonresponse in Surveys*. 2004, New York: John Wiley & Sons Inc.
36. van Buuren, S., *Flexible imputation of missing data*". 2018, Boca Raton: Chapman & Hall/CRC Interdisciplinary Statistics.
37. R Core Team, *R: a Language and Environment for Statistical Computing*. 2021, R Foundation for Statistical Computing.: Vienna, Austria.
38. Meng, X.-L. and Rubin, D. B., *Performing Likelihood Ratio Tests with Multiply-Imputed Data Sets*. Biometrika, 1992. **79**(1).
39. Aydin, S., Crone, M. R., Siebelink, B. M., Vermeiren, R., Numans, M. E., and Westenberg, P. M., *Recognition of anxiety disorders in children: a cross-sectional vignette-based survey among general practitioners*. BMJ Open, 2020. **10**(4): p. e035799.
40. Russell, G., Rodgers, L. R., and Ford, T., *The strengths and difficulties questionnaire as a predictor of parent-reported diagnosis of autism spectrum disorder and attention deficit hyperactivity disorder*. PLoS One, 2013. **8**(12): p. e80247.
41. Salayev, K. A. and Sanne, B., *The strengths and difficulties questionnaire (SDQ) in autism spectrum disorders*. International Journal on Disability and Human Development, 2017. **16**(3).
42. Moons, K. G., de Groot, J. A., Linnet, K., Reitsma, J. B., and Bossuyt, P. M., *Quantifying the added value of a diagnostic test or marker*. Clin Chem, 2012. **58**(10): p. 1408-17.
43. Whitmyre, E. D., Adams, L. M., Defayette, A. B., Williams, C. A., and Esposito-Smythers, C., *Is the focus of community-based mental health treatment consistent with adolescent psychiatric diagnoses?* Child Youth Serv Rev, 2019. **103**: p. 247-254.
44. Scott, J., Leboyer, M., Hickie, I., Berk, M., Kapczynski, F., Frank, E., et al., *Clinical staging in psychiatry: a cross-cutting model of diagnosis with heuristic and practical value*. Br J Psychiatry, 2013. **202**(4): p. 243-5.
45. Mash, E. J. and Hunsley, J., *Evidence-based assessment of child and adolescent disorders: issues and challenges*. J Clin Child Adolesc Psychol, 2005. **34**(3): p. 362-79.
46. Bennett, S. D., Heyman, I., Coughtrey, A. E., Buszewicz, M., Byford, S., Dore, C. J., et al., *Assessing feasibility of routine identification tools for mental health disorder in neurology clinics*. Arch Dis Child, 2019. **104**(12): p. 1161-1166.

47. Priebe, S., McCabe, R., Bullenkamp, J., Hansson, L., Lauber, C., Martinez-Leal, R., et al., *Structured patient-clinician communication and 1-year outcome in community mental healthcare: cluster randomised controlled trial*. Br J Psychiatry, 2007. **191**: p. 420-6.
48. Kazdin, A. E., *Evidence-based assessment for children and adolescents: issues in measurement development and clinical application*. J Clin Child Adolesc Psychol, 2005. **34**(3): p. 548-58.