



Universiteit
Leiden
The Netherlands

Comprehensive measurement of long-term outcomes and costs of rehabilitation in patients with stroke

Meijeren-Pont, W. van

Citation


Meijeren-Pont, W. van. (2023, February 22). *Comprehensive measurement of long-term outcomes and costs of rehabilitation in patients with stroke*. Retrieved from <https://hdl.handle.net/1887/3564442>

Version: Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3564442>

Note: To cite this publication please use the final published version (if applicable).



Chapter 4

The long-term course of participation in stroke patients with paid employment

van Meijeren-Pont W | van Velzen JM | Volker G | Arwert HJ | Meesters JJL | de Kloet AJ
| van Bennekom CAM | Vliet Vlieland TPM | Tamminga SJ | Oosterveer DM |
on behalf of the SCORE-study group.

Submitted

Abstract

Background

Knowledge on long-term participation for patients with paid employment at the time of stroke is scarce.

Aims/Objectives

Describe the characteristics and the course of participation (concerning paid employment and overall participation) in patients who did and did not remain in paid employment.

Material and Methods

Patients with paid employment at the time of stroke completed questions on work up to 30 months after starting rehabilitation, and the Utrecht Scale for Evaluation of Rehabilitation-Participation (USER-P, Frequency, Restrictions and Satisfaction scales) up to 24 months. Baseline characteristics of patients with and without paid employment at 30 months were compared using Fisher's exact tests and Mann-Whitney U tests. USER-P scores over time were analysed using linear mixed models.

Results

Of the 170 included patients (median age 54.2 interquartile range 11.2 years; 40% women) 50.6% reported paid employment at 30 months. Those returning to work reported at baseline more working hours, better quality of life and communication, were more often self-employed and in an office job. The USER-P scores did not change statistically significantly over time.

Conclusions

About half of the stroke patients remained in paid employment.

Significance

Optimizing interventions for returning to work and achieving meaningful participation outside of employment seem desirable.

Introduction

Stroke is a common and serious medical condition¹ often leading to impairments in physical and emotional functioning, cognition, and communication²⁻⁴, negatively influencing participation in society⁵.

Regarding the course of participation on the longer term after stroke, the literature is scarce. Nevertheless, the relatively few available studies with a longer duration of follow-up showed that, despite improvements in particular in the first year, a considerable proportion of patients with stroke still experience restrictions in various aspects of participation on the longer term⁶⁻⁹. These restrictions in participation include the domain employment⁹, since approximately a quarter of the patients is younger than 65 years at the time of stroke and thus part of the labour force⁸. A review of Treger et al.¹⁰ demonstrated differences in the proportion of stroke patients that return to work between countries ranging from 14% in Germany to 73% in France and Portugal. Other reviews also mentioned wide ranges of return to work: 4.0-90.9% with a pooled summary estimate of 67.4% two years post stroke⁹, 11% three months after rehabilitation to 85% seven years post stroke¹¹, or 0% 0-3 years post stroke to 100%, with an average of 44%^{12,13}. These differences may not only reflect different international differences such as retirement age or social security systems, but may also be related to the inclusion of different stroke populations (population-based, hospital-based, rehabilitation-based), different definitions or assessment of employment status, and different follow-up durations.

Overall, it must be noted that most studies on the course of employment status report on one specific time point after stroke, usually not beyond one year, and do not describe the course of returning to work over time. Moreover, most of the studies did not report on aspects of participation other than return to paid employment, whereas participation in other meaningful activities is very important as well, both in patients who do and do not return to work.

Given that knowledge gap, the aim of the present study was to describe the long-term employment outcomes and overall participation in a Dutch cohort of stroke patients who received multidisciplinary rehabilitation and who had paid employment at the time of stroke. More specifically, the study aimed, in patients with paid employment at the time of stroke, a) to explore differences in characteristics of patients who did and did not return to work at 30 months; b) to describe the proportion of patients with paid employment and on partial or full-time sick leave over time as well as their use of employment adaptations and support; and c) to describe the course of participation in patients who did and did not remain in the work force.

Materials and methods

Study design

This study was part of the Stroke Cohort Outcomes of REhabilitation (SCORE) study¹⁴, a longitudinal cohort study that was executed from March 2014 until December 2019 at Basalt, a multidisciplinary rehabilitation facility in the Netherlands.

The Medical Ethical Committee of the Leiden University Medical Center (protocol number NL465321.058.13) approved the SCORE study, that is registered at the International Clinical Trial Registry Platform (<https://trialsearch.who.int/Default.aspx>: NTR4293). The current study on the long-term course of participation is reported in accordance with the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) guidelines¹⁵.

Setting

In the Netherlands, after an average of six days of hospital admission for stroke, approximately 14% of the patients are referred to inpatient multidisciplinary rehabilitation in a rehabilitation facility, 15% are discharged to inpatient geriatric rehabilitation and 71% of the patients are discharged home¹⁶. Some of the patients who are discharged to their homes are referred to outpatient multidisciplinary rehabilitation in a rehabilitation facility¹⁶. As compared with geriatric rehabilitation, the population of stroke patients admitted to multidisciplinary rehabilitation is composed of the younger patients, who were more active before stroke and have complex impairments⁸.

With respect to the Dutch legislation and social security system related to sick leave and work disability, it is compulsory for anyone that has paid employment with an employment contract to be insured under the Dutch Employee Insurance Schemes. This insurance obligates employers to continue to pay (a percentage of) the salary when an employee is fully or partly sick-listed during the first two years. In addition, during this period employers should do all they can to ensure that the sick employee returns to work as quickly as possible in a responsible way, including providing (temporary) modified work within the own company or elsewhere when necessary¹⁷. When the employee stays disabled and sick-listed for work for more than two years, the employee's 'ability to work' is examined. When this 'ability to work' is not present anymore, the employee receives a benefit of the Dutch government and the employer is allowed to terminate the employment contract of the employee. In case of self-employment this legislation does not apply; return to work is the patient's own responsibility, and it depends on his or her private insurance for sick leave and work disability whether or not he or she receives a benefit during sick leave and when there is no ability to work.

Patients

Consecutive stroke patients starting with inpatient or outpatient rehabilitation in the multidisciplinary rehabilitation facility were invited by their rehabilitation physician to participate in the SCORE study when they: 1) were 18 years or older; 2) had a first or recurrent stroke less than six months ago; 3) had no dementia or psychiatric disorder; and 4) were able to complete questionnaires in Dutch. Eligible patients who were willing to participate were only included after they provided written informed consent.

The current analysis concerned a subset of patients who had paid employment at the time of stroke, were aged <66 years (retirement age in the Netherlands in 2019) 30 months after start of rehabilitation (T30) and completed the questionnaire related to paid employment at T30.

4

Assessments

Sociodemographic and clinical characteristics

Sociodemographic and clinical characteristics were recorded at the start of rehabilitation, i.e. baseline. Age, sex and stroke type (i.e. ischemic or haemorrhagic stroke) were extracted from the patients' medical file. A questionnaire was used to assess educational level and living situation. Comorbidities were determined by the Dutch Life Situation Cohort Questionnaire, comprising 16 chronic diseases¹⁸. The Barthel Index was completed only for patients receiving inpatient rehabilitation. The Barthel Index is a nurse-reported measurement instrument that measures functional independence. It yields a score between 0 and 20, with higher scores indicating more independence¹⁹.

Employment prior to stroke and at follow-up

A questionnaire about paid employment prior to stroke was completed at baseline and included the following questions: type of contract (permanent, temporary, self-employed, other), amount of working hours per week according to contract, type of occupation (office job, service job or industrial/manual job) and managerial position (yes/no).

At 6 (T6), 12 (T12), 18 (T18), 24 (T24) and 30 (T30) months after baseline, patients were asked whether they had paid employment (yes/no), defined as having an employment contract or being self-employed, regardless of being actually working or not (because of partial or full sick leave).

If patients indicated that they were in paid employment, an additional questionnaire was completed. They were asked whether they were actually working and/or were on partial or full sick leave. This questionnaire also comprised questions on the occurrence of employment adaptations (changes of tasks, working hours, function/position, work accommodations, or a change of employer) and support related to return to work (work-related support from

employer/supervisor, occupational physician, rehabilitation center or other), all over the past 6 months, in yes/no format.

Other Patient Reported Outcome Measures (PROMs)

Apart from the questionnaire concerning paid employment, the EuroQol-5 Dimensions-3 Levels (EQ-5D-3L)²⁰ and four domains of the Stroke Impact Scale (SIS) version 3.0²¹ were completed at baseline. The Utrecht Scale for Evaluation of Rehabilitation-Participation (USER-P)²² was completed at T6, T12 and T24.

The EQ-5D-3L was used to measure health-related quality of life²⁰. It comprises the following five dimensions: mobility, self-care, usual activities, pain/discomfort and anxiety/depression. Each dimension has 3 levels of severity: no problems, some problems, and extreme problems. The patient is asked to indicate his/her health state by ticking the box next to the most appropriate statement in each of the five dimensions. The resulting index ranges from -0.33 (serious problems on all five dimensions) to 1 (perfect health)²³. Next to this index, the EQ-5D-3L comprised a vertical visual analogue scale (VAS), ranging from 0 ('worst imaginable health state') to 100 ('best imaginable health state') to quantify the patient's self-rated health status²⁰.

The SIS is a stroke-specific health status measure, that assesses several domains²¹. Items are scored on a 5-point Likert scale and transformed to a score ranging from 0-100 for each domain, with higher scores indicating better functioning on that specific domain. The domains 'Communication' and 'Memory and thinking' were administered in all patients. In April 2015, the domains 'Mobility' and 'Mood and emotions' were added.

The USER-P is a measure that is based on the International Classification of Functioning, Disability and Health (ICF) and assesses objective and subjective participation²². It consists of 32 items divided into three scales: Frequency, Restrictions, and Satisfaction. The Frequency scale consists of four items on vocational activities ('paid work', 'unpaid work', 'education', 'household duties'), scored in hours per week ranging from 0 (not at all) to 5 (36 hours or more); and seven items on leisure and social activities, scored in frequency in the last four week ranging from 0 (not at all) to 5 (19 times or more). The Restrictions scale consists of 11 items on activities that may be restricted due to a health condition, including one item about 'paid work, unpaid work or education'. The perceived difficulty in performing the activity is rated on a four-point scale, ranging from 0 (not possible) to 3 (without difficulty). A 'not applicable' option is available for every item and can be used if the item is not relevant to the patient or if experienced restrictions are not related to the patient's health condition. The Satisfaction scale includes ten items on satisfaction with vocational, leisure and social activities and relationships. Items are rated on a scale from 0 (very dissatisfied) to 5 (very satisfied). For the items 'paid work, unpaid work or education' and 'your relationship with your partner' a 'not

'applicable' option is available. The sum score of each scale is based on all applicable items and is converted to a 0–100 scale, with higher scores indicating better participation (more time spent/higher frequency, less restrictions, higher satisfaction)²².

Statistical analysis

Data analyses were performed in IBM SPSS version 25 (IBM Corp: Armonk, NY, USA 2013). For all statistical analyses a two-sided *p* value of ≤ 0.05 was considered statistically significant. Data are presented as numbers (n) with percentages (%), as means with standard deviations (SD) or as medians and interquartile ranges (IQR) depending on their nature and their distribution. The Kolmogorov-Smirnov test was used to assess whether or not continuous variables were normally distributed.

Baseline sociodemographic, clinical and employment characteristics and PROMs of included patients were compared with those of patients who had paid employment at the time of stroke and were still <66 years old at T30, but who did not complete the questionnaire related to paid employment at T30 and were therefore excluded. For this comparison Fisher's exact tests and Mann-Whitney U tests were used. Baseline sociodemographic, clinical and employment characteristics and PROMs of patients with paid employment at T30 were compared with those of patients without paid employment at T30 using Fisher's exact tests and Mann-Whitney U tests, where appropriate.

The proportion of patients with paid employment was computed as the number of patients reporting they had paid employment at that time point divided by the number of patients completing the questionnaire on work at that time point. Only for patients with paid employment, the proportions of patients who were on sick leave, who had specific work adaptations and who received work-related support for each follow-up time point were calculated.

With respect to participation, the scores of each USER-P scale at T6, T12 and T24 were compared between patients with and without paid employment using Mann Whitney U tests. In order to make a fair comparison between patients with and without paid employment, additional analyses were done with the scores of each USER-P scale without the items concerning employment. For the Frequency scale it concerned omitting the items 'paid work' and 'education', as the latter is described as 'only training courses taken in the context of your paid work or to help you obtain paid work'. For the Restrictions and Satisfaction scales only the item 'paid work, unpaid work or education' was omitted. The minimum number of completed items for the Frequency scale for the first four items was set on two instead of three, and for the Satisfaction scale this was set on five instead of six.

In addition, to evaluate whether or not USER-P scores of each scale changed over time, linear mixed models were used. Analyses were done with both the complete USER-P scale scores and the scale scores without the items related to work as dependent variables. Paid employment was included in the model as being employed at T24 (yes/no). Time was the independent variable, and also an interaction term between time and paid employment at T24 was added to the model to analyse whether the slope of the change over time was different in patients with and without paid employment.

Results

Between March 2014 and December 2019, 836 patients were included in the SCORE study. Of these patients, 620 reported whether they had paid employment or not at the time of stroke: 348 (41.6%) patients reported they had paid employment at the time of stroke. Of these patients, 288 were younger than 66 years old at T30, of whom 170 (59%) completed the questionnaire related to paid employment at T30.

These 170 patients were included in the current analyses (Table 1). Their median age was 54.2 (IQR 11.2) years and 68 (40.0%) of them were female. The included patients did not statistically significantly differ from the 118 patients who had paid employment at the time of stroke and were still younger than 66 at T30, but who did not complete the employment questionnaire at T30 (Online resource 1).

Characteristics of patients with and without paid employment at T30

At T30, 86 patients (50.6%) reported to be in paid employment. Table 1 shows that compared to those who did not remain in the work force, patients with paid employment at T30 had statistically significantly more working hours and better EQ-5D-3L and SIS Communication scores at baseline. In addition, they were more often self-employed (versus permanent contract $p = 0.015$; versus temporary contract $p = 0.053$; versus other $p = 0.004$) and had more often an office job (versus service job $p = 0.026$; versus industrial/manual job $p = 0.013$).

Table 1. Baseline characteristics of stroke patients receiving multidisciplinary rehabilitation who had paid employment at the time of stroke.

| | n 170 | Included in the current analyses 86 | With paid employment at T30 84 | n 84 | Without paid employment at T30 | p value* |
|--|------------------------|--|---|-----------------------|---|--------------------------|
| Sociodemographic characteristics | | | | | | |
| Age in years | 170 | 54.2 (11.2) | 86 | 52.7 (9.3) | 84 | 56.4 (13.2) 0.087 |
| Female sex | 170 | 68 (40.0%) | 86 | 30 (34.9%) | 84 | 38 (45.2%) 0.210 |
| Low education level | 167 | 46 (27.1%) | 85 | 20 (23.5%) | 82 | 26 (31.7%) 0.299 |
| Living alone | 169 | 30 (17.8%) | 86 | 13 (15.1%) | 83 | 17 (20.5%) 0.423 |
| Clinical characteristics | | | | | | |
| Ischemic stroke | 167 | 126 (75.4%) | 84 | 63 (75.0%) | 83 | 63 (75.9%) 1.000 |
| Number of comorbidities | 131 | 1.0 (1.0) | 64 | 1.0 (2.0) | 67 | 1.0 (1.0) 0.054 |
| Barthel Index at start rehabilitation ¹ | 92 | 17.0 (9.0) | 44 | 17.0 (8.0) | 48 | 17.0 (9.0) 0.803 |
| Employment characteristics prior to stroke | | | | | | |
| Type of contract | 170 | 131 (77.1%) | 86 | 64 (74.4%) | 84 | 67 (79.8%) 0.008 |
| Permanent | | 12 (7.1%) | | 5 (5.8%) | | 7 (8.3%) |
| Temporary | | 20 (11.8%) | | 16 (18.6%) | | 4 (4.8%) |
| Self-employed | | 7 (4.1%) | | 1 (1.2%) | | 6 (7.1%) |
| Other | | | | | | |
| Number of working hours according to contract | 169 | 36.0 (11.0) | 86 | 38.0 (4.0) | 83 | 35.0 (16.0) 0.001 |
| Type of occupation | 155 | 67 (43.2%) | 79 | 43 (54.4%) | 76 | 24 (31.6%) 0.014 |
| Office job | | 51 (32.9%) | | 22 (27.8%) | | 29 (38.3%) |
| Service job | | 37 (23.9%) | | 14 (17.7%) | | 23 (30.3%) |
| Industrial or manual job | | 18 (11.7%) | | 13 (16.5%) | | 5 (6.7%) 0.079 |
| Managerial position | 154 | | | | | |
| Patient Reported Outcome Measures | | | | | | |
| EQ-5D-3L index | 151 | 0.78 (0.26) | 77 | 0.81 (0.21) | 74 | 0.77 (0.34) 0.008 |
| EQ-5D-3L VAS | 159 | 65.0 (26.0) | 81 | 70.0 (20.0) | 78 | 60.0 (31.0) 0.003 |
| SIS Communication | 161 | 92.2 (25.0) | 84 | 96.1 (17.9) | 77 | 89.3 (28.6) 0.035 |
| SIS Mobility | 84 | 84.7 (38.2) | 39 | 91.7 (36.1) | 45 | 83.3 (52.8) 0.207 |
| SIS Memory and thinking | 163 | 85.7 (25.0) | 84 | 85.7 (24.1) | 79 | 82.1 (32.1) 0.132 |
| SIS Mood and emotions | 84 | 79.2 (23.6) | 39 | 80.6 (19.4) | 45 | 75.0 (23.6) 0.183 |

Dichotomous variables are described as numbers with percentages (%) and continuous variables as medians with interquartile ranges; *p values are given of Fisher Exact Tests or Mann-Whitney U Tests, when appropriate.

¹For inpatients only

Abbreviations: EQ-5D-3L EuroQoL-5 Dimensions-3 Levels; SIS Stroke Impact Scale; VAS visual analogue scale.

Table 2. Employment outcomes at all follow-up time point of patients who had paid employment at the time of stroke.

| | n | 6 months | n | 12 months | n | 18 months | n | 24 months | n | 30 months |
|---|----------|-----------------|----------|------------------|----------|------------------|----------|------------------|----------|------------------|
| Employment | 159 | | 158 | | 153 | | 154 | | 170 | |
| No paid employment | | 25 (15.7%) | | 25 (15.8%) | | 38 (24.8%) | | 64 (41.6%) | | 84 (49.4%) |
| Paid employment | | 134 (84.3%) | | 133 (84.2%) | | 115 (75.2%) | | 90 (58.4%) | | 86 (50.6%) |
| Presence of sick leave when employed¹ | 93 | | 88 | | 81 | | 62 | | 50 | |
| Working without sick leave | | 9 (9.7%) | | 30 (37.5%) | | 43 (53.1%) | | 50 (80.6%) | | 41 (82.0%) |
| Partial sick leave | | 27 (29.0%) | | 35 (39.8%) | | 21 (25.9%) | | 7 (11.3%) | | 5 (10.0%) |
| Full sick leave | | 57 (61.3%) | | 23 (26.1%) | | 17 (21.0%) | | 5 (8.1%) | | 4 (8.0%) |
| Employment adaptations when employed¹ | 92 | | 133 | | 114 | | 90 | | 86 | |
| No employment adaptations | | 53 (57.6%) | | 67 (50.4%) | | 65 (57.0%) | | 57 (63.3%) | | 74 (86.0%) |
| Work tasks/activities | | 24 (26.1%) | | 32 (24.1%) | | 22 (19.3%) | | 13 (14.4%) | | 7 (8.1%) |
| Working hours | | 24 (26.1%) | | 38 (28.6%) | | 23 (20.2%) | | 9 (10.0%) | | 8 (9.3%) |
| Work function/position | | 3 (3.3%) | | 11 (8.3%) | | 9 (7.9%) | | 5 (5.6%) | | 5 (5.8%) |
| Work accommodations (e.g. devices) | | 5 (5.4%) | | 12 (9.0%) | | 6 (5.3%) | | 5 (5.6%) | | 1 (1.2%) |
| Change in employer | | 0 (0.0%) | | 3 (2.3%) | | 4 (3.5%) | | 3 (3.3%) | | 1 (1.2%) |
| Employment-related support¹ | 92 | | 133 | | 114 | | 90 | | 86 | |
| No employment-related support | | 34 (37.0%) | | 55 (41.4%) | | 65 (57.0%) | | 64 (71.1%) | | 80 (93.0%) |
| Employer/supervisor | | 34 (37.0%) | | 53 (39.8%) | | 29 (25.4%) | | 10 (11.1%) | | 4 (4.7%) |
| Occupational physician | | 38 (41.3%) | | 55 (41.4%) | | 36 (31.6%) | | 16 (17.8%) | | 4 (4.7%) |
| Rehabilitation centre | | 26 (28.3%) | | 23 (17.3%) | | 4 (3.5%) | | 3 (3.3%) | | 1 (1.2%) |
| Other | | 5 (5.4%) | | 10 (7.5%) | | 15 (13.2%) | | 4 (4.4%) | | 1 (1.2%) |

Variables are described as numbers with percentages (%).

¹In the previous six months in the patients who had paid employment at that specific follow-up time point; several answers were possible when patients reported that there were employment adaptations or that they received employment-related support.

Paid employment over time

Table 2 shows the employment status of the 170 participants over time. The proportions of patients reporting that they were employed decreased, in particular between T18 and T24, with eventually 50.6% of patients reporting paid employment at T30. Only few patients reported changing jobs. The individual courses of patients of employment status are described in Online Resource 2.

Patients reporting paid employment could also be on sick leave partially or fully. Although the proportions of patients in paid employment decreased over time, among those with paid employment, the percentage of patients reporting that they were working without sick leave increased from 9.7% at T6 to 82.0% at T30. It must be noted that at the various follow-up time points, only 58.1%-70.4% of patients reporting paid employment provided information on sick leave.

Employment adaptations and support

Table 2 also provides insight into the implementation of employment adaptations and the support from the employer or health professionals with respect to return to work. It appeared that overall changes in tasks and activities and changes in working hours were the most frequently reported employment adaptations. With regard to support the guidance from the employer/supervisor and occupational physician were reported more often than that from the rehabilitation center or other sources. Like the questions on sick leave, the response rates to the questions on employment adaptations and support at the various time points were varying between 68.7%-100.0%.

Participation over time

Table 3 shows the scores of all three USER-P scales of the total group of patients and separately for patients either reporting or not reporting paid employment at T6, T12 and T24.

Regarding the differences of USER-P scale scores between patients with and without paid employment, there were no statistically significant differences at T6, whereas at T12 patients reporting paid employment had significantly better scores for the USER-P Frequency and Restrictions scales ($p < 0.05$) and at T24 for all three USER-P scales (all $p < 0.001$). With respect to USER-P scale scores over time, there were no statistically significant changes over time, neither in the total, nor within the subgroups of patients with or without paid employment at T24 (Online Resource 3).

When leaving out the items concerning employment, again at T6 no statistically significant differences in USER-P scale scores were seen between patients who did and did not report paid employment at that time point. At T12 only the difference for the Restrictions scale remained. At T24, the scores for the Restrictions and Satisfaction scales were statistically significantly better for patients with paid employment, whereas the Frequency scale score was not statistically significantly different. Regarding the course of the Frequency scale score, its scores diminished over time for patients with paid employment ($\beta -1.74$, 95%CI $-2.96 - -0.52$, $p = 0.005$), but not in patients without paid employment at 24 months.

Table 3. USER-P in stroke patients with and without paid employment up to 24 months after the start of rehabilitation.

| | n | 6 months | p value* | n | 12 months | p value* | n | 24 months | p value* |
|---|-----|-------------|----------|-----|-------------|--------------|-----|-------------|------------------|
| USER-P Frequency | | | | | | | | | |
| <i>All items</i> | | | | | | | | | |
| All patients | 159 | 33.6 (16.4) | | 160 | 33.6 (14.8) | | 158 | 31.4 (16.3) | |
| Paid employment | 131 | 34.6 (17.5) | 0.182 | 133 | 34.6 (15.0) | 0.001 | 90 | 36.0 (17.0) | <0.001 |
| No paid employment | 25 | 29.2 (13.0) | | 25 | 28.9 (16.3) | | 61 | 29.3 (10.4) | |
| <i>Without item 'Paid work' and 'education'</i> | | | | | | | | | |
| All patients | 159 | 34.3 (15.7) | | 160 | 33.1 (14.8) | | 158 | 32.9 (16.8) | |
| Paid employment | 131 | 34.3 (16.4) | 0.470 | 133 | 33.6 (14.3) | 0.799 | 90 | 31.8 (17.9) | 0.697 |
| No paid employment | 25 | 33.6 (17.3) | | 25 | 33.3 (23.6) | | 61 | 34.3 (13.6) | |
| USER-P Restrictions | | | | | | | | | |
| <i>All items</i> | | | | | | | | | |
| All patients | 161 | 83.3 (33.0) | | 157 | 87.5 (27.3) | | 158 | 87.9 (33.3) | |
| Paid employment | 133 | 83.3 (33.2) | 0.197 | 130 | 89.4 (27.9) | 0.005 | 90 | 96.8 (19.0) | <0.001 |
| No paid employment | 25 | 76.7 (43.3) | | 25 | 74.1 (35.3) | | 61 | 70.0 (30.8) | |
| <i>Without item 'Paid work, unpaid work or education'</i> | | | | | | | | | |
| All patients | 161 | 86.7 (33.3) | | 157 | 90.0 (30.0) | | 158 | 90.0 (29.3) | |
| Paid employment | 133 | 88.9 (33.3) | 0.284 | 130 | 92.6 (26.7) | 0.032 | 90 | 96.7 (18.5) | <0.001 |
| No paid employment | 25 | 80.0 (42.1) | | 25 | 74.1 (34.6) | | 61 | 73.3 (29.6) | |
| USER-P Satisfaction | | | | | | | | | |
| <i>All items</i> | | | | | | | | | |
| All patients | 158 | 72.2 (27.1) | | 157 | 72.5 (25.8) | | 155 | 72.5 (27.5) | |
| Paid employment | 130 | 72.2 (26.9) | 0.713 | 130 | 72.5 (26.3) | 0.093 | 90 | 77.6 (26.5) | <0.001 |
| No paid employment | 25 | 75.0 (33.8) | | 25 | 69.4 (29.5) | | 58 | 65.6 (27.8) | |
| <i>Without item 'Paid work, unpaid work or education'</i> | | | | | | | | | |
| All patients | 158 | 75.0 (25.7) | | 158 | 75.0 (25.3) | | 158 | 75.0 (26.2) | |
| Paid employment | 130 | 75.0 (25.0) | 0.813 | 131 | 75.0 (26.7) | 0.064 | 90 | 77.8 (26.9) | <0.001 |
| No paid employment | 25 | 75.0 (32.5) | | 25 | 69.4 (29.3) | | 61 | 69.4 (26.4) | |

Variables are described as medians with interquartile ranges; *p values are given of Mann-Whitney U Tests comparing patients with and without paid employment.

Abbreviations: USER-P Utrecht Scale for Evaluation of Rehabilitation-Participation.

Discussion

This study on the long-term course of employment outcomes and overall participation in patients with paid employment pre-stroke receiving multidisciplinary rehabilitation, found that half of them reported paid employment at 30 months after starting rehabilitation. The proportion of patients that had paid work was highest at six months with a marked decrease between 18 and 24 months after start of rehabilitation. These results reflect the Dutch social security system, where patients who are employed but sick-listed are entitled to a two-year period of (partial) salary payment and possible re-integration.

Baseline characteristics of employment, namely self-employment, a higher number of working hours and having an office job were associated with having paid employment at T30. In addition, the patients remaining in the work force reported better quality of life and less impact of their stroke on communication at baseline.

With respect to participation that is not employment related, patients who reported paid employment experienced less restrictions and were more satisfied than patients who did not. However, frequencies of participation outside of employment did not differ and decreased with time in those who retained work.

4

Our study showed a decrease in proportions of patients reporting paid employment that seems in contrast to previous studies such as that of Saeki et al.²⁴, that demonstrate an increase of patients that return to work over time. Nevertheless this contrast is not an actual contrast, because looking at the proportions of patients that reported paid employment and actually worked the same increase over time is seen.

In our study, half of the patients returned to paid employment at 30 months, but it is difficult to directly compare this result with previous studies, in part due to methodological differences. Therefore, and as mentioned in the introduction, estimated proportions of stroke patients returning to work varied largely⁹⁻¹². Nevertheless, our finding is in the same range as the proportions seen in a previous Dutch cross-sectional, hospital-based study including patients aged 18-65 years at 2-5 years post-stroke, where 39% returned to work²⁵. The patients of that study were younger and more often had an ischemic stroke than the patients in our study, but the proportions females and patients with a low level of education were comparable. In addition, our results were in the same range of a review which calculated a pooled summary estimate of return to work two years post-stroke of 67.4%. Overall, the heterogeneity in study methodology seen in the studies on this topic underlines the need for international consensus on how to best define and assess employment status in clinical and epidemiological studies in stroke patients^{10,26,27}.

This quantitative study did not elaborate on why patients were not able to return to paid employment. Depending on the patient's health status, the work situation, and the social security system, the work status of patients may vary largely within and across patients, with possible combinations of either or not working fully or partially and either or not being on fulltime or parttime sick leave, and either or not receiving a fulltime or parttime disability pension. For a detailed description an individual interview or an extensive questionnaire is needed.

Regarding the association of baseline characteristics with long-term paid employment, our findings are in general in line with previous literature. Regarding work characteristics, previous

literature in particular demonstrated that white collar occupation was beneficial for return to work compared to blue collar occupation (24,28). Our study found that self-employment and more working hours at baseline were also associated with return to work.

Regarding stroke characteristics, previous studies found that the presence of aphasia was negatively associated with return to work^{28,29}, which is in line with the observation in our study that the SIS Communication score was lower in patients who did not return to work.

Moreover, it has been found previously that normal muscle strength, absence of apraxia and more independence in activities of daily life measured with the Barthel Index were positively related to return to work^{24,29}, while other studies showed no influence of stroke severity³⁰. In our study, better scores for the EQ-5D-3L, which involves questions about mobility and activities of daily life, were associated with paid employment at T30. However this was not true for the SIS Mobility nor for the Barthel Index, perhaps because of low number of patients for whom these outcomes were known.

This study found that at all time points, a considerable proportion of patients reported employment adaptions in the previous six months, with only few changing jobs. The need for reductions in working hours and employment modifications because of changes in abilities due to stroke are previously mentioned in literature¹⁰. However, our results are hard to compare with those from other studies, as we did not record the cumulative, overall changes from baseline onwards. However, by recording adaptations over the previous six months, we were able to demonstrate that the occurrence of adaptations in those with paid employment decreased with time. It remains unclear to what extent this finding can be interpreted as a decreasing need and successful work integration over time.

Support from the employer and occupational physician were the most often reported sources of help. Although we have no cumulative figures, the findings at six months can be interpreted on their own, where it is striking that less than half of the patients reported support from their employer or occupational physician. These results may be flattened by the reporting of self-employed patients, but nevertheless may indicate that there is room for improvement, in particular given the far-reaching legal responsibility to support the return to work process in the Netherlands²⁶. Work-directed interventions in combination with education/coaching were shown previously to be effective regarding return to work³¹. It could be considered to include these interventions more consistently by the employer, occupational physician or rehabilitation center.

Considering participation outside of paid employment, it was striking that although frequencies were comparable, patients without paid employment experience more restrictions and less satisfaction with this participation. A previous study demonstrated that stroke patients retain predominately more sedentary and home-based activities and fewer physically demanding

and community based activities³². This might be more the case for patients without paid employment, explaining the difference in restrictions and satisfaction. Indeed, a need for well-founded, proven effective interventions for achieving meaningful participation outside of employment has been mentioned previously³³. It was suggested that this might require different types of support at various stages after stroke³⁴ and should take into account the social support system and other environmental factors, such as transportation^{35,36}.

Strengths and limitations

4

A strength of this study is the long-term, prospective design and the comprehensive assessment of both employment and participation. The computation of the USER-P scale scores with and without work-related items allowed a fair comparison on the perception of participation of patients who remained in paid employment and who did not. Limitations include the relatively small sample size, inclusion from only one rehabilitation facility and missing items in questionnaires of patients in the study. Moreover, the study population was selected based on their completion of the 30-months assessment. Although their characteristics at baseline did not differ from those who did not complete the study, selection bias cannot be ruled out. Another limitation was that the definition of 'having paid employment' could reflect different situations, including being actually at work or being fully or partly sick-listed. Although we aimed to gather detailed information from all patients, the precise working situation was missing for a proportion of patients. Finally, the results from the present study are influenced by the Dutch context and might therefore not be applicable to other countries with different legislation, social security and health systems.

In conclusion, the results of the present study suggest that there are windows of opportunity to improve the participation outcomes for patients in paid employment at the time of stroke receiving rehabilitation, both in those who do and do not remain in the work force, by implementing more consistently effective work-directed interventions and interventions for achieving meaningful participation outside of employment.

References

1. Feigin VL, Krishnamurthi RV, Parmar P, et al. Update on the Global Burden of Ischemic and Hemorrhagic Stroke in 1990-2013: The GBD 2013 Study. *Neuroepidemiology*. 2015;45(3):161-176.
2. Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: a systematic review. *Lancet Neurol*. 2009;8(8):741-754.
3. Lazar RM, AK Boehme. Aphasia as a predictor of stroke outcome. *Curr Neurol Neurosci Rep*. 2017;17(11):83.
4. van Rijssbergen MW, Mark RE, de Kort PL, et al. Subjective cognitive complaints after stroke: a systematic review. *J Stroke Cerebrovasc Dis*. 2014;23(3):408-420.
5. Verberne DPJ, Post MWM, Köhler S, et al. Course of social participation in the first 2 years after stroke and its associations with demographic and stroke-related factors. *Neurorehabil Neural Repair*. 2018;32(9):821-833.
6. van Velzen JM, van Bennekom CA, Edelaar MJ, et al. How many people return to work after acquired brain injury?: a systematic review. *Brain Inj*. 2009;23(6):473-488.
7. Törnbom K, Lundälv J, Sunnerhagen KS. Long-term participation 7-8 years after stroke: Experiences of people in working-age. *PLoS One*. 2019;14(3):e0213447.
8. de Boer AR, van Dis I, Visseren FLJ, Vaartjes I, Bots ML. Hart- en vaatziekten in Nederland 2019, cijfers over incidentie, prevalentie, ziekte en sterfte. Den Haag: Hartstichting; 2019.
9. Duong P, Sauvé-Schenk K, Egan MY, et al. Operational definitions and estimates of return to work poststroke: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2019;100(6):1140-1152.
10. Treger I, Shames J, Giaquinto S, et al. Return to work in stroke patients. *Disabil Rehabil*. 2007;29(17):1397-1403.
11. Wozniak MA, Kittner SJ. Return to work after ischemic stroke: a methodological review. *Neuroepidemiology*. 2002;21(4):159-166.
12. Daniel K, Wolfe CD, Busch MA, et al. What are the social consequences of stroke for working-aged adults? A systematic review. *Stroke*. 2009;40(6):e431-440.
13. Guzik A, Kwolek A, Družbicki M, et al. Return to work after stroke and related factors in Poland and abroad: A literature review. *Work*. 2020;65(2):447-462.
14. Groeneveld IF, Meesters JJL, Arwert HJ, et al. Onderzoeksopzet met analyse van structuur, proces en uitkomsten. Praktijkvariatie in de CVA revalidatie (Research design of an analysis of structure, processes and outcomes. Practice variation in stroke rehabilitation). *Nederlands Tijdschrift voor Revalidatiegeneeskunde*. 2015;37:134-137. Dutch.
15. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol*. 2008;61(4):344-349.
16. van Eeden M, van Heugten C, van Mastrigt GA, et al. The burden of stroke in the Netherlands: estimating quality of life and costs for 1 year poststroke. *BMJ Open*. 2015;5(11):e008220.
17. Wet verbetering poortwachter (Internet). Den Haag: 2001 (cited 2022 nov 28). Available from: <https://wetten.overheid.nl/BWBR0013063/2008-11-01>
18. Sociaal en Cultureel Planbureau - SCP; Centraal Bureau voor de Statistiek - CBS (Internet). Culturele Veranderingen in Nederland (CV) en SCP Leefsituatie Index (SLI) 2018. (cited 2022 June 24) Available from: http://www.scp.nl/Onderzoek/Bronnen/Beknopte_onderzoeksbeschrijvingen/Permanent_onderzoek_naar_de_leefsituatie_POLS

19. Quinn TJ, Langhorne P, Stott DJ. Barthel index for stroke trials: development, properties, and application. *Stroke*. 2011;42(4):1146-1151.

20. Lamers LM, Stalmeier PFM, McDonnell J, et al. Kwaliteit van leven meten in economische evaluaties: het Nederlands EQ-5D-tarief (Measuring quality of life in economic evaluations: the Dutch EQ-5D tariff). *Ned Tijdschr Geneesk*. 2005;149:1574-1578.

21. Duncan PW, Bode RK, Min Lai S, et al. Rasch analysis of a new stroke-specific outcome scale: the Stroke Impact Scale. *Arch Phys Med Rehabil*. 2003;84(7):950-963.

22. van der Zee CH, Priesterbach AR, van der Dussen L, et al. Reproducibility of three self-report participation measures: The ICF Measure of Participation and Activities Screener, the Participation Scale, and the Utrecht Scale for Evaluation of Rehabilitation-Participation. *J Rehabil Med*. 2010;42(8):752-757.

23. Hunger M, Sabariego C, Stollenwerk B, et al. Validity, reliability and responsiveness of the EQ-5D in German stroke patients undergoing rehabilitation. *Qual Life Res*. 2012;21(7):1205-1216.

24. Saeki S, Ogata H, Okubo T, et al. Return to work after stroke. A follow-up study. *Stroke*. 1995;26(3):399-401.

25. Arwert HJ, Schults M, Meesters JJL, et al. Return to Work 2-5 Years After Stroke: A Cross Sectional Study in a Hospital-Based Population. *J Occup Rehabil*. 2017;27(2):239-246.

26. Ashley KD, Lee LT, Heaton K. Return to Work Among Stroke Survivors. *Workplace Health Saf*. 2019;67(2):87-94.

27. Van Patten R, Merz ZC, Mulhauser K, et al. Multivariable prediction of return to work at 6-month follow-up in patients with mild to moderate acute stroke. *Arch Phys Med Rehabil*. 2016;97(12):2061-2067.

28. Wang YC, Kapellusch J, Garg A. Important factors influencing the return to work after stroke. *Work*. 2014;47(4):553-559.

29. Black-Schaffer RM, Osberg JS. Return to work after stroke: development of a predictive model. *Arch Phys Med Rehabil*. 1990;71(5):285-290.

30. Radford K, Grant MI, Sinclair EJ, et al. Describing return to work after stroke: a feasibility trial of 12-month outcomes. *J Rehabil Med*. 2020;52(4):jrm00048.

31. Donker-Cools BH, Daams JG, Wind H, et al. Effective return-to-work interventions after acquired brain injury: A systematic review. *Brain Inj*. 2016;30(2):113-131.

32. Tse T, Lentin P, Douglas J, et al. Understanding activity participation 3-months after stroke: a mixed methodology study. *Disabil Rehabil*. 2020;44(12):2868-2878.

33. Engel-Yeger B, Tse T, Josman N, et al. Scoping review: the trajectory of recovery of participation outcomes following stroke. *Behav Neurol*. 2018;2018:5472018.

34. Norlander A, Iwarsson S, Jönsson AC, et al. Participation in social and leisure activities while reconstructing the self: understanding strategies used by stroke survivors from a long-term perspective. *Disabil Rehabil*. 2022;44(16):4284-4292.

35. Malinowsky C, Olofsson A, Nyman A, et al. Patterns of participation: Facilitating and hindering aspects related to places for activities outside the home after stroke. *Scand J Occup Ther*. 2020;27(3):204-212.

36. Erler KS, Sullivan V, Mckinnon S, et al. Social support as a predictor of community participation after stroke. *Front Neurol*. 2019;10:1013.

Online Resource 1. Baseline characteristics of stroke patients receiving multidisciplinary rehabilitation who had paid employment at the time of stroke

| | n 170 | Included in the current analyses | n 118 | Excluded in the current analyses | p-value* |
|--|--------------------------|--|-------------|--|-------------|
| Sociodemographic characteristics | | | | | |
| Age in years | 169 | 54.2 (11.2) | 115 | 53.9 (13.2) | 0.839 |
| Female sex | 170 | 68 (40.0%) | 118 | 51 (43.2%) | 0.627 |
| Low education level | 167 | 46 (27.5%) | 116 | 43 (37.1%) | 0.093 |
| Living alone | 169 | 30 (17.8%) | 115 | 29 (25.2%) | 0.138 |
| Clinical characteristics | | | | | |
| Ischemic stroke | 167 | 126 (75.4%) | 117 | 90 (76.9%) | 0.888 |
| Number of comorbidities | 131 | 1.0 (1.0) | 97 | 1.0 (2.0) | 0.832 |
| Barthel Index at start rehabilitation ¹ | 92 | 17.0 (9.0) | 70 | 17.0 (9.0) | 0.494 |
| Employment characteristics | | | | | |
| Type of contract | Permanent | 170 | 131 (77.1%) | 118 | 84 (71.2%) |
| | Temporary | | 12 (7.1%) | | 11 (9.3%) |
| | Self-employed | | 20 (11.8%) | | 16 (13.6%) |
| | Other | | 7 (4.1%) | | 7 (5.9%) |
| Number of working hours according to contract | | 169 | 36.0 (11.0) | 116 | 36.0 (16.0) |
| Type of occupation | Office job | | 67 (43.2%) | | 37 (38.1%) |
| | Service job | 155 | 51 (32.9%) | 97 | 27 (27.8%) |
| | Industrial or manual job | | 37 (23.9%) | | 33 (34.0%) |
| Managerial position | | 154 | 18 (11.7%) | 98 | 11 (11.2%) |
| Patient Reported Outcome Measures | | | | | |
| EQ-5D-3L index | | 151 | 0.78 (0.26) | 108 | 0.76 (0.35) |
| EQ-5D-3L VAS | | 159 | 65.0 (26.0) | 108 | 64.0 (25.0) |
| SIS Communication | | 161 | 92.2 (25.0) | 109 | 89.3 (26.8) |
| SIS Mobility ² | | 84 | 84.7 (38.2) | 80 | 86.1 (29.9) |
| SIS Memory and thinking | | 163 | 85.7 (25.0) | 111 | 78.6 (35.7) |
| SIS Mood and emotions | | 84 | 79.2 (23.6) | 82 | 77.8 (22.2) |

Dichotomous variables are described as numbers with percentages (%) and continuous variables as medians with interquartile ranges; *p-values are given of Fisher Exact Tests or Mann-Whitney U Tests, when appropriate.

¹For inpatients only

²Added later to the set of questionnaires

Abbreviations: EQ-5D-3L EuroQoL-5 Dimensions-3 Levels; SIS Stroke Impact Scale; VAS visual analogue scale.

Online Resource 2. Overview of employment status at the different measurement moments of stroke patients with paired measurements.

| | n 126 | 6 months | 12 months | 18 months | 24 months | 30 months |
|--|------------|-------------|--------------|--------------|--------------|--------------|
| Group 1 Paid employment at all measurement moments | | | | | | |
| | 54 (42.9%) | + | + | + | + | + |
| Group 2 Paid employment over time | | | | | | |
| Paid employment at 12, 18, 24 and 30 months | 5 (4.0%) | - | + | + | + | + |
| Paid employment at 18, 24 and 30 months | 0 (0.0%) | - | - | + | + | + |
| Paid employment at 24 and 30 months | 2 (1.6%) | - | - | - | + | + |
| Paid employment at 30 months | 0 (0.0%) | - | - | - | - | + |
| Group 3 No longer paid employment over time | | | | | | |
| No longer paid employment at 12, 18, 24 and 30 months | 6 (4.8%) | + | - | - | - | - |
| No longer paid employment at 18, 24 and 30 months | 9 (7.1%) | + | + | - | - | - |
| No longer paid employment at 24 and 30 months | 20 (15.9%) | + | + | - | - | - |
| No longer paid employment at 30 months | 12 (9.5%) | + | + | + | - | - |
| Group 4 Having paid employment fluctuates over time | | | | | | |
| Paid employment at 6, 18, 24 and 30 months | 1 (0.8%) | + | - | + | + | + |
| Paid employment at 6, 12, 24 and 30 months | 4 (3.2%) | + | + | - | + | + |
| Paid employment at 6, 12, 18 and 30 months | 1 (0.8%) | + | + | - | - | + |
| Paid employment at 6 and 18 months | 1 (0.8%) | + | - | + | - | - |
| Paid employment at 12 months | 2 (1.6%) | - | + | - | - | - |
| Group 5 No longer paid employment after baseline | | | | | | |
| | 9 (7.1%) | - | - | - | - | - |

+ means paid employment at that measurement moment

- means no paid employment at that measurement moment

Online Resource 3. Linear mixed model results of USER-P scales over time

| | Complete USER-P scale | | | USER-P scale without items paid employment | | |
|----------------------------------|-----------------------|--------------|------------------|--|---------------|------------------|
| | β | 95% CI | p-value | β | 95% CI | p-value |
| USER-P Frequency scale | | | | | | |
| Time | -0.62 | -1.97 – 0.74 | 0.370 | 0.45 | -1.03 – 1.93 | 0.550 |
| Paid employment at T24 | 7.81 | 3.31 – 12.31 | 0.001 | 6.77 | 1.82 – 11.71 | 0.007 |
| Time*Paid employment at T24 | 0.52 | -1.23 – 2.28 | 0.557 | -2.19 | -4.11 – -0.27 | 0.026 |
| USER-P Restrictions scale | | | | | | |
| Time | 1.19 | -0.87 – 3.26 | 0.257 | 1.41 | -0.59 – 3.41 | 0.166 |
| Paid employment at T24 | 16.54 | 9.28 – 23.80 | <0.001 | 15.24 | 8.16 – 22.32 | <0.001 |
| Time*Paid employment at T24 | 0.60 | -2.08 – 3.29 | 0.659 | -0.19 | -2.78 – 2.41 | 0.888 |
| USER-P Satisfaction scale | | | | | | |
| Time | 0.45 | -1.50 – 2.39 | 0.652 | 0.09 | -1.84 – 2.02 | 0.930 |
| Paid employment at T24 | 12.78 | 5.48 – 20.07 | 0.001 | 12.44 | 5.12 – 19.77 | 0.001 |
| Time*Paid employment at T24 | 0.31 | -2.17 – 2.80 | 0.804 | 0.10 | -2.38 – 2.59 | 0.935 |

Abbreviations: CI confidence interval; T24 time measurement point 24 months after start of the rehabilitation; USER-P Utrecht Scale for Evaluation of Rehabilitation–Participation.