



Universiteit
Leiden
The Netherlands

Cosmological forecasts on thermal axions, relic neutrinos, and light elements

Giare, W.; Renzi, F.; Melchiorri, A.; Mena, O.; Valentino, E. di

Citation

Giare, W., Renzi, F., Melchiorri, A., Mena, O., & Valentino, E. di. (2022). Cosmological forecasts on thermal axions, relic neutrinos, and light elements. *Monthly Notices Of The Royal Astronomical Society*, 511(1), 1373-1382. doi:10.1093/mnras/stac126

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3567503>

Note: To cite this publication please use the final published version (if applicable).

Cosmological forecasts on thermal axions, relic neutrinos, and light elements

William Giaré¹,^{*} Fabrizio Renzi², Alessandro Melchiorri¹, Olga Mena³ and Eleonora Di Valentino⁴

¹Physics Department and INFN, Università di Roma ‘La Sapienza’, Ple Aldo Moro 2, I-00185 Rome, Italy

²Lorentz Institute for Theoretical Physics, Leiden University, PO Box 9506, NL-2300 RA Leiden, the Netherlands

³IFIC, Universidad de Valencia–CSIC, E-46071 Valencia, Spain

⁴Institute for Particle Physics Phenomenology, Department of Physics, Durham University, Durham DH1 3LE, UK

Accepted 2022 January 13. Received 2021 December 10; in original form 2021 October 8

ABSTRACT

One of the targets of future cosmic microwave background (CMB) and baryon acoustic oscillation measurements is to improve the current accuracy in the neutrino sector and reach a much better sensitivity on extra dark radiation in the early Universe. In this paper, we study how these improvements can be translated into constraining power for well-motivated extensions of the standard model of elementary particles that involve axions thermalized before the quantum chromodynamics (QCD) phase transition by scatterings with gluons. Assuming a fiducial Lambda cold dark matter cosmological model, we simulate future data for Stage-IV CMB-like and Dark Energy Spectroscopic Instrument (DESI)-like surveys and analyse a mixed scenario of axion and neutrino hot dark matter. We further account also for the effects of these QCD axions on the light element abundances predicted by big bang nucleosynthesis. The most constraining forecasted limits on the hot relic masses are $m_a \lesssim 0.92$ eV and $\sum m_\nu \lesssim 0.12$ eV at 95 per cent Confidence Level, showing that future cosmic observations can substantially improve the current bounds, supporting multimessenger analyses of axion, neutrino, and primordial light element properties.

Key words: cosmic background radiation – cosmological parameters – dark matter – early Universe – cosmology: observations.

1 INTRODUCTION

The most compelling solution to the strong CP problem in quantum chromodynamics (QCD) would require the Lagrangian of the standard model of elementary particles to be invariant under an additional global $U(1)_{PQ}$ (Peccei–Quinn) symmetry, spontaneously broken at some energy scale f_a (Nanopoulos 1973; Weinberg 1973, 1975; Belavin et al. 1975; Callan, Dashen & Gross 1976; Jackiw & Rebbi 1976; Peccei & Quinn 1977a, b; Wilczek 1978; Kim 1979; Shifman, Vainshtein & Zakharov 1980; Dine, Fischler & Srednicki 1981; Peccei 1989, 2008; Berezhiani & Khlopov 1991; Berezhiani, Sakharov & Khlopov 1992). The result is an associated pseudo-Nambu–Goldstone boson (PNGB), the *axion* (Weinberg 1978; Kim 1979, 1987; Shifman et al. 1980; Dine et al. 1981; Cheng 1988; Peccei 1989, 2008; Sikivie 2008; Marsh 2016; Di Luzio et al. 2020).

Axions could be copiously produced in the early Universe via both thermal and non-thermal processes (Kibble 1976; Vilenkin 1981; Kibble, Lazarides & Shafi 1982; Sikivie 1982; Vilenkin & Everett 1982; Abbott & Sikivie 1983; Dine & Fischler 1983; Preskill, Wise & Wilczek 1983; Huang & Sikivie 1985; Linde 1985; Seckel & Turner 1985; Davis 1986; Linde & Lyth 1990; Lyth 1990; Vilenkin & Shellard 2000; Hannestad, Mirizzi & Raffelt 2005; Hannestad et al. 2007, 2008, 2010; Melchiorri, Mena & Slosar 2007; Archidiacono et al. 2013; Giusarma et al. 2014;

Archidiacono et al. 2015; Di Valentino et al. 2015, 2016). Those produced non-thermally behave as cold dark matter, while thermal axions (produced by interactions with other particles of the standard model) contribute to the hot dark matter component of the Universe. In what follows, we shall concentrate on the thermal axion scenario.

In the thermal production mechanism, axions can be described by two parameters – the axion coupling constant f_a and the axion mass m_a – related as

$$m_a = \frac{f_\pi m_\pi}{f_a} \frac{\sqrt{R}}{1+R} \simeq 0.6 \text{ eV} \times \frac{10^7 \text{ GeV}}{f_a}, \quad (1)$$

where $R \doteq m_u/m_d \simeq 0.553 \pm 0.043$ is the up-to-down quark mass ratio and $f_\pi \simeq 93$ MeV is the pion decay constant (Zyla et al. 2020).

As long as thermal axions remain relativistic particles, they behave as extra dark radiation, contributing to the effective number of relativistic degrees of freedom N_{eff} , modifying the damping tail of the cosmic microwave background (CMB) temperature angular power spectrum, and changing two important scales at recombination: the sound horizon and the Silk damping scale. Furthermore, the primordial abundances of light elements predicted by the big bang nucleosynthesis (BBN) are also sensitive to extra light species since larger values of N_{eff} would increase the expansion rate of the Universe, leading to a higher freeze-out temperature for weak interactions and implying a higher fraction of primordial helium.

Conversely, when thermal axions become non-relativistic particles, they leave cosmological signatures that are very similar to those originated by massive neutrinos. These axions therefore contribute to

* E-mail: william.giare@gmail.com

the (hot) dark matter component, suppressing structure formation at scales smaller than their free-streaming scale and leaving an imprint on the CMB temperature anisotropies, via the early integrated Sachs–Wolfe effect. So, a large degeneracy between the axion and the total neutrino masses is typically expected.

Updated bounds on thermal axion masses should be then obtained within a realistic mixed hot dark matter scenario, including also massive neutrinos.¹ For the thermal axions decoupling before the QCD phase transition, the most constraining bounds are $m_a < 7.46$ eV and $\sum m_\nu < 0.114$ eV, both at 95 per cent Confidence Level (CL, hereafter) (Giaré et al. 2021). For axions decoupling after the QCD phase transition, these bounds can be improved to $m_a < 0.91$ eV and $\sum m_\nu < 0.105$ eV, both at 95 per cent CL.² These limits were obtained by means of the final release of *Planck* 2018 temperature and polarization data (Aghanim et al. 2020a) in combination with the other most recent (CMB-independent) cosmological observations. While these bounds are able to probe a significant range of the parameter space allowed by direct axion searches, the current constraining power on the total variation of the effective number of relativistic species due to extra dark radiation ($\Delta N_{\text{eff}} \lesssim 0.4$ at 95 per cent CL) represents an important limitation as it is not accurate enough to reveal the presence of thermal axions produced before the QCD transition. These axions would lead to $\Delta N_{\text{eff}} \lesssim 0.1$, which lies well below the present sensitivity to ΔN_{eff} . In this regard, the next generation of CMB experiments is expected to significantly increase the constraints on N_{eff} . In particular, a Stage-IV CMB (CMB-S4) experiment will increase the accuracy on extra dark radiation by almost an order of magnitude, $\Delta N_{\text{eff}} \lesssim 0.06$ (Abazajian et al. 2016), opening the possibility of robustly constraining the mass of QCD axions also for this thermal channel. In addition, future observations of large-scale structure will lead to highly accurate baryon acoustic oscillation (BAO) data, expected to provide an enormous improvement on the neutrino mass bound (Font-Ribera et al. 2014). Interestingly, the combination of CMB-S4 and Dark Energy Spectroscopic Instrument (DESI) should be able to constrain the sum of the neutrino masses at the 2σ level with a precision of $\sigma(\sum m_\nu) \sim 16$ meV (Abazajian et al. 2015; Xu et al. 2021) for $m_\nu \sim 0.06$ eV.

In this work, focusing exclusively on QCD axions produced *before* the QCD phase transition, we analyse realistic mixed hot dark matter scenarios that include also massive neutrinos. The aim is to study the improvement in the constraining power on hot relics expected by the next-generation CMB and BAO observations. We also investigate and discuss the implications for BBN light element primordial abundances up to beryllium-7.

The paper is organized as follows. In Section 2, we review the thermalization processes involved in the axion production. In Section 3, we describe the numerical methods adopted to simulate

the data used for the forecasts. In Section 4, we present and discuss our results. Finally, in Section 5 we present our conclusions.

2 AXION THERMALIZATION

The axion contribution to the effective number of relativistic degrees of freedom

$$\Delta N_{\text{eff}} \simeq \frac{4}{7} \left[\frac{43}{4 g_{\star}(T_d)} \right]^{4/3} \quad (2)$$

depends on the temperature T_d at which axions decouple from the thermal bath via the number of entropic degrees of freedom $g_{\star}(T)$. Notice that the decoupling temperature can be estimated solving the usual freeze-out condition

$$H(T_d) = \Gamma(T_d), \quad (3)$$

where $\Gamma(T)$ is the axion interaction rate and $H(T)$ is the expansion rate, which, in a radiation-dominated Universe, is given by

$$H(T_d) = \sqrt{\frac{4\pi^3}{45} g_{\star}(T_d)} \left(\frac{T_d^2}{M_{\text{pl}}} \right), \quad (4)$$

with $M_{\text{pl}} \simeq 1.22 \times 10^{19}$ GeV the Planck mass and $g_{\star}(T_d) = g_{\star}^{\text{SM}}(T_d) + 1$ the standard model relativistic degrees of freedom with the additional contribution from the axion. On the other hand, considering only the two-body processes with cross-sections $\sigma_i = \sigma(p_i a \leftrightarrow p_j p_k)$ and with all the particles in thermal equilibrium, the axion interaction rate reads (Melchiorri et al. 2007)

$$\Gamma \propto \sum_i n_i \langle v \sigma_i \rangle, \quad (5)$$

where n_i is the number density of p_i , $v \simeq 1$ the relativistic velocity, and the brackets refer to the thermal average.

After decoupling ($T < T_d$), axions maintain a thermal distribution, which basically remains unaffected by other phenomena occurring in the plasma and therefore the current axion number density is simply given by

$$n_a = \frac{g_{\star}(T_0)}{g_{\star}(T_d)} \times \frac{n_\gamma}{2}, \quad (6)$$

with $n_\gamma \simeq 411 \text{ cm}^{-3}$ the present photon density and $g_{\star}(T_0) \simeq 3.91$ the current number of entropic degrees of freedom (we recall that before the neutrino freeze-out $g_{\star} = g_{\star}$).

In the early Universe, there are several processes that can keep the axion in thermal equilibrium with the standard model thermal bath and, depending on whether axions decouple before or after the QCD phase transition ($T_{\text{QCD}} \simeq 160$ MeV), different thermal channels should be considered (D’Eramo et al. 2018; Ferreira & Notari 2018; Arias-Aragón et al. 2021; D’Eramo, Hajkarim & Yun 2021a, b; D’Eramo & Yun 2021; Ferreira, Notari & Rompineve 2021). If axions decouple before the QCD phase transition ($T_d > T_{\text{QCD}}$), the most relevant thermalization channel is provided by the axion–gluon scattering (Hannestad et al. 2005; Melchiorri et al. 2007; Hannestad et al. 2007, 2008, 2010; Archidiacono et al. 2013; Giusarma et al. 2014; Archidiacono et al. 2015; Di Valentino et al. 2015, 2016). Indeed in any QCD axion model, axions necessarily couple to gluons via a model-independent coupling $\alpha_s/(8\pi f_a) a G \tilde{G}$ required to solve the strong CP problem. In this case, the relevant processes for axion thermalization are

- (i) $a + q \leftrightarrow g + q$ and $a + \bar{q} \leftrightarrow g + \bar{q}$,
- (ii) $a + g \leftrightarrow q + \bar{q}$, and
- (iii) $a + g \leftrightarrow g + g$,

¹As robustly indicated by oscillation experiments (de Salas et al. 2018, 2021), neutrinos should be regarded as massive particles and cosmology provides a powerful (albeit indirect) means to constrain their mass (Bond, Efstathiou & Silk 1980; Doroshkevich et al. 1980; Giusarma et al. 2016; Vagnozzi et al. 2017, 2018a, b; De Salas et al. 2018; Vagnozzi 2021; Hagstotz et al. 2021). However, a quite strong correlation between $\sum m_\nu$ and H_0 is typically observed with possible implications for the cosmological tensions (Capozzi et al. 2021; Di Valentino & Melchiorri 2021).

²Note anyway that in the latter case the upper bound on the axion mass is mostly due to the limitation of the range of validity of chiral perturbation theory and weakly depends on data sensitivity.

and the decoupling temperature for this axion population is estimated to be (Di Luzio et al. 2020)

$$T_d \simeq 12.5 \frac{\sqrt{g_*(T)} f_a^2}{\alpha_s^3 M_{\text{pl}}} \text{ GeV}. \quad (7)$$

If instead axions decouple after the QCD phase transition ($T_d < T_{\text{QCD}}$), processes involving pions and nucleons must be considered. In practice, however, nucleons are so rare in the early Universe with respect to pions that the only relevant process is the axion–pion interaction $\pi + \pi \leftrightarrow \pi + a$. In this case, a leading-order computation in chiral perturbation theory predicts the interaction rate (Di Luzio et al. 2020; Di Luzio, Martinelli & Piazza 2021)

$$\Gamma_{a\pi}^{\text{LO}} \simeq 0.215 C_{a\pi}^2 \frac{T^5}{f_a^2 f_\pi^2} h_{\text{LO}} \left(\frac{m_\pi}{T} \right), \quad (8)$$

where $h(x)$ is a rapidly decreasing function of its arguments [and normalized to $h(0) = 1$], and $C_{a\pi}$ is the axion–pion coupling, which is given by

$$C_{a\pi} = \frac{1}{3} \left(\frac{m_d - m_u}{m_u + m_d} + c_d^0 - c_u^0 \right). \quad (9)$$

The axion–pion coupling is a model-dependent quantity, sensitive to the nature of axion–fermion interactions via the axion–quark couplings c_d^0 and c_u^0 . Therefore, the thermal production of axions via pion scattering is strongly model-dependent since the relation between the axion mass and the (decoupling) temperature changes according to the axion–pion interaction strength and, consequently, the thermal production could range between relatively large thermal abundances and negligible ones, depending on the precise value of $C_{a\pi}$. Furthermore, the perturbative approach can only be extended up to temperatures $T_d \lesssim 62 \text{ MeV}$ since above this limit the next-to-leading order corrections become of the same order as the leading order part ($\Gamma_{a\pi}^{\text{NLO}} \simeq \Gamma_{a\pi}^{\text{LO}}$) (Di Luzio et al. 2021). Consequently, no reliable bounds on axion masses can be derived until robust lattice QCD techniques provide a precise answer for any given model in these temperature ranges. In this work, we ignore this model-dependent thermal channel and we exclusively focus on the axion–gluon scatterings.

3 METHOD

In this section, we describe the method followed for our forecasted analyses, focusing on future CMB and BAO observations. In particular, we simulate future data for a CMB-S4-like (Abazajian et al. 2016) observatory and for a DESI-like (Levi et al. 2013; Font-Ribera et al. 2014) BAO survey. These probes are expected to provide scientific results in the next few years and have been carefully designed to improve the constraints on the neutrino sector and other forms of dark radiation in a significant way (Font-Ribera et al. 2014; Abazajian et al. 2015; Abazajian et al. 2016; Abazajian & Heeck 2019). Finally, we also address the effect of additional thermal species on the observational prediction of BBN on light element abundances up to beryllium-7.

All our forecasted data sets make use of the COBAYA software (Torrado & Lewis 2021). The code allows us to build synthetic realization of cosmological data for both CMB and BAO observations and test them again on a given cosmological model. The parameter posteriors have been sampled using the Markov chain Monte Carlo (MCMC) algorithm developed for COSMOMC (Lewis & Bridle 2002; Lewis 2013). The predictions of the theoretical observational probes are calculated using the latest version of the cosmological Boltzmann integrator code CAMB (Lewis, Challinor & Lasenby 2000; Howlett

et al. 2012). To include the effect of the axion–gluon coupling as an additional form of dark radiation, we have modified the CAMB package accordingly to the description detailed in the previous section. The strength of the coupling and its effect on the neutrino sector are functions only of the axion mass that we include as an additional cosmological parameter in our analyses.

To complete this picture, one needs to choose a fiducial cosmological model to build the forecasted data. We perform our forecasts using values of the parameters that are in agreement with the latest *Planck* 2018 constraints for a Lambda cold dark matter (Λ CDM) scenario (Aghanim et al. 2020b). In particular, we choose the following values for the standard six cosmological parameters: $n_s = 0.965$, $\omega_b = 0.0224$, $\omega_c = 0.12$, $H_0 = 67.4$, $\tau = 0.05$, $A_s = 2.1 \times 10^{-9}$, $N_{\text{eff}} = 3.046$, $\sum m_\nu = 0.06 \text{ eV}$, and $m_a = 0 \text{ eV}$. These values are commonly used in the forecasts available in the literature, and, therefore, for the sake of comparison, are the most convenient and useful ones, despite the fact that none of these previous works have considered m_a as a parameter to be constrained.

3.1 CMB-S4 forecasts

We build our forecast for future CMB observations using a well-established and robust method that is now a common practice in cosmological analyses. Using the fiducial model introduced earlier, we compute the angular power spectra of temperature C_ℓ^{TT} , E , and B polarization $C_\ell^{EE, BB}$ and cross-temperature-polarization C_ℓ^{TE} anisotropies. Then, we consider an experimental noise for the temperature angular spectra of the form (Perotto et al. 2006)

$$N_\ell = w^{-1} \exp(\ell(\ell+1)\theta^2/8 \ln 2), \quad (10)$$

where θ is the FWHM angular resolution and w^{-1} is the experimental sensitivity in units of $\mu\text{K arcmin}$. The polarization noise is derived assuming $w_p^{-1} = 2w^{-1}$ (one detector measures two polarization states). The simulated spectra are compared with theoretical ones using the following likelihood $\mathcal{L}$ (Perotto et al. 2006; Cabass et al. 2016):

$$-2 \ln \mathcal{L}_{\text{CMB}} = \sum_\ell (2\ell+1) f_{\text{sky}} \left(\frac{D_\ell}{|C_\ell|} + \ln \frac{|C_\ell|}{|\hat{C}_\ell|} - 3 \right), \quad (11)$$

where $\hat{C}$ and C are the theoretical and simulated spectra (plus noise), respectively, and are defined by

$$|C_\ell| = C_\ell^{TT} C_\ell^{EE} C_\ell^{BB} - (C_\ell^{TE})^2 C_\ell^{BB}, \quad (12)$$

$$|\hat{C}_\ell| = \hat{C}_\ell^{TT} \hat{C}_\ell^{EE} \hat{C}_\ell^{BB} - (\hat{C}_\ell^{TE})^2 \hat{C}_\ell^{BB}, \quad (13)$$

while D is

$$D_\ell = \hat{C}_\ell^{TT} C_\ell^{EE} C_\ell^{BB} + C_\ell^{TT} \hat{C}_\ell^{EE} C_\ell^{BB} + C_\ell^{TT} C_\ell^{EE} \hat{C}_\ell^{BB} - C_\ell^{TE} (C_\ell^{TE} \hat{C}_\ell^{BB} + 2C_\ell^{TE} C_\ell^{BB}). \quad (14)$$

In this study we construct synthetic realizations of CMB data for only one experimental configuration, namely CMB-S4 (see e.g. Abazajian et al. 2016), using $\theta = 3 \text{ arcmin}$ and $w = 1 \mu\text{K arcmin}$. The range of multipoles is $5 \leq \ell \leq 3000$ and the sky coverage of the 40 per cent $f_{\text{sky}} = 0.4$. We do not include CMB lensing derived from trispectrum data.

3.2 DESI (BAO) forecasts

For the future BAO data set, we consider the DESI experiment (Levi et al. 2013). As a tracer for BAO observations, we employ the volume-




Figure 1. The fiducial BAO data sets employed in our forecasts. Error bars refer to 3σ CL uncertainties.

averaged distance defined as

$$D_V(z) \equiv \left(\frac{(1+z)^2 D_A(z)^2 cz}{H(z)} \right)^{\frac{1}{3}}, \quad (15)$$

where D_A is the angular diameter distance and $H(z)$ the Hubble parameter. Assuming the fiducial model described previously, we compute the theoretical values of the ratio D_V/r_s for several redshifts in the range $z = [0.15-1.85]$, where r_s is sound horizon at the photon–baryon decoupling epoch. The uncertainties on D_V/r_s are calculated propagating those for D_A/r_s and $H(z)$ reported in Font-Ribera et al. (2014). The simulated BAO data are compared with the theoretical D_V/r_s values through a multivariate Gaussian likelihood:

$$-2 \ln \mathcal{L}_{\text{BAO}} = \sum (\boldsymbol{\mu} - \hat{\boldsymbol{\mu}}) \mathbf{C}^{-1} (\boldsymbol{\mu} - \hat{\boldsymbol{\mu}})^T, \quad (16)$$

where $\boldsymbol{\mu}$ and $\hat{\boldsymbol{\mu}}$ are the vectors containing the simulated and theoretical values of D_V/r_s at each redshift and $\mathbf{C}$ is their simulated covariance matrix.

It would also be possible to forecast BAO data considering D_A/r_s and $H(z)$ as independent measurements, allowing for stronger constraints. However, some small tension ($\sim 1\sigma$) has been identified between the current constraints from D_A/r_s and $H(z)$ (Addison et al. 2018). Given the difficulty of properly accounting for this small tension between D_A/r_s and $H(z)$, we decided to follow the approach of Allison et al. (2015) and employ the volume-averaged distance for our BAO forecasts. The resulting data set is the same as that obtained in Di Valentino et al. (2018; see also their table 2); a plot representing the forecasted data set is presented in Fig. 1.

3.3 BBN primordial light element abundances

BBN is a cornerstone of the hot big bang cosmology which explains the formation of the first light nuclei (from H up to ${}^7\text{Li}$) by a solid understanding of the nuclear interactions involved in the production of elements.

The set of differential equations that regulate those interactions in the primordial plasma can be solved numerically (Pisanti et al. 2008; Consiglio et al. 2018; Pitrou et al. 2018; Gariazzo et al. 2022) after neutrino decoupling ($T \gtrsim 1$ MeV) up to the end of BBN ($T \sim 10$ keV), yielding the total abundance of primordial elements in terms of their ratio with respect to the hydrogen abundance. BBN provides a natural laboratory to probe new physical scenarios of the early Universe and its predictions can be compared with the primordial abundances of light elements inferred by astrophysical

and cosmological observations. Given current uncertainties, BBN predictions and primordial light element measurements show good agreement (Mossa et al. 2020; Pisanti et al. 2021; Pitrou et al. 2021).³ Notice also that since the BBN epoch ends before recombination, its outcome does not have any impact on the recombination epoch or else on the CMB power spectra. In other words, recombination and BBN are two independent and complementary probes that can be combined to check the consistency of particle interactions in the early Universe. What is relevant for the CMB is the BBN prediction for the helium abundance, which can be used to estimate the baryon density through a simple formula:

$$\Omega_b h^2 = \frac{1 - 0.007125 Y_p^{\text{BBN}}}{273.279} \left(\frac{T_{\text{CMB}}}{2.7255 \text{ K}} \right)^3 \eta_{10}, \quad (17)$$

where $\eta_{10} \doteq 10^{10} n_b/n_\gamma$ is the photon–baryon ratio today, T_{CMB} is the CMB temperature at the present time, and $Y_p^{\text{BBN}} \doteq 4n_{\text{He}}/n_b$ is the helium nucleon fraction defined as the ratio of the helium-4 to the baryon density one. Furthermore, BBN depends on the expansion rate $H(z)$, which sets the value of the temperature of the Universe during the radiation epoch via a function of the radiation density:

$$H(z) \simeq \frac{8\pi G}{3} \rho_{\text{rad}} \simeq \frac{7\pi G}{3} N_{\text{eff}} \left(\frac{4}{11} \right)^{4/3} \rho_\gamma, \quad (18)$$

making BBN a very powerful tool to constrain the total number of relativistic species via $H(z)$.

In this work, we made use of the code PARTHENOPE (Gariazzo et al. 2022). Using the values of N_{eff} , τ_n (the neutron lifetime⁴), and η_{10} (or equivalently $\Omega_b h^2$), the code computes the value of Y_p^{BBN} and other light element abundances. To include the BBN code predictions in our MCMC analysis, we follow the same procedure used by the Planck Collaboration (Aghanim et al. 2020b). Namely, we fix the neutron lifetime and create an interpolation grid varying as $\Omega_b h^2$ and $\Delta N_{\text{eff}} = N_{\text{eff}} - 3.045$ within a given range. We choose these ranges to be $\Delta N_{\text{eff}} \in [-3; 3]$ and $\Omega_b h^2 \in [0.0073; 0.033]$. The neutron lifetime is fixed to $\tau_n = 879.4$ s, corresponding to the latest measurement reported by the Particle Data Group ($\tau_n = 879.4 \pm 0.6$ s) (Zyla et al. 2020).⁵

4 RESULTS

In this section, we discuss the results obtained with the forecasting method presented in the previous section. As a baseline model,

³Nevertheless, there are some discrepancies in the observed abundance of ${}^7\text{Li}$ [which is a factor of ~ 2 smaller than those measured from low-metallicity stars (Cyburt, Fields & Olive 2008; Fields 2011)] and in that of the primordial deuterium [which exhibits a 1.8σ discrepancy with the CMB + BAO value (Pitrou et al. 2021)].

⁴It is worth noting that the interaction rates used in BBN codes assume a prior knowledge of τ_n , which sets the efficiency of nuclear reactions. Therefore, BBN abundances are significantly affected even by a small change in the precise value of this parameter.

⁵This estimate of the neutron lifetime is derived averaging over a large number of measurements. However, beam-only and bottle-only experiments show a 4σ discrepancy in measuring the neutron lifetime, leading, respectively, to $\tau_n = 888.0 \pm 2.0$ s and $\tau_n = 879.2 \pm 0.6$ s (see also the discussion in Salvati et al. 2016). Interestingly, independent constraints can be derived by CMB data only, but these limits are not accurate enough to disentangle the two results ($\tau_n = 851 \pm 60$ s). Nevertheless, the neutron lifetime discrepancy is beyond the scope of this work and we therefore fix its value to that of Zyla et al. (2020), even if this could produce a systematic error in N_{eff} (Capparelli et al. 2018). Other theoretical uncertainties that can affect the axion contribution to the relativistic degrees of freedom in the early Universe are briefly discussed in Appendix A.

Table 1. Results for the Λ CDM + $m_a + \sum m_\nu$ cosmological model and the primordial light element abundances. The constraints on the parameters are at 68 per cent CL, while upper bounds are quoted at 95 per cent CL. We make use of the PARTHENOPE package to compute the BBN predictions.

Parameter	Fiducial value	CMB-S4	CMB-S4 + DESI
$\Omega_b h^2$	0.0224	$0.022\,420 \pm 0.000\,034$	$0.022\,422 \pm 0.000\,034$
$\Omega_c h^2$	0.12	0.12066 ± 0.00062	$0.120\,19 \pm 0.000\,32$
H_0 (km s $^{-1}$ Mpc $^{-1}$)	67.4	$66.94^{+0.63}_{-0.57}$	67.39 ± 0.24
τ	0.05	0.0508 ± 0.0026	0.0508 ± 0.0025
$\log(10^{10} A_S)$	3.044	3.0491 ± 0.0049	3.0478 ± 0.0047
n_s	0.965	0.9647 ± 0.0021	0.9659 ± 0.0017
$\sum m_\nu$ (eV)	0.06	<0.183	<0.122
m_a (eV)	0.0	<1.60	<0.924
Y_p^{BBN}	0.247	$0.247\,268^{+0.000\,052}_{-0.000\,085}$	$0.247\,244^{+0.000\,030}_{-0.000\,042}$
10^5D/H	2.514	2.5211 ± 0.0065	2.5200 ± 0.0063
10^7T/H	0.808	0.8104 ± 0.0022	0.8101 ± 0.0021
10^5He 3/H	1.032	$1.033\,74 \pm 0.000\,95$	$1.033\,58 \pm 0.000\,94$
10^{10}Li 7/H	4.67	4.670 ± 0.015	4.671 ± 0.015
10^{10}Be 7/H	4.40	4.396 ± 0.031	4.398 ± 0.015

we employ an extension of the standard cosmological model that includes both neutrinos and axions as thermal massive relics. We refer to it with Λ CDM + $\sum m_\nu + m_a$. Within this model, we study the improvement on the bounds of QCD axions achievable by future CMB and BAO experiments. As mentioned previously, thermal axions also contribute as additional relativistic species prior to recombination, increasing the value of N_{eff} and thus leading to modification of the standard BBN predictions. Therefore, we also take into account the effect of additional thermal species on the observational predictions of BBN light element abundances. Finally, we shall also compare the constraints on the hot relics and on the helium nucleon fraction Y_p^{BBN} achievable with our simulated data sets without employing the BBN code, testing the dependence of our results on the assumptions adopted for the BBN sector and proving their robustness.

4.1 Mixed hot dark matter: axions and neutrinos

Table 1 summarizes the results obtained from our forecasting methods for future CMB and BAO experiments, while Fig. 2 shows the 68 per cent and 95 per cent CL contour plots for different cosmological parameters.

Using our forecasting data for future CMB-S4 observations, we derive the 95 per cent CL upper bounds on thermal relics of $\sum m_\nu < 0.183$ eV and $m_a < 1.60$ eV. These values should be compared with those derived in Giarè et al. (2021) for the same cosmological model, exploiting the last CMB data release provided by the Planck Collaboration. In particular, one can appreciate that future CMB-S4 measurements are expected to improve the current bounds on the axion–gluon interaction scenario by a factor of ~ 5 , while we estimate the improvement in the constraining power on the neutrino sector to be ~ 2 . This enhancement in the constraining power on thermal relic masses is mostly due to the much higher sensitivity to the effective number or relativistic degrees of freedom N_{eff} expected from future CMB measurements.⁶

⁶We recall that, while the current *Planck* data lead to a 95 per cent CL upper limit of $\Delta N_{\text{eff}} \lesssim 0.4$, future CMB-S4-like experiments are expected to bring this upper limit down by a factor of ~ 10 , resulting in a much more tighter limit on dark radiation, $\Delta N_{\text{eff}} \lesssim 0.06$, and eV-scale light relics (DePorzio et al. 2021).

Notice that, due to the degeneracy between the axion and the neutrino masses discussed in the introductory section, the contours in $(\sum m_\nu, m_a)$ show a clear anticorrelation. Furthermore, these two parameters show very similar degeneracies with other cosmological parameters such as H_0 , σ_8 , and Ω_m . It is well known that hot thermal particles suppress structure formation at small scales and therefore galaxy clustering information becomes crucial to set bounds on the amount of dark matter in the form of these relics. As discussed in Giarè et al. (2021), the largest impact on CMB bounds on hot relics arises from the inclusion of the large-scale structure information from BAO measurements. For this reason, here, together with the likelihood for future CMB-S4 observations, we consider also a likelihood for future BAO measurements from the DESI-like experiment. Combining our simulated CMB-S4 and DESI forecasts, we obtain a further improvement in the cosmological constraining power for thermal relics, reaching the 95 per cent CL limits $m_a < 0.924$ eV and $\sum m_\nu < 0.122$ eV. In this case, these bounds can be compared with those obtained for current *Planck*+BAO real data (Giarè et al. 2021), observing an improvement of a factor of ~ 8 and ~ 1.5 in the sensitivity to the axion and neutrino masses, respectively.

Our results clearly state that future cosmological observations can substantially improve the current constraints on m_a , exploring a much larger range of the parameter space than currently allowed for QCD thermal axions and reaching the sub-eV mass range. Conversely, when axions are included in the picture as additional thermal species, the possibility to detect the expected minimum neutrino mass of 0.06 eV is no longer there, and only upper bounds, close to the inverted mass ordering prediction, can be derived.

4.2 Primordial abundances of light elements

Thermal axions contribute to the effective number of relativistic degrees of freedom, modifying the expansion rate at the radiation epoch and affecting, indirectly, the canonical BBN predictions. Even though the latest results of the Planck Collaboration place tight bounds on both the baryon density ($\Omega_b h^2 = 0.0224 \pm 0.0001$) and N_{eff} (limiting the amount of additional relativistic degrees of freedom to $\Delta N_{\text{eff}} \lesssim 0.4$), the impact of axions on the helium fraction is extremely small and the *Planck* uncertainties on $\Omega_b h^2$ are still too large to provide robust theoretical predictions on the helium abundance in the presence of the axion. However, the next generation




Figure 2. Marginalized 2D and 1D posteriors for different cosmological parameters obtained from the forecasting data and methods.

of CMB and BAO observations will substantially improve the bounds on the baryon energy density by a factor of 2, strongly reducing the theoretical uncertainties on Y_p^{BBN} , and, possibly, allowing us to test signatures of the axion in the primordial abundances. Fig. 3 shows a comparison between CMB-S4 and *Planck* in determining the helium fraction. In particular, we show the theoretical helium fraction predictions in the $\Lambda\text{CDM} + m_a + \sum m_\nu$ cosmological model as a function of the axion mass (or, equivalently, as a function of the axion contribution to ΔN_{eff}) together with the 2D marginalized posterior distribution obtained for the CMB-S4 and CMB-S4+DESI simulated data. Note that the BBN predictions introduce a strong correlation between the axion mass and the helium fraction (Y_p^{BBN}) that, combined with the substantial improvement in the constraining power expected by CMB-S4 and DESI, suggests that the BBN could be a useful tool to make predictions on hot axions and that astrophysical measurements of the primordial fraction of helium could be used as an independent test together with the cosmological observations. For this reason, we included all the BBN light elements in our analysis. We provide the 68 per cent CL results for the other light elements up to beryllium-7 in Table 1. It should be noted,

however, that these results are derived without considering the experimental error in the measurement of the neutron lifetime τ_n . This error could dominate the total error budget, enlarging the theoretical uncertainties on the BBN predictions for Y_p^{BBN} by a factor of ~ 3 [$\Delta Y_p^{\text{BBN}}(\Delta \tau_n) \simeq 0.00012$], producing the same effect as an extra dark radiation component (Capparelli et al. 2018). Consequently a large degeneracy between the axion mass and the neutron lifetime is expected and this effect may change the correlations between the primordial helium fraction and the axion mass. For this reason, to prove the robustness of our results on hot massive relics, it is mandatory to follow also a very conservative approach and study the impact of additional hot relics on the abundances of primordial elements without assuming the BBN theoretical predictions but leaving all the parameters varying in uninformative flat priors. We therefore analyse a cosmological model where, together with axions and massive neutrinos, we also include the abundance of primordial helium as an additional free parameter. We refer to this model as $\Lambda\text{CDM} + m_a + \sum m_\nu + Y_p^{\text{BBN}}$ and report the results obtained with our CMB and BAO forecasting data in Table 2. In this case, the 68 per cent and 95 per cent CL marginalized contours in the plane




Figure 3. Theoretical helium fraction predictions in the Λ CDM + $m_a + \sum m_\nu$ cosmological model. The black solid line represents the helium fraction as a function of the axion mass (with the corresponding ΔN_{eff} on the top axis) obtained by fixing $\Omega_b h^2 = 0.0224$. The green (grey) region represents the 3σ uncertainties on Y_p^{BBN} by CMB-S4 (*Planck*). The vertical lines are the 95 per cent CL upper limits on m_a from current cosmological data (Giarè et al. 2021) and from CMB and BAO future experiments obtained in this work, together with the respective 68 per cent and 95 per cent CL contours.

Table 2. Results for the Λ CDM + $m_a + \sum m_\nu + Y_p^{\text{BBN}}$ case (i.e. leaving the helium nucleon fraction as a free parameter of the model, without assuming the BBN theoretical predictions). The constraints on parameters are at 68 per cent CL, while the quoted upper bounds are at 95 per cent CL.

Parameter	Fiducial value	CMB-S4	CMB-S4 + DESI
$\Omega_b h^2$	0.0224	$0.022\,399 \pm 0.000\,050$	$0.022\,406 \pm 0.000\,050$
$\Omega_c h^2$	0.12	$0.120\,70 \pm 0.000\,62$	$0.120\,20 \pm 0.000\,31$
H_0 (km s $^{-1}$ Mpc $^{-1}$)	67.4	$66.90^{+0.65}_{-0.57}$	67.37 ± 0.24
τ	0.05	0.0506 ± 0.0026	0.0507 ± 0.0025
$\log(10^{10} A_s)$	3.044	3.0486 ± 0.0049	3.0474 ± 0.0047
n_s	0.965	0.9635 ± 0.0034	0.9649 ± 0.0030
$\sum m_\nu$ (eV)	0.06	<0.183	<0.120
m_a (eV)	0.0	<1.63	<0.991
Y_p^{BBN}	0.247	$0.2458^{+0.0057}_{-0.0058}$	$0.2460^{+0.0057}_{-0.0058}$

(m_a , Y_p^{BBN}) are shown in Fig. 4. On removing the BBN predictions, the strong positive correlation between the axion mass and the helium fraction Y_p^{BBN} is relaxed as well. Furthermore, the bounds on the helium fraction are much less constraining, with 68 per cent CL bounds of $Y_p^{\text{BBN}} = 0.2458^{+0.0057}_{-0.0058}$ and $Y_p^{\text{BBN}} = 0.2460^{+0.0057}_{-0.0058}$ for CMB-S4 and CMB-S4+DESI, respectively.

On the other hand, the constraints on hot dark matter are basically unchanged. Exploiting our forecasting data for future CMB-S4 observations, we can still derive the 95 per cent CL upper bounds $m_a < 1.63$ eV and $\sum m_\nu < 0.183$ eV for axions and neutrinos, respectively. The upper limit on the total neutrino mass is exactly the same as that derived including the BBN code as well as the upper bound on the total axion mass. Similarly, combining future CMB-S4 and BAO data the upper bound on neutrino masses is unchanged

($\sum m_\nu < 0.120$ eV at 95 per cent CL), while the upper bound on axions is only slightly worsened to $m_a < 0.991$ eV at 95 per cent CL. These results prove that the impact of the BBN uncertainties on axion and neutrino masses is negligible and therefore the extraction of both m_a and m_ν does not rely on the assumptions adopted for the neutrino lifetime.

5 DISCUSSION AND CONCLUSIONS

One of the declared targets of the next-generation CMB and BAO cosmic observations is to improve the current constraining power on the neutrino sector, reaching a much better sensitivity both on their total mass and on extra dark radiation, severely constraining additional contribution to the number of relativistic degrees of




Figure 4. Marginalized 2D posteriors in the plane (m_a, Y_p^{BBN}) when the BBN predictions are relaxed and the helium fraction is considered as a free parameter.

freedom in the early Universe. In this study, we have translated the expected experimental improvements into constraining power for well-motivated physical extensions of the standard model of elementary particles, involving QCD axions as the solution to the strong CP problem. In particular, we have focused on axions produced thermally before the QCD phase transition via scattering with free gluons and analysed mixed hot dark matter scenarios that include also neutrinos as additional massive relics. Assuming a fiducial Λ CDM cosmological model and following robust numerical procedures, we have simulated forecasting data for future CMB and BAO observations, focusing in particular on CMB-S4-like and DESI-like experiments. Exploiting these simulated data, we have studied the improvements in the cosmological constraints on axion and neutrino mixed hot dark matter scenarios. We have shown that future CMB-S4 measurements are expected to improve the current cosmological bounds on axion–gluon interactions by a factor of ~ 5 . The improvement in the constraining power on the neutrino sector is ~ 2 , and the expected 95 per cent CL upper bounds on hot relic masses are $\sum m_\nu < 0.183$ eV and $m_a < 1.60$ eV. Since hot thermal particles such as axions and neutrinos suppress structure formation at small scales, galaxy clustering information becomes crucial to set bounds on the amount of hot dark matter in the form of thermal relics. For this reason, we combine the CMB-S4 simulated data with our likelihood for future BAO DESI-like experiments, showing that in this case the bounds on thermal relics can be further improved to $m_a < 0.924$ eV and $\sum m_\nu < 0.122$ eV, both at 95 per cent CL. Interestingly, future cosmic observers can substantially improve the current constraints on m_a , reaching the sub-eV mass range. Conversely, when axions are included in the picture as additional thermal species, a 2σ detection of the neutrino mass for a fiducial value of $\sum m_\nu = 0.06$ eV (which is another declared target of the next-generation CMB and BAO experiments) is excluded. Only upper bounds (close to the inverted mass ordering prediction) can be derived in the former case. Finally, since thermal axions also contribute as additional relativistic species prior to recombination, increasing the value of N_{eff} and thus leading to a modification of the standard BBN predictions, we have also taken into account the effect of an additional thermal species on the observational prediction of BBN on light element abundances up

to beryllium-7. We have compared the constraints on hot relics achievable within our simulated data sets with and without employing the PARTHENOPE BBN code for computing the theoretical predictions of the different abundances of primordial elements. We have shown that our results do not rely strongly on the assumptions adopted in the BBN sector, definitively proving their robustness. We conclude supporting and underlying the relevance of multimessenger searches of axions, neutrinos, and primordial light element measurements. Indeed, cosmology-independent limits on the axion and neutrino masses, combined with precise astrophysical measurements of light elements, may provide an important cosmological test for checking the BBN predictions. On the other hand, future cosmic observations should also be able to probe scenarios with hot axions with masses $m_a \gtrsim 1$ eV and a missing piece of evidence would constrain the axion mass at the sub-eV level, favouring the normal ordering as the one governing the mass pattern of neutral fermions. In the same multimessenger spirit, future cosmology-independent probes of neutrino masses (i.e. future terrestrial double beta decay and/or long baseline neutrino experiments) will play, even if indirectly, a crucial role in axion searches. Complementarity between astrophysical and cosmological axion searches has been carefully explored in a very recent study by Green, Guo & Wallisch (2021), which appeared while this manuscript was being completed. Our results agree fairly well with those obtained by the authors of the aforementioned reference.

ACKNOWLEDGEMENTS

We thank Francesco D’Eramo, Seokhoon Yun, Sadra Hajkarim, and Eoin O Colgain for useful comments and suggestions.

WG and AM were supported by ‘Theoretical Astroparticle Physics’ (TASP), iniziativa specifica INFN. FR acknowledges support from the NWO and the Dutch Ministry of Education, Culture and Science (OCW), and from the D-ITP consortium, a programme of the NWO that is funded by the OCW. OM was supported by the Spanish grants FPA2017-85985-P and PROMETEO/2019/083, and the European ITN project HIDEEN (H2020-MSCA-ITN-2019//860881-HIDEEN). EDV acknowledges the support of the Addison-Wheeler Fellowship awarded by the Institute of Advanced Study at Durham University. In this work, we made use of the following PYTHON packages that are not mentioned in the text: SCIPY (Virtanen et al. 2020) for numerical sampling of the statistical distributions involved in our data analysis; GetDist (Lewis 2019), a tool for the analysis of MCMC samples that employs MATPLOTLIB (Hunter 2007) for the realization of the plots in the paper; and NUMPY (Oliphant 2006) for numerical linear algebra.

DATA AVAILABILITY

The simulated data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Abazajian K. N., Heeck J., 2019, *Phys. Rev. D*, 100, 075027
- Abazajian K. N. et al., 2015, *Astropart. Phys.*, 63, 66
- Abazajian K. N. et al., 2016, preprint ([arXiv:1610.02743](https://arxiv.org/abs/1610.02743))
- Abbott L. F., Sikivie P., 1983, *Phys. Lett. B*, 120, 133
- Addison G. E., Watts D. J., Bennett C. L., Halpern M., Hinshaw G., Weiland J. L., 2018, *ApJ*, 853, 119
- Aghanim N. et al., 2020a, *A&A*, 641, A1
- Aghanim N. et al., 2020b, *A&A*, 641, A6
- Allison R., Caucal P., Calabrese E., Dunkley J., Louis T., 2015, *Phys. Rev. D*, 92, 123535

- Archidiacono M., Hannestad S., Mirizzi A., Raffelt G., Wong Y. Y. Y., 2013, *J. Cosmol. Astropart. Phys.*, 10, 020
- Archidiacono M., Basse T., Hamann J., Hannestad S., Raffelt G., Wong Y. Y. Y., 2015, *J. Cosmol. Astropart. Phys.*, 05, 050
- Arias-Aragón F., D'Eramo F., Ferreira R. Z., Merlo L., Notari A., 2021, *J. Cosmol. Astropart. Phys.*, 03, 090
- Belavin A. A., Polyakov A. M., Schwartz A. S., Tyupkin Y. S., 1975, *Phys. Lett. B*, 59, 85
- Bereziani Z. G., Khlopov M. Y., 1991, *Z. Phys. C*, 49, 73
- Bereziani Z. G., Sakharov A. S., Khlopov M. Y., 1992, *Sov. J. Nucl. Phys.*, 55, 1063
- Bond J. R., Efstathiou G., Silk J., 1980, *Phys. Rev. Lett.*, 45, 1980
- Cabass G., Pagano L., Salvati L., Gerbino M., Giusarma E., Melchiorri A., 2016, *Phys. Rev. D*, 93, 063508
- Callan C. G. Jr, Dashen R. F., Gross D. J., 1976, *Phys. Lett. B*, 63, 334
- Capozzi F., Di Valentino E., Lisi E., Marrone A., Melchiorri A., Palazzo A., 2021, *Phys. Rev. D*, 104, 083031
- Capparelli L., Di Valentino E., Melchiorri A., Chluba J., 2018, *Phys. Rev. D*, 97, 063519
- Cheng H.-Y., 1988, *Phys. Rep.*, 158, 1
- Consiglio R., de Salas P. F., Mangano G., Miele G., Pastor S., Pisanti O., 2018, *Comput. Phys. Commun.*, 233, 237
- Cybart R. H., Fields B. D., Olive K. A., 2008, *J. Cosmol. Astropart. Phys.*, 11, 012
- Davis R. L., 1986, *Phys. Lett. B*, 180, 225
- DePorzio N., Xu W. L., Muñoz J. B., Dvorkin C., 2021, *Phys. Rev. D*, 103, 023504
- D'Eramo F., Yun S., 2021, preprint ([arXiv:2111.12108](https://arxiv.org/abs/2111.12108))
- D'Eramo F., Ferreira R. Z., Notari A., Bernal J. L., 2018, *J. Cosmol. Astropart. Phys.*, 11, 014
- D'Eramo F., Hajkarim F., Yun S., 2021a, preprint ([arXiv:2108.04259](https://arxiv.org/abs/2108.04259))
- D'Eramo F., Hajkarim F., Yun S., 2021b, *J. High Energy Phys.*, 10, 224
- De Salas P. F., Gariazzo S., Mena O., Ternes C. A., Tórtola M., 2018, *Front. Astron. Space Sci.*, 5, 36
- de Salas P. F., Forero D. V., Gariazzo S., Martínez-Miravé P., Mena O., Ternes C. A., Tórtola M., Valle J. W. F., 2021, *J. High Energy Phys.*, 02, 071
- Dine M., Fischler W., 1983, *Phys. Lett. B*, 120, 137
- Dine M., Fischler W., Srednicki M., 1981, *Phys. Lett. B*, 104, 199
- Di Luzio L., Giannotti M., Nardi E., Visinelli L., 2020, *Phys. Rep.*, 870, 1
- Di Luzio L., Martinelli G., Piazza G., 2021, *Phys. Rev. Lett.*, 126, 241801
- Di Valentino E., Melchiorri A., 2021, preprint ([arXiv:2112.02993](https://arxiv.org/abs/2112.02993))
- Di Valentino E., Gariazzo S., Giusarma E., Mena O., 2015, *Phys. Rev. D*, 91, 123505
- Di Valentino E., Giusarma E., Lattanzi M., Mena O., Melchiorri A., Silk J., 2016, *Phys. Lett. B*, 752, 182
- Di Valentino E., Holz D. E., Melchiorri A., Renzi F., 2018, *Phys. Rev. D*, 98, 083523
- Doroshkevich A. G., Zeldovich Y. B., Sunyaev R. A., Khlopov M., 1980, *Sov. Astron. Lett.*, 6, 252
- Ferreira R. Z., Notari A., 2018, *Phys. Rev. Lett.*, 120, 191301
- Ferreira R. Z., Notari A., Rompineve F., 2021, *Phys. Rev. D*, 103, 063524
- Fields B. D., 2011, *Annu. Rev. Nucl. Part. Sci.*, 61, 47
- Font-Ribera A., McDonald P., Mostek N., Reid B. A., Seo H.-J., Slosar A., 2014, *J. Cosmol. Astropart. Phys.*, 05, 023
- Gariazzo S., de Salas P. F., Pisanti O., Consiglio R., 2022, *Comput. Phys. Commun.*, 271, 108205
- Ghodbane N., Martyn H.-U., 2002, preprint ([arXiv:hep-ph/0201233](https://arxiv.org/abs/hep-ph/0201233))
- Giarè W., Di Valentino E., Melchiorri A., Mena O., 2021, *MNRAS*, 505, 2703
- Giusarma E., Di Valentino E., Lattanzi M., Melchiorri A., Mena O., 2014, *Phys. Rev. D*, 90, 043507
- Giusarma E., Gerbino M., Mena O., Vagnozzi S., Ho S., Freese K., 2016, *Phys. Rev. D*, 94, 083522
- Green D., Guo Y., Wallisch B., 2021, preprint ([arXiv:2109.12088](https://arxiv.org/abs/2109.12088))
- Hagstotz S., de Salas P. F., Gariazzo S., Gerbino M., Lattanzi M., Vagnozzi S., Freese K., Pastor S., 2021, *Phys. Rev. D*, 104, 123524
- Hannestad S., Mirizzi A., Raffelt G., 2005, *JCAP*, 07, 002
- Hannestad S., Mirizzi A., Raffelt G. G., Wong Y. Y. Y., 2007, *J. Cosmol. Astropart. Phys.*, 08, 015
- Hannestad S., Mirizzi A., Raffelt G. G., Wong Y. Y. Y., 2008, *J. Cosmol. Astropart. Phys.*, 04, 019
- Hannestad S., Mirizzi A., Raffelt G. G., Wong Y. Y. Y., 2010, *J. Cosmol. Astropart. Phys.*, 08, 001
- Howlett C., Lewis A., Hall A., Challinor A., 2012, *J. Cosmol. Astropart. Phys.*, 1204, 027
- Huang M. C., Sikivie P., 1985, *Phys. Rev. D*, 32, 1560
- Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
- Jackiw R., Rebbi C., 1976, *Phys. Rev. Lett.*, 37, 172
- Kibble T. W. B., 1976, *J. Phys. A*, 9, 1387
- Kibble T. W. B., Lazarides G., Shafi Q., 1982, *Phys. Rev. D*, 26, 435
- Kim J. E., 1979, *Phys. Rev. Lett.*, 43, 103
- Kim J. E., 1987, *Phys. Rep.*, 150, 1
- Levi M. et al., 2013, preprint ([arXiv:1308.0847](https://arxiv.org/abs/1308.0847))
- Lewis A., 2013, *Phys. Rev. D*, 87, 103529
- Lewis A., 2019, preprint ([arXiv:1910.13970](https://arxiv.org/abs/1910.13970))
- Lewis A., Bridle S., 2002, *Phys. Rev. D*, 66, 103511
- Lewis A., Challinor A., Lasenby A., 2000, *ApJ*, 538, 473
- Linde A. D., 1985, *Phys. Lett. B*, 158, 375
- Linde A. D., Lyth D. H., 1990, *Phys. Lett. B*, 246, 353
- Lyth D. H., 1990, *Phys. Lett. B*, 236, 408
- Marsh D. J. E., 2016, *Phys. Rep.*, 643, 1
- Melchiorri A., Mena O., Slosar A., 2007, *Phys. Rev. D*, 76, 041303
- Mossa V. et al., 2020, *Nature*, 587, 210
- Nanopoulos D. V., 1973, *Lett. Nuovo Cimento*, 8S2, 873
- Oliphant T., 2006, NumPy: A Guide to NumPy. Trelgol Publishing, USA
- Peccei R. D., 1989, in Jarlskog C., ed., CP Violation, Advanced Series on Directions in High Energy Physics, Vol. 3. World Scientific, Singapore, p. 501
- Peccei R. D., 2008, in Kuster M., ed., Axons, Lecture Notes in Physics, Vol. 741. Springer-Verlag, Berlin, p. 3
- Peccei R. D., Quinn H. R., 1977a, *Phys. Rev. D*, 16, 1791
- Peccei R. D., Quinn H. R., 1977b, *Phys. Rev. Lett.*, 38, 1440
- Perotto L., Lesgourgues J., Hannestad S., Tu H., Wong Y. Y. Y., 2006, *J. Cosmol. Astropart. Phys.*, 10, 013
- Pisanti O., Cirillo A., Esposito S., Iocco F., Mangano G., Miele G., Serpico P. D., 2008, *Comput. Phys. Commun.*, 178, 956
- Pisanti O., Mangano G., Miele G., Mazzella P., 2021, *J. Cosmol. Astropart. Phys.*, 04, 020
- Pitrou C., Coc A., Uzan J.-P., Vangioni E., 2018, *Phys. Rep.*, 754, 1
- Pitrou C., Coc A., Uzan J.-P., Vangioni E., 2021, *MNRAS*, 502, 2474
- Preskill J., Wise M. B., Wilczek F., 1983, *Phys. Lett. B*, 120, 127
- Salvati L., Pagano L., Consiglio R., Melchiorri A., 2016, *J. Cosmol. Astropart. Phys.*, 03, 055
- Schwarz D. J., 2003, *Ann. Phys.*, 12, 220
- Seckel D., Turner M. S., 1985, *Phys. Rev. D*, 32, 3178
- Shifman M. A., Vainshtein A. I., Zakharov V. I., 1980, *Nucl. Phys. B*, 166, 493
- Sikivie P., 1982, *Phys. Rev. Lett.*, 48, 1156
- Sikivie P., 2008, in Kuster M., ed., Axons, Lecture Notes in Physics, Vol. 741. Springer-Verlag, Berlin, p. 19
- Torrado J., Lewis A., 2021, *J. Cosmol. Astropart. Phys.*, 5, 057
- Vagnozzi S., 2021, *J. Cosmol. Astropart. Phys.*, 2021, 057
- Vagnozzi S., Giusarma E., Mena O., Freese K., Gerbino M., Ho S., Lattanzi M., 2017, *Phys. Rev. D*, 96, 123503
- Vagnozzi S., Brinckmann T., Archidiacono M., Freese K., Gerbino M., Lesgourgues J., Sprenger T., 2018a, *J. Cosmol. Astropart. Phys.*, 09, 001
- Vagnozzi S., Dhawan S., Gerbino M., Freese K., Goobar A., Mena O., 2018b, *Phys. Rev. D*, 98, 083501
- Vilenkin A., 1981, *Phys. Rev. D*, 24, 2082
- Vilenkin A., Everett A. E., 1982, *Phys. Rev. Lett.*, 48, 1867
- Vilenkin A., Shellard E. P. S., 2000, Cosmic Strings and Other Topological Defects. Cambridge Univ. Press, Cambridge
- Virtanen P. et al., 2020, *Nat. Methods*, 17, 261
- Weinberg S., 1973, *Phys. Rev. Lett.*, 31, 494

- Weinberg S., 1975, *Phys. Rev. D*, 11, 3583
 Weinberg S., 1978, *Phys. Rev. Lett.*, 40, 223
 Wilczek F., 1978, *Phys. Rev. Lett.*, 40, 279
 Xu W. L., DePorzio N., Muñoz J. B., Dvorkin C., 2021, *Phys. Rev. D*, 103, 023503
 Zyla P. A. et al., 2020, *Prog. Theor. Exp. Phys.*, 2020, 083C01

APPENDIX: DEGREES OF FREEDOM IN THE EARLY UNIVERSE

The results derived in this paper assume a minimal extension of the standard model that includes QCD axions thermalized at high temperatures, prior to the QCD phase transition. However, little is known about the number of degrees of freedom above the QCD epoch and particularly around the electroweak transition. In many extensions of the standard model of particles, such as supersymmetry, additional degrees of freedom in the early Universe are in principle allowed, possibly changing the contributions of QCD axions to the different cosmological observables and, consequently, our forecasted bounds.

In this appendix, we briefly discuss what happens to our forecasted limits in many-degrees-of-freedom models of the early Universe. In particular, we consider minimal supersymmetrical extensions of the standard model (MSSM) (Ghodbane & Martyn 2002) where, above the electroweak scale, the relativistic degrees of freedom are more than doubled since all supersymmetric partners can eventually be excited. More precisely, when the temperature is larger than all particle masses, within the MSSM we have $g_{\star}^{\text{MSSM}} = 228.75$ (Schwarz 2003), while within the SM we have $g_{\star}^{\text{SM}} = 106.75$.

Retracing our analysis in such scenarios, it is clear that if the axion decouples from the thermal bath at sufficiently high temperatures, its residual number density is suppressed by the additional entropic degrees of freedom and consequently also its contribution to the effective number of relativistic degrees of freedom will be smaller. However, although the supersymmetric model considered here has light superpartners compared with the other models proposed in the literature, the lightest supersymmetric particle in the picture is the lighter neutralino with a mass of ~ 95 GeV. Therefore, the




Figure A1. Axion contribution to ΔN_{eff} in a minimal supersymmetrical extension of the standard model.

additional supersymmetric degrees of freedom can be excited only at temperatures of the order of the electroweak scale ($T \gtrsim 100$ GeV) that, by equation (7), correspond to tiny axion masses $m_a \lesssim 0.5$ eV. As shown in Fig. A1, these mass values are beyond (even though close to) the sensitivity expected by future cosmological and astrophysical experiments. None the less, these effects can be relevant to the abundance of primordial helium predicted by the BBN. Indeed, in the limit of high decoupling temperatures ($T \gg 100$ GeV), the contribution of the axion to the effective number of relativistic degrees of freedom is reduced from $\Delta N_{\text{eff}} \simeq 0.027$ to $\Delta N_{\text{eff}} \lesssim 0.01$ (see also Fig. A1). This difference changes the theoretical predictions shown in Fig. 3 for $Y_{\text{p}}^{\text{BBN}}$ versus m_a . Therefore, we recommend caution when applying our forecasted limits on primordial abundances to scenarios that involve additional degrees of freedom in the early Universe.

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.