



Universiteit
Leiden
The Netherlands

Comparative analysis of right ventricular strain in Fabry cardiomyopathy and sarcomeric hypertrophic cardiomyopathy

Meucci, M.C.; Lillo, R.; Lombardo, A.; Lanza, G.A.; Bootsma, M.; Butcher, S.C.; ... ; Graziani, F.

Citation

Meucci, M. C., Lillo, R., Lombardo, A., Lanza, G. A., Bootsma, M., Butcher, S. C., ... Graziani, F. (2022). Comparative analysis of right ventricular strain in Fabry cardiomyopathy and sarcomeric hypertrophic cardiomyopathy. *European Heart Journal - Cardiovascular Imaging*. doi:10.1093/ehjci/jeac151







Version: Publisher's Version

License: [Creative Commons CC BY-NC 4.0 license](https://creativecommons.org/licenses/by-nc/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3567710>

Note: To cite this publication please use the final published version (if applicable).

Comparative analysis of right ventricular strain in Fabry cardiomyopathy and sarcomeric hypertrophic cardiomyopathy

Maria Chiara Meucci ^{1,2}, Rosa Lillo^{2,3}, Antonella Lombardo^{2,4}, Gaetano A. Lanza^{2,4}, Marianne Bootsma¹, Steele C. Butcher ^{1,5}, Massimo Massetti^{2,4}, Raffaele Manna ^{2,6}, Jeroen J. Bax ^{1,7}, Filippo Crea ^{2,4}, Nina Ajmone Marsan¹, and Francesca Graziani ^{4*}

¹Department of Cardiology, Leiden University Medical Center, Albinusdreef 2 2300 RC, Leiden, The Netherlands; ²Catholic University of the Sacred Heart, Largo F. Vito 1, Rome 00168, Italy; ³Department of Emergency Medicine, Fondazione Policlinico Universitario A. Gemelli IRCCS, Largo A. Gemelli 8, 00168 Rome, Italy; ⁴Department of Cardiovascular Medicine, Fondazione Policlinico Universitario A. Gemelli IRCCS, Largo A. Gemelli 8, 00168 Rome, Italy; ⁵Department of Cardiology, Royal Perth Hospital, Victoria Square, 6000 Perth WA, Australia; ⁶Institute of Internal Medicine, Periodic Fever and Rare Diseases Research Centre, Fondazione Policlinico A. Gemelli IRCCS, Largo A. Gemelli 8, 00168 Rome, Italy; and ⁷Heart Center, University of Turku and Turku University Hospital, Kiinamyllynkatu 4-8, 20521 Turku, Finland

Received 9 March 2022; revised 22 June 2022; accepted 15 July 2022

Aims

To perform a comparative analysis of right ventricle (RV) myocardial mechanics, assessed by 2D speckle-tracking echocardiography (2D-STE), between patients with Fabry disease and patients with sarcomeric disease.

Methods and results

Patients with Fabry cardiomyopathy (FC) ($n = 28$) were compared with patients with sarcomeric hypertrophic cardiomyopathy (HCM), matched for degree of left ventricle hypertrophy (LVH) and demographic characteristics ($n = 112$). In addition, patients with Fabry disease and no LVH [phenotype-negative carriers of pathogenic α -galactosidase gene mutations (GLA LVH-)] ($n = 28$) were compared with age and sex-matched carriers of sarcomeric gene mutations without LVH [Phenotype-negative carriers of pathogenic sarcomeric gene mutations (Sarc LVH-)] ($n = 56$). Standard echocardiography and 2D-STE were performed in all participants. Despite a subtle impairment of RV global longitudinal strain (RV-GLS) was common in both groups, patients with FC showed a more prominent reduction of RV free wall longitudinal strain (RV-FWS) and lower values of difference between RV-FWS and RV-GLS (Δ RV strain), in comparison to individuals with HCM ($P < 0.001$ and $P = 0.002$, respectively). RV-FWS and Δ RV strain demonstrated an independent and additive value in discriminating FC from HCM, over the presence of symmetric LVH, systolic anterior motion of the mitral valve and RV hypertrophy. Similar results were found in GLA LVH- patients: they had worse RV-FWS and lower values of Δ RV strain as compared to Sarc LVH- patients (both $P < 0.001$).

Conclusion


Patients with FC show a specific pattern of RV myocardial mechanics, characterized by a larger impairment of RV-FWS and lower Δ RV strain in comparison to patients with HCM, which may be helpful in the differential diagnosis between these two diseases.

* Corresponding author. Tel: +39 0630154187; Fax: +39 063055535. E-mail: francesca.graziani@policlinicogemelli.it

© The Author(s) 2022. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Graphical Abstract



Keywords

Fabry disease • hypertrophic cardiomyopathy • strain analysis • speckle-tracking echocardiography • right ventricle

Introduction

Fabry disease is a rare lysosomal storage disorder, often misdiagnosed and with poor outcome if left untreated. Cardiac involvement is frequent and represents the strongest determinant of prognosis.¹ Left ventricular hypertrophy (LVH) is the hallmark of Fabry cardiomyopathy (FC), which may be indistinguishable from sarcomeric hypertrophic cardiomyopathy (HCM) using conventional echocardiography.¹

Right ventricular hypertrophy (RVH) is found in a consistent proportion of patients with FC,^{2–4} while is less frequent and usually milder in HCM.^{5–7} In both diseases, right ventricular (RV) systolic function is typically normal when assessed by conventional echocardiography.^{2–8} Two-dimensional speckle-tracking echocardiography (2D-STE) allows to detect subtle RV impairment in different clinical settings, including FC^{9,10} and HCM.^{7,11,12} However, potential differences between FC and HCM in terms of RV strain analysis have never been investigated. The aim of the current study was to perform a comparative analysis of RV myocardial mechanics assessed by 2D-STE in: (i) patients with sarcomeric HCM vs. patients with FC and (ii) phenotype-negative carriers of pathogenic sarcomeric gene mutations (Sarc LVH-) vs. phenotype-negative carriers of pathogenic α -galactosidase (GLA) gene mutations (GLA LVH-).

Methods

Study population

We retrospectively screened two cohorts of: (i) patients with Fabry disease ($n = 74$), evaluated at the Fondazione Policlinico A. Gemelli IRCCS, Rome, Italy between January 2010 and January 2021 and (ii) patients with pathogenic sarcomeric mutations ($n = 293$), evaluated at the Leiden University Medical Center (LUMC), Leiden, The Netherlands during the same period.

As currently recommended,¹ the diagnosis of Fabry disease was based on the measurement of the enzymatic activity of α -galactosidase A in leucocytes (for male patients) and confirmed by genetic testing demonstrating pathogenic variants in the GLA gene (for all patients). In the second study cohort, HCM-causing mutations were identified by direct DNA sequencing of sarcomeric genes, performed through clinical evaluation (patients with LVH or family history of HCM). Genotypic information for both study populations are reported in [Supplementary data online, Table S1](#).

Exclusion criteria from the present study were: age <18 years ($n = 14$), obstructive form (defined as maximum resting gradient >30 mmHg and/or maximum provokable gradient >50 mmHg, $n = 33$), insufficient quality of echocardiographic images ($n = 38$), known coronary artery disease ($n = 8$).

Both populations were divided in two groups according to left ventricular (LV) wall thickness: patients with LVH (defined as LV thickness ≥ 13 mm) were considered as those with an overt cardiac phenotype (FC and HCM). Patients without LVH (LV wall thickness < 13 mm) were regarded as genetic positive but phenotype negative (GLA LVH- and Sarc LVH-).¹³

For the purpose of the present study, FC patients were compared with HCM patients matched for maximal LV wall thickness and demographic characteristics in a 1:4 ratio. In addition, GLA LVH- patients were compared with age and sex-matched Sarc LVH- individuals in a 1:2 ratio.

The Fabry population was previously enrolled in ongoing clinical and echocardiographic registries (Fabry Outcome Survey, Fabry Registry and Follow Me Registry) and all patients signed a dedicated informed consent. Due to the retrospective design of this study, the Medical Ethical Committee of the LUMC declared that no formal ethical approval was needed and waived the need for written informed consent. The study was conducted in accordance with the principles of the Helsinki Declaration.

Conventional echocardiographic assessment

Comprehensive echocardiographic examinations were performed with the patients at rest in the left lateral decubitus position using commercially available ultrasound systems (Toshiba Artida™, Toshiba Medical System, Tokio, Japan or Philips EPIQ CVx, Philips Medical Systems, Andover, Massachusetts, USA in the Policlinico A. Gemelli; E7, E9 and E95 system, General Electric Vingmed, Horten, Norway or Philips EPIQ 7, Philips Medical Systems, Andover, Massachusetts, USA in the LUMC). Two-dimensional, colour, pulsed-wave, and continuous-wave Doppler images were acquired from the parasternal, apical and subcostal views. All images were digitally stored for offline analyses, which were performed by an experienced operator (M.C.M.), in both study centres.

From a short-axis view at basal, mid, and apical levels the maximal LV end-diastolic wall thickness was assessed. LV linear dimensions were measured from parasternal long-axis views, and LV mass was calculated according to the Devereux formula and indexed for body surface area.¹⁴ The asymmetry of LVH was quantified using the ratio between the thickness of the interventricular septum and the posterior wall (IVS-PW thickness ratio). Chamber quantification, LV ejection fraction (LVEF) and diastolic function were assessed according to the most recent guidelines.^{14,15}

A comprehensive assessment of RV geometry, size and systolic function was performed according to current recommendations.¹⁶ RV free wall thickness was derived from 2D echocardiography using the subcostal view; RV hypertrophy was defined as RV wall thickness > 5 mm. RV end-diastolic and end-systolic area were measured from a RV focused apical four-chamber view. RV systolic function was evaluated using tricuspid annular plane systolic excursion (TAPSE) by M-mode, tricuspid annular peak systolic velocity (RV S' velocity) by pulsed-wave tissue Doppler imaging (TDI) and RV fractional area change (RVFAC). TAPSE < 17 mm, RV S' velocity < 9.5 cm/s and RVFAC $< 35\%$ were considered suggestive of RV systolic dysfunction.¹⁶

The presence of systolic anterior movement (SAM) of the mitral valve apparatus was visually assessed from the parasternal and apical long-axis views. The peak velocity of the tricuspid regurgitant jet was derived using continuous-wave Doppler, from a RV focused apical four-chamber view.¹⁶ Finally, the severity of mitral regurgitation and tricuspid regurgitation was evaluated using a multiparametric approach, as recommended.

Two-dimensional speckle-tracking echocardiographic analysis

2D-STE analysis was performed offline using commercially available, vendor-independent, dedicated software, 2D Cardiac Performance Analysis© by TomTec-Arena™ (TomTec Imaging Systems, Unterschleissheim, Germany). Images from the apical 4- and 2-chamber and long-axis views were used for the assessment of LV global longitudinal strain (LV-GLS), while the RV focused apical four-chamber view was used to obtain RV global longitudinal strain (RV-GLS) and RV free wall longitudinal strain (RV-FWS).^{14,17} Images with frame rates between 50 and 90 were selected for 2D-STE analysis. The endocardial border was traced from an end-systolic frame by using a point-and-click approach. The region of interest was defined by the software and manually adapted to include the entire myocardial thickness. Then, the myocardium was automatically divided in six segments in each view. For the assessment of RV strain, the average values of the longitudinal peak systolic strain from the three segments of the free wall (RV-FWS) and from all six segments of the free wall and septal wall of the RV (RV-GLS) were calculated.^{17,18} For the assessment of LV-GLS, the average of the longitudinal peak systolic values obtained from the 17 LV segments in the three apical views was evaluated.¹⁴ In the present study, the values of strain measurements are reported in absolute values. According to current evidence,¹⁸ RV-GLS $< 20\%$ and RV-FWS $< 23\%$ were considered as impaired RV systolic function. In addition, the difference between RV-FWS and RV-GLS, referred to as Δ RV strain, was investigated as marker of the equilibrium of RV mechanical properties. RV strain is typically higher in the free wall as compared to the septum and a reference range of $5 \pm 2\%$ for Δ RV strain has been reported in healthy subjects.¹⁸

Statistical analysis

Normally-distributed continuous variables are presented as mean \pm standard deviation whereas non-normally distributed data are presented as median and interquartile range. Categorical variables are expressed as frequencies and percentages. Comparison of clinical and echocardiographic characteristics between groups was performed by the unpaired Student's *t*-test (for normally-distributed continuous variables), Mann-Whitney *U* test (for non-normally distributed continuous variables) and χ^2 test or the Fisher's exact test, as appropriate (for categorical variables). Multiple comparisons of continuous variables were tested with Bonferroni correction. The correlation between RV-FWS and RV free wall thickness was assessed using the Spearman's method. Univariable and multivariable binary logistic regression analyses were performed to investigate the association between echocardiographic characteristics and FC (vs. HCM). Variables with a significant correlation at univariable analysis ($P < 0.05$) were included in the multivariable analysis. The odds ratio (OR) and 95% confidence intervals (CIs) were calculated for each variable. The goodness of fit of the univariable and multivariable regression models was evaluated by calculating Harrell's C statistic with 95% CI. Since Δ RV strain has not been previously investigated in the context of cardiomyopathies with hypertrophic phenotype, a penalized spline curve was fitted to specifically characterize the association between FC and Δ RV strain. The spline curve displayed the changes of FC probability (expressed as percentage) across a range of Δ RV strain values and, based on this analysis, a threshold value of Δ RV strain to predict FC was proposed. Additionally, receiver-operating characteristics (ROC) curve analyses were performed to determine the optimal cut-offs of RV-FWS and

Δ RV strain for discriminating FC from HCM. Different measures of diagnostic accuracy were computed for the relevant thresholds of RV-FWS and Δ RV strain.

Twenty random individuals (10 from each study cohort) were selected for the evaluation of intra- and inter-observer variability of RV strain parameters. Excellent agreement was defined by an intraclass correlation coefficient > 0.9 , whereas good agreement was defined by a value between 0.75 and 0.90.

All tests were two-sided and P -values < 0.05 were considered statistically significant. Statistical analysis was performed using SPSS version 25.0 (IBM Corporation, Armonk, NY, USA) and R version 4.0.1 (R Foundation for Statistical Computing, Vienna, Austria).

Results

Phenotype-positive patients

Clinical and echocardiographic characteristics of FC and HCM patients are summarized in *Table 1*. A total of 140 patients with LVH (28 FC and 112 HCM) were included.

FC and HCM patients were similar for most of the echocardiographic parameters, such as LV and RV dimensions, LVEF, LV-GLS as well as parameters of LV diastolic function. In accordance with the selection criteria, the degree of LVH was similar between the two groups, but FC patients showed a more symmetric pattern of LVH, as expressed by a significantly lower IVS-PW thickness ratio ($P < 0.001$). Mitral SAM (without LV outflow tract gradient) was more frequently observed in HCM patients than FC patients ($P = 0.010$). FC patients had significantly increased RV wall thickness as compared to HCM patients ($P < 0.001$). Accordingly, the prevalence of RVH was higher in the FC group (93% vs. 44%, $P < 0.001$). The assessment of RV systolic function using conventional echocardiographic parameters did not show significant differences between the two groups and TAPSE, RVFAC and RV S' were within the normal range in most of the patients. On the contrary, RV strain analysis revealed a subclinical RV systolic impairment in a sizeable proportion of subjects in both groups. Specifically, RV-GLS was impaired ($< 20\%$) in 61% of FC patients and 46% of HCM patients, with no significant difference between the two populations ($P = 0.151$). Conversely, abnormal RV-FWS ($< 23\%$) was more common in FC individuals than HCM patients (68% vs. 27%, $P < 0.001$), with significantly lower values of Δ RV strain in FC vs. HCM ($P = 0.002$). *Figure 1* shows the comparison of RV parameters in FC vs. HCM. *Figure 2* illustrates the proportion of patients with RV systolic dysfunction based on different echocardiographic parameters. Notably, there were no significant differences in RV strain parameters between patients with disease-causing mutations of the MYBPC3 gene (72% of the HCM patients) and patients with other sarcomeric gene mutations (see [Supplementary data online, Table S2](#)). In the overall population, RV-FWS was moderately correlated with RV wall thickness (correlation coefficient = 0.370; $P < 0.001$).

Diagnostic value of RV mechanics to discriminate FC vs. HCM

At univariate logistic regression analysis, a significant correlation was found between FC and the following echocardiographic variables: IVS-PW thickness ratio, mitral SAM, RV wall thickness, RV-GLS,

RV-FWS and Δ RV strain (see [Supplementary data online, Table S3](#)). For collinearity reasons, different multivariable logistic regression models including IVS-PW thickness ratio, mitral SAM, RV wall thickness and one parameter of RV mechanics were built (*Table 2*). After correcting for IVS-PW thickness ratio, mitral SAM and RV wall thickness, RV-GLS was not significantly associated with FC ($P = 0.141$). Conversely, both RV-FWS and Δ RV strain retained an independent association with FC ($P = 0.005$ and $P = 0.002$, respectively). Notably, the likelihood ratio test demonstrated a significant improvement in the predictive value with the addition of RV-FWS or Δ RV strain to the baseline model ($P = 0.003$ and $P = 0.005$, respectively). The multivariate model incorporating Δ RV strain yielded the highest increment in the C statistic (0.881, 95% CI: 0.809–0.953). Therefore, a spline curve was specifically performed to characterize the association between Δ RV strain and FC. *Figure 3* demonstrates a linear increase of FC probability with lower values of Δ RV strain; based on the fitted spline curve, the threshold value of 4% was chosen to discriminate FC vs. HCM. This cut-off value (Δ RV strain $< 4\%$) was also confirmed by the ROC analysis [sensitivity 68%, specificity 65%, positive predictive value (PPV) 33%, negative predictive value (NPV) 89%] (see [Supplementary data online, Figure S1](#)). The ROC analysis built for RV-FWS identified a value of 23.3% as the best threshold for discriminating FC vs. HCM (sensitivity 75%, specificity 72%, PPV 40%, NPV 92%). Notably, this threshold corresponded approximately to the lower limit of normality for RV-FWS (23.0%), derived from the literature (see [Supplementary data online, Figure S1 and Table S4](#)).

Phenotype-negative patients

A total of 84 patients without LVH were included (28 GLA LVH- and 56 Sarc LVH-) (*Table 3*). As expected, most of the conventional echocardiographic parameters were within the normal range and did not significantly differ between the two groups, except for the E/e' ratio that was higher in the Sarc LVH- group ($P = 0.001$). GLA LVH- patients showed significantly lower values of LV-GLS ($P < 0.001$). No significant differences were found in conventional parameters of RV systolic function between the two populations. However, when assessing RV systolic function by strain analysis, GLA LVH- patients showed significantly reduced RV-FWS ($P < 0.0001$) with lower values of Δ RV strain ($P < 0.001$), as compared to Sarc LVH- patients (*Figure 4*). Impaired RV-FWS values (defined as $< 23\%$) were found in 21% of GLA LVH- patients, as compared to 5% of Sarc LVH- patients ($P = 0.025$).

Reproducibility

The intra- and inter-observer reproducibility of RV strain parameters is summarized in [Supplementary data online, Table S5](#). The agreement was excellent for RV-GLS and RV-FWS and good for Δ RV strain measurements.

Discussion

In the present study we performed a comparative analysis of RV longitudinal mechanics assessed by 2D-STE in FC vs. HCM, as well as in genotype positive-phenotype negative patients. The original and relevant finding of the current study is that the impairment of RV




Figure 1 Box plots showing echocardiographic parameters of RV systolic function in phenotype-positive patients. Lower, middle, and upper hinges of the box correspond to the 25th, 50th, and 75th percentiles. The upper and lower whiskers extend from the hinge to the largest and smallest value, respectively, no further than 1.5 times the interquartile range of the hinge.




Figure 2 Prevalence of RV systolic dysfunction according to different echocardiographic parameters in phenotype-positive patients. The following thresholds were used to define RV systolic dysfunction: TAPSE <17 cm; RV S' velocity <9.5 m/s; RVFAC <35%; RV-GLS <20%; RV-FWS <23%.

Phenotype-positive patients

RVH is considered a 'red flag' for HCM phenocopies.¹⁹ RVH has been reported with a prevalence ranging between 40 and 70% in patients with FC,²⁻⁴ and its presence has been shown to correlate with LVH, global disease severity and increasing age, suggesting that it is a feature of advanced stages of FC.²⁻⁴ In HCM, RVH is usually less frequent and milder as compared to storage/infiltrative cardiomyopathies⁵⁻⁷ but is still correlated with clinical and echocardiographic features of severe disease, especially LVH.⁵⁻⁷ In both diseases, RVH is typically associated with normal RV systolic function when assessed by standard echocardiography.^{3,4,6-8} However, the complex RV shape and geometry weaken the accuracy of conventional parameters of RV systolic function and recent studies on RV strain analysis revealed that this novel technique can unveil a common subclinical impairment of RV mechanics in both HCM and FC patients.^{7,9-12} In the present study, RV strain impairment was

confirmed in a sizeable proportion of the cohort and was more frequent in FC than in HCM patients with similar degree of LVH. However, data on segmental variability of RV strain in cardiomyopathies with hypertrophic phenotype are still limited.¹⁷ In patients with FC, we previously reported that both RV-GLS and RV-FWS are more impaired in comparison to patients with Fabry disease in the pre-hypertrophic stage and healthy controls, while the equilibrium of RV mechanical properties was preserved, as demonstrated by similar values of Δ RV strain.¹⁰ Intriguingly, in the present study, FC patients showed a more prominent reduction of RV-FWS and lower values of Δ RV strain than HCM patients, adjusting for the potential confounding effect of the degree of LVH.

A previous work by Militaru *et al.*²⁰ showed a larger impairment of RV-FWS in 20 FC patients vs. 20 HCM patients with similar age and degree of LVH. The present study extended these findings in a larger and more homogenous population (only patients with genetically-

Table 3 Clinical and echocardiographic characteristics of the phenotype-negative population

	Total LVH- (N = 84)	Sarc LVH- (N = 56)	GLA LVH- (N = 28)	P-value*
Demographic characteristics				
Age (years)	39 ± 15	37 ± 15	40 ± 15	0.337
Female (n,%)	57 (68)	38 (68)	19 (68)	1.000
Hypertension (n,%)	15 (18)	9 (16)	6 (21)	0.546
Echocardiographic characteristics				
LV mass index (g/m ²)	80 (68–93)	80 (69–92)	78 (66–93)	0.708
LVEDV index (mL/m ²)	49 (43–54)	51 (45–56)	47 (40–52)	0.256
LVEF (%)	63 ± 5	63 ± 6	64 ± 3	0.258
LV-GLS (%)	21.8 (19.9–23.2)	22.3 (20.6–23.7)	20.2 (18.5–22.0)	0.002
LAV index (mL/m ²)	28 (23–32)	27 (23–33)	29 (24–30)	1.000
E/A ratio	1.3 (1.1–1.7)	1.3 (1.1–1.5)	1.6 (1.3–1.9)	0.013
E/E' ratio	7.1 (5.8–8.6)	7.9 (6.0–9.3)	6.3 (5.4–7.3)	0.001
RVWT (mm)	3.5 (3.0–4.0)	3.5 (3.0–4.0)	3.6 (3.2–4.0)	0.747
RVEDA (cm ²)	16.4 (14.2–19.3)	16.9 (14.1–19.3)	15.1 (14.5–18.6)	0.407
RVESA (cm ²)	8.3 (6.5–9.7)	8.3 (6.5–9.7)	8.0 (6.3–10.6)	0.859
RAV index (mL/m ²)	14 (11–19)	15 (11–19)	14 (11–19)	0.772
TAPSE (mm)	24 ± 3	24 ± 3	23 ± 3	0.745
RVFAC (%)	51 ± 6	51 ± 6	51 ± 6	0.817
RV S' velocity (cm/s)	13.4 ± 2.2	13.2 ± 3.0	13.5 ± 1.5	0.669
RV-GLS (%)	24.9 (22.2–27.4)	25.3 (22.9–28.0)	23.5 (21.8–26.0)	0.049
RV-FWS (%)	29.2 (26.6–32.6)	30.3 (27.5–33.4)	26.6 (23.9–29.6)	<0.001
ΔRV strain (%)	4.3 (2.8–5.9)	5.0 (3.3–7.0)	2.9 (2.0–4.3)	<0.001
TR jet velocity (m/s)	2.1 (2.0–2.3)	2.1 (2.0–2.3)	2.1 (2.0–2.4)	0.989

*Comparison of echocardiographic parameters was performed applying a Bonferroni correction. *P* values <0.0028 were considered statistically significant (18 comparisons) and are shown in bold type.

LAV, left atrial volume; LVEDV, LV end diastolic volume; RAV, right atrial volume; RVEDA, RV end-diastolic area; RVESA, RV end-systolic area; RVWT, RV wall thickness; TR, tricuspid regurgitation.




Figure 4 Box plots showing parameters of LV and RV strain in phenotype-negative patients. Lower, middle, and upper hinges of the box correspond to the 25th, 50th, and 75th percentiles. The upper and lower whiskers extend from the hinge to the largest and smallest value, respectively, no further than 1.5 times the interquartile range of the hinge.

patients with Fabry disease showed reduced TDI systolic velocities and LV-GLS values as compared to healthy controls.^{25–27} In the current study, patients with Fabry disease in the pre-hypertrophic stage

showed worse LV-GLS values as compared to patients with sarcomeric mutations, even if LV-GLS was within the normal range in both populations. As regards the RV, the results obtained in the

- phenotypes and final diagnosis. A position statement from the ESC working group on myocardial and pericardial diseases. *Eur Heart J* 2013;**34**:1448–58.
20. Militaru S, Jurcuț R, Adam R, Roșca M, Ginghina C, Popescu BA. Echocardiographic features of Fabry cardiomyopathy-comparison with hypertrophy-matched sarcomeric hypertrophic cardiomyopathy. *Echocardiography* 2019;**36**:2041–9
 21. Niemann M, Liu D, Hu K, Herrmann S, Breunig F, Strotmann J et al. Prominent papillary muscles in Fabry disease: a diagnostic marker? *Ultrasound Med Biol* 2011;**37**:37–43.
 22. Labombarda F, Saloux E, Milesi G, Bienvenu B. Loss of base-to-apex circumferential strain gradient: a specific pattern of Fabry cardiomyopathy? *Echocardiography* 2017;**34**:504–10.
 23. Nagueh SF, McFalls J, Meyer D, Hill R, Zoghbi WA, Tam JW et al. Tissue Doppler imaging predicts the development of hypertrophic cardiomyopathy in subjects with subclinical disease. *Circulation* 2003;**108**:395–8.
 24. Williams LK, Misurka J, Ho CY, Chan WX, Agmon Y, Seidman C et al. Multilayer myocardial mechanics in genotype-positive left ventricular hypertrophy-negative patients with hypertrophic cardiomyopathy. *Am J Cardiol* 2018;**122**:1754–60.
 25. Pieroni M, Chimenti C, Ricci R, Sale P, Russo MA, Frustaci A. Early detection of Fabry cardiomyopathy by tissue Doppler imaging. *Circulation* 2003;**107**:1978–84.
 26. Saccheri MC, Cianciulli TF, Lax JA, Gagliardi JA, Cáceres GL, Quarin AE et al. Two-dimensional speckle tracking echocardiography for early detection of myocardial damage in young patients with Fabry disease. *Echocardiography* 2013;**30**:1069–77.
 27. Lu DY, Huang WM, Wang WT, Hung SC, Sung SH, Chen CH et al. Reduced global longitudinal strain as a marker for early detection of Fabry cardiomyopathy. *Eur Heart J Cardiovasc Imaging* 2022;**23**:487–95.