



Universiteit
Leiden
The Netherlands

Modulating energy metabolism: pathophysiological aspects and novel interventions

Straat, M.E.

Citation

Straat, M. E. (2023, March 16). *Modulating energy metabolism: pathophysiological aspects and novel interventions*. Retrieved from <https://hdl.handle.net/1887/3571820>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3571820>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 7

Circadian control of brown adipose tissue

Maaïke E. Straat^{1,2}, Rick Hogenboom^{1,2}, Mariëtte R. Boon^{1,2}, Patrick C.N. Rensen^{1,2}, Sander Kooijman^{1,2}

1. Department of Medicine, Division of Endocrinology, Leiden University Medical Center, Leiden, the Netherlands.
2. Eindhoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, the Netherlands

Biochim Biophys Acta Mol Cell Biol Lipids 2021, 1866(8):158961.

ABSTRACT

Disruption of circadian (~24 hour) rhythms is associated with an increased risk of cardiometabolic diseases. Therefore, unravelling how circadian rhythms are regulated in different metabolic tissues has become a prominent research focus. Of particular interest is brown adipose tissue (BAT), which combusts triglyceride-derived fatty acids and glucose into heat and displays a circannual and diurnal rhythm in its thermogenic activity. In this review, the genetic, neuronal and endocrine generation of these rhythms in BAT is discussed. In addition, the potential risks of disruption or attenuation of these rhythms in BAT, and possible factors influencing these rhythms, are addressed.

INTRODUCTION

Organisms have dealt with daily and seasonal fluctuations in their environment ever since the earth has been habited. To synchronize with the daily variations of the environment, an evolutionary conserved endogenous time-keeping system exists. This system orchestrates an approximate 24-hour oscillatory pattern in intracellular and behavioral processes, which is referred to as the circadian rhythm (*circa* = approximately, *dies* = day). The importance of the circadian rhythm in energy metabolism in mammals is illustrated by numerous studies showing that disruption of the circadian rhythm through genetic, environmental or behavioral factors can contribute to increased risk of developing cardiometabolic diseases. For example, shift work is associated with metabolic syndrome, type 2 diabetes mellitus and cardiovascular diseases (1-4). Therefore, unravelling how circadian rhythms are regulated in different metabolic tissues has become a major research focus. An important metabolic organ is brown adipose tissue (BAT), which can combust triglyceride (TG)-derived fatty acids (FAs) and glucose into heat. BAT exhibits both circannual (seasonal) and circadian (daily) rhythms in its thermogenic activity, which we will discuss in the next sections of this review. In addition, we will discuss the potential risks of circadian disruptions or loss of amplitude in circadian rhythms of BAT as occurs for instance during aging. Finally, we will describe the potential implications of rhythms in BAT activity as a therapeutic means to promote thermogenic activity, which is relevant in the battle against cardiometabolic diseases.

GENERATION OF BIOLOGICAL RHYTHMS

Circadian rhythms are present in almost every cell of the body including neurons in the brain and adipocytes in the various adipose tissue depots, across all species. Generation of circadian rhythms is a complex process. In principle, these rhythms are self-sustained molecular systems, however, they are also entrained to the environment via external cues called 'zeitgebers' (literally 'time-givers'), such as light exposure and food intake, the latter being especially relevant in our current society. In addition, they are subject to a hierarchical architecture, with neurons in the suprachiasmatic nucleus (SCN) receiving photic information serving as the master synchronizer of the peripheral circadian clocks (5, 6). A more detailed description of the various processes involved is outlined below.

Two key processes determine the generation of the endogenous cellular circadian rhythms: 1) activation and repression of key clock components entrenched in the transcriptional-translational feedback loop (TTFL), and 2) post-translational modifications of proteins involved in the biological clock (5, 7, 8). In mammals, the

TTFL mainly consists of two interconnected feedback loops that collectively determine and regulate the near-24-hour-rhythm, namely the core loop and the stabilizing loop (**Figure 1**). In the core loop, the two transcription factors BMAL1 and CLOCK dimerize and translocate to the nucleus. This BMAL1:CLOCK complex initiates the transcription of repressor genes: two cryptochrome members (*CRY1* and *CRY2*) and three periods (*PER1*, *PER2* and *PER3*). The corresponding proteins repress BMAL1 and CLOCK, thereby stopping their own gene transcription. BMAL1 and CLOCK remain suppressed until CRY and PER are degraded, after which the cycle starts again. In the stabilizing loop, the BMAL1:CLOCK complex drives the expression of nuclear receptors (*REV-ERBa* and *REV-ERBb*) and retinoid-related orphan receptors (*RORa*, *RORb* and *RORy*), of which the proteins control the expression of *BMAL1*. Whilst REV-ERB proteins repress transcription of *BMAL1*, RORs promote the expression of *BMAL1*. In addition to the TTFL, post-translational modifications of core clock proteins determine the robustness of the circadian rhythm (9-12). Together, the clock genes and clock proteins form the molecular clock, which cycles around in an approximate 24-hour pattern. The clock proteins can regulate transcription of tissue-specific target genes, resulting in endogenous circadian rhythms in many tissue-specific genes and proteins.

Although this molecular clock machinery is self-sustaining, vertebrates show a hierarchical composition in the time-keeping network. That is, they accommodate a clock-regulating center in the brain (6). This so-called 'master clock' is located in the neurons of the SCN, a small structure within the anterior hypothalamus. The SCN receives photic information from the retina about the external light/dark cycle and communicates this information to distinct brain areas and peripheral tissues through synaptic and non-synaptic mechanisms, such as paracrine and endocrine signaling. Thereby, the SCN synchronizes peripheral circadian clocks to external environmental information (6). In addition, the SCN also receives and integrates information about the internal environment to fine-tune its output. The importance of the SCN is illustrated by studies showing that SCN-ablated rodents do not exhibit circadian rhythmicity in endocrine aspects, such as the adrenal corticosterone rhythm, (13) and drinking behavior (14), suggesting that the SCN reigns over circadian physiology and behavior. In addition, neural grafts from the SCN region restore circadian rhythms in arrhythmic hamsters whose own SCN had been ablated. Moreover, experiments in which the SCN was transplanted from mutated short-period hamsters to long-period hamsters showed that the restored rhythms always adopt the period of the donor (15). The importance of the SCN has been further demonstrated in humans. For instance, patients treated for nonfunctioning pituitary macroadenomas often suffer from a disturbed sleep-wake rhythm, probably due to dysfunction of the adjacent located SCN (16, 17).




Figure 1. The molecular clock. After dimerizing and translocating into the nucleus, the BMAL1:CLOCK complex initiates transcription of three periods (PER1, PER2 and PER3) and two cryptochrome members (CRY1 and CRY2). PER and CRY proteins in turn repress BMAL1 and CLOCK, thereby negatively regulating their own gene transcription. In addition, the BMAL1:CLOCK complex drives the expression of nuclear receptors (REV-ERBa and REV-ERBβ) and retinoid-related orphan receptors (RORα, RORβ and RORγ). REV-ERB proteins repress transcription of BMAL1, RORs promote the expression of BMAL1. The clock proteins in turn initiate transcription of tissue-specific target genes.

The photoperiod (*i.e.*, the amount of daylight perceived) serves as the primary zeitgeber for the SCN. However, to adapt to other environmental fluctuations, endogenous molecular clocks respond to several other zeitgebers that can foster chronobiological homeostasis by transiently extending or reducing intracellular oscillations. A rhythm that is synchronized with the day/night cycle by zeitgebers is formally called a diurnal rhythm and may or may not be a circadian rhythm. Next to the photoperiod, zeitgebers include temperature, physical activity, and eating/drinking patterns. The rhythm of feeding behavior, for example, acts as a powerful synchronizer for peripheral clocks in the liver, muscle, and adipose tissue, while leaving the SCN in most cases unaffected (18, 19). The result is that many organs exhibit a unique set of genes that are expressed in a circadian fashion, ranging from 3% of genes in the hypothalamus up to 16% in the liver. In mice, in total, 43% of all known genes exhibit a circadian profile anywhere in the body (20). To date, there is no consensus on how the circadian clock exactly induces

tissue-specific rhythmic gene expression, but there are suggestions that clock genes rely on tissue-specific transcription factors (21, 22) as well as tissue-specific enhancer-enhancer interactions to facilitate DNA binding (23).

Together, the central and peripheral clocks have profound impact on biological processes. Notably, many aspects of metabolism are under stringent circadian control in order to maintain homeostatic balance during day and night and therefore promote survival (5, 24). Timing of zeitgeber information is thus crucial for optimal clock responses and therewith metabolic health, but is also challenged in our modern society. A significant proportion of the population is involved in shift work to fulfil the demands of our 24/7 economy, resulting in misalignment of rhythms in energy metabolism and the sleep-wake cycle. In addition, technological developments such as electrical lights, computers and smartphones have made it possible for humans to do virtually anything they want whenever they want. However, the extensive use of these devices in the evening or night often results in an extension of the photoperiod affecting sleep quality (25). Exposure to dim light at night is already sufficient to shift behavioral and metabolic rhythms, induce weight gain, and disturb the rhythm in body temperature (26). Another example is that late-night-snacking profoundly disrupts peripheral oscillators (27) and is associated with adverse metabolic and cardiovascular outcomes (28, 29). Knowledge on how the circadian clock is regulated in specific tissues and what the implications are on health outcomes is been recognized to be of major importance for the prevention and optimal treatment of disease. In the next sections we will specifically focus on the circadian mechanisms regulating the activity of BAT.

BROWN ADIPOSE TISSUE

Adipose tissue functions as an endocrine organ and plays a central role in energy metabolism. Most adipocytes are classified as white adipocytes, residing in white adipose tissue (WAT) depots located at various locations throughout our body. These adipocytes are characterized by large unilocular lipid droplets that contain TG to provide fuel to other organs during fasting. In contrast, brown adipocytes within BAT and beige/brite adipocytes scattered throughout WAT are characterized by multiple small lipid droplets and abundant mitochondria that uniquely contain uncoupling protein 1 (UCP1). In the presence of UCP1, brown adipocytes can convert chemical energy into heat, a process referred to as non-shivering thermogenesis.

The most potent activator of thermogenesis in BAT is cold exposure. Cold leads to activation of the sympathetic nervous system (**Figure 2**). In mice, norepinephrine (NE) activates BAT via the β 3-adrenergic receptor (30). The regulation of human BAT is less

clear although recent data indicate a crucial involvement of β 2-adrenergic receptor (31). In both cases, cyclic adenosine monophosphate (cAMP) is produced intracellularly that activates protein kinase A (PKA). PKA phosphorylates the cAMP response element-binding protein (CREB) to activate the thermogenic gene program and furthermore phosphorylates hormone sensitive lipase (HSL) to liberate FA from intracellularly stored TG. The released FA can either allosterically activate UCP1 or be used as a substrate for oxidation. Oxidation of reducing equivalents, mainly nicotinamide adenine dinucleotide (NADH), using the terminal electron acceptor oxygen in the inner mitochondrial membrane, produces a proton motive force. In the presence of activated UCP1 within the inner mitochondrial membrane, protons run back along the gradient, resulting in heat production instead of ATP generation (30, 32).




Figure 2. Regulation of thermogenesis in brown adipocytes. Norepinephrine binds to the β -adrenergic receptor. As a consequence, cyclic adenosine monophosphate (cAMP) is produced intracellularly, which activates protein kinase A (PKA). PKA phosphorylates the cAMP response element-binding protein (CREB) to activate the thermogenic gene program. In addition, PKA phosphorylates hormone sensitive lipase (HSL) to lipolyze intracellularly stored triglycerides (TG). The released fatty acids can either activate UCP1 or be used as a substrate for oxidation. In addition, brown adipocytes can take up fatty acids, from TG-rich lipoproteins, and glucose from the bloodstream, both for use as substrate for oxidation.

Besides its key role in thermoregulation, some studies suggest that BAT contributes to the preservation of the energy balance. This phenomenon is called 'diet-induced

thermogenesis' and is most likely an adaptive response to overfeeding to prevent excessive levels of circulating glucose and lipids (33, 34). The fact that BAT takes up large amounts of glucose and TG-derived FA upon cold- or diet-induced thermogenesis, can be used to visualize and quantify metabolic BAT activity. For mice, we (35, 36) and others (37) have used radiolabeled [^{14}C]deoxyglucose (^{14}C -DG) and glycerol tri[^3H]oleate (^3H -TO), to quantify glucose and TG-derived FA uptake, respectively. In humans, the presence of active BAT was originally identified by the use of [^{18}F]fluorodeoxyglucose (^{18}F -FDG) and detection by position emission tomography combined with X-ray computed tomography (PET-CT) (38-41). Magnetic resonance imaging (MRI) has been used as an alternative technique to assess BAT volume and activity based on changes in TG content of the tissue (42), and infrared thermography as alternative technique to non-invasively monitor temperature of the neck and suprascapular area (43). All these techniques have pros and cons, also depending on body composition and insulin sensitivity status of the subject. Most likely a combination of different techniques including future lipid-based tracers is needed to measure true metabolic BAT activity in humans.

The promotion of BAT function and the browning of WAT have emerged as promising therapeutic targets to increase energy expenditure and counteract weight gain. Human BAT volume and activity as determined by ^{18}F -FDG uptake after cold exposure inversely correlate with fat mass (40). Furthermore, repeated or prolonged cold exposure reduces fat mass in humans (44). In addition, the presence of active BAT also inversely associates with diabetes incidence and fasting glucose levels (45, 46), whereas cold exposure has been shown to improve glucose tolerance and insulin sensitivity (47). A recent study analyzed 134,529 ^{18}F -FDG PET-CT scans in thermoneutral condition and showed that individuals with detectable BAT suffered significantly less from cardiovascular diseases than individuals without detectable BAT (48). Moreover, in mice the uptake of TG-derived FA by activated BAT is accompanied by enhanced hepatic TG-rich lipoprotein (TRL) remnant clearance and promotion of reverse cholesterol transport through increased formation and functionality of high-density lipoproteins (HDL). The combined effect is reversal of dyslipidemia and attenuation of atherosclerosis development (35, 49, 50). Questions remain, however, to what extent classic BAT can be activated in humans and whether research should focus on browning of WAT. WAT browning is an intriguing concept given the large amount of WAT present in (obese) individuals. In addition, a clear distinction between classic brown and beige/brite fat likely does not exist in humans (38, 51). Nonetheless, irrespective of what the appearance of the thermogenic tissues, accumulating evidence suggests that thermogenic adipocytes in human adipose depots can be recruited and/or (trans)differentiated in response to specific stimuli (52).

Thus, BAT is a thermogenic organ with roles in temperature regulation and energy metabolism in mammals. Since outdoor temperatures and the need for energy fluctuate throughout the year and even on a daily basis it is not surprising that BAT displays a circannual and diurnal rhythm, on which we will focus in the following paragraphs.

CIRCANNUAL AND DIURNAL RHYTHMS IN BAT

Circannual rhythms in BAT

Preparations for the cold winter in small mammals presumably starts in fall with enhanced lipid accumulation in WAT and BAT (53). This is followed by an increase in the thermogenic capacity through elevation in the number of mitochondria in BAT during winter (54). Such adaptations of adipose tissue to the environment are likely critical for the survival of small mammals, nonetheless, also in humans the prevalence of metabolically active (*i.e.*, ^{18}F -FDG-positive) BAT is higher in winter as compared to the summer months (38, 48). Indeed, ten days of cold exposure recruits BAT in humans (55). However, not only daytime outdoor temperature, but also day length of the day preceding ^{18}F -FDG PET-CT scanning are found to be predictive for the glucose uptake by BAT (56). Interesting enough, it has even been reported that a stronger correlation exists between the presence of active BAT and photoperiod than with outdoor temperature (57). Based on these data it is tempting to speculate that seasonal changes in photoperiod may be a way to prepare the body for upcoming changes in temperature. In line with this notion, experimental exposure to a short photoperiod of 8h prevents weight gain in field voles (58) and stimulates BAT growth and thermogenic potential in hamsters (59, 60). Conversely, exposing mice to a prolonged photoperiod diminishes metabolic BAT activity, as evidenced by strongly reduced uptake of TG-derived FA and glucose by BAT, resulting in increased lipid deposition in WAT (61).

Changes in temperature and photoperiod may thus both be responsible for the initiation of seasonal adaptive changes in BAT morphology, and this also seems to have implications for cold-induced thermogenic activity, which is markedly increased in humans during wintertime (39). Collectively, this suggests that when BAT is considered as a therapeutic target, seasonal changes in the basal state of BAT may be an important determinant of the treatment outcome.

Diurnal rhythms in BAT

In addition to seasonal changes, also diurnal rhythms are present in BAT. Murine BAT is characterized by circadian gene expression of clock proteins as well as of proteins involved in adipose function and thermogenesis (62). Using RNA-sequencing, about 8% of all protein coding genes in BAT were found to fluctuate in a twenty-four hour cycle,

twice as many as in WAT (20). These rhythms in gene expression most likely translate to tissue function, as ^{18}F -FDG uptake by murine BAT also exhibits a rhythm within the course of the light-dark cycle, peaking 3h before the start of the dark/active phase of mice (63). In line with these data, it was recently reported that primary human brown adipocytes in synchronized cell cultures display a diurnal rhythm in insulin-stimulated glucose uptake (64). This rhythm in glucose uptake is superimposed by a rhythm in the expression of glucose transporter protein 4 (*GLUT4*) and *UCP1*. Interestingly, these circadian expressions are also maintained in cultured human BAT explants, irrespective of the co-incubation with adrenergic or glucocorticoid antagonists, hinting to cell-autonomous mechanisms being involved.

In addition to glucose, the uptake of TG-derived FA by BAT displays a high-amplitude diurnal rhythm, at least in mice (65). BAT shows the highest uptake of TG-derived FA at the onset of the active period, which coincides with high expression of lipoprotein lipase (LPL) and low expression of angiopoietin-like 4 (ANGPTL4), which is a potent LPL inhibitor. The diurnal rhythm in BAT activity determines the rate at which lipids can be cleared from the circulation, thereby regulating the daily rhythm in plasma lipid concentrations. In mice as well as humans, postprandial lipid excursions are nearly absent at waking which may at least partly be due to higher BAT activity at this time of day.

Collectively, these studies show that BAT is more active near the start of the active period in rodents and likely also in humans. Again, it is tempting to speculate about the reasons for this pattern. BAT activity likely relates to body temperature, which reaches its lowest point in the middle of the inactive phase (66). At that point non-shivering thermogenesis might be required to increase body temperature until waking. In the next section we will discuss how diurnal rhythms in BAT are generated.

CIRCADIAN MECHANISMS REGULATING BAT ACTIVITY

Clock genes and BAT function

Many clock gene-mutant mouse strains show changes in BAT morphology and/or function, underscoring the importance of the circadian clock in BAT. Whole-body *Clock* mutant mice are hyperphagic and obese when fed a high fat diet. Additionally, they have many features of metabolic syndrome, including hyperlipidemia (67), and their BAT is characterized by excessive lipid accumulation and loss of its multilocular appearance. Global and adipose-specific *Bmal1* knockout typically potentiate diet-induced obesity (68). Brown adipocytes in these animals are characterized by enlarged lipid droplets (69), but also by increased expression of thermogenic genes (69, 70). The net effect on thermogenesis is likely context dependent with one study reporting decreased cold-

tolerance (70), while others reported no (69, 71) or even improved cold-tolerance (72). Mechanistically, BMAL1 suppresses brown adipogenesis via direct transcriptional control of key components of the transforming growth factor beta (TGF- β) pathway (72). BMAL1 is also known to facilitate transcription of PER2, but PER2 actually stimulates BAT function through interaction with peroxisome proliferator-activated receptor alpha (PPAR α) (73).

BAT function is improved in mice lacking REV-ERB α , related to the repressing activity of REV-ERB α on *Ucp1* gene expression. Consequently, UCP1 expression and cold-tolerance exhibit a diurnal rhythm, both of which are in antiphase with REV-ERB α (74). More recently, it was shown that BAT-specific REV-ERB α deficiency was sufficient to elevate UCP1 levels (75). REV-ERB α shares nuclear targets with ROR α , with opposite regulation of *Bmal1* expression, while both REV-ERB α and ROR α show repressive effects on *Ucp1* expression. Homozygous staggerer (*sg/sg*) mice, featuring a global ROR α dysfunction, are lean and protected from diet-induced obesity explained by a remarkable increase in the thermogenic gene program including *Ucp1* expression in BAT and WAT (76, 77).

Circadian sympathetic innervation of BAT

The sympathetic nervous system is the dominant driver of thermogenic activity in BAT. Interestingly, the autonomic nervous system itself follows a circadian pattern as well. SCN neurons connect with neurons regulating sympathetic activity in BAT of Siberian hamsters (78) and injection of glutamate into the SCN increases BAT thermogenesis in rats (79). To what extent these connections translate into actual circadian release of NE in BAT is, to the best of our knowledge, not known. Nevertheless, circadian disruption upon continuous light exposure was found to reduce sympathetic outflow towards BAT (61) and sympathetic denervation abolishes rhythmic TG-derived FA uptake by BAT (65), although this latter observation should be interpreted with caution as denervation by itself has quite a dramatic effect on BAT activity.

Interestingly, chronic cold exposure (4°C) was reported to strongly increase the amplitude of circadian rhythm in genes controlling *de novo* lipogenesis in BAT of mice, including *Srebf1* and *Mlxipl*, encoding SREBP-1 and ChREBP respectively (75, 80). *De novo* lipogenesis in BAT is required for the adaptation to chronic cold exposure (75). On the other hand, during thermoneutrality (30°C), ChREBP-dependent lipogenesis was shown to cause lipid accumulation in inactive BAT (80). Depletion of SREBP-1 hinders maintenance of body temperature specifically during the light phase when mice were housed at 4°C (75). The question remains, however, to what extent this is a direct effect of (circadian) sympathetic nervous system activity. In addition, it is not clear why *Srebp1* and *Mlxipl* expression is specifically stimulated in the dark phase. Possibly, nutrient availability during the dark phase, combined with *de novo* lipogenesis helps the mouse to prepare for the resting phase when BAT activity is needed to maintain body temperature.

Thus, circadian disruption negatively affects BAT function, possibly via the sympathetic nervous system, and sympathetic activity increases the amplitude of genes involved in BAT functioning. On the other hand, it seems that adrenergic input is dispensable for circadian BAT activity since mice lacking β -ARs (β -less mice) have low expression of *Ucp1* at room temperature, but the circadian rhythmicity of *Ucp1* and clock genes in BAT is maintained (81). Similarly, housing mice at thermoneutrality, coinciding with minimal to no sympathetic outflow to BAT, leaves circadian rhythmic heat production intact (74). The latter indicates that apart from circadian sympathetic outflow other factors must be involved in the generation of the diurnal rhythm in BAT.

Circadian release of hormones and BAT function

Many hormones exhibit circadian or diurnal rhythms and quite a few of them have been implicated in BAT function, either direct or indirect via modulating sympathetic outflow (82). In this review, we will focus on two (types of) hormones that are well-known for their contribution to the circadian control of metabolism, namely glucocorticoids (GC) and melatonin.

Glucocorticoids

The release of GCs, mainly represented by corticosterone in mice and cortisol in humans, is controlled through the hypothalamus-pituitary-adrenal (HPA) axis and shows a robust rhythm peaking at the onset of the active phase. The promoter regions of the clock genes *PER1* and *PER2* (83), as well as those of many genes involved in BAT function (84), contain glucocorticoid receptor (GR)- and mineralocorticoid receptor (MR)-responsive elements. Clock proteins can physically interact with GR to regulate transcription in peripheral tissues, as CLOCK represses GR action (85), CRY transactivates GRs (86), and the recently newly identified BMAL:CLOCK inhibitor CHRONO represses GR expression (87).

In BAT, GCs typically promote adipogenesis including uptake of TG-derived FA and glucose (88) but simultaneously repress the thermogenic gene program (89), resulting in increased nutrient storage. Surprisingly, exogenous adrenocorticotrophic hormone (ACTH), which promotes GC release as part of the HPA axis, was found to directly promote BAT activity (90), although the question remains whether endogenous ACTH reaches levels that are sufficiently high to elicit this effect. A recent study also suggested species-specific differences in the regulation of UCP1 expression by GCs, with repression of transcriptional activity in brown adipocytes of mice and stimulation in human cells. Despite these data it is very unlikely that GCs will ever be used to promote BAT activity in humans, because long term treatment was found to suppress BAT activity in both mice and humans (91), possibly due to a strong overflow in nutrient influx and

the necessity to store these. In addition, GCs are well-known to cause multiple negative metabolic side effects (92, 93).

Adrenalectomy increases BAT activity in mice, while rhythmic activity is maintained (65), indicating that at least the peak at waking is not required for circadian BAT activity. On the other hand, continuous high dose corticosterone treatment not only reduces BAT activity but also decreases the amplitude of circadian activity. However, in this type of experiments it is not possible to discriminate between effects caused by absence of rhythmicity or by changes in GC levels. We (94) and others (95) have therefore implanted low-dose corticosterone-containing pellets. These pellets increase circulating GC levels just above the threshold at which the negative feedback on the HPA axis kicks in shutting off endogenous GC production, resulting in flattened corticosterone levels without exceeding the normal levels. Flattening of corticosterone rhythms resulted in lipid accumulation in BAT, while TG-derived FA uptake was decreased, due to alterations in rhythmic LPL expression (94). Interestingly, these effects were independent of GR expression by the brown adipocyte and rather explained by reduced sympathetic outflow at waking (94).

Melatonin

The SCN controls timing of melatonin production by the pineal gland and it thus directly transfers information about circadian cycles. Accordingly, melatonin shows a robust circadian rhythm with peak levels during night time. On exposure to (day)light melatonin levels rapidly drop to almost undetectable levels (96). In addition, melatonin secretion varies over the year with prolonged secretion during long dark winter nights (*i.e.* short photoperiods) as compared to short summer nights (*i.e.* long photoperiods) (97). Siberian hamsters, transferred from a long photoperiod to a short photoperiod, exhibit a naturally-occurring reversal in obesity. This is mediated by the duration of melatonin secretion and subsequent activation of the sympathetic nervous system (98). Consequently, pinealectomized rats show reduced UCP1 levels (99) and an impaired BAT response to either HFD or cold (100), phenotypes that could be reversed by exogenous melatonin administration.

Melatonin may promote BAT activity through different ways. Besides stimulating sympathetic activity, it also inhibits ACTH-mediated GC secretion (101). In addition, melatonin receptors are not only expressed in the brain but also in metabolic organs and cells including adipocytes (102). An early study already showed that melatonin can promote BAT growth in hamsters (103). A more recent study showed comparable effects in patients with melatonin deficiency, who show increased BAT volume and activity as measured with PET-MRI after 3-months oral melatonin administration (104). Of note, most inbred mouse strains, including the widely used C57BL/6 mice,

are naturally deficient for melatonin because serotonin is not acetylated by serotonin N-acetyltransferase (arylalkylamine N-acetyltransferase, AANAT) to generate melatonin (105). Nevertheless, as we have repeatedly demonstrated that BAT activity has a circadian rhythm in C57BL/6J mice (61, 65), we may conclude that although melatonin is able to modulate circadian BAT activity, it is dispensable.




Figure 3. Proposed model of how several circadian factors influence diurnal rhythms in BAT. A) The SCN contains a direct innervation pathway to BAT via the sympathetic nervous system (SNS), releasing norepinephrine locally that binds to β -adrenergic receptors. In addition, rhythmic secretion of glucocorticoids stimulates fatty acids and glucose uptake by the brown adipocyte but conversely inhibits thermogenesis. The SCN also promotes synthesis of melatonin by the pineal gland, which signals through melatonin receptors and enhances thermogenesis in brown adipocytes. Clock genes are also involved. Sympathetic innervation represses REV-ERB in brown adipocytes, which inhibits thermogenesis. ROR and BMAL1 suppress thermogenesis, while PER and CLOCK probably stimulate thermogenesis. B) The circadian rhythm in BAT thermogenesis coincides with the rhythm of SNS activity and is in anti-phase with the activity of REV-ERBa. Cortisol peaks at the beginning of the active phase, while melatonin peaks during the inactive phase.

RELEVANCE OF BAT RHYTHMS: WHEN ARE INTERVENTIONS USEFUL AND MOST EFFICIENT?

It should be clear by now that circadian disruptions affect BAT activity and that BAT is characterized by pronounced circannual and diurnal rhythms with respect to thermogenic capacity and activity. This raises at least three questions: 1) What factors attenuate or disturb the circadian rhythm in BAT? 2) Can we prevent metabolic problems associated with attenuated or disturbed circadian rhythms by stimulating BAT? 3) Should we consider circadian cycles in BAT during therapy, otherwise known as chronotherapy? For example, when pharmacologically stimulating BAT, we could question whether the time of the day is a crucial determinant. Below we will discuss these questions over a few examples.

Aging

During aging, there is an increased chance of developing pathologies, including cardiovascular diseases, type 2 diabetes mellitus, and cancer. Aging is associated with deterioration of the biological clock, including a reduced amplitude in the rhythms of hormones, body temperature, and processes that regulate the sleep-wake cycle.

During aging in both humans and mice BAT mass and activity are reduced, and browning of white fat is decreased (38, 106). Transcriptome analysis covering a 24-hour period of BAT in aged mice revealed 1,021 differentially expressed genes including elevated amplitude of *Cry1* and *Rev-erba* expression (107). Comparison of the rhythmic lipidome of BAT in young and old mice shows that lipids tend to accumulate with age, yet the number of oscillating lipids reduces and the time of the day at which lipids reach their peak shifts (108). Mechanistically there are many possibilities by which age can affect BAT function, including reduced sympathetic outflow and dysfunction of mitochondria, and therefore it will probably be challenging to disentangle the various contributors from that of altered circadian rhythmicity. Nevertheless, it is tempting to speculate that restoring rhythmic BAT activity may promote healthy aging. Accordingly, several studies have described improved BAT activity in long-lived animals (109-112) and that transplantation of BAT from young to old mice reverses the aging phenotype (112).

Timing of food intake

The time of the day is an important determinant for the fate of consumed nutrients. Interesting enough, diet-induced thermogenesis (*i.e.* the postprandial increase in energy expenditure) is known to exhibit a day-night rhythm, with the highest response in the early morning (113-115). This may help explain why meal timing affects body weight regulation and why shift work is a risk factor for obesity (116-118). The role of BAT in diet-

induced thermogenesis is somewhat controversial, however, the response is higher in persons who have detectable ^{18}F -FDG uptake by BAT compared to those without (119). Following a meal, ^{18}F -FDG uptake (120), oxygen consumption, and blood flow in human BAT acutely increase (121). In mice, the time of the day also determines to what extent TG-derived FA are taken up by BAT, which is reflected in postprandial TG levels (65).

Regardless of the contribution of BAT to diet-induced thermogenesis, the diurnal rhythm in diet-induced thermogenesis provides the scientific basis for the concept of time-restricted feeding. Mice fed a high fat diet only during the dark phase show reduced weight gain compared to *ad libitum* fed mice, an effect accompanied by increased and rhythmic UCP1 expression and whole body energy expenditure (122-124). In human studies without instructions on caloric intake, time-restricted eating (*i.e.* only eating during an either self- or by the researchers selected 6-9 hour window) reduces body weight and other risk factors of cardiovascular disease and type 2 diabetes in obese individuals with/without metabolic syndrome (125-127). Independent of weight loss, early time-restricted food intake with a feeding period between 9 a.m. and 3 p.m. was also found to improve insulin sensitivity in men with prediabetes (128). Surprisingly, a very recent randomized controlled trial did not find any metabolic benefit of late time-restricted eating between 12 p.m. and 8 p.m. (129), indicating that more research is needed to identify who may or may not benefit from such intervention and what time-window would be, if at all, effective.

Timing of cold exposure and therapeutic interventions

Chronotherapy is a relatively new field of research that is based on the premise that administering medications or other interventions at specific times of the circadian cycle will minimize side effects and/or maximize therapeutic effects. As mentioned before, cold exposure is the dominant driver of BAT activity and mice tolerate cold considerably better at the end of the dark phase than at the end of the light phase (74, 130). To the best of our knowledge, no study has addressed this issue in humans thus far. Nonetheless, night time cold acclimation was demonstrated to result in a pronounced increase in BAT abundance accompanied by improved insulin sensitivity (131). Similarly, the question whether the effects of pharmacological interventions targeting BAT are time-dependent has not been addressed in humans nor in mice. With respect to circannual rhythms, cold-induced thermogenic activity is markedly increased in humans during wintertime (39, 56, 57), indicating that the basal state is important.

Very interestingly, over the last years various small molecules have been developed to interfere with key clock proteins. Two agonistic compounds, SR9011 and SR9009, targeting REV-ERBs were found to improve energy homeostasis in diet-induced obese

mice (132). Treatment with SR1555, an inverse agonist of ROR γ , induces UCP1 expression in BAT of diet-induced obese mice (133). To what extent these molecules promote thermogenic activity in BAT and can be applied to reverse circadian disturbances in humans is yet to be discovered.

CONCLUSIONS AND FUTURE DIRECTIONS

The discovery of metabolically active BAT in human adults a decade ago led to interest in BAT as a therapeutic target to combat metabolic and cardiovascular diseases. Whilst animal studies show great promise, it is unclear as yet to what extent BAT or browned WAT contributes to energy expenditure in humans. Nonetheless, UCP1 positive adipocytes are present in human adults and are expected to have, even in small amounts, an impact on energy homeostasis in the long term. Accumulating evidence suggests that rodent and human BAT possesses clear circannual and diurnal rhythms, regulated through processes summarized in **Figure 3**. These rhythms should be considered when measuring BAT function and possibly when therapeutically targeting BAT, although more research is needed to confirm this. On the other hand, disruption of circadian rhythms through genetic clock deficiency in mice, prolonged light exposure, (mimicking) shift work, and other ways have been associated with metabolic and cardiovascular problems, often accompanied by deteriorated BAT function. The exact contribution of BAT dysfunction to these phenotypes is not known and probably difficult to address, but unravelling the affected mechanistic pathways may lead to the discovery of novel therapeutic targets to combat cardiometabolic diseases.

ACKNOWLEDGMENTS

Illustrations were created with BioRender.com.

FUNDING

This work was supported by the Netherlands CardioVascular Research Initiative: ‘the Dutch Heart Foundation, Dutch Federation of University Medical Centers, the Netherlands Organization for Health Research and Development and the Royal Netherlands Academy of Sciences’ [CVON2017-20 GENIUS-II] to PCNR (Patrick C.N. Rensen). MRB (Mariëtte R. Boon) is supported by the Dutch Diabetes Research Foundation Fellowship [grant number 2015.81.1808] and SK (Sander Kooijman) by the Dutch Heart Foundation (2017T016).

REFERENCES

1. Pietroiusti A, Neri A, Somma G, Coppeta L, Iavicoli I, Bergamaschi A, et al. Incidence of metabolic syndrome among night-shift healthcare workers. *Occup Environ Med*. 2010;67(1):54-7.
2. Schiavo-Cardozo D, Lima MM, Pareja JC, Geloneze B. Appetite-regulating hormones from the upper gut: disrupted control of xenin and ghrelin in night workers. *Clin Endocrinol (Oxf)*. 2013;79(6):807-11.
3. Knutsson A, Akerstedt T, Jonsson BG, Orth-Gomer K. Increased risk of ischaemic heart disease in shift workers. *Lancet*. 1986;2(8498):89-92.
4. Kroenke CH, Spiegelman D, Manson J, Schernhammer ES, Colditz GA, Kawachi I. Work characteristics and incidence of type 2 diabetes in women. *Am J Epidemiol*. 2007;165(2):175-83.
5. Albrecht U. Timing to perfection: the biology of central and peripheral circadian clocks. *Neuron*. 2012;74(2):246-60.
6. Dibner C, Schibler U, Albrecht U. The mammalian circadian timing system: organization and coordination of central and peripheral clocks. *Annu Rev Physiol*. 2010;72:517-49.
7. Albrecht U, Eichele G. The mammalian circadian clock. *Curr Opin Genet Dev*. 2003;13(3):271-7.
8. Partch CL, Green CB, Takahashi JS. Molecular architecture of the mammalian circadian clock. *Trends in Cell Biology*. 2014;24(2):90-9.
9. Siepka SM, Yoo SH, Park J, Song W, Kumar V, Hu Y, et al. Circadian mutant Overtime reveals F-box protein FBXL3 regulation of cryptochrome and period gene expression. *Cell*. 2007;129(5):1011-23.
10. Eide EJ, Woolf MF, Kang H, Woolf P, Hurst W, Camacho F, et al. Control of mammalian circadian rhythm by CKepsilon-regulated proteasome-mediated PER2 degradation. *Mol Cell Biol*. 2005;25(7):2795-807.
11. Tamaru T, Hirayama J, Isojima Y, Nagai K, Norioka S, Takamatsu K, et al. CK2alpha phosphorylates BMAL1 to regulate the mammalian clock. *Nat Struct Mol Biol*. 2009;16(4):446-8.
12. Sahar S, Zocchi L, Kinoshita C, Borrelli E, Sassone-Corsi P. Regulation of BMAL1 protein stability and circadian function by GSK3beta-mediated phosphorylation. *PLoS One*. 2010;5(1):e8561.
13. Moore RY, Eichler VB. Loss of a circadian adrenal corticosterone rhythm following suprachiasmatic lesions in the rat. *Brain Res*. 1972;42(1):201-6.
14. Stephan FK, Zucker I. Circadian rhythms in drinking behavior and locomotor activity of rats are eliminated by hypothalamic lesions. *Proc Natl Acad Sci U S A*. 1972;69(6):1583-6.
15. Ralph MR, Foster RG, Davis FC, Menaker M. Transplanted suprachiasmatic nucleus determines circadian period. *Science*. 1990;247(4945):975-8.
16. Joustra SD, Kruijssen E, Verstegen MJ, Pereira AM, Biermasz NR. Determinants of altered sleep-wake rhythmicity in patients treated for nonfunctioning pituitary macroadenomas. *J Clin Endocrinol Metab*. 2014;99(12):4497-505.

17. Joustra SD, Thijs RD, van den Berg R, van Dijk M, Pereira AM, Lammers GJ, et al. Alterations in diurnal rhythmicity in patients treated for nonfunctioning pituitary macroadenoma: a controlled study and literature review. *Eur J Endocrinol.* 2014;171 (2):217-28.
18. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev.* 2000;14 (23):2950-61.
19. Marchant EG, Mistlberger RE. Anticipation and entrainment to feeding time in intact and SCN-ablated C57BL/6j mice. *Brain Res.* 1997;765 (2):273-82.
20. Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB. A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci U S A.* 2014;111 (45):16219-24.
21. Meireles-Filho ACA, Bardet AF, Yanez-Cuna JO, Stampfel G, Stark A. cis-regulatory requirements for tissue-specific programs of the circadian clock. *Curr Biol.* 2014;24 (1):1-10.
22. Zhang Y, Fang B, Emmett MJ, Damle M, Sun Z, Feng D, et al. GENE REGULATION. Discrete functions of nuclear receptor Rev-erbalpha couple metabolism to the clock. *Science.* 2015;348 (6242):1488-92.
23. Beytebiere JR, Trott AJ, Greenwell BJ, Osborne CA, Vitet H, Spence J, et al. Tissue-specific BMAL1 cistromes reveal that rhythmic transcription is associated with rhythmic enhancer-enhancer interactions. *Genes Dev.* 2019;33 (5-6):294-309.
24. Lamia KA, Storch KF, Weitz CJ. Physiological significance of a peripheral tissue circadian clock. *Proc Natl Acad Sci U S A.* 2008;105 (39):15172-7.
25. Wright KP, Jr., McHill AW, Birks BR, Griffin BR, Rusterholz T, Chinoy ED. Entrainment of the human circadian clock to the natural light-dark cycle. *Curr Biol.* 2013;23 (16):1554-8.
26. Borniger JC, Maurya SK, Periasamy M, Nelson RJ. Acute dim light at night increases body mass, alters metabolism, and shifts core body temperature circadian rhythms. *Chronobiol Int.* 2014;31 (8):917-25.
27. Archer SN, Laing EE, Moller-Levet CS, van der Veen DR, Bucca G, Lazar AS, et al. Mistimed sleep disrupts circadian regulation of the human transcriptome. *Proc Natl Acad Sci U S A.* 2014;111 (6):E682-91.
28. Morris CJ, Purvis TE, Hu K, Scheer FA. Circadian misalignment increases cardiovascular disease risk factors in humans. *Proc Natl Acad Sci U S A.* 2016;113 (10):E1402-11.
29. Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci U S A.* 2009;106 (11):4453-8.
30. Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev.* 2004;84 (1):277-359.
31. Blondin DP, Nielsen S, Kuipers EN, Severinsen MC, Jensen VH, Miard S, et al. Human Brown Adipocyte Thermogenesis Is Driven by beta2-AR Stimulation. *Cell Metab.* 2020;32 (2):287-300 e7.
32. Nicholls DG, Locke RM. Thermogenic mechanisms in brown fat. *Physiol Rev.* 1984;64 (1):1-64.
33. Kozak LP. Brown fat and the myth of diet-induced thermogenesis. *Cell Metab.* 2010;11 (4):263-7.

34. Peterson CM, Orooji M, Johnson DN, Naraghi-Pour M, Ravussin E. Brown adipose tissue does not seem to mediate metabolic adaptation to overfeeding in men. *Obesity (Silver Spring)*. 2017;25 (3):502-5.
35. Berbee JF, Boon MR, Khedoe PP, Bartelt A, Schlein C, Worthmann A, et al. Brown fat activation reduces hypercholesterolaemia and protects from atherosclerosis development. *Nat Commun*. 2015;6:6356.
36. Khedoe PP, Hoeke G, Kooijman S, Dijk W, Buijs JT, Kersten S, et al. Brown adipose tissue takes up plasma triglycerides mostly after lipolysis. *J Lipid Res*. 2015;56 (1):51-9.
37. Bartelt A, Bruns OT, Reimer R, Hohenberg H, Ittrich H, Peldschus K, et al. Brown adipose tissue activity controls triglyceride clearance. *Nature Medicine*. 2011;17 (2):200-5.
38. Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, et al. Identification and importance of brown adipose tissue in adult humans. *N Engl J Med*. 2009;360 (15):1509-17.
39. Saito M, Okamatsu-Ogura Y, Matsushita M, Watanabe K, Yoneshiro T, Nio-Kobayashi J, et al. High incidence of metabolically active brown adipose tissue in healthy adult humans: effects of cold exposure and adiposity. *Diabetes*. 2009;58 (7):1526-31.
40. van Marken Lichtenbelt WD, Vanhomerig JW, Smulders NM, Drossaerts JM, Kemerink GJ, Bouvy ND, et al. Cold-activated brown adipose tissue in healthy men. *N Engl J Med*. 2009;360 (15):1500-8.
41. Virtanen KA, Lidell ME, Orava J, Heglind M, Westergren R, Niemi T, et al. Functional brown adipose tissue in healthy adults. *N Engl J Med*. 2009;360 (15):1518-25.
42. Abreu-Vieira G, Sardjoe Mishre ASD, Burakiewicz J, Janssen LGM, Nahon KJ, van der Eijk JA, et al. Human Brown Adipose Tissue Estimated With Magnetic Resonance Imaging Undergoes Changes in Composition After Cold Exposure: An in vivo MRI Study in Healthy Volunteers. *Front Endocrinol (Lausanne)*. 2019;10:898.
43. Boon MR, Bakker LE, van der Linden RA, Pereira Arias-Bouda L, Smit F, Verberne HJ, et al. Supraclavicular skin temperature as a measure of 18F-FDG uptake by BAT in human subjects. *PLoS One*. 2014;9 (6):e98822.
44. Yoneshiro T, Aita S, Matsushita M, Kayahara T, Kameya T, Kawai Y, et al. Recruited brown adipose tissue as an antiobesity agent in humans. *J Clin Invest*. 2013;123 (8):3404-8.
45. Lee P, Greenfield JR, Ho KK, Fulham MJ. A critical appraisal of the prevalence and metabolic significance of brown adipose tissue in adult humans. *Am J Physiol Endocrinol Metab*. 2010;299 (4):E601-6.
46. Ouellet V, Routhier-Labadie A, Bellemare W, Lakhali-Chaieb L, Turcotte E, Carpentier AC, et al. Outdoor temperature, age, sex, body mass index, and diabetic status determine the prevalence, mass, and glucose-uptake activity of 18F-FDG-detected BAT in humans. *J Clin Endocrinol Metab*. 2011;96 (1):192-9.
47. Chondronikola M, Volpi E, Borsheim E, Porter C, Annamalai P, Enerback S, et al. Brown adipose tissue improves whole-body glucose homeostasis and insulin sensitivity in humans. *Diabetes*. 2014;63 (12):4089-99.
48. Becher T, Palanisamy S, Kramer DJ, Eljalby M, Marx SJ, Wibmer AG, et al. Brown adipose tissue is associated with cardiometabolic health. *Nature Medicine*. 2021.

49. Bartelt A, John C, Schaltenberg N, Berbee JFP, Worthmann A, Cherradi ML, et al. Thermogenic adipocytes promote HDL turnover and reverse cholesterol transport. *Nat Commun.* 2017;8:15010.
50. Hoeke G, Kooijman S, Boon MR, Rensen PC, Berbee JF. Role of Brown Fat in Lipoprotein Metabolism and Atherosclerosis. *Circ Res.* 2016;118 (1):173-82.
51. Shinoda K, Luijten IH, Hasegawa Y, Hong H, Sonne SB, Kim M, et al. Genetic and functional characterization of clonally derived adult human brown adipocytes. *Nat Med.* 2015;21 (4):389-94.
52. Herz CT, Kiefer FW. Adipose tissue browning in mice and humans. *J Endocrinol.* 2019;241 (3):R97-R109.
53. Burlington RF, Therriault DG, Hubbard RW. Lipid changes in isolated brown fat cells from hibernating and aroused thirteen-lined ground squirrels (*Citellus tridecemlineatus*). *Comp Biochem Physiol.* 1969;29 (1):431-7.
54. Johansson BW, Senturia JB. Seasonal variations in the physiology and biochemistry of the European hedgehog (*Erinaceus europaeus*) including comparisons with non-hibernators, guinea-pig and man. *Acta Physiol Scand Suppl.* 1972;380:1-159.
55. van der Lans AA, Hoeks J, Brans B, Vijgen GH, Visser MG, Vosselman MJ, et al. Cold acclimation recruits human brown fat and increases nonshivering thermogenesis. *J Clin Invest.* 2013;123 (8):3395-403.
56. Bahler L, Deelen JW, Hoekstra JB, Holleman F, Verberne HJ. Seasonal influence on stimulated BAT activity in prospective trials: a retrospective analysis of BAT visualized on 18F-FDG PET-CTs and 123I-mIBG SPECT-CTs. *J Appl Physiol (1985).* 2016;120 (12):1418-23.
57. Au-Yong IT, Thorn N, Ganatra R, Perkins AC, Symonds ME. Brown adipose tissue and seasonal variation in humans. *Diabetes.* 2009;58 (11):2583-7.
58. Krol E, Duncan JS, Redman P, Morgan PJ, Mercer JG, Speakman JR. Photoperiod regulates leptin sensitivity in field voles, *Microtus agrestis*. *J Comp Physiol B.* 2006;176 (2):153-63.
59. McElroy JF, Wade GN. Short photoperiod stimulates brown adipose tissue growth and thermogenesis but not norepinephrine turnover in Syrian hamsters. *Physiol Behav.* 1986;37 (2):307-11.
60. Wiesinger H, Heldmaier G, Buchberger A. Effect of photoperiod and acclimation temperature on nonshivering thermogenesis and GDP-binding of brown fat mitochondria in the Djungarian hamster *Phodopus s. sungorus*. *Pflugers Arch.* 1989;413 (6):667-72.
61. Kooijman S, van den Berg R, Ramkisoensing A, Boon MR, Kuipers EN, Loeff M, et al. Prolonged daily light exposure increases body fat mass through attenuation of brown adipose tissue activity. *Proc Natl Acad Sci USA.* 2015;112 (21):6748-53.
62. Zvonic S, Ptitsyn AA, Conrad SA, Scott LK, Floyd ZE, Kilroy G, et al. Characterization of peripheral circadian clocks in adipose tissues. *Diabetes.* 2006;55 (4):962-70.
63. van der Veen DR, Shao J, Chapman S, Leevy WM, Duffield GE. A diurnal rhythm in glucose uptake in brown adipose tissue revealed by in vivo PET-FDG imaging. *Obesity (Silver Spring).* 2012;20 (7):1527-9.
64. Lee P, Bova R, Schofield L, Bryant W, Dieckmann W, Slattery A, et al. Brown Adipose Tissue Exhibits a Glucose-Responsive Thermogenic Biorhythm in Humans. *Cell Metab.* 2016;23 (4):602-9.

65. van den Berg R, Kooijman S, Noordam R, Ramkisoensing A, Abreu-Vieira G, Tambyrajah LL, et al. A Diurnal Rhythm in Brown Adipose Tissue Causes Rapid Clearance and Combustion of Plasma Lipids at Wakening. *Cell Rep.* 2018;22 (13):3521-33.
66. Kräuchi K. How is the circadian rhythm of core body temperature regulated? *Clinical Autonomic Research.* 2002;12 (3):147-9.
67. Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, et al. Obesity and metabolic syndrome in circadian Clock mutant mice. *Science.* 2005;308 (5724):1043-5.
68. Shi SQ, Ansari TS, McGuinness OP, Wasserman DH, Johnson CH. Circadian disruption leads to insulin resistance and obesity. *Curr Biol.* 2013;23 (5):372-81.
69. Li S, Yu Q, Wang GX, Lin JD. The biological clock is regulated by adrenergic signaling in brown fat but is dispensable for cold-induced thermogenesis. *PLoS One.* 2013;8 (8):e70109.
70. Chang L, Xiong W, Zhao X, Fan Y, Guo Y, Garcia-Barrio M, et al. Bmal1 in Perivascular Adipose Tissue Regulates Resting-Phase Blood Pressure Through Transcriptional Regulation of Angiotensinogen. *Circulation.* 2018;138 (1):67-79.
71. Paschos GK, Ibrahim S, Song WL, Kunieda T, Grant G, Reyes TM, et al. Obesity in mice with adipocyte-specific deletion of clock component Arntl. *Nat Med.* 2012;18 (12):1768-77.
72. Nam D, Guo B, Chatterjee S, Chen MH, Nelson D, Yechoor VK, et al. The adipocyte clock controls brown adipogenesis through the TGF-beta and BMP signaling pathways. *J Cell Sci.* 2015;128 (9):1835-47.
73. Chappuis S, Ripperger JA, Schnell A, Rando G, Jud C, Wahli W, et al. Role of the circadian clock gene Per2 in adaptation to cold temperature. *Mol Metab.* 2013;2 (3):184-93.
74. Gerhart-Hines Z, Feng D, Emmett MJ, Everett LJ, Loro E, Briggs ER, et al. The nuclear receptor Rev-erbalpha controls circadian thermogenic plasticity. *Nature.* 2013;503 (7476):410-3.
75. Adlanmerini M, Carpenter BJ, Remsberg JR, Aubert Y, Peed LC, Richter HJ, et al. Circadian lipid synthesis in brown fat maintains murine body temperature during chronic cold. *Proc Natl Acad Sci U S A.* 2019;116 (37):18691-9.
76. Lau P, Tuong ZK, Wang SC, Fitzsimmons RL, Goode JM, Thomas GP, et al. Roralpha deficiency and decreased adiposity are associated with induction of thermogenic gene expression in subcutaneous white adipose and brown adipose tissue. *Am J Physiol Endocrinol Metab.* 2015;308 (2):E159-71.
77. Monnier C, Auclair M, Le Cam G, Garcia MP, Antoine B. The nuclear retinoid-related orphan receptor RORalpha controls circadian thermogenic programming in white fat depots. *Physiol Rep.* 2018;6 (8):e13678.
78. Bamshad M, Song CK, Bartness TJ. CNS origins of the sympathetic nervous system outflow to brown adipose tissue. *Am J Physiol.* 1999;276 (6):R1569-78.
79. Amir S, Shizgal P, Rompre PP. Glutamate injection into the suprachiasmatic nucleus stimulates brown fat thermogenesis in the rat. *Brain Res.* 1989;498 (1):140-4.
80. Schlein C, Fischer AW, Sass F, Worthmann A, Tödter K, Jaeckstein MY, et al. Endogenous Fatty Acid Synthesis Drives Brown Adipose Tissue Involution. *Cell Reports.* 2021;34 (2):108624.
81. Razzoli M, Emmett MJ, Lazar MA, Bartolomucci A. beta-Adrenergic receptors control brown adipose UCP-1 tone and cold response without affecting its circadian rhythmicity. *FASEB J.* 2018;32 (10):5640-6.

82. Kooijman S, van den Heuvel JK, Rensen PCN. Neuronal Control of Brown Fat Activity. *Trends Endocrinol Metab.* 2015;26 (11):657-68.
83. Yamamoto T, Nakahata Y, Tanaka M, Yoshida M, Soma H, Shinohara K, et al. Acute physical stress elevates mouse period1 mRNA expression in mouse peripheral tissues via a glucocorticoid-responsive element. *J Biol Chem.* 2005;280 (51):42036-43.
84. Luijten IHN, Cannon B, Nedergaard J. Glucocorticoids and Brown Adipose Tissue: Do glucocorticoids really inhibit thermogenesis? *Mol Aspects Med.* 2019;68:42-59.
85. Nader N, Chrousos GP, Kino T. Circadian rhythm transcription factor CLOCK regulates the transcriptional activity of the glucocorticoid receptor by acetylating its hinge region lysine cluster: potential physiological implications. *FASEB J.* 2009;23 (5):1572-83.
86. Lamia KA, Papp SJ, Yu RT, Barish GD, Uhlénhaut NH, Jonker JW, et al. Cryptochromes mediate rhythmic repression of the glucocorticoid receptor. *Nature.* 2011;480 (7378):552-6.
87. Goriki A, Hatanaka F, Myung J, Kim JK, Yoritaka T, Tanoue S, et al. A novel protein, CHRONO, functions as a core component of the mammalian circadian clock. *PLoS Biol.* 2014;12 (4):e1001839.
88. van den Heuvel JK, Boon MR, van Hengel I, Peschier-van der Put E, van Beek L, van Harmelen V, et al. Identification of a selective glucocorticoid receptor modulator that prevents both diet-induced obesity and inflammation. *Br J Pharmacol.* 2016;173 (11):1793-804.
89. Strack AM, Bradbury MJ, Dallman MF. Corticosterone decreases nonshivering thermogenesis and increases lipid storage in brown adipose tissue. *Am J Physiol.* 1995;268 (1 Pt 2):R183-91.
90. van den Beukel JC, Grefhorst A, Quarta C, Steenbergen J, Mastroberardino PG, Lombes M, et al. Direct activating effects of adrenocorticotrophic hormone (ACTH) on brown adipose tissue are attenuated by corticosterone. *FASEB J.* 2014;28 (11):4857-67.
91. Ramage LE, Akyol M, Fletcher AM, Forsythe J, Nixon M, Carter RN, et al. Glucocorticoids Acutely Increase Brown Adipose Tissue Activity in Humans, Revealing Species-Specific Differences in UCP-1 Regulation. *Cell Metab.* 2016;24 (1):130-41.
92. Schäcke H, Döcke WD, Asadullah K. Mechanisms involved in the side effects of glucocorticoids. *Pharmacol Ther.* 2002;96 (1):23-43.
93. Chotiyarnwong P, McCloskey EV. Pathogenesis of glucocorticoid-induced osteoporosis and options for treatment. *Nature Reviews Endocrinology.* 2020;16 (8):437-47.
94. Kroon J, Schilperoord M, In Het Panhuis W, van den Berg R, van Doeselaar L, Verzijl CRC, et al. A physiological glucocorticoid rhythm is an important regulator of brown adipose tissue function. *Mol Metab.* 2021;47:101179.
95. Tholen S, Kovary K, Rabiee A, Bielczyk-Maczyńska E, Yang W, Teruel MN. Flattened circadian glucocorticoid oscillations cause obesity due to increased lipid turnover and lipid uptake. *bioRxiv.* 2020:2020.01.02.893081.
96. Wurtman RJ, Axelrod J, Phillips LS. Melatonin Synthesis in the Pineal Gland: Control by Light. *Science.* 1963;142 (3595):1071-3.
97. Reiter RJ. Pineal melatonin: cell biology of its synthesis and of its physiological interactions. *Endocr Rev.* 1991;12 (2):151-80.

98. Ryu V, Zarebidaki E, Albers HE, Xue B, Bartness TJ. Short photoperiod reverses obesity in Siberian hamsters via sympathetically induced lipolysis and Browning in adipose tissue. *Physiol Behav.* 2018;190:11-20.
99. Buonfiglio D, Parthimos R, Dantas R, Cerqueira Silva R, Gomes G, Andrade-Silva J, et al. Melatonin Absence Leads to Long-Term Leptin Resistance and Overweight in Rats. *Front Endocrinol (Lausanne).* 2018;9:122.
100. de Souza CAP, Gallo CC, de Camargo LS, de Carvalho PVV, Olescuck IF, Macedo F, et al. Melatonin multiple effects on brown adipose tissue molecular machinery. *J Pineal Res.* 2019;66 (2):e12549.
101. Torres-Farfan C, Richter HG, Rojas-Garcia P, Vergara M, Forcelledo ML, Valladares LE, et al. mt1 Melatonin receptor in the primate adrenal gland: inhibition of adrenocorticotropin-stimulated cortisol production by melatonin. *J Clin Endocrinol Metab.* 2003;88 (1):450-8.
102. Brydon L, Petit L, Delagrangé P, Strosberg AD, Jockers R. Functional expression of MT2 (Mel1b) melatonin receptors in human PAZ6 adipocytes. *Endocrinology.* 2001;142 (10):4264-71.
103. Heldmaier G, Hoffmann K. Melatonin stimulates growth of brown adipose tissue. *Nature.* 1974;247 (5438):224-5.
104. Halpern B, Mancini MC, Bueno C, Barcelos IP, de Melo ME, Lima MS, et al. Melatonin Increases Brown Adipose Tissue Volume and Activity in Patients With Melatonin Deficiency: A Proof-of-Concept Study. *Diabetes.* 2019;68 (5):947-52.
105. Roseboom PH, Namboodiri MA, Zimonjic DB, Popescu NC, Rodriguez IR, Gastel JA, et al. Natural melatonin 'knockdown' in C57BL/6J mice: rare mechanism truncates serotonin N-acetyltransferase. *Brain Res Mol Brain Res.* 1998;63 (1):189-97.
106. Goncalves LF, Machado TQ, Castro-Pinheiro C, de Souza NG, Oliveira KJ, Fernandes-Santos C. Ageing is associated with brown adipose tissue remodelling and loss of white fat browning in female C57BL/6 mice. *Int J Exp Pathol.* 2017;98 (2):100-8.
107. Sutton GM, Ptitsyn AA, Floyd ZE, Yu G, Wu X, Hamel K, et al. Biological aging alters circadian mechanisms in murine adipose tissue depots. *Age (Dordr).* 2013;35 (3):533-47.
108. Held NM, Buijink MR, Elfrink HL, Kooijman S, Janssens GE, Vaz FM, et al. Aging selectively dampens oscillation of lipid abundance in white and brown adipose tissue. *bioRxiv.* 2020:2020.09.04.283226.
109. Darcy J, McFadden S, Fang Y, Huber JA, Zhang C, Sun LY, et al. Brown Adipose Tissue Function Is Enhanced in Long-Lived, Male Ames Dwarf Mice. *Endocrinology.* 2016;157 (12):4744-53.
110. Li Y, Knapp JR, Kopchick JJ. Enlargement of interscapular brown adipose tissue in growth hormone antagonist transgenic and in growth hormone receptor gene-disrupted dwarf mice. *Exp Biol Med (Maywood).* 2003;228 (2):207-15.
111. Ortega-Molina A, Efeyan A, Lopez-Guadamillas E, Munoz-Martin M, Gomez-Lopez G, Canamero M, et al. Pten positively regulates brown adipose function, energy expenditure, and longevity. *Cell Metab.* 2012;15 (3):382-94.
112. Vatner DE, Zhang J, Oydanich M, Guers J, Katsyuba E, Yan L, et al. Enhanced longevity and metabolism by brown adipose tissue with disruption of the regulator of G protein signaling 14. *Aging Cell.* 2018;17 (4):e12751.

113. Morris CJ, Garcia JI, Myers S, Yang JN, Trienekens N, Scheer FA. The Human Circadian System Has a Dominating Role in Causing the Morning/Evening Difference in Diet-Induced Thermogenesis. *Obesity (Silver Spring)*. 2015;23 (10):2053-8.
114. Richter J, Herzog N, Janka S, Baumann T, Kistenmacher A, Oltmanns KM. Twice as High Diet-Induced Thermogenesis After Breakfast vs Dinner On High-Calorie as Well as Low-Calorie Meals. *J Clin Endocrinol Metab*. 2020;105 (3).
115. Romon M, Edme JL, Boulenguez C, Lescroart JL, Frimat P. Circadian variation of diet-induced thermogenesis. *Am J Clin Nutr*. 1993;57 (4):476-80.
116. Bandin C, Scheer FA, Luque AJ, Avila-Gandia V, Zamora S, Madrid JA, et al. Meal timing affects glucose tolerance, substrate oxidation and circadian-related variables: A randomized, crossover trial. *Int J Obes (Lond)*. 2015;39 (5):828-33.
117. Bo S, Musso G, Beccuti G, Fadda M, Fedele D, Gambino R, et al. Consuming more of daily caloric intake at dinner predisposes to obesity. A 6-year population-based prospective cohort study. *PLoS One*. 2014;9 (9):e108467.
118. Wehrens SMT, Christou S, Isherwood C, Middleton B, Gibbs MA, Archer SN, et al. Meal Timing Regulates the Human Circadian System. *Curr Biol*. 2017;27 (12):1768-75 e3.
119. Hibi M, Oishi S, Matsushita M, Yoneshiro T, Yamaguchi T, Usui C, et al. Brown adipose tissue is involved in diet-induced thermogenesis and whole-body fat utilization in healthy humans. *Int J Obes (Lond)*. 2016;40 (11):1655-61.
120. Vosselman MJ, Brans B, van der Lans AA, Wiert R, van Baak MA, Mottaghy FM, et al. Brown adipose tissue activity after a high-calorie meal in humans. *Am J Clin Nutr*. 2013;98 (1):57-64.
121. M UD, Saari T, Raiko J, Kudomi N, Maurer SF, Lahesmaa M, et al. Postprandial Oxidative Metabolism of Human Brown Fat Indicates Thermogenesis. *Cell Metab*. 2018;28 (2):207-16 e3.
122. Chaix A, Lin T, Le HD, Chang MW, Panda S. Time-Restricted Feeding Prevents Obesity and Metabolic Syndrome in Mice Lacking a Circadian Clock. *Cell Metab*. 2019;29 (2):303-19 e4.
123. Hatori M, Vollmers C, Zarrinpar A, DiTacchio L, Bushong EA, Gill S, et al. Time-restricted feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet. *Cell Metab*. 2012;15 (6):848-60.
124. Sherman H, Genzer Y, Cohen R, Chapnik N, Madar Z, Froy O. Timed high-fat diet resets circadian metabolism and prevents obesity. *FASEB J*. 2012;26 (8):3493-502.
125. Chow LS, Manoogian ENC, Alvear A, Fleischer JG, Thor H, Dietsche K, et al. Time-Restricted Eating Effects on Body Composition and Metabolic Measures in Humans who are Overweight: A Feasibility Study. *Obesity (Silver Spring)*. 2020;28 (5):860-9.
126. Hutchison AT, Regmi P, Manoogian ENC, Fleischer JG, Wittert GA, Panda S, et al. Time-Restricted Feeding Improves Glucose Tolerance in Men at Risk for Type 2 Diabetes: A Randomized Crossover Trial. *Obesity (Silver Spring)*. 2019;27 (5):724-32.
127. Wilkinson MJ, Manoogian ENC, Zadourian A, Lo H, Fakhouri S, Shoghi A, et al. Ten-Hour Time-Restricted Eating Reduces Weight, Blood Pressure, and Atherogenic Lipids in Patients with Metabolic Syndrome. *Cell Metab*. 2020;31 (1):92-104 e5.
128. Sutton EF, Beyl R, Early KS, Cefalu WT, Ravussin E, Peterson CM. Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes. *Cell Metab*. 2018;27 (6):1212-21 e3.

129. Lowe DA, Wu N, Rohdin-Bibby L, Moore AH, Kelly N, Liu YE, et al. Effects of Time-Restricted Eating on Weight Loss and Other Metabolic Parameters in Women and Men With Overweight and Obesity: The TREAT Randomized Clinical Trial. *JAMA Intern Med.* 2020.
130. Talan MI, Tatelman HM, Engel BT. Cold tolerance and metabolic heat production in male C57BL/6J mice at different times of day. *Physiol Behav.* 1991;50 (3):613-6.
131. Lee P, Smith S, Linderman J, Courville AB, Brychta RJ, Dieckmann W, et al. Temperature-acclimated brown adipose tissue modulates insulin sensitivity in humans. *Diabetes.* 2014;63 (11):3686-98.
132. Solt LA, Wang Y, Banerjee S, Hughes T, Kojetin DJ, Lundasen T, et al. Regulation of circadian behaviour and metabolism by synthetic REV-ERB agonists. *Nature.* 2012;485 (7396):62-8.
133. Chang MR, He Y, Khan TM, Kuruvilla DS, Garcia-Ordonez R, Corzo CA, et al. Antiobesity Effect of a Small Molecule Repressor of RORgamma. *Mol Pharmacol.* 2015;88 (1):48-56.

