



**Universiteit
Leiden**
The Netherlands

Definition of a structured training curriculum for robot-assisted radical cystectomy with intracorporeal ileal conduit in male patients: a Delphi consensus study led by the ERUS educational board

Dell'Oglio, P.; Turri, F.; Larcher, A.; D'Hondt, F.; Sanchez-Salas, R.; Bochner, B.; ... ; YAU Working Grp Robot-Assisted Sur

Citation

Dell'Oglio, P., Turri, F., Larcher, A., D'Hondt, F., Sanchez-Salas, R., Bochner, B., ... Mottrie, A. (2022). Definition of a structured training curriculum for robot-assisted radical cystectomy with intracorporeal ileal conduit in male patients: a Delphi consensus study led by the ERUS educational board. *European Urology Focus*, 8(1), 160-164.
doi:10.1016/j.euf.2020.12.015

Version: Accepted Manuscript

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3567627>

Note: To cite this publication please use the final published version (if applicable).



available at www.sciencedirect.com
journal homepage: www.europeanurology.com/eufocus



Brief Correspondence

Definition of a Structured Training Curriculum for Robot-assisted Radical Cystectomy with Intracorporeal Ileal Conduit in Male Patients: A Delphi Consensus Study Led by the ERUS Educational Board

Paolo Dell'Oglio^{a,b,c,†,*}, Filippo Turri^{b,d,†}, Alessandro Larcher^e, Frederiek D'Hondt^{b,c}, Rafael Sanchez-Salas^f, Bernard Bochner^g, Joan Palou^h, Robin Westonⁱ, Abolfazl Hosseini^j, Abdullah E. Canda^k, Jørgen Bjerggaard^l, Giovanni Cacciamani^m, Kasper Ørding Olsen^l, Inderbir Gill^m, Thierry Piechautⁿ, Walter Artibani^o, Pim J. van Leeuwen^p, Arnulf Stenzl^q, John Kelly^r, Prokar Dasgupta^s, Carl Wijburg^t, Justin W. Collins^{b,r}, Mihir Desai^m, Henk G. van der Poel^p, Francesco Montorsi^e, Peter Wiklund^u, Alexandre Motttrie^{b,c}, on behalf of the ERUS Educational Working Group and the YAU Working Group on Robot-assisted Surgery

^a Department of Urology, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; ^b ORSI, Academy, Melle, Belgium; ^c Department of Urology, Onze Lieve Vrouw Hospital, Aalst, Belgium; ^d Department of Urology, University of Modena & Reggio Emilia, Modena, Italy; ^e Department of Urology, Division of Oncology, Urological Research Institute, IRCCS Ospedale San Raffaele, Milan, Italy; ^f Department of Urology, Institut Mutualiste Montsouris, Paris, France; ^g Urology Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY, USA; ^h Department of Urology, Fundació Puigvert, Autonomous University of Barcelona, Barcelona, Spain; ⁱ Department of Urology, Royal Liverpool University Hospital, Liverpool, UK; ^j Department of Urology, Karolinska University Hospital, Stockholm, Sweden; ^k Department of Urology, School of Medicine, Koc University, Istanbul, Turkey; ^l Department of Clinical Medicine—Department of Urology, Aarhus University Hospital, Aarhus, Denmark; ^m USC Institute of Urology, and Catherine and Joseph Aresty Department of Urology, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA; ⁿ Department of Urology, Clinique Saint Augustin, Bordeaux, France; ^o Department of Urology, Azienda Ospedaliera Universitaria Integrata, University of Verona, Verona, Italy; ^p Department of Urology, The Netherlands Cancer Institute, Amsterdam, The Netherlands; ^q Department of Urology, Eberhard Karls University, Tübingen, Germany; ^r Department of Urology, University College London Hospital, London, UK; ^s Department of Urology, Guy's Hospital, King's Health Partners, London, UK; ^t Department of Urology, Robotic Surgery, Rijnstate Hospital, Arnhem, The Netherlands; ^u Department of Urology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[†] These authors contributed equally.

* Corresponding author. Department of Urology, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; ORSI, Academy, Melle, Belgium; Department of Urology, Onze Lieve Vrouw Hospital, Aalst, Belgium. Tel. +3407981232; Fax: +390226437298.
E-mail address: paolo.delloglio@gmail.com (P. Dell'Oglio).

Article info

Article history:

Accepted December 18, 2020

Associate Editor: Derya Tilki

Keywords:

Curriculum
Learning curve
Radical cystectomy
Robot-assisted surgery
Training

Abstract

Robot-assisted radical cystectomy (RARC) continues to expand, and several surgeons start training for this complex procedure. This calls for the development of a structured training program, with the aim to improve patient safety during RARC learning curve. A modified Delphi consensus process was started to develop the curriculum structure. An online survey based on the available evidence was delivered to a panel of 28 experts in the field of RARC, selected according to surgical and research experience, and expertise in running training courses. Consensus was defined as $\geq 80\%$ agreement between the responders. Overall, 96.4% experts completed the survey. The structure of the RARC curriculum was defined as follows: (1) theoretical training; (2) preclinical simulation-based training: 5-d simulation-based activity, using models with increasing complexity (ie, virtual reality, and dry- and wet-laboratory exercises), and nontechnical skills training session; (3) clinical training: modular console activity of at least 6 mo at the host center (a RARC case was divided into 11 steps and steps of similar complexity were grouped into five modules); and (4) final evaluation: blind review of a video-recorded RARC case. This structured training pathway will guide a starting surgeon from the first steps of RARC toward independent completion of a full procedure. Clinical implementation is urgently needed.

Patient summary: Robot-assisted radical cystectomy (RARC) is a complex procedure. The first structured training program for RARC was developed with the goal of aiding surgeons to overcome the learning curve of this procedure, improving patients' safety at the same time.

© 2020 European Association of Urology. Published by Elsevier B.V. All rights reserved.

Radical cystectomy (RC) represents the gold standard for surgical management of organ-confined muscle-invasive bladder cancer and refractory non-muscle-invasive disease [1]. Owing to the surgical morbidity related to the open RC approach [2], the minimally invasive technique has been introduced in an attempt to improve patient outcomes. In the last decades, robot-assisted RC (RARC) increased exponentially, overtaking open RC [3]. Despite RARC being probably the most complex urological procedure with a non-negligible learning curve [4] and high complication rate, no training program is currently available to assist surgeons during their learning process. To overcome this issue, we developed the first structured training program for RARC, with the ultimate goal of improving patient's safety during the learning phase of the surgeon.

The structure of the curriculum was defined relying on a modified Delphi consensus process [5]. Using Google Forms (<https://www.google.com/intl/it/forms/about/>), a web-based survey (Supplementary material) based on the available evidence was generated and delivered to a panel of 28 experts in the field of RARC. Experts were selected according to their extensive experience in robotic surgery, the impact and number of their publications on robotic surgery, and their surgical training and expertise in running training courses in the field of RARC. For simplicity, the web-based survey focused exclusively on RARC for male patients with intracorporeal ileal conduit. Consensus was defined as $\geq 80\%$ agreement between responders. Frequency and proportions were used to describe the outcomes of the survey.

The response rate in the first Delphi round was 96.4% (27/28). Consensus was reached in several areas assessed (Table 1). Unanimous agreement was reached on the statement that the adoption of a standardized curriculum for training can improve clinical outcomes during RARC

learning curve and the established structure of the robot-assisted radical prostatectomy (RARP) curriculum [6] should be applied to the RARC curriculum. The panel agreed on the eligibility criteria for a trainee who should have previous experience with RARP, defined as full surgical autonomy in performing RARP, regardless of the number of procedures performed. Moreover, the trainee should have a minimum table-side assistance experience of 10 RARCs before beginning the curriculum. The structure of the RARC curriculum was divided into four main phases (Fig. 1). The first phase consisted of *theoretical training*, including e-learning and case observation. The second phase consisted of *5-d preclinical simulation-based training*, including simulation-based activity using models with increasing complexity (ie, virtual reality, and dry- and wet-laboratory exercises) and nontechnical skills training session. In the dry laboratory, synthetic hydrogel models are used to improve suturing skills and simulate ureteroileal and bowel anastomosis. In the wet laboratory, cadaveric canine and living porcine models are used for simulation of the full case procedure and of vascular and bowel injury. Each trainee's performance is assessed in an objective fashion relying on a set of five Da Vinci skills simulator exercises [7] before (baseline assessment) and after (final assessment) the training session. The third phase consisted of *clinical modular training* of at least 6 mo at the host center. A RARC case was divided into 11 steps, defined as discrete segmental units of surgery in chronological order. There was high agreement regarding the progression of the trainee through the different steps that must follow a modular pattern according to the level of complexity of each step. Therefore, each step's complexity was ranked using a scale from 1 (easy) to 5 (complex), and steps with similar complexity were grouped together into five modules, defined as a unit of the same

Table 1 – Key statements of the modified Delphi consensus process for the definition of a structured training curriculum for robot-assisted radical cystectomy.

Domain	Item	Consensus
Overview	Clinical outcomes during RARC learning curve can be improved by the adoption of a standardized curriculum for training	100% (27/27)
	The established structure of the robot-assisted radical prostatectomy curriculum should be applied to the RARC curriculum	100% (27/27)
Eligibility	Before beginning the RARC curriculum, the candidate should have previous experience with robot-assisted radical prostatectomy	81.5% (22/27)
	Before beginning the RARC curriculum, the candidate should have a minimum table-side assistance experience of 10 RARCs	81.5% (22/27)
	To be eligible, a center should already have an ERUS host center certification	81.5% (22/27)
	A total of 30 annual RARC cases is the minimum volume for a center to be eligible as a RARC curriculum host center	81.5% (22/27)
Preclinical simulation-based training	Preclinical simulation-based training will have the structure presented in Fig. 1	89% (24/27)
	The baseline assessment and virtual reality simulation exercises already established for the robot-assisted radical prostatectomy curriculum should be applied to the RARC curriculum	89% (24/27)
	The dry-lab exercises with synthetic hydrogel models should include the following:	96.3% (26/27)
	Ureteroileal anastomosis	85.2% (23/27)
	Bowel anastomosis	
	The animal model for the wet-lab exercises should be a cadaveric canine model and a living porcine model	81.5% (22/27)
	The wet-lab exercises should include the following:	92.6% (25/27)
	Case simulation on animal model	81.5% (22/27)
	Vascular injury	81.5% (22/27)
	Bowel injury	92.6% (25/27)
	Bowel stapling	81.5% (22/27)
	Vessel sealer management	
	Nontechnical skills training should include the following:	92.6% (25/27)
	Decision-making	96.3% (26/27)
	Emergency scenario	85.2% (23/27)
	Team training	
Clinical modular training	RARC steps were defined as follows:	100% (27/27)
	I Patient positioning and trocar placement	
	II Identification and isolation of the ureters	
	III Incision of the Douglas pouch, freeing of the seminal vesicles, transection of vasa deferentia, opening of Denonvilliers, and preparation of the space between the prostate and the rectum. Opening of the endopelvic fascia	
	IV Dissection and division of the bladder vascular pedicles. Lateral dissection of the prostate (with or without nerve sparing)	
	V Cutting of puboprostatic ligaments, cutting of Santorini plexus with oversewing, and cutting of the urethra (avoiding tumor spillage by placing a clip on proximal urethra)	
	VI Bagging of bladder specimen	
	VII Extended pelvic lymph node dissection (can be performed after step II and before step III)	
	VIII Tunnelization of the left ureter behind the sigmoid	
	IX Harvesting of the bowel segment and ileoileal anastomosis	
	X Ureteroileal anastomosis	
	XI Stoma creation (RARC with ileal conduit)	
	RARC was divided into five modules according to the complexity of each step using a scale from I (easy) to V (complex):	100% (27/27)
	Module I: steps I and VI	
	Module II: steps II and III	
	Module III: steps V, VII, and XI	
	Module IV: steps IV and VIII	
	Module V: steps IX and X	
	Progression of the trainee through the different steps must follow a modular pattern according to the level of complexity of each step	96.3% (26/27)
Final assessment	The duration of the modular training activity should be 6–12 mo	100% (27/27)
	The final assessment should be based on the evaluation of an index video by certified independent examiners in a blind-review process	92.6% (25/27)
	A new procedure-specific scoring system should be developed for the RARC curriculum, based on objective metrics defining the steps to be completed and the errors to be avoided	89% (24/27)
	The final assessment should be based on a procedure-specific scale focusing on the following steps: (1) dissection of bladder pedicles, (2) harvesting of the ileal segment and ileoileal anastomosis, and (3) ureteroileal anastomosis	81.5% (22/27)

ERUS = European Association of Urology Robotic Urology Section; RARC = robot-assisted radical cystectomy.

complexity regardless of chronological order. The fourth phase consisted of the *final evaluation*, based on the blind review of a video-recorded RARC case.

The increasing adoption of robot-assisted surgery in urological field also for RC calls for the development of standardized and validated training programs in order to

improve patient outcomes during the learning process of this complex surgical procedure. The increasing focus on patients' safety is bringing surgical robotic training more and more out of the operating room and has forced mentors to move away from the Halstedian surgical training program of "see one, do one, teach one," where patients are




Fig. 1 – Structure of the European Association of Urology Robotic Urology Section curriculum for robot-assisted radical cystectomy (RARC) defined by the modified Delphi consensus process.

inevitably exposed to an increased risk of suboptimal outcomes. Under these premises, we developed the first available structured curriculum for RARC with intracorporeal ileal conduit. This structured training program is in line with those proposed by the European Association of Urology Robotic Urology Section (ERUS) for RARP [6] and robot-assisted partial nephrectomy [5]. *Theoretical and preclinical simulation-based training* comprises the key elements. The

latter allows replacement of a real patient with virtual, synthetic, and animal models in order to improve technical skills of trainees and shorten their learning curve for a specific procedure. It is required before clinical training in the operating theater because it was shown to improve surgical performance significantly [7]. It is of note that high agreement was reached by the panel regarding the importance of including nontechnical skills training into the

preclinical simulation-based phase, underlying that appropriate communication/confirmatory feedback with bedside assistant, anesthetist, and theater staff are mandatory to ensure safe robotic surgical performance [8]. The core of all ERUS curricula is the *modular console training*, where an interaction between mentor and trainee is mandatory to allow the progression through modules of increasing complexity. The role of the mentor is pivotal to the success of this process, and appropriate experience and skills are vital: the ability to share expertise often is not innate, and the increasing need for training of mentors is being recognized [9]. Another key element is the *final evaluation*. Measurement of outcomes and performance-level verification become imperative. The panel reached agreement that a new specific scoring system should be developed for the RARC curriculum, based on objective and validated performance metrics, defining the steps to be completed and the errors to be avoided, to track the progression of the trainee and ensure that defined benchmarks of skills will be reached before the trainee progresses to the next level of difficulty (proficiency-based progression training) [10].

Despite its novelty, this study is not devoid of limitations. First, it was developed for male candidates for RARC with intracorporeal ileal conduit. Consequently, this curriculum is not applicable to female candidates for RC and cannot be used to assist surgeons during their learning process for RARC with neobladder. Second, no pilot clinical validation was provided that will ultimately show whether this ERUS curriculum will allow progression through the learning curve without a detrimental effect on the patient's clinical outcomes [5].

The first structured training pathway for RARC was developed based on an international consensus between experts. This curriculum will drive a surgeon from the first steps of RARC toward independent completion of a full procedure. Clinical implementation of this curriculum is urgently needed.

Author contributions: Paolo Dell'Oglio had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Dell'Oglio, Turri, Larcher, Mottrie.

Acquisition of data: Dell'Oglio, Turri.

Analysis and interpretation of data: Dell'Oglio, Turri.

Drafting of the manuscript: Dell'Oglio, Turri.

Critical revision of the manuscript for important intellectual content: D'Hondt, Sanchez-Salas, Bochner, Palou, Weston, Hosseini, Canda, Bjerggaard, Cacciamani, Olsen, Gill, Piechaud, Artibani, van Leeuwen, Stenzl, Kelly, Dasgupta, Wijburg, Collins, Desai, van der Poel, Montorsi, Wiklund, Mottrie.

Statistical analysis: Dell'Oglio.

Obtaining funding: None.

Administrative, technical, or material support: Mottrie.

Supervision: D'Hondt, van der Poel, Montorsi, Wiklund, Mottrie.

Other: None.

Financial disclosures: Paolo Dell'Oglio certifies that all conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript (eg, employment/affiliation, grants or funding, consultancies, honoraria, stock ownership or options, expert testimony, royalties, or patents filed, received, or pending), are the following: None.

Funding/Support and role of the sponsor: None.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.euf.2020.12.015>.

References

- [1] European Association of Urology. Guidelines on muscle-invasive and metastatic bladder cancer. 2020 <https://uroweb.org/guideline/bladder-cancer-muscle-invasive-and-metastatic/>
- [2] Lawrentschuk N, Colombo R, Hakenberg OW, et al. Prevention and management of complications following radical cystectomy for bladder cancer. *Eur Urol* 2010;57:983–1001.
- [3] Zamboni S, Soria F, Mathieu R, et al. Differences in trends in the use of robot-assisted and open radical cystectomy and changes over time in peri-operative outcomes among selected centres in North America and Europe: an international multicentre collaboration. *BJU Int* 2019;124, 656–64.
- [4] Dell'Oglio P, Mazzone E, Lambert E, et al. The effect of surgical experience on perioperative and oncological outcomes after robot-assisted radical cystectomy with intracorporeal urinary diversion: evidence from a referral centre with extensive experience in robotic surgery. *Eur Urol Focus*. In press. <https://doi.org/10.1016/j.euf.2020.01.016>.
- [5] Larcher A, De Naeyer G, Turri F, et al. The ERUS curriculum for robot-assisted partial nephrectomy: structure definition and pilot clinical validation. *Eur Urol* 2019;75:1023–31.
- [6] Volpe A, Ahmed K, Dasgupta P, et al. Pilot validation study of the European Association of Urology robotic training curriculum. *Eur Urol* 2015;68:292–9.
- [7] Larcher A, Turri F, Bianchi L, et al. Virtual reality validation of the ERUS simulation-based training programmes: results from a high-volume training centre for robot-assisted surgery. *Eur Urol* 2019;75:885–7.
- [8] Collins JW, Dell'Oglio P, Hung AJ, Brook NR. The importance of technical and non-technical skills in robotic surgery training. *Eur Urol Focus* 2018;4, 674–6.
- [9] Collins JW, Levy J, Stefanidis D, et al. Utilising the Delphi process to develop a proficiency-based progression train-the-trainer course for robotic surgery training. *Eur Urol* 2019;75:775–85.
- [10] Gallagher A. Proficiency-based progression simulation training for more than an interesting educational experience. *J Musculoskelet Surg Res* 2018;2:139–41.