



Universiteit
Leiden
The Netherlands

Destructive inflammatory reaction after an autologous retinal pigment epithelium and choroid transplantation: no detection of an auto-immune response

Romunde, S.H.M. van; Vergouwen, D.P.C.; Iacovello, D.; Roelen, D.L.; Verdijk, R.M.; Berge, J.C.E.M. ten; ... ; Meurs, J.C. van

Citation

Romunde, S. H. M. van, Vergouwen, D. P. C., Iacovello, D., Roelen, D. L., Verdijk, R. M., Berge, J. C. E. M. ten, ... Meurs, J. C. van. (2022). Destructive inflammatory reaction after an autologous retinal pigment epithelium and choroid transplantation: no detection of an auto-immune response. *Journal Of Ophthalmic Inflammation And Infection*, 12(1).
doi:10.1186/s12348-022-00305-2

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3486990>

Note: To cite this publication please use the final published version (if applicable).

RESEARCH

Open Access



Destructive inflammatory reaction after an autologous retinal pigment epithelium and choroid transplantation: no detection of an auto-immune response

Saskia H. M. van Romunde^{1*}, Daphne P. C. Vergouwen^{2,3}, Daniela Iacovello⁴, Dave L. Roelen⁵, Robert M. Verdijk^{1,6,7}, Josianne C. E. M. ten Berge², Grazia Pertile⁴, Marco W. J. Schreurs³ and Jan C. van Meurs^{1,2}

Abstract

Purpose: Five patients who underwent uncomplicated retinal pigment epithelium (RPE)-choroid transplantation for neovascular age-related macular degeneration developed a destructive inflammatory reaction causing subretinal fluid accumulation and extensive RPE atrophy in the graft. We hypothesized that this inflammation could be caused by an auto-immune response against the graft, resulting in circulating auto-antibodies. The aim of our study was to examine a potential autoimmune origin, which would allow a more targeted therapy approach.

Methods: Five above-mentioned patients and four control groups of five patients each were included: 1) after uncomplicated RPE-choroid transplantation, 2) after full macular translocation, 3) treated with anti-vascular endothelial growth factor, and 4) healthy controls. Histopathology of rejected graft tissue was performed using standard procedures. Presence of RPE-choroid autoantibodies in serum was examined by indirect immunofluorescence and Western blot, and human leukocyte antigen (HLA) typing was performed.

Results: Histopathological examination of an explanted graft showed infiltration of T-lymphocytes and macrophages in the choroid and RPE, and an increased number of B-cell lymphocytes were found in the choroid. Indirect immunofluorescence showed weak RPE-choroid autoantibody immunoreactivity in three patients of different groups. Western blot did not show specific RPE-choroid autoantibody immunoreactivity and no difference of HLA genotypes between the groups was found.

Conclusions: Although local mononuclear inflammatory cell infiltration and a high number of B-lymphocytes were observed in an explanted graft, we did not detect serological evidence of an autoimmune origin of the postoperative inflammation using direct immunofluorescence and Western Blot. Alternatively, the graft failure may have been caused by local innate inflammation, triggered by breakdown of tolerance. Based on our current findings of this small study group, we have no rationale to pursue therapies targeted towards autoreactive graft failure. More research is needed to confirm our findings.

Keywords: Age-related macular degeneration, Antibodies, Western blot, Immunofluorescence, Submacular surgery

Introduction

Neovascular age-related macular degeneration (AMD) is a common eye disease in the elderly population that is usually treated with intraocular injections of anti-vascular endothelial growth factor (anti-VEGF)

*Correspondence: s.vanromunde@oogziekenhuis.nl

¹ Department of Vitreoretinal surgery, Rotterdam Eye Hospital, Schiedamsse Vest 180 – 3011BH, Rotterdam, the Netherlands
Full list of author information is available at the end of the article

[1]. Sometimes anti-VEGF is less effective, for example in case of a retinal pigment epithelial (RPE) tear, a large subretinal hemorrhage, or subretinal fibrosis. In these patients, surgery can be an alternative treatment option. In RPE-choroid transplantation and full macular translocation (FMT), the choroidal neovascularization (CNV) is surgically removed and the macula is provided with extrafoveal RPE in two different ways. Specifically, in RPE-choroid transplantation, a sheet of RPE and choriocapillaris is transplanted from the midperiphery to the macula, while in FMT the fovea is rotated to an extrafoveal area after counterrotation of the globe [2, 3].

In the Rotterdam Eye Hospital (REH) and Ospedale Sacro Cuore (OSC) RPE-choroid transplantation for AMD has been performed in more than 400 patients [3, 4]. An unusual postoperative course occurred several months after uncomplicated surgery in five patients, which was characterized by accumulation of subretinal fluid (SRF), development of hyperreflective spots (HRS) and diffuse RPE atrophy, suggesting a destructive inflammatory reaction.

The eye is unique in its ways to suppress inflammation. Several passive and active systems include the blood-retinal barrier, production of immunomodulatory cytokines, and suppression by regulatory T-cells [5]. Possibly, these systems were disrupted due to surgical stress or ischemic injury. According to the 'danger theory', distressed or injured cells can evoke an inflammatory response via damage-associated molecular patterns (DAMPs), activating a T-cell response [6]. We hypothesize that in the patients with the indicated inflammatory reaction after surgery, the transplanted RPE cells were recognized as 'dangerous' by antigen presenting cells, leading to the activation of an adaptive autoreactive T-cell response, and subsequent a B-cell mediated autoimmune response, illustrated by production of autoantibodies and eventually degradation of the graft.

Autoimmunity in several diseases is associated with certain human leukocyte antigen (HLA) genotypes, for example HLA-DR15 and multiple sclerosis [7]. Similarly, there could be an HLA predisposition in patients with the postoperative inflammation after RPE-choroid transplantation when its origin would be autoimmune.

The aim of this study was to investigate the potential role of an adaptive autoimmune response in the five patients that developed the inflammatory reaction after RPE-choroid transplantation, by identifying the presence of RPE-choroid reactive autoantibodies and evaluating a possible HLA-based genetic predisposition. An autoimmune mechanism would allow us to target specific factors in the autoimmune response, e.g. with monoclonal antibody therapy.

Main text

Materials and methods

This case-control study was conducted according to the Declaration of Helsinki and the Good Clinical Practice guidelines. The study was approved by the local ethical committee (trial number: NL72704.078.20).

Patients

Five patients that developed SRF, HRS, and diffuse RPE atrophy after uncomplicated RPE-choroid transplantation were included at the REH (Rotterdam, the Netherlands), or OSC (Negrar (Verona), Italy). Considering that presence of auto-antibodies has been described previously in AMD patients and even after laser photocoagulation [8, 9] four control groups were included each consisting of five patients: 1) patients that underwent uncomplicated RPE-choroid transplantation, 2) patients that underwent uncomplicated FMT, 3) AMD patients treated with anti-VEGF injections, and 4) healthy age-matched controls. Exclusion criteria were a history of uveitis, diabetic retinopathy, and a systemic autoimmune disease.

Data collection

Retrospectively, the following data were collected from medical files of the included patients with the inflammatory reaction: demographics, ophthalmic history, general history, indication for surgery, intra- and post-operative complications, and best-corrected visual acuity (BCVA) before surgery, after surgery and at last visit. Spectral Domain – optical coherence tomography (SD-OCT) images, fundus photography, fluorescein angiography images and fundus autofluorescence (FAF) were collected. From the subjects of the control groups, demographics, ocular history and general history were reported. After informed consent was obtained, peripheral blood was collected providing both serum, and mononuclear cell samples for laboratory tests. The samples were stored at a temperature of -80°C until analyses.

Histopathology

In one patient the RPE-choroid graft was explanted because of a recurrent submacular hemorrhage and histopathologic examination was performed. The choroidal explant tissue was fixated in neutral buffered-formalin 10% and after processing, the formalin fixed paraffin embedded block was cut at $4\mu\text{m}$ sections which were mounted on slides for histochemical and immunohistochemical staining. Routine staining protocols were used for standard haematoxylin and eosin staining using an automated staining system (HE600, Ventana Medical Systems, Tucson, AZ, USA). Immunohistochemistry was performed with an automated, validated and accredited

staining system (Ventana Benchmark ULTRA, Ventana Medical Systems, Tucson, AZ, USA) using ultraview universal DAB detection kit. In brief, following deparaffinisation and heat-induced antigen retrieval, the tissue samples were incubated according to their optimized time with the antibody of interest. Incubation was followed by haematoxylin II counter stain for 8 minutes and then a blue coloring reagent for 8 minutes according to the manufacturer's instructions (Ventana). Tonsil tissue was used as a positive control for all antibodies.

Indirect immunofluorescence

Serum anti-RPE-choroid antibodies and anti-retinal antibodies were determined as described previously, with slight optimization of the protocol [10]. In short, biochip primate retinal slides (Euroimmun, Lübeck, Germany) were incubated with patients' serum for 30 minutes at room temperature (RT) (dilution 1:100). After incubation, the slides were washed with phosphate-buffered saline (PBS) for 15 minutes and incubated with goat anti-human IgG Cy5 (ab97172; Abcam, Cambridge, UK) for 30 minutes at RT. Tissue slides were washed again with PBS for 15 minutes, embedded in glycerol and covered with a coverslip. Slides were analysed and photographed using an Axioplan2 fluorescence microscope with AxioCam MRm at 160x magnification and an exposure time of 6 seconds (Zeiss, Thornwood, NY, USA). In case of a weak result the procedure was repeated with a serum dilution 1:50 and 1:25. PBS and serum negative for anti-nuclear antibodies (ANA) were used as negative references. Recombinant rabbit anti-RPE65 antibody (ab231782; dilution 1:250; Abcam, Cambridge, UK), recombinant anti-collagen type 1 (ab138492; dilution 1:500; Abcam, Cambridge, UK), and serum of an anti-retinal antibody positive, and ANA-positive subject (dilution 1:100) were used as positive references.

Western blot protein extracts

Human RPE and choroid tissue extracts for Western blot were prepared as described previously, implementing slight changes in the protocol [11]. RPE and choroid were obtained from a healthy donor eye post-mortem. After careful removal of the retina, RPE and choroid were dissected and subsequently homogenized in RIPA buffer with Halt protease inhibitor cocktail (Thermo Fisher Scientific, Waltham, USA) for 30 minutes. After homogenization the samples were centrifuged at 17000G at 4°C for 15 minutes, the supernatants were separated, and protein yield was assessed by colorimetric Bradford assay. In addition to the human donor RPE-choroid protein extract, an RPE cell line was used to provide an RPE protein extract [12]. ARPE-19 cells were cultured in culture flasks in DME/F (HyClone, Logan, UT) containing 10%

fetal calf serum, and 1% penicillin/streptomycin. Two times (72 h and 24 h before the harvest) 50 µg/ml ascorbic acid and 200 µM L-Proline were added to the cell culture, and thrombin (5 U/ml) was added 48 h before the harvest. Confluent grown ARPE19 cells were harvested and subsequently homogenized in cold RIPA buffer with Halt protease inhibitor cocktail (Thermo Fisher Scientific, Waltham, USA). After homogenization, the samples were centrifuged at 17000G at 4°C for 15 minutes, the supernatants were separated, and protein yield was assessed by colorimetric Bradford assay. Extracts were stored at a temperature of – 80°C until further use.

Western blot procedure

Protein extract were diluted in RIPA buffer, with 4x Laemmli sample buffer (Bio-rad, Hercules, California, USA), and incubated at 95°C for 5 minutes. The lysate was loaded on 4–12% mini-PROTEAN TGX Stain-Free Precast Gels (Bio-rad, Hercules, California, USA), and transferred to nitrocellulose membranes. Membranes were blocked with trix-buffered saline (TBS) containing 3% bovine serum albumine (BSA), and subsequently incubated with diluted samples (serum 1:100 TBS-Tween 0,1% containing 3% BSA, other antibodies as mentioned) for 1 hour at RT. Membranes were washed multiple times using an automated laboratory workstation (auto-LIA 48; Fujirebio, Japan), and incubated with goat anti-rabbit IgG (IRDye 680; dilution 1:4000; Licor, Lincoln, NE, USA), or goat anti-Human IgG (IRDye 680; dilution 1:15000; Licor, Lincoln, NE, USA) for 30 minutes. Visualization was performed with the imaging system Odyssey CLx (Thermo Fisher Scientific, Waltham, USA). Recombinant rabbit anti-RPE65 antibody, anti-recoverin antibody, and recombinant anti-collagen type I antibody were used as references.

Human leukocyte antigen typing

DNA was collected from whole blood using a QIAamp DNA Blood Mini kit, or using PAXgene blood tubes and a PAXgene Blood DNA Kit (Qiagen, Hilden, Germany), following manufacturer protocol. HLA class I and II typing was performed on all samples except for the five healthy participants. The procedure was performed as described by van Sonderen et al. [13] The HLA type of the study subjects was compared to the frequency in 5604 Dutch healthy blood donors [13].

Statistical analyses

Because of the small number of patients, and the explorative nature of the study, no statistical analysis was performed. The data have been evaluated descriptively.

Table 1 Characteristics of patients with inflammatory reaction after retinal pigment epithelial (RPE)-choroid transplantation

Pt No.	Sex	Age	Eye	Ocular history	Indication for surgery	Onset ^a	BCVA preop	BCVA postop	BCVA inflammation	Blood withdrawal ^b
1	M	92	OS	None	RPE tear	3	20/200	20/40	20/63	15 & 42
2	F	84	OD	Submacular hemorrhage treated with vitrectomy, gas and r-tPA	RPE tear	8	20/125	20/125	20/200	13 & 37
3	F	86	OS	Glaucoma, phaco	RPE tear	5	20/100	20/100	20/200	0 & 24
4	M	71	OD	Glaucoma treated with Baerveldt implant, anti-VEGF	Fibrotic scar	2	20/40	20/40	20/1250	106
5	F	73	OS	Anti-VEGF	Fibrotic scar	3	20/100	20/40	20/80	43

^a Onset of inflammatory reaction after RPE-choroid transplantation in months

^b Time between inflammation and blood withdrawal in months

Anti-VEGF Anti-vascular endothelial growth factor, BCVA Best-corrected visual acuity, RPE Retinal pigment epithelium, r-tPA Recombinant tissue plasminogen activator

Results

The demographics and ocular history of the five patients that developed an inflammatory reaction after RPE-choroid transplantation are shown in Table 1. None of the patients had a history of uveitis, or systemic autoimmune disease. Patient 1–3 were operated at REH and patient 4 and 5 were operated at OSC. On ophthalmic examination, there were no cells in the anterior or posterior segment. The onset of the inflammatory reaction after surgery ranged from 2 to 8 months. Figure 1 shows the changes of the OCT image, including SRF, HRS, and diffuse RPE atrophy. The latter is also clearly illustrated on FAF. Patient 2 developed a recurrent CNV with a large hemorrhage 2 years after primary surgery (Fig. 2). The graft was explanted and a second autologous graft was placed under the fovea. Interestingly, with a follow up period of 3 years, the inflammatory reaction did not recur in the new graft.

Histopathologic assessment of the explanted first graft showed mononuclear inflammatory cell infiltration of both helper- and cytotoxic T-lymphocytes and macrophages in the RPE and choroid, based on increased presence of cells expressing CD3, CD4, CD8 and CD68, respectively. An increased number of B-lymphocytes, of which a few were plasma cells, were seen only in the choroid but not in the RPE, based on the presence of cells expressing CD20 and CD138, respectively. Patient 1–3 were treated with peribulbar steroid injection and patient 4 and 5 with topical steroids after the inflammatory reaction. In all patients SRF resolved after several months. The blood samples were withdrawn 0–106 months after the inflammatory reaction. In patient 1–3 a second sample was taken 24–42 months after the inflammation.

Demographics of the control groups are shown in Table 2. The groups RPE-choroid transplantation, anti-VEGF injection, and healthy participants were recruited at the REH and patients that underwent FMT were recruited at OSC.

Autoantibody detection by indirect immunofluorescence

Negative controls, positive anti-RPE65 and anti-retinal antibodies confirmed the validity of autoantibody detection by indirect immunofluorescence of ocular tissue (Fig. 3A–C). There was no strong immunoreactivity against RPE-choroid or other ocular structures observed in the serum of any of the subjects. In three patients there was weak fluorescence of the RPE and the photoreceptors: patient number 4 of the inflammation group, one patient in the FMT group, and one patient in the uncomplicated RPE-choroid transplantation group

(Fig. 3D–F). The fluorescence of the RPE increased with higher serum concentrations (1:50 and 1:25), but the signal remained weak. In one FMT patient, one patient after uncomplicated RPE-choroid transplantation and two patients treated with anti-VEGF only, various intraretinal structures (photoreceptor layer, ganglion cell layer, inner plexiform layer) showed weak fluorescence with increasing intensity with higher serum concentrations. The presence of antibodies in both disease and control groups indicates a lack of specific autoantibody immunoreactivity.

Autoantibody detection by Western blot

The results of the Western blots performed using the donor RPE-choroid tissue lysate and protein extract of the RPE-cell line are illustrated in Fig. 4. In both Western blots anti-RPE65 was positive and anti-recoverin was negative, validating the protein content of the lysates. Anti-collagen type 1 was positive with high intensity light exposure in the Western blot using donor tissue lysate, indicating the presence of choroidal tissue. This Western blot showed complete absence of potential autoantibody immunoreactivity in all patients. The positive signals at ~40 kDa and ~60 kDa could be explained by the presence of immunoglobulins in human donor RPE-choroid tissue. The Western blot using the RPE cell line demonstrated no evident immunoreactivity in the inflammation group as well. There were several weak signals mostly in the group of uncomplicated RPE-choroid transplantation. For the most part these signals did not seem to be specific. The patients with weak immunoreactivity on Western blot did not match those patients with weak immunoreactivity on indirect immunofluorescence, indicating lack of relevance of these observations.

HLA typing

There was no overrepresentation of an HLA class I or class II type in patients that developed the inflammatory reaction as compared to the included controls and to the general population. No significant differences could be found between the inflammation group, the control groups, and the healthy subjects. See Supplementary Table S1 for complete results.

Discussion

The aim of our study was to examine a potential autoimmune origin of RPE-choroid transplant failure. Our initial hypothesis is supported by the presence of a mononuclear inflammatory infiltrate, including T-cells, in the failed graft and a high number of B-lymphocytes in the choroid. No serological evidence, however, of a systemic




Fig. 1 Imaging of patients with inflammatory reaction after RPE-choroid transplantation. Patient 1–5 after retinal pigment epithelium (RPE) – choroid transplantation. **A** Optical coherence tomography (OCT)-image taken 2–6 weeks after surgery show an intact layer of transplanted RPE. **B** OCT-image during inflammation showing subretinal fluid and diffuse atrophy of the transplanted RPE. **C** Fundus autofluorescence during inflammation illustrating a leopard-like pattern of atrophy

humoral response was detected using indirect immunofluorescence as well as Western blot analyses in patients who developed an inflammatory reaction after uncomplicated RPE-choroid transplantation. In addition, we were unable to identify a specific HLA association as possible genetic predisposition for autoimmune mediated graft failure.

The role of humoral autoimmunity in graft failure, activated by T-helper cells, has become more clear in recent years [14]. Presence of autoantibodies has been found after allogeneic and autologous transplantation [14–17]. Along the same line, we hypothesized that the inflammatory reaction was associated with an adaptive response of autoimmune origin resulting in the formation of detectable autoantibodies against RPE-choroid tissue and overrepresentation of a certain HLA type. The results of our study, however, do not support our theory.

The appearance of the graft and findings on multimodal imaging, i.e. SRF, HRS, and RPE atrophy, implies a destructive inflammatory process. It is common to notice a thin layer of SRF underneath the graft, which disappears when revascularisation is established [18]. However, in the presented patients the fluid appeared in a later phase and persisted for several months. HRS are believed to represent migrated RPE cells or activated microglial cells [19]. The latter cells have an important role in the innate immune system, and possibly in the pathogenesis of several retinal diseases including AMD [20, 21]. Postoperative RPE atrophy in the graft is highly prevalent after RPE-choroid transplantation [3]. Normally it initiates at the edge of the graft and slowly expands towards the centre. Expansion is more rapid in areas with a large gap between the native RPE and transplanted RPE [22]. However, the sudden onset and the involvement of the entire graft suggest an inflammation instead of a degenerative process. Moreover, the histopathological examination of the explanted graft showed infiltration of mononuclear inflammatory cells.

There are several possible explanations for our results. The immune reaction may have been targeted specifically against injured RPE. According to the danger theory, DAMP molecules induce inflammation to eliminate the injured cells [6, 23]. In that case, we may not expect any adaptive immune reactivity against healthy RPE-choroid tissue. Although some antibody responses persist lifelong, others last only a short time [14]. Possibly, autoantibodies against the transplanted tissue were produced in a concentration too low to detect in serum or the serum was sampled too long after the actual autoantibody production. Concentrations may have been higher




Fig. 2 Imaging and histopathologic findings in a patient with an inflammatory reaction after RPE-choroid transplantation. Imaging (A-E) and histopathologic examination (F-I) of patient 2. Fundus shows a leopard-like pattern on the graft (A). At the time of inflammation fluorescein angiography shows no recurrence (B) and indocyanine green angiography shows perfusion of the graft (C). OCT-image during inflammation shows subretinal fluid and hyperreflective spots (arrow heads) (D). No inflammatory reaction occurred in the second RPE-choroid transplant 21 months after re-transplantation (E). Histopathology of the explanted graft demonstrates presence of CD3 positive T-lymphocytes (F), showing both CD4 positive T-helper phenotype (G), and CD8 positive cytotoxic T cell phenotype (H), and CD68 positive macrophages (I) in the choroid and RPE. An increased number of CD20 positive B-cells (J), of which a few were CD138 positive plasma cells (K), was seen in the choroid but not in the RPE

Table 2 Characteristics of the control groups

	RPE-choroid transplantation (n = 5)	Full macular translocation (n = 5)	Anti-VEGF injection (n = 5)	Healthy participants (n = 5)
Male, no (%)	3 (60%)	3 (60%)	3 (60%)	3 (60%)
Mean age (SD, range)	71 (5, 65–78)	82 (3, 77–84)	84 (5, 78–90)	72 (4, 66–77)

Anti-VEGF Anti-vascular endothelial growth factor, RPE Retinal pigment epithelium, SD Standard deviation

either locally at the inflammation site or earlier during the failure process. Unfortunately, we did not have any ocular fluids or tissues available to examine local deposition of autoantibodies. Alternatively, it is possible that only a local T-cell mediated inflammation was causing SRF and damage to the RPE of the graft. Studying autoreactive T-cells is highly challenging in this matter, therefore we first investigated the indirect consequence

of an autoreactive response by T-cells, using detection of autoantibodies. Without further evidence for auto-reactivity we believe it is not feasible to further study antigen-specific T-cells. Moreover, absence of inflammation after transplantation of a second graft in one patient implies that there would be no autoimmune origin. Still, it remains unclear what triggered the inflammation, since




Fig. 3 Indirect immunofluorescence. Indirect immunofluorescence of references (A-C) and serum of patients (dilution 1:100) with weak fluorescence of the retinal pigment epithelium (RPE) and photoreceptors (*arrow heads*) (D-F). Photographs were taken at 160x magnification and 6 seconds exposure time. **A** healthy control. **B** Recombinant rabbit anti-RPE65 antibody. **C** anti-recoverin antibody. **D** patient with inflammation after RPE-choroid transplantation. **E** patient that underwent full macular translocation. **F** patient after uncomplicated RPE-choroid transplantation. CH = choroid, PR = photoreceptor, RPE = retinal pigment epithelium

the surgery and initial postoperative course in all five patients were uncomplicated.


Detection of autoantibodies is generally used for diagnostic purposes when autoimmune retinopathy is suspected. In the last decade, the interest for ocular autoantibodies has broadened. There is growing evidence for the involvement of autoantibodies in the pathogenesis of other ocular diseases, including AMD [24]. It has been demonstrated that autoantibodies to macular tissue are significantly more prevalent and show stronger autoreactivity in AMD patients compared to age-matched controls [25]. Patel et al. found a higher titre of anti-retinal antibodies in dry and exudative AMD patients compared to controls (94%, 82% and 9%, respectively) [9]. In agreement with the literature, we noticed more immunofluorescent reactivity in the RPE layer and neuroretina in AMD patients compared to healthy controls in our study. Drusen containing complement factors and IgG immune complexes add to the suggestion that immune-related processes play a role in the pathogenesis of AMD [26]. The question whether such autoantibody-driven immunoreactivity is pathogenic or merely an epiphenomenon of tissue destruction is still largely unanswered. Further research into this topic is important because autoimmune biomarkers might be useful for diagnoses, prognoses and perhaps as a target for AMD treatment [24].

This study has several limitations. Evidently, the number of patients is too low to draw definitive conclusions. Autoantibody concentration may have been insufficient for detection due to the long period between onset

of inflammation and blood sample collection in some patients. The validity of antiretinal antibody testing has been questioned previously because of false-positive binding [27]. Probably, the reported low validity may be influenced by lack of standardization of the laboratory assessment [10, 28]. In our study, all assessments were performed in one laboratory according to previously described protocols. The autoimmune origin of the observed inflammation could be primarily T-cell mediated. Unfortunately, we were unable to specifically examine anti-RPE-choroid T-cell immunoreactivity due to technical limitation. Although the low number of patients precludes any conclusion, the lack of HLA type association with the inflammatory reaction may argue against autoreactive T-cell involvement.

Conclusion

Assessment with indirect immunofluorescence and Western blot of RPE-choroid reactive autoantibodies and HLA typing did not show evidence for an autoimmune origin of the inflammatory complication, in spite of the observed numerous B-cells in the choroid and the local tissue infiltration by T-cells. Therefore, in this series of patients, we found no rationale to pursue more specific and off-label targeted immunomodulatory therapies such as anti-B-cell antibodies. Possibly the inflammation was caused by a local innate inflammation only, triggered by a breakdown of tolerance and weakening of the blood-retinal barrier. Moreover, there are important drawbacks



including a small sample size and limitations of the laboratory assessments. Although we were unable to pinpoint the cause in our series of patients, we consider it important to recognize and study this complication, and to establish a diagnostic set-up, as we may encounter this phenomenon more often in the future. Though submacular surgery for AMD is rarely performed since the introduction of anti-VEGF, there is a growing interest in

the transplantation of induced pluripotent stem cells and embryonic stem cell derived RPE [29].

Abbreviations

ANA: Antinuclear antibodies; Anti-VEGF: Anti-vascular endothelial growth factor; BCVA: Best-corrected visual acuity; BSA: Bovine serum albumine; CNV: Choroidal neovascularization; DAMP: Damage-associated molecular pattern; FAF: Fundus autofluorescence; FMT: Full macular translocation; HLA: Human leukocyte antigen; HRS: Hyperreflective spots; PBS: Phosphate-buffered saline; REH: Rotterdam Eye Hospital; RPE: Retinal pigment epithelium; RT:

Room temperature; OSC: Hospital Sacro Cuore – Don Calabria; SD-OCT: Spectral-domain optical coherence tomography; SRF: Subretinal fluid; TBS: Trix-buffered Saline.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12348-022-00305-2>.

Additional file 1: Supplementary Table S1. Human leukocyte antigen (HLA) results.

Acknowledgements

The authors would like to thank Aniki Rothova for her critical revision of the manuscript.

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

SvR assisted in laboratory assessments, collected and analysed the data and drafted the manuscript. DV performed the laboratory assessment, analysed the data and revised the manuscript. DI collected the blood samples and revised the manuscript. DR performed the HLA gene typing and revised the manuscript. RV performed the histopathological examination of the explanted graft, prepared the donor tissue for Western blot, and revised the manuscript. JtB contributed to design of the work and revised the manuscript, GP enabled the acquisition of the blood samples and revised the manuscript, MS contributed to the design of the work, to the interpretation of the results and revised the manuscript, JvM had a major contribution in the conception of the study, enabled acquisition of the blood samples and revised the manuscript. All authors read and approved the final manuscript.

Funding

This study was sponsored by Combined Ophthalmic Research Rotterdam (CORR).

Declarations

Ethics approval and consent to participate

Ethical approval was obtained by the Medische Ethische Toetings Commissie (METC) Erasmus MC. Trial number: NL72704.078.20.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Vitreoretinal surgery, Rotterdam Eye Hospital, Schiedamse Vest 180 – 3011BH, Rotterdam, the Netherlands. ²Department of Ophthalmology, Erasmus MC, University Medical Center Rotterdam, Doctor Molewaterplein 40, 3015, GD, Rotterdam, the Netherlands. ³Department of Immunology, Erasmus MC, University Medical Center Rotterdam, Doctor Molewaterplein 40, 3015, GD, Rotterdam, the Netherlands. ⁴Department of Ophthalmology, IRCCS Sacro Cuore Don Calabria, Viale Rizzardi, 437024 Negrar, Italy. ⁵Department of Immunology, Leiden University Medical Center, Albinusdreef 2, 2333, ZA, Leiden, the Netherlands. ⁶Department of Pathology, Section Ophthalmic Pathology, Erasmus MC, University Medical Center, Doctor Molewaterplein 40, 3015, GD, Rotterdam, the Netherlands. ⁷Department of Pathology, Leiden University Medical Center, Albinusdreef 2, 2333, ZA, Leiden, the Netherlands.

Received: 13 March 2022 Accepted: 11 August 2022

Published online: 26 August 2022

References

1. Rofagha S, Bhisitkul RB, Boyer DS, Sadda SR, Zhang K, SEVEN-UP Study Group (2013) Seven-year outcomes in ranibizumab-treated patients in ANCHOR, MARINA, and HORIZON: a multicenter cohort study (SEVEN-UP) ophthalmology. *Ophthalmology* 120(11):2292–2299. <https://doi.org/10.1016/j.ophtha.2013.03.046>
2. van Romunde SHM, Polito A, Bertazzi L, Guerriero M, Pertile G (2015) Long-term results of full macular translocation for choroidal neovascularization in age-related macular degeneration. *Ophthalmology* 122(7):1366–1374. <https://doi.org/10.1016/j.ophtha.2015.03.012>
3. van Romunde SHM, Polito A, Peroglio Deiro P, Guerriero M, Pertile G (2019) Retinal pigment epithelium-choroid graft with a peripheral retinotomy for exudative age-related macular degeneration: long-term outcome. *Retina* 39(2):288–295. <https://doi.org/10.1097/IAE.00000000000001945>
4. van Zeeburg EJ, Maaijwee KJ, Missotten TO, Heimann H, van Meurs JC (2012) A free retinal pigment epithelium-choroid graft in patients with exudative age-related macular degeneration: results up to 7 years. *Am J Ophthalmol* 153(1):120–7.e2. <https://doi.org/10.1016/j.ajo.2011.06.007>
5. Mochizuki M, Sugita S, Kamoi K (2013) Immunological homeostasis of the eye. *Prog Retin Eye Res* 33:10–27. <https://doi.org/10.1016/j.preteyeres.2012.10.002>
6. Matzinger P (2002) The danger model: a renewed sense of self. *Science* 296(5566):301–305. <https://doi.org/10.1126/science.1071059>
7. Dendrou CA, Petersen J, Rossjohn J, Fugger L (2018) HLA variation and disease. *Nat Rev Immunol* 18(5):325–339. <https://doi.org/10.1038/nri.2017.143>
8. Rahi AH, Addison DJ (1983) Autoimmunity and the outer retina. *Trans Ophthalmol Soc U K* 103(Pt 4):428–437
9. Patel N, Ohbayashi M, Nugent AK, Ramchand K, Toda M, Chau KY, Bunce C, Webster A, Bird AC, Ono SJ, Chong V (2005) Circulating anti-retinal antibodies as immune markers in age-related macular degeneration. *Immunology* 115(3):422–430. <https://doi.org/10.1111/j.1365-2567.2005.02173.x>
10. Ten Berge JC, Schreurs MW, Vermeer J, Meester-Smoor MA, Rothova A (2016) Prevalence and clinical impact of antiretinal antibodies in uveitis. *Acta Ophthalmol* 94:282–288. <https://doi.org/10.1111/aos.12939>
11. Kuiper JJ, Emmelot ME, Rothova A, Mutis T (2013) Interleukin-17 production and T helper 17 cells in peripheral blood mononuclear cells in response to ocular lysate in patients with birdshot chorioretinopathy. *Mol Vis* 19:2606–2614
12. Bastiaans J, van Meurs JC, van Holten-Neelen C, Nagtzaam NM, van Hagen PM, Chambers RC, Hooijkaas H, Dik WA (2013) Thrombin induces epithelial-mesenchymal transition and collagen production by retinal pigment epithelial cells via autocrine PDGF-receptor signaling. *Invest Ophthalmol Vis Sci* 54(13):8306–8314. <https://doi.org/10.1167/iov.13-12383>
13. van Sonderen A, Roelen DL, Stoop JA, Verdijk RM, Haasnoot GW, Thijs RD, Wirtz PW, Schreurs MW, Claas FH, Sillevs Smitt PA, Titulaer MJ (2017) Anti-LGI1 encephalitis is strongly associated with HLA-DR7 and HLA-DRB4. *Ann Neurol* 81(2):193–198. <https://doi.org/10.1002/ana.24858>
14. Siu JHY, Motallebzadeh R, Pettigrew GJ (2020) Humoral autoimmunity after solid organ transplantation: germinal ideas may not be natural. *Cellular Immunol* 354:104131. <https://doi.org/10.1016/j.cellimm.2020.104131>
15. Zorn E, See SB (2019) Is there a role for natural antibodies in rejection following transplantation? *Transplantation* 103(8):1612–1619. <https://doi.org/10.1097/TP.0000000000002743>
16. Ali KF, San Martin VT, Stevens T, Walsh RM, Bottino R, Trucco M, Hatipoglu B (2018) Autoimmunity in autologous islet transplantation. *OBM Transplant* 2(3). <https://doi.org/10.21926/obm.transplant.1803014>
17. Bellin MD, Moran A, Wilhem JJ, O'Brien TD, Gottlieb PA, Yu L, Dunn TB (2015) Development of autoimmune-mediated β cell failure after total pancreatectomy with autologous islet transplantation. *Am J Transplant* 15(7):1991–1994. <https://doi.org/10.1111/ajt.13216>
18. van Zeeburg EJ, Cereda MG, van der Schoot J, Pertile G, van Meurs JC (2011) Early perfusion of a free RPE-choroid graft in patients with exudative macular degeneration can be imaged with spectral domain-OCT. *Invest Ophthalmol Vis Sci* 52(8):5881–5886. <https://doi.org/10.1167/iov.11-7245>
19. Vujosevic S, Berton M, Bini S, Casciano M, Cavarzeran F, Midena E (2016) Hyperreflective retinal spots and visual function after anti-vascular

- endothelial growth factor treatment in center-involving diabetic macular edema. *Retina* 36(7):1298–1308. <https://doi.org/10.1097/IAE.0000000000000912>
20. Streilein JW, Ma N, Wenkel H, Ng TF, Zamiri P (2002) Immunobiology and privilege of neuronal retina and pigment epithelium transplants. *Vis Res* 42(4):487–495. [https://doi.org/10.1016/s0042-6989\(01\)00185-7](https://doi.org/10.1016/s0042-6989(01)00185-7)
 21. Ma W, Zhao L, Wong WT (2012) Microglia in the outer retina and their relevance to pathogenesis of age-related macular degeneration. *Adv Exp Med Biol* 723:37–42. https://doi.org/10.1007/978-1-4614-0631-0_6
 22. Pertile G, Mete M, Peroglio Deiro A, Guerriero M, Sartore M, Alfano M, Polito A (2018) New insights into the development and progression of geographic atrophy after full thickness autologous choroidal graft. *Invest Ophthalmol Vis Sci* 59(4):AMD93–AMD103. <https://doi.org/10.1167/iov.18-24229>
 23. Relja B, Land WG (2020) Damage-associated molecular patterns in trauma. *Eur J Trauma Emerg Surg* 46(4):751–775. <https://doi.org/10.1007/s00068-019-01235-w>
 24. Morohoshi K, Goodwin AM, Ohbayashi M, Ono SJ (2009) Autoimmunity in retinal degeneration: autoimmune retinopathy and age-related macular degeneration. *J Autoimmun* 33(3–4):247–254. <https://doi.org/10.1016/j.jaut.2009.09.003>
 25. Iannaccone A, Giorgianni F, New DD, Hollingsworth TJ, Umfress A, Alhateem AH, Neeli I, Lenchik NI, Jennings BJ, Calzada JI, Satterfield S, Mathews D, Diaz RI, Harris T, Johnson KC, Charles S, Kritchevsky SB, Gerling IC, Beranova-Giorgianni S, Radic MZ (2015) Health ABC study (2015) circulating autoantibodies in age-related macular degeneration recognize human macular tissue antigens implicated in autophagy, immunomodulation, and protection from oxidative stress and apoptosis. *Plos One* 10(12). <https://doi.org/10.1371/journal.pone.0145323>
 26. Hageman GS, Luthert PJ, Victor Chong NH, Johnson LV, Anderson DH, Mullins RF (2001) An integrated hypothesis that considers drusen as biomarkers of immune-mediated processes at the RPE-Bruch's membrane interface in aging and age-related macular degeneration. *Prog Retin Eye Res* 20(6):705–732. [https://doi.org/10.1016/s1350-9462\(01\)00010-6](https://doi.org/10.1016/s1350-9462(01)00010-6)
 27. Chen JJ, McKeon A, Greenwood TM, Flanagan EP, Bhatti MT, Dubey D, Pulido JS, Iezzi R, Smith WM, Sen HN, Gordon LK, Pittcock SJ (2021) Clinical utility of antiretinal antibody testing. *JAMA Ophthalmol* 139(6):658–662. <https://doi.org/10.1001/jamaophthalmol.2021.0651>
 28. Forooghian F, Macdonald IM, Heckenlively JR, Héon E, Gordon LK, Hooks JJ, Detrick B, Nussenblatt RB (2008) The need for standardization of antiretinal antibody detection and measurement. *Am J Ophthalmol* 146(4):489–495. <https://doi.org/10.1016/j.ajo.2008.05.046>
 29. Mandai M, Watanabe A, Kurimoto Y, Hiram Y, Morinaga C, Daimon T, Fujihara M, Akimaru H, Sakai N, Shibata Y, Terada M, Nomiya Y, Tanishima S, Nakamura M, Kamao H, Sugita S, Onishi A, Ito T, Fujita K, Kawamata S, Go MJ, Shinohara C, Hata KI, Sawada M, Yamamoto M, Ohta S, Ohara Y, Yoshida K, Kuwahara J, Kitano Y, Amano N, Umekage M, Kitaoka F, Tanaka A, Okada C, Takasu N, Ogawa S, Yamanaka S, Takahashi M (2017) Autologous induced stem-cell-derived retinal cells for macular degeneration. *N Engl J Med* 376(11):1038–1046. <https://doi.org/10.1056/NEJMoa1608368>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
