



Universiteit
Leiden
The Netherlands

Proactive care programs in the emergency department: effectiveness and feasibility

Loon, M. van

Citation

Loon, M. van. (2023, April 13). *Proactive care programs in the emergency department: effectiveness and feasibility*. Retrieved from <https://hdl.handle.net/1887/3593961>


Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3593961>

Note: To cite this publication please use the final published version (if applicable).





CHAPTER 6

Frequencies and reasons for unplanned emergency department return visits by older adults

Merel van Loon-van Gaalen
Ilje E. Voshol
M. Christien van der Linden
Jacobijn Gussekloo
Roos C. van der Mast

Submitted ■

ABSTRACT

Background

As unplanned Emergency Department (ED) return visits (URVs) are associated with adverse health outcomes in older adults, many EDs have initiated post-discharge interventions to reduce URVs. Unfortunately, most interventions fail to reduce URVs, including telephone follow-up after ED discharge, investigated in a recent trial. To understand why these interventions were not effective, we analyzed patient and ED visit characteristics and reasons for URVs within 30 days for patients aged ≥ 70 years.

Methods

Data was used from a randomized controlled trial, investigating whether telephone follow-up after ED discharge reduced URVs compared to a satisfaction survey call. Only observational data from control group patients were used. Patient and index ED visit characteristics were compared between patients with and without URVs. Two independent researchers determined the reasons for URVs and categorized them into: patient-related, illness-related, new complaints and other reasons. Associations were examined between the number of URVs per patient and the categories of reasons for URVs.

Results

Of the 1659 control group patients, 222 (13.4%) had at least one URV within 30 days. Male sex, ED visit in the 30 days before the index ED visit, triage category “urgent”, longer length of ED stay, urinary tract problems, and dyspnea were associated with URVs. Of the 222 patients with an URV, 31 (14%) returned for patient-related reasons, 95 (43%) for illness-related reasons, 76 (34%) for a new complaint and 20 (9%) for other reasons. URVs of patients who returned ≥ 3 times were mostly illness-related (72%).

Conclusion

As the majority of patients had an URV for illness-related reasons or new complaints, these data fuel the discussion as to whether URVs can or should be prevented.

INTRODUCTION

With demographic change, there is an increase in Emergency Department (ED) presentations by patients aged 70 years and older worldwide.¹ Up to 25% of these older patients have an unplanned ED return visit (URV) within one month.²⁻⁶ Since URVs in older adults are associated with adverse health outcomes, they are often viewed as negative.^{3,7} Therefore, many EDs have initiated post-discharge interventions in order to reduce URVs.^{8,9}

Many post-ED discharge intervention programs are focused on older patients at high risk for hospital return. However, prediction tools that have been developed to identify patients at risk have poor predictive accuracy, contain different predictors, and are often not suitable for clinical use.^{4,10-13} However, all previous studies consistently report that the majority of older adults who return to the ED suffer from chronic and often comorbid health conditions, functional dependency or cognitive problems.^{2,10,14,15} In addition, several (psycho)social factors, such as living alone, lack of social support and uncertainty about the health condition, as well as insufficient understanding or provision of discharge information are found to be associated with URVs in older adults.^{2,6,7,11,14-19}

Several of these predicting factors could be addressed through specific interventions, such as patient education and community follow-up by a geriatric nurse. However, systematic reviews evaluating the effects of post-discharge interventions initiated in the ED have found that many were not effective in reducing ED re-attendances.^{8,9} In a pragmatic randomized controlled trial, our research group also failed to find a beneficial effect of a transitional care program, consisting of post-ED discharge telephone follow-up for older adults, on the reduction of unplanned hospital admissions and URVs within 30 days after ED discharge.²⁰

In order to understand why these interventions are not effective in reducing URVs, more insight is needed into the reasons why older patients return to the ED. Therefore, we investigated the frequencies, associated patient and ED visit characteristics and reasons for URVs within 30 days after the index ED visit among patients aged ≥ 70 years. In addition, we examined whether specific categories of reasons for URVs were associated with the number of URVs per patient.

METHODS

Study design and setting

For this study, we used data from a pragmatic randomized controlled trial (RCT). The research question of this RCT was whether a telephone follow-up call reduces unplanned hospitalizations and URVs within 30 days of ED discharge, compared to

a satisfaction survey call. The trial was conducted in the EDs of Haaglanden Medical Center (HMC), a non-academic teaching hospital in the Netherlands, from February 1, 2018 to July 1, 2019. In this RCT, 3175 patients were allocated to either the intervention (n=1516) or the control (n=1659) group, according to the month of their ED visit; patients included in odd months received an intervention telephone call to identify post-discharge problems and to offer additional information, and patients included in even months received a satisfaction survey telephone call.²⁰ The Medical Ethics Review Committee of HMC waived the necessity for formal approval of the study as it closely followed routine care (METC Zuidwest Holland, nr. 17-028).

Participants

For this study, only observational data from control group patients were used to exclude a possible effect of the intervention telephone follow-up call. Patients aged ≥ 70 years who were discharged from one of the EDs of HMC to an unassisted living environment were eligible for inclusion. Exclusion criteria were: admission to the hospital, discharge to a nursing home, another care facility or assisted living environment, and planned follow-up appointment at an outpatient clinic or at the ED within 24 hours.²⁰

Data collection and measurements

Unplanned ED return visits (URVs)

Data on ED return visits were collected from the electronic hospital system (EHS). ED return visits that could not be foreseen were defined as URVs.²¹ The index ED visit was the first ED visit during the study period that was followed by a telephone call.

Baseline data

We collected baseline data that were associated with URVs in previous studies, including demographics (age,^{4,10,13} gender,^{4,5,10} whether or not living alone,^{2,3,11,22}) and ED visit characteristics (mode of arrival, Manchester Triage System triage urgency level,²³ chief complaint, ED length of stay^{2,10,11,24}). We also used data concerning level of ED crowding at discharge, measured by the National Emergency Department Overcrowding Scale (NEDOCS).²⁵ Data were abstracted from the EHS by an information technology specialist, who was not involved in the study.²⁰

Determination of reasons for URVs

Prior to the start of the study, reasons for URVs were defined and categorized, based on findings in the literature (see Supplementary file 1).^{4,7,15,19,26} Two investigators (MvLvG and IEV), both medical doctors, independently determined and categorized the reason for each URV by reviewing the emergency medical records (EMRs). In case of disagreement, the EMR was reviewed and reasons for ED return were discussed until consensus was achieved. In case of no agreement, the EMR was reviewed by a

third investigator (MCvdL) for the final decision. This study method has been used in previous studies on URVs.^{16,26-28} During analyses, we found that only few URVs were categorized as physician-related, system-related or not classifiable. Therefore, these three categories have been merged into the “other reasons” category. This resulted in the following four main categories: 1. patient-related reasons, 2. illness-related reasons, 3. new complaints and 4. other reasons (see Supplementary file 1). The study was conducted in adherence to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement.²⁹

Statistical analysis

Categorical data are presented as numbers and percentages. Continuous data were skewed and therefore presented as median and interquartile ranges (IQR). Differences in characteristics of patients with and without URVs were analyzed using χ^2 -tests and univariable logistic regression.

The χ^2 -test was used to examine the association between number of URVs per patient and the categories of reasons for URVs. Odds ratios (ORs) were calculated with 95% confidence intervals (95% CIs). If a patient had multiple URVs during the 30-day follow up period, only the first URV was included to determine the reason for unplanned return and to assess associations between patient and index ED visit characteristics and occurrence of an URV. To investigate whether specific categories of reasons for URVs were associated with the number of URVs per patient, all URVs within 30 days after the index ED visit were included in the analysis.

Inter-rater reliability regarding the initial determination of reasons and categories of URVs was measured with Cohen’s kappa coefficient.

Statistical analyses were performed using the Statistical Package for the Social Sciences (IBM Corp. Released 2019. IBM SPSS Statistics, Version 26.0. Armonk, New York, USA).

RESULTS

Of the 1659 control group patients, 222 (13.4%) had at least one URV within 30 days. The total number of URVs within 30 days was 279.

Patient and ED visit characteristics associated with URVs

Table 1 shows the differences in baseline patient and index ED visit characteristics between patients with and without an URV. In univariate analysis, the following factors were associated with an URV within 30 days: male sex, ED visit in the 30 days before the index ED visit, triage category “urgent”, longer length of ED stay, and the chief complaints “urinary tract problems” and “dyspnea”.

Table 1. Baseline patient and index Emergency Department (ED) visit characteristics of patients with and without an URV

	Unplanned ED return visit (URV) ≤30 days		
	Yes (n=222)	No (n=1437)	OR (95% CI)
<u>Demographics</u>			
Age in years, median (IQR)	78 (73-83)	78 (73-83)	1.0 (1.0-1.0) [#]
Male sex, n (%)	106 (47.7)	588 (40.9)	1.3 (1.0-1.8)
Living without partner, n (%) [*]	83 (42.3)	413 (37.7)	1.2 (0.9-1.7)
<u>Characteristics of index ED visit</u>			
Mode of referral, n (%)			
- Self-referral	52 (23.4)	307 (21.4)	1.1 (0.8-1.6)
- General practitioner	65 (29.3)	485 (33.8)	0.8 (0.6-1.1)
- Medical specialist	44 (19.8)	246 (17.1)	1.2 (0.8-1.7)
ED visit ≤ 30 days before index visit, n (%)	45 (20.3)	164 (11.4)	2.0 (1.4-2.8)
Arrival by ambulance, n (%)	79 (35.6)	476 (33.1)	1.1 (0.8-1.5)
Triage category urgent, n (%) ^{**}	168 (76.4)	999 (70.0)	1.4 (1.0-1.9)
ED visit at daytime, n (%)	153 (68.9)	1009 (70.2)	0.9 (0.7-1.3)
Length of ED stay (minutes), median (IQR)	179 (128-242)	151 (106-204)	1.0 (1.0-1.1) [§]
NEDOCS at discharge ≥ 60, n (%) [^]	66 (34.6)	425 (32.9)	1.1 (0.8-1.5)
<u>Chief complaint, n (%)</u>			
- Urinary tract problems	16 (7.2)	47 (3.3)	2.3 (1.3-4.1)
- Headache or neurological problems	10 (4.5)	55 (3.8)	1.2 (0.6-2.4)
- Wounds	11 (5.0)	76 (5.3)	0.9 (0.5-1.8)
- Abdominal pain	16 (7.2)	76 (5.3)	1.4 (0.8-2.4)
- Syncope or palpitations	8 (3.6)	90 (6.3)	0.6 (0.3-1.2)
- Dyspnea	29 (13.1)	116 (8.1)	1.7 (1.1-2.6)
- Malaise	19 (8.6)	131 (9.1)	0.9 (0.6-1.5)
- Chest pain	28 (12.6)	177 (12.3)	1.0 (0.7-1.6)
- Limb complaints	37 (16.7)	299 (20.8)	0.8 (0.5-1.1)
- Fall or trauma	48 (10.7)	358 (13.1)	0.8 (0.6-1.1)
- Other complaints	17 (7.7)	152 (10.6)	0.7 (0.4-1.2)

[#] In univariable logistic regression model^{*} Living condition unknown in 26 patients with URV and in 341 patients without URV^{**} Triage category urgent: red, orange and yellow according to Manchester Triage System. Triage category

missing in 2 patients with URV and in 10 patients without URV

[§] Per 10 minutes increase in length of stay[^] If the NEDOCS at discharge is ≥ 60, the ED is considered to be busy. NEDOCS at discharge was missing in 31 patients with URV and 144 patients without URV, due to technical malfunction of electronic hospital system on days that patients were discharged from the ED.

ED Emergency Department, NEDOCS National Emergency Department Overcrowding Scale, IQR Interquartile Range, n number, URV unplanned emergency department return visit

Reasons for unplanned ED return

Figure 1 shows the number of URVs per reason for return. Patient-related reasons for URVs were found in 31 (14%) of the 222 patients with one or more URVs. The two most frequently occurring patient-related reasons for URVs were non-compliance with

discharge instructions (n=7), and worrying about health (n=19). Illness-related reasons for URVs were found in 95 (43%) of the 222 patients, of which recurrent complaints/disease (n=28) and progression of disease (n=38) were the two largest subgroups. A new complaint was the reason for URV in 76 (34%) of the 222 patients, and 20 (9%) out of the 222 patients had an URV for other reasons. Within the latter category, 6 of the 20 patients were misdiagnosed during the index ED visit, resulting in inappropriate treatment. Other physician-related and system-related reasons occurred in <2% of the 222 patients. Five URVs could not be classified and were therefore coded as “undefined”.




Figure 1. Reasons for unplanned Emergency Department (ED) return visits (n=222), divided into four categories. *ED*, emergency department; *GP*, general practitioner

Multiple URVs and reasons for ED return

Of the 222 patients with URVs, 176 (79.2%) had one URV, 39 (17.6%) had two URVs and 7 (3.2%) had three or more URVs within 30 days (Table 2). Most URVs in patients with one or two URVs were illness-related (40.9% and 46.2%, respectively) or because of a new complaint (38.0% and 30.8%, respectively). Patients with three or more URVs also returned mainly for illness-related reasons (72.0%), followed by patient-related reasons (24.0%), while new complaints were less common (4.0%).

■ **Table 2.** Association between the number of URVs per patient and per category of reasons for URVs

	Number of URVs per patient			Total
	1	2	≥ 3	
Number of patients, n (%)	176 (79.2)	39 (17.6)	7 (3.2)	222
Total number of URVs, n (%)	176 (63.1)	78 (28.0)	25 (9.0)	279
<u>Category reasons for URV:</u>				
Patient-related, n (%)	22 (12.5)	10 (12.8)	6 (24.0)	38
Illness-related, n (%)	72 (40.9)	36 (46.2)	18 (72.0)	126
New complaint, n (%)	67 (38.0)	24 (30.8)	1 (4.0)	92
Other, n (%)	15 (8.5)	8 (10.3)	0 (0.0)	23

ED, emergency department; n, number; URV, unplanned emergency department return visit

Inter-rater reliability regarding assessment of reasons for URVs

The inter-rater reliability after initial independent determination and categorization of the reasons for the URVs, measured with Cohen’s kappa coefficient, was 0.57. All disagreements concerning the determination of the reasons for URVs were solved by discussion between the two researchers and hence, the judgement of a third researcher for the final decision was not needed.

DISCUSSION

In this study, we found that 222 of the 1659 (13.4%) older adults had at least one URV within 30 days after being discharged from the ED. Of them, 171 (77%) returned for medical reasons, including 95 (43%) for problems related to the same illness of the index ED visit and 76 (34%) for a new medical complaint unrelated to the presenting problem of the index ED visit. URVs for patient-related reasons occurred in only 31 (14%) patients. Also, patients with three or more URVs returned mainly for problems related to the same illness of the index ED visit.

The URV rate in our study was comparable with URV rates among older adults reported in other studies.^{2-4,10,30} We also found that male sex,^{2,4,14,15} an ED visit in the 30 days before the index ED visit,^{2,10,11,14} triage category “urgent”, and a longer length of ED stay²⁴ were more common in patients with an URV. In accordance with other studies, we found that the chief complaints “urinary tract problems”²⁶ and “dyspnea”^{15,19,26,31} were associated with URVs.

Although transitional care programs that focus on patient education and post-discharge support may have a positive effect on the patient’s capacity for self-care, disease control, and perceived support,³²⁻³⁴ the limited number of patient-related URVs found in our study may explain why many of these programs do not reduce URVs. Our finding that most older adults returned to the ED for illness-related reasons or new problems indicates that the majority of these patients needed diagnostic work-

up of their health problem and/or acute care. This fuels the discussion of whether URVs can and need to be prevented. If the aim is to divert older patients from the ED, it will have to be sorted out where else diagnostic work-ups can be performed and patients can receive the necessary (acute) care outside the ED. This will depend on the organization of the health care system and should therefore be investigated locally. An example is the organization of an acute geriatric community hospital for older adults.³⁵ On the other hand, as the ED is organized and equipped to conduct targeted diagnostic work-ups and deliver acute care, it may be more feasible to make existing EDs more senior-friendly by applying the initiatives already described.³⁶⁻³⁹ Interventions focusing on close collaboration between primary care, hospital care, and community services may be more successful in reducing unplanned ED visits for older adults than interventions involving only the ED. Within these collaborations, it may be easier to deliver the best care for the patient at the most suitable location. It would be interesting to explore such collaborations in future studies.^{40,41}

STRENGTHS AND LIMITATIONS

We were able to compare an extensive set of patient and ED visit characteristics between patients with and without URVs. Although previous studies mentioned reasons for URVs in older adults, this is one of the few studies that investigated the frequencies of the different reasons for URVs in older adults.^{2,7} Data were prospectively collected and derived from the hospital database to diminish confounding by recall bias.

Some limitations, however, could be considered. The reasons for URVs were defined and categorized prior to the start of the study and based on explicit criteria, used in previous studies. However, the reasons for URVs were determined retrospectively. By having the URVs assessed by two independent researchers, we tried to comply with the classification criteria as much as possible. The Cohen's kappa coefficient of 0.57, reflecting a moderate inter-rater reliability of the categorization system, may be a limitation.

Furthermore, not all data about health determinants that are associated with hospital return were available. Finally, this study was conducted in two EDs of a non-academic hospital in the Netherlands. The findings may not be generalizable to all EDs. However, two studies, one conducted in a Dutch academic ED and one in two Australian large referral hospital EDs, reported comparable percentages of URVs for illness-related and patient-related reasons,⁷ and for new complaints.²

CONCLUSION

In this study, most older patients returned unplanned to the ED for medical reasons, whereas URVs for patient-related reasons, such as uncertainty about health or misunderstanding of discharge instructions, were less common. These findings may explain why many transitional care programs that focus on patient education and post-discharge support are ineffective in reducing URVs. In addition, the results suggest that most patients who return to the ED require urgent care. This fuels the discussion as to whether URVs can or need to be prevented.

REFERENCES

1. Samaras N, Chevalley T, Samaras D, Gold G. Older patients in the emergency department: a review. *Ann Emerg Med* 2010;56:261-269.
2. Arendts G, Fitzhardinge S, Pronk K, Hutton M, Nagree Y, Donaldson M. Derivation of a nomogram to estimate probability of revisit in at-risk older adults discharged from the emergency department. *Intern Emerg Med* 2013;8:249-254.
3. Aminzadeh F, Dalziel WB. Older adults in the emergency department: a systematic review of patterns of use, adverse outcomes, and effectiveness of interventions. *Ann Emerg Med* 2002;39:238-247.
4. de Gelder J, Lucke JA, de Groot B, et al. Predictors and Outcomes of Revisits in Older Adults Discharged from the Emergency Department. *J Am Geriatr Soc* 2018;66:735-741.
5. Graf CE, Giannelli SV, Herrmann FR, et al. Identification of older patients at risk of unplanned readmission after discharge from the emergency department - comparison of two screening tools. *Swiss Med Wkly* 2012;141:w13327.
6. Hastings SN, Barrett A, Weinberger M, et al. Older patients' understanding of emergency department discharge information and its relationship with adverse outcomes. *J Patient Saf* 2011;7:19-25.
7. Driesen B, Merten H, Wagner C, Bonjer HJ, Nanayakkara PWB. Unplanned return presentations of older patients to the emergency department: a root cause analysis. *BMC Geriatr* 2020;20:365. <https://doi.org/10.1186/s12877-020-01770-x>
8. Crede SH, O'Keeffe C, Mason S, et al. What is the evidence for the management of patients along the pathway from the emergency department to acute admission to reduce unplanned attendance and admission? An evidence synthesis. *BMC Health Serv Res* 2017;17:355.
9. Hughes JM, Freiermuth CE, Shepherd-Banigan M, et al. Emergency Department Interventions for Older Adults: A Systematic Review. *J Am Geriatr Soc* 2019;67:1516-1525.
10. Lowthian J, Straney LD, Brand CA, et al. Unplanned early return to the emergency department by older patients: the Safe Elderly Emergency Department Discharge (SEED) project. *Age Ageing* 2016;45:255-261.
11. McCusker J, Cardin S, Bellavance F, Belzile E. Return to the emergency department among elders: patterns and predictors. *Acad Emerg Med* 2000;7:249-259.
12. Hastings SN, Purser JL, Johnson KS, Sloane RJ, Whitson HE. Frailty predicts some but not all adverse outcomes in older adults discharged from the emergency department. *J Am Geriatr Soc* 2008;56:1651-1657.
13. LaMantia MA, Platts-Mills TF, Biese K, et al. Predicting hospital admission and returns to the emergency department for elderly patients. *Acad Emerg Med* 2010;17:252-259.
14. Šteinmiller J, Routasalo P, Suominen T. Older people in the emergency department: a literature review. *Int J Older People Nurs* 2015;10:284-305.
15. Sheikh S. Risk Factors Associated with Emergency Department Recidivism in the Older Adult. *West J Emerg Med* 2019;20:931-938.
16. Keith KD, Bocka JJ, Kobernick MS, Krome RL, Ross MA. Emergency department revisits. *Ann Emerg Med* 1989;18:964-968.

17. Rising KL, Padrez KA, O'Brien M, Hollander JE, Carr BG, Shea JA. Return visits to the emergency department: the patient perspective. *Ann Emerg Med* 2015;65:377-386.e373.

18. Nielsen LM, Gregersen Østergaard L, Maribo T, Kirkegaard H, Petersen KS. Returning to everyday life after discharge from a short-stay unit at the Emergency Department-a qualitative study of elderly patients' experiences. *Int J Qual Stud Health Well-being* 2019;14:1563428.

19. Trivedy CR, Cooke MW. Unscheduled return visits (URV) in adults to the emergency department (ED): a rapid evidence assessment policy review. *Emerg Med J* 2015;32:324-329.

20. van Loon-van Gaalen M, van der Linden MC, Gussekloo J, van der Mast RC. Telephone follow-up to reduce unplanned hospital returns for older emergency department patients: A randomized trial. *J Am Geriatr Soc* 2021;69:3157-3166.

21. Landrum L, Weinrich S. Readmission data for outcomes measurement: identifying and strengthening the empirical base. *Qual Manag Health Care* 2006;15:83-95.

22. McCusker J, Healey E, Bellavance F, Connolly B. Predictors of repeat emergency department visits by elders. *Acad Emerg Med* 1997;4:581-588.

23. Mackway-Jones K MJ, Windle J. *Emergency triage*. London: BMJ Publishing Group, 2005.

24. Verelst S, Pierloot S, Desruelles D, Gillet JB, Bergs J. Short-term unscheduled return visits of adult patients to the emergency department. *J Emerg Med* 2014;47:131-139.

25. Weiss SJ, Derlet R, Arndahl J, et al. Estimating the degree of emergency department overcrowding in academic medical centers: results of the National ED Overcrowding Study (NEDOCS). *Acad Emerg Med* 2004;11:38-50.

26. van der Linden MC, Lindeboom R, de Haan R, et al. Unscheduled return visits to a Dutch inner-city emergency department. *Int J Emerg Med* 2014;7:23.

27. Pereira L, Choquet C, Perozziello A, et al. Unscheduled-return-visits after an emergency department (ED) attendance and clinical link between both visits in patients aged 75 years and over: a prospective observational study. *PLoS One* 2015;10:e0123803.

28. Wu CL, Wang FT, Chiang YC, et al. Unplanned emergency department revisits within 72 hours to a secondary teaching referral hospital in Taiwan. *J Emerg Med* 2010;38:512-517.

29. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370:1453-1457.

30. Biese KJ, Busby-Whitehead J, Cai J, et al. Telephone Follow-Up for Older Adults Discharged to Home from the Emergency Department: A Pragmatic Randomized Controlled Trial. *J Am Geriatr Soc* 2018;66:452-458.

31. Hardy M, Cho A, Stavig A, et al. Understanding Frequent Emergency Department Use Among Primary Care Patients. *Popul Health Manag* 2018;21:24-31.

32. Smith S, Mitchell C, Bowler S. Standard versus patient-centred asthma education in the emergency department: a randomised study. *Eur Respir J* 2008;31:990-997.

33. Lowthian JA, Lennox A, Curtis A, et al. HOspitals and patients WoRking in Unity (HOW R U?): telephone peer support to improve older patients' quality of life after emergency department discharge in Melbourne, Australia-a multicentre prospective feasibility study. *BMJ Open* 2018;8(6):e020321.
34. Leppin AL, Gionfriddo MR, Kessler M, et al. Preventing 30-day hospital readmissions: a systematic review and meta-analysis of randomized trials. *JAMA Intern Med* 2014;174:1095-1107.
35. Ribbink ME, Macneil-Vroomen JL, van Seben R, Oudejans I, Buurman BM. Investigating the effectiveness of care delivery at an acute geriatric community hospital for older adults in the Netherlands: a protocol for a prospective controlled observational study. *BMJ Open* 2020;10(3):e033802.
36. American College of Emergency Physicians; American Geriatrics Society; Emergency Nurses Association; Society for Academic Emergency Medicine; Geriatric Emergency Department Guidelines Task Force. Geriatric emergency department guidelines. *Ann Emerg Med* 2014;63:e7-e25.
37. Fox MT, Persaud M, Maimets I, et al. Effectiveness of acute geriatric unit care using acute care for elders components: a systematic review and meta-analysis. *J Am Geriatr Soc* 2012;60:2237-2245.
38. Mooijaart SP, Lucke JA, Brabrand M, Conroy S, Nickel CH. Geriatric emergency medicine: time for a new approach on a European level. *Eur J Emerg Med* 2019;26:75-76.
39. Blomaard LC, de Groot B, Lucke JA, et al. Implementation of the acutely presenting older patient (APOP) screening program in routine emergency department care : A before-after study. *Z Gerontol Geriatr* 2021;54:113-121.
40. Hendry A. Creating an Enabling Political Environment for Health and Social Care Integration. *Int J Integr Care* 2016;16:7.
41. Gomes B, Calanzani N, Curiale V, McCrone P, Higginson IJ. Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers. *Cochrane Database Syst Rev* 2013;Cd007760.

Supplementary file 1. Definitions of reasons used to analyze unplanned emergency department return visits (URV) and categorization of the reasons.

Reasons for ED return per category	Definition
<u>Patient-related ED return</u>	
Non-compliance	There is evidence in the medical records that the patient did not follow instructions. The patient returned to the ED for the same problem.
Left without being seen	The patient was registered in the ED but left before being seen by a physician. The patient returned to the ED for the same problem.
Left against medical advice	The patient was seen by a physician and left the ED against medical advice. The patient returned to the ED for the same problem.
Patient was instructed to visit own GP	The patient was instructed to return to the GP for re-evaluation but did not go and returned to the ED instead.
Psychiatric disorder / substance abuse	The patient has a psychiatric disorder and/or uses drugs or alcohol, which causes him/her to repeatedly visit the ED for the same or similar problems. Mentally, the patient is in a chronic stable state.
Worrying about health	The patient's anxiety caused him/her to return to the ED for the same or similar problem. After re-evaluation in the ED, there was no change in diagnosis or treatment and medical management consisted of reassurance only.
Patient refuses admission or treatment	The patient refused the treatment advised by the treating physician during the index ED visit or refused hospital admission. The patient returned to the ED for the same problem.
<u>Illness-related ED return</u>	
Recurrent complaints/disease	The patient was diagnosed and treated appropriately during the index ED visit, with resolution of symptoms, but later returned with a second exacerbation of the disease or with recurrence of the same or similar problem.
Complication	The patient was diagnosed and treated appropriately during the index ED visit, but returned to the ED because of a complication of the disease or side effect of treatment (e.g., allergic drug reaction).
Progression of disease	The medical records reveal that the patient was treated appropriately at the index ED visit and that admission was not indicated. Appropriate follow-up was arranged, but the patient's disease or problem got worse, and he/she returned to the ED as instructed.
Failure of adequate treatment	The patient was diagnosed and treated appropriately during the index ED visit, but the symptoms did not resolve, neither progressed (e.g., persistent pain due to fracture despite adequate use of pain medication). The patient returned to the ED because of persistent complaints.
<u>New complaint</u>	The patient returned to the ED with a new complaint, which was different from the disease or complaint presented at the index ED visit and not determined as a complication or different presentation of the disease, presented during the index ED visit.

<i>Other</i>	
<i>Physician-related ED return</i>	
Treatment error	The physician made the right diagnosis during the index ED visit, but made an error in treatment. The patient returned to the ED for the same or similar problem or progression of disease.
No painkillers prescribed	The disease or injury warranted pain medication but no prescription or advice for the use of pain medication was given. The patient returned primarily because of continued pain.
Misdiagnosis	Medical record review reveals a diagnosis or problem missed by the physician who saw the patient during the index ED visit. The patient returned to the ED for the same problem.
Discharged while admission indication	Medical record review reveals a hospital admission indication, considering the severity of the patient's complaints, but the physician judged that admission was not indicated. The patient returned to the ED because of the severity of the complaints.
<i>System-related ED return</i>	
Not admitted due to lack of hospital capacity	Hospital admission was indicated, but the patient was sent home due to lack of hospital admission capacity. The patient returned to the ED for the same problem.
Not called for follow-up	The patient did not receive a follow-up appointment within the time limit set upon discharge after the index ED visit, due to system-related reasons (e.g., miscommunication, waiting list). The patient returned to the ED for the same problem or progression of disease.
<i>Undefined ED return</i>	The reason for the patient's return cannot be classified in one of the other reasons for an URV.

ED, Emergency Department; *e.g.*, *exempli gratia*; meaning “for example”; *GP*, General Practitioner; *URVs*, Unplanned emergency department return visits