



Universiteit  
Leiden  
The Netherlands

## **The peroxidase activity of cytochrome c-550 from *Paracoccus versutus***

Diederix, R.E.M.; Ubbink, M.; Canters, G.W.

### **Citation**

Diederix, R. E. M., Ubbink, M., & Canters, G. W. (2001). The peroxidase activity of cytochrome c-550 from *Paracoccus versutus*. *European Journal Of Biochemistry*, 268(15), 4207-4216.  
doi:10.1046/j.1432-1327.2001.02335.x

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3618209>

**Note:** To cite this publication please use the final published version (if applicable).

## The peroxidase activity of cytochrome *c*-550 from *Paracoccus versutus*

Rutger E. M. Diederix, Marcellus Ubbink and Gerard W. Canters

Gorlaeus Laboratories, Leiden Institute of Chemistry, Leiden University, the Netherlands

Next to their natural electron transport capacities, *c*-type cytochromes possess low peroxidase and cytochrome P-450 activities in the presence of hydrogen peroxide. These catalytic properties, in combination with their structural robustness and covalently bound cofactor make cytochromes *c* potentially useful peroxidase mimics. This study reports on the peroxidase activity of cytochrome *c*-550 from *Paracoccus versutus* and the loss of this activity in presence of H<sub>2</sub>O<sub>2</sub>. The rate-determining step in the peroxidase reaction of cytochrome *c*-550 is the formation of a reactive intermediate, following binding of peroxide to the haem iron. The reaction rate is very low compared to horseradish peroxidase (approximately one millionth), because of the poor accessibility of the haem iron for H<sub>2</sub>O<sub>2</sub>, and the lack of a base catalyst such as the distal His of the

peroxidases. This is corroborated by the linear dependence of the reaction rate on the peroxide concentration up to at least 1 M H<sub>2</sub>O<sub>2</sub>. Steady-state conversion of a reducing substrate, guaiacol, is preceded by an activation phase, which is ascribed to the build-up of amino-acid radicals on the protein. The inactivation kinetics in the absence of reducing substrate are mono-exponential and shown to be concurrent with haem degradation up to 25 mM H<sub>2</sub>O<sub>2</sub> (pH 8.0). At still higher peroxide concentrations, inactivation kinetics are biphasic, as a result of a remarkable protective effect of H<sub>2</sub>O<sub>2</sub>, involving the formation of superoxide and ferrocycytochrome *c*-550.

**Keywords:** cytochrome *c*; peroxidase; protein radicals; haem; oxidation.

Peroxidases are haem containing enzymes that efficiently catalyse substrate oxidations using hydrogen peroxide [1,2]. Peroxidases can function as catalysts in a variety of oxidation reactions on a broad spectrum of substrates and their potential use is therefore considerable. This is more so because they utilize the 'clean' oxidant H<sub>2</sub>O<sub>2</sub> [2]. Unfortunately, peroxidases are prone to inactivation during normal turnover. This inherent instability is poorly understood and it is important to investigate the mechanism of inactivation because it is currently the main restriction to commercial application of peroxidases and peroxidase mimics [2–4].

Peroxidase activity is inherent to many haem-proteins besides peroxidases. It has been detected in, e.g. haemoglobins and myoglobins, cytochrome *c* and microperoxidases [5–10]. The latter are small peptides derived from extensive proteolysis of cytochrome *c*, which have contained a covalently bound haem moiety [8,11]. In some cases protein modification has resulted in enhanced activity [12–14]. Understanding the peroxidase properties of *c*-type cytochromes is particularly interesting, because these are

very stable proteins that remain highly soluble even under conditions of extreme heat, acidity and basicity. Importantly, the covalent linkage, via thioether bonds, of their haem prosthetic group to the protein matrix prevents dissociation of the catalytic moiety from the protein. These properties render cytochromes *c* excellent candidates to be used as peroxidase mimics.

We therefore set out to understand the peroxidase activity and inactivation kinetics of cytochrome *c*-550 from *Paracoccus versutus*. Such a study is a prerequisite to ultimately improve these properties in cytochrome *c*-550 by utilizing protein engineering and other means. Eventually, such studies might direct the way to how to suppress the H<sub>2</sub>O<sub>2</sub>-driven inactivation that currently hampers the application of peroxidases or peroxidase mimics as 'green chemistry catalysts'. Cytochrome *c*-550, which was selected for this study, is a member of the Class I cytochromes *c*, a class which contains also the archetypal mitochondrial cytochromes *c* and many other, bacterial cytochromes [15]. *P. versutus* cytochrome *c*-550 has a strong homology to the well-studied cytochrome *c*<sub>2</sub> from *Paracoccus denitrificans* [15–17]. *P. versutus* cytochrome *c*-550 has been studied extensively [16,18–22], and we could take advantage of an excellent heterologous expression system, allowing site-directed mutagenesis with high protein yields [22].

The present work shows that cytochrome *c*-550 has peroxidase activity. An activation phase precedes maximal rate of turnover. This activation involves an oxidative process centered on the protein. The catalytic characteristics of cytochrome *c*-550 are similar to those of the microperoxidases, although the reaction rate is more than 1000-fold slower. However, the rate of inactivation of cytochrome *c*-550 by H<sub>2</sub>O<sub>2</sub> is also very low, and the protein stays active at very high peroxide concentrations. In the

Correspondence to M. Ubbink, Gorlaeus Laboratories, Leiden Institute of Chemistry, Leiden University, PO Box 9502, 2300 RA Leiden, the Netherlands. Fax: + 31 71 5274593, Tel.: + 31 71 5274628, E-mail: m.ubbink@chem.leidenuniv.nl

Abbreviations: ABTS, 2,2'-azino-bis (3-ethylbenzo-6-thiazolinesulfonic acid); M100K cytochrome *c*-550, site-directed mutant of *P. versutus* cytochrome *c*-550 in which the iron co-ordinating methionine is replaced by lysine;  $K_m^{per}$ , apparent Michaelis constant for hydrogen peroxide; MP-8, microperoxidase-8.

Note: a website is available at <http://wwwchem.leidenuniv.nl/metprot> (Received 2 April 2001, revised 1 June 2001, accepted 4 June 2001)

absence of reductants the inactivation of the protein obeys first-order kinetics, except at high concentrations of peroxide where the inactivation is biphasic.

## EXPERIMENTAL PROCEDURES

### Expression and purification of cytochrome *c*-550

Cytochrome *c*-550 was heterologously expressed in *P. denitrificans* strain 2131 containing the pEG400.Tv1 plasmid [22]. Typically, 3 L of culture was prepared for expression of cytochrome *c*-550. The growth medium was BHI (GibcoBRL), supplemented with 50  $\mu\text{g}\cdot\text{mL}^{-1}$  streptomycin. The cells were cultured in 2 L conical flasks (500 mL culture per flask) and shaken at 300 r.p.m. After 24 h of growth at 30 °C, the  $D_{660}$  was measured, and the cells were harvested by centrifugation at 11 300 g. The cell pellet was resuspended to a calculated  $D_{660}$  of 200 in an ice-cold solution of 1 M sucrose, 1 mM EDTA, 50 mM Tris, pH 8.0 (Tes). Solid lysozyme (50 mg, Sigma) was added and the cell suspension was stirred for 20 min at room temperature and centrifuged. The cell pellet was resuspended and this procedure was repeated once. The pellet was resuspended again in ice-cold Milli Q water, using four times the volume used with Tes. After stirring for 20 min, the resultant protoplasts were centrifuged at 27 200 g. The supernatants containing cytochrome *c*-550 were pooled. Further purification was performed as described [16]. The average yield of pure protein was 12  $\text{mg}\cdot\text{L}^{-1}$  culture. The protein was judged to be more than 95% pure by its UV-visible spectrum and by SDS/PAGE. The spectra of both ferrous and ferric cytochrome *c*-550 were identical to those previously published [18].

### Activity assays

The assays were typically performed in 1.5 mL plastic cuvettes. The reactions were started by the addition of  $\text{H}_2\text{O}_2$  (Merck). The  $\text{H}_2\text{O}_2$  stock solution was freshly made and its concentration was verified spectrophotometrically ( $\epsilon_{240} = 39.4 \text{ M}^{-1}\cdot\text{cm}^{-1}$  [23]). The hydrogen donor was guaiacol (i.e. 2-methoxyphenol, Sigma), diluted from a fresh aqueous stock solution (100 mM). It was assumed that the coloured product, tetraguaiacol, is the result of four one-electron oxidations [24]. The formation of tetraguaiacol ( $\epsilon_{470} = 26.6 \text{ mM}^{-1}\cdot\text{cm}^{-1}$  [25]) was followed with a Shimadzu UVPC-200 spectrophotometer fitted with a thermostat. The reactions were performed at 25 °C in 100 mM sodium phosphate buffer, pH 8.0 (unless stated otherwise). In some cases, the reaction was monitored by using an Applied Photophysics SX.18 MV stopped-flow apparatus. Other reagents used were mannitol (Brocacef, the Netherlands), superoxide dismutase, horse heart cytochrome *c*, ABTS (all from Sigma), and M100K cytochrome *c*-550 (kindly provided by Mrs Ing. Gertrud Warmerdam). In a number of experiments, directed at investigating the activation process preceding the steady-state reaction, a mixture was employed containing 10 mM  $\text{H}_2\text{O}_2$  and 5  $\mu\text{M}$  cytochrome *c*-550. After a fixed amount of time the  $\text{H}_2\text{O}_2$  was removed by addition of  $\approx 1200$  units of buffer-equilibrated catalase attached to 4% cross-linked beaded agarose (Sigma) to 1 mL of reaction mixture. The catalase was removed after about 2 min by filtration

(0.2  $\mu\text{m}$  Acrodisc, Gelman Sciences). Activity was then measured by addition of  $\text{H}_2\text{O}_2$ /guaiacol and the UV/visible spectrum recorded. Controls were taken without  $\text{H}_2\text{O}_2$ , cytochrome *c*-550 or both, respectively.

### Inactivation assays

Cytochrome *c*-550 was preincubated with  $\text{H}_2\text{O}_2$  for a recorded time. The activity remaining after this time was measured by adding  $\text{H}_2\text{O}_2$ /guaiacol. The concentrations of reactants in the preincubation mix were varied, but in the end-mix the conditions were 10 mM guaiacol, and either 50 or 100 mM  $\text{H}_2\text{O}_2$ . The spectral changes of cytochrome *c*-550 upon addition of  $\text{H}_2\text{O}_2$  in the absence of guaiacol were monitored by using the stopped-flow apparatus with a photodiode array (320–1000 nm).

## RESULTS


### Peroxidase activity

When cytochrome *c*-550, guaiacol and  $\text{H}_2\text{O}_2$  were mixed together, the orange-coloured tetraguaiacol was formed. When cytochrome *c*-550 or  $\text{H}_2\text{O}_2$  was left out, no formation of tetraguaiacol was observed. The rate of production of tetraguaiacol depends linearly on the cytochrome *c*-550 concentration (0.1–10  $\mu\text{M}$  cytochrome *c*-550). When mannitol (100 mM) or when superoxide dismutase was added, no change in rate was observed. Therefore hydroxyl or superoxide radicals, respectively, are not involved in the reaction, thus excluding Haber–Weiss or Fenton chemistry. The activity was negligible after a vigorous cytochrome *c*-550 inactivation procedure, involving a long (> 60 min) preincubation with  $\text{H}_2\text{O}_2$ .

Although guaiacol can reduce ferricytochrome *c*-550, and thus is oxidized in the absence of  $\text{H}_2\text{O}_2$ , the peroxidase activity of cytochrome *c*-550 is not due to a 'pseudo-peroxidase' cycle where cytochrome *c*-550 cycles between its  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$  oxidation states. This is concluded from the observation that, when the reaction is in steady state, ferrocycytochrome *c*-550 is not observed, while the reaction between cytochrome *c*-550 and  $\text{H}_2\text{O}_2$  is the rate-limiting step in the oxidation of guaiacol (as will be shown later). This means that ferric cytochrome *c*-550 and not ferrous cytochrome *c*-550 is the relevant reaction partner with  $\text{H}_2\text{O}_2$ . In addition, when ABTS was used as the reducing substrate, peroxidase activity is also observed (ABTS does not reduce ferricytochrome *c*-550). Finally, peroxidase activity was also observed for M100K cytochrome *c*-550. The redox potential of M100K cytochrome *c*-550 is 329 mV lower (298 K, pH 7.0) than the wild-type cytochrome *c*-550 value [20], and it is not reduced by guaiacol.

### The peroxidase assay

A typical peroxidase assay with cytochrome *c*-550 and guaiacol gives a product formation curve with four phases, as depicted in Fig. 1. An initial activation phase (I in Fig. 1) is followed by a steady-state phase, represented by the straight part of the curve (II). Then the curve levels off (III) and ultimately a decrease in absorption is seen (IV). The last two features can be ascribed to the inherent instability of the reaction product (tetraguaiacol), in combination with




**Fig. 1.** A typical tetra-guaiacol formation curve. The four phases (I–IV) are described in the text. Conditions were 0.2  $\mu$ M cytochrome *c*-550, 50 mM  $\text{H}_2\text{O}_2$  and 20 mM guaiacol.

catalyst inactivation [7,25,26]. The rate of the steady-state reaction was determined by taking the maximum of the first derivative of the product formation curve. The length of the steady-state phase (II), as judged from the width of the first derivative maximum, depends strongly on conditions such


as  $[\text{H}_2\text{O}_2]$  and pH. Ideally, the length of the steady-state phase is  $> 10$  s, but frequently it was shorter.

### The activation phase

The presence of a lag period (phase I) suggests that cytochrome *c*-550 is activated by  $\text{H}_2\text{O}_2$ . The length of this activation phase was assessed in terms of an activation rate constant,  $k_{\text{act}}$ , which was obtained by fitting the first derivative of the product formation curve to a single exponential (Fig. 2A,B). The length of the activation phase depends on the concentration of  $\text{H}_2\text{O}_2$  (Fig. 2A). The activation phase is also shortened by increased concentrations of cytochrome *c*-550 (Fig. 2B). When cytochrome *c*-550 is preincubated for 30 s with  $\text{H}_2\text{O}_2$  before addition of guaiacol, the activation phase is absent (Fig. 3A). When an aliquot is removed at the end of the activation phase and transferred to a fresh mixture of substrates, no activation phase is seen. When the  $\text{H}_2\text{O}_2$  is removed after a 30-s preincubation time by addition of agarose-immobilized catalase, the protein exhibits a diminished activation phase (Fig. 3B), and remains activated for at least 30 min. The UV/visible spectrum of the activated form is unchanged



**Fig. 2.** Rate of activation of cytochrome *c*-550 as a function of (A)  $\text{H}_2\text{O}_2$  and (B) cytochrome *c*-550. The value of the activation rate constant,  $k_{\text{act}}$ , was obtained by fitting the first derivative of the product formation curve to a single exponential. Conditions: (A) 25 mM guaiacol and 1  $\mu$ M cytochrome *c*-550 and (B) 25 mM guaiacol 5 mM  $\text{H}_2\text{O}_2$ , cytc550, cytochrome *c*-550.



**Fig. 3.** Tetra-guaiacol formation without (solid line) and with (dotted line) a 30-s preincubation with 10 mM  $\text{H}_2\text{O}_2$  (A) and tetra-guaiacol formation after a 30-s preincubation in absence of  $\text{H}_2\text{O}_2$  (solid line) and in the presence of 10 mM  $\text{H}_2\text{O}_2$  (dotted line) after which agarose-immobilized catalase was added (B). (A) Activity was assayed at 10 mM  $\text{H}_2\text{O}_2$ , 10 mM guaiacol and  $< 4$   $\mu$ M cytochrome *c*-550. (B) Conditions of the assay after removal of the catalase were 5 mM  $\text{H}_2\text{O}_2$ , 25 mM guaiacol and  $< 1$   $\mu$ M cytochrome *c*-550.

with respect to native ferricytochrome *c*-550. The activation appears to be partly reversible (diminished length of the activation phase) and does not originate from changes in the haem-environment or iron co-ordination, because the UV-visible spectrum is normally sensitive to such changes. The peroxidase activity of M100K cytochrome *c*-550, in which the axial Met is replaced by Lys [20], exhibits an activation phase just like the wild-type cytochrome *c*-550, corroborating the conclusion that axial co-ordination plays no role in the activation process.

### Dependence on guaiacol and hydrogen peroxide

When the concentration of guaiacol is between 15 and 30 mM, the rate of the steady-state peroxidase reaction (phase II in Fig. 1) of cytochrome *c*-550 depends linearly on the concentration of H<sub>2</sub>O<sub>2</sub>. This linear dependence holds even up to 1 M H<sub>2</sub>O<sub>2</sub> (Fig. 4). The rate also depends linearly on the concentration of cytochrome *c*-550, and therefore the oxidation rate of guaiacol due to the peroxidase activity of cytochrome *c*-550 follows the bimolecular rate law [Eqn (1)]:

$$v = k_{\text{obs}}[\text{cytochrome } c\text{-550}][\text{H}_2\text{O}_2] \quad (1)$$

The value of the bimolecular rate constant  $k_{\text{obs}}$  is  $43.4 \pm 0.8 \text{ M}^{-1} \cdot \text{s}^{-1}$  (25 mM guaiacol, 100 mM sodium phosphate, pH 8.0, 25 °C). The H<sub>2</sub>O<sub>2</sub> dependence of the peroxidase activity of horse heart cytochrome *c* was also probed. Similar to cytochrome *c*-550, no saturation is seen up to 1 M H<sub>2</sub>O<sub>2</sub> (Fig. 4). Its bimolecular rate constant  $k_{\text{obs}}$  is  $4.1 \pm 0.1 \text{ M}^{-1} \cdot \text{s}^{-1}$  (25 mM guaiacol, 100 mM sodium phosphate, pH 8.0). The complete absence of saturation by H<sub>2</sub>O<sub>2</sub> is contrary to earlier reports where  $K_{\text{m}}^{\text{per}}$  values varying from 25 to 65 mM have been reported for horse heart cytochrome *c* [27–30]. However, in our hands it was not possible to reproduce these results, and a linear dependence of the rate on the concentration of H<sub>2</sub>O<sub>2</sub> was found with both guaiacol (Fig. 4) and ABTS as reducing substrates (not shown).

The guaiacol concentration has a limited effect on the rate, even down to 50 μM guaiacol. The steady-state




Fig. 4. Dependence of the steady-state peroxidase reaction rate as a function of H<sub>2</sub>O<sub>2</sub> for cytochrome *c*-550 (○) and horse heart cytochrome *c* (●). Conditions: 5 μM cytochrome *c* and 25 mM guaiacol.




Fig. 5. Inverse slope ( $V_{\text{max}}/K_{\text{m}}^{\text{per}}$ ) of plots of  $1/\text{rate}$  vs.  $1/[\text{H}_2\text{O}_2]$  as a function of [guaiacol]. [Cytochrome *c*-550] was 1 μM.

reaction rate is reduced however, by increasing concentrations of guaiacol when the H<sub>2</sub>O<sub>2</sub> concentration is low. This can be attributed to substrate inhibition by guaiacol, and in fact serves as evidence for a ping-pong mechanism [31]. When the dependence of the steady-state reaction rate on [H<sub>2</sub>O<sub>2</sub>] is plotted in a double reciprocal plot ( $1/v$  vs.  $1/[\text{H}_2\text{O}_2]$ ) at various guaiacol concentrations, parallel lines are seen above 15 mM guaiacol (not shown). Such parallel lines again are evidence for a ping-pong mechanism, in which the association with H<sub>2</sub>O<sub>2</sub> is rate limiting [31]. The inverse slope of these parallel lines represents  $V_{\text{max}}/K_{\text{m}}^{\text{per}}$  in Michaelis–Menten formalism (which reduces to  $k_{\text{obs}}[\text{cytochrome } c\text{-550}]$  for  $K_{\text{m}}^{\text{per}} \gg [\text{H}_2\text{O}_2]$ ). The thus calculated values of  $V_{\text{max}}/K_{\text{m}}^{\text{per}}$  are plotted against guaiacol concentration in Fig. 5. This shows that the value of  $V_{\text{max}}/K_{\text{m}}^{\text{per}}$ , or  $k_{\text{obs}}[\text{cytochrome } c\text{-550}]$ , is constant above 15 mM guaiacol. The reduction of this value at lower guaiacol concentrations can be attributed to catalyst inactivation. When reducing substrate is lacking, inactivation can become more prevalent for peroxidases and peroxidase models [26,27,32]. This would pose no problem if true initial rates were measured, but in our assay initial rates can not be measured because cytochrome *c*-550 exhibits an activation phase in its peroxidase activity (see above). Thus the steady-state rate is measured after a certain time, and when the concentration of guaiacol is low, relatively more cytochrome *c*-550 is inactivated at the moment the rate is measured.

### pH dependence of the peroxidase activity

As shown in Fig. 6, the peroxidase activity of cytochrome *c*-550 depends strongly on pH. The steady-state rate increases with increasing pH up to pH 10.0, after which a decrease is seen. Horse heart cytochrome *c* exhibits a stable rate value from pH 5.0–7.0, after which a decline in activity is seen (Fig. 6). This agrees well with earlier observations [27].

### Inactivation of cytochrome *c*-550 by hydrogen peroxide

The rate of inactivation of cytochrome *c*-550 in the absence of guaiacol was determined as described in Experimental procedures. Two types of inactivation behaviour can be distinguished, depending on the concentration of H<sub>2</sub>O<sub>2</sub>.




Fig. 6. Dependence of the steady-state peroxidase reaction rate as a function of pH for cytochrome *c*-550 (○) and horse heart cytochrome *c* (●). Conditions: 1  $\mu$ M cytochrome *c*, 10 mM H<sub>2</sub>O<sub>2</sub> and 25 mM guaiacol.

When preincubated with up to about 25 mM H<sub>2</sub>O<sub>2</sub>, the peroxidase activity of cytochrome *c*-550 decreases in time as a single exponential (Fig. 7, open circles). In addition, the UV-visible spectrum of cytochrome *c*-550 bleaches when exposed to up to 25 mM H<sub>2</sub>O<sub>2</sub>. The loss of the Soret band intensity exactly matches the decrease in activity (Fig. 7, solid line), and both can be fitted to the same single exponential function (not shown). The inactivation rate does not depend on the concentration of cytochrome *c*-550 between 0.5 and 5  $\mu$ M. Up to 25 mM H<sub>2</sub>O<sub>2</sub>, the value of the exponential rate constant of inactivation ( $k_i$ ) depends linearly on the H<sub>2</sub>O<sub>2</sub> concentration. Note that with short preincubation times, the activity is somewhat increased. During this short preincubation, inactivation has relatively little effect and nearly all protein is activated at the start of the activity assay (no activation phase). In the absence of a




Fig. 7. The normalized absorption change at 410 nm in the presence of 20 mM H<sub>2</sub>O<sub>2</sub> as a function of time (solid line) and the normalized steady-state peroxidase reaction rate of 0.5  $\mu$ M cytochrome *c* as a function of preincubation time with 20 mM H<sub>2</sub>O<sub>2</sub> (open circles). Normalization of the peroxidase reaction rate data was performed by taking the amplitude of a first-order exponential fit of the data into account. Activity was assayed with 10 mM H<sub>2</sub>O<sub>2</sub>, 10 mM guaiacol and <0.25  $\mu$ M cytochrome *c*-550.




Fig. 8. Rate of the steady-state peroxidase reaction as a function of the preincubation time with H<sub>2</sub>O<sub>2</sub>. In (A), [cytochrome *c*-550] was 1  $\mu$ M in the preincubation mix and [H<sub>2</sub>O<sub>2</sub>] was varied (○, 50 mM; ●, 100 mM; □, 200 mM). In B, [H<sub>2</sub>O<sub>2</sub>] was 50 mM in the preincubation mix and [cytochrome *c*-550] was varied (○, 10  $\mu$ M; ●, 5  $\mu$ M; □, 2.5  $\mu$ M; ■, 1  $\mu$ M; △, 0.6  $\mu$ M). Activity was assayed in 10 mM guaiacol, 50 mM H<sub>2</sub>O<sub>2</sub>, and <0.5  $\mu$ M cytochrome *c*-550. cytc550, cytochrome *c*-550.

preincubation, the protein is being activated during the activation phase but simultaneously the activated form is reduced by the substrate. In the latter case, not all the protein will become activated, but rather an equilibrium is established. This is why the activity seems to increase slightly at short incubation times.

Above 50 mM H<sub>2</sub>O<sub>2</sub>, the inactivation behavior is markedly changed; the activity decreases in a biphasic fashion. A fast exponential phase is followed by a slower exponential phase (Fig. 8A,B). At even higher H<sub>2</sub>O<sub>2</sub> concentrations, the second phase is reached sooner, and the rate constants of both phases are lower (Fig. 8A). The same effect is seen when the concentration of cytochrome *c*-550 is lowered in the premix (Fig. 8B). So, paradoxically, the more peroxide and the less protein present, the higher the residual activity is after a certain period of incubation. Control measurements in the presence of mannitol show that hydroxyl radicals hardly affect the inactivation. However, when superoxide dismutase is added the overall inactivation is increased (Fig. 9A). Thus, superoxide is formed when cytochrome *c*-550 is incubated with high [H<sub>2</sub>O<sub>2</sub>]. Superoxide is a potent reductant of cytochrome *c*-550 [33] and may be responsible for the formation of ferrocycytochrome *c*-550. By using fast



As in natural peroxidases, the reducing substrate might be expected to react with the (high-valent) haem. However, considering that protein radicals are so easily found in peroxidase mimics such as myoglobin and cytochrome *c* (see above), it is possible that guaiacol is oxidized via one of these radicals. Unstabilized protein radicals can be transferred easily and rapidly between different sites on a protein [51–54], and thus also to the two surface exposed tyrosines of cytochrome *c*-550 [16,17], which might be involved in guaiacol oxidation. Studies on the myoglobin/ $\text{H}_2\text{O}_2$  catalysed oxygenation and oxidation reactions provide evidence that a protein radical is indeed the true oxidizing species in such types of reactions [54–61].

### The rate-determining step in the peroxidase reaction of cytochrome *c*-550

The rate-determining step in the peroxidase activity of cytochrome *c*-550 is the formation of a high-potential intermediate. This rate-determining step is characterized by ping-pong kinetics with a high apparent Michaelis constant for  $\text{H}_2\text{O}_2$ , allowing a description of the rate in terms of a bimolecular rate-law (Eqn 1). In this respect cytochrome *c*-550 is similar to microperoxidase-8 (MP-8) [24], even though the bimolecular rate constant  $k_{\text{obs}}$  is approximately 1000-fold larger for MP-8 than for cytochrome *c*-550 [24] (this work). The latter observation seems to reflect the larger accessibility of the haem iron in MP-8 (which is fully exposed to the solvent) [8,37–39], compatible with the observed lower  $K_{\text{m}}^{\text{per}}$  for MP-8 [62].

### The pH dependence of the peroxidase activity of cytochrome *c*-550 and horse heart cytochrome *c*

Cytochrome *c*-550 also resembles MP-8 in the pH dependence of its peroxidase activity. The pH dependence on the activity of horse heart cytochrome *c* and cytochrome *c*-550 is shown in Fig. 6. The curve of the latter compares well with that of MP-8: a steady increase in activity with pH leading to a maximum, and a drop at even higher pH [9,24]. Metal-replacement studies showed that the pH dependence of MP-8 could be explained by assuming that the  $\text{pK}_{\text{a}}$  of  $\text{H}_2\text{O}_2$  is lowered by the metal [9]. The drop in rate with MP-8 at more alkaline pH has been attributed to product instability and ligation of hydroxide to the iron [8,9,24], although the latter has been disputed [63]. A similar reasoning can be used to explain the pH dependence of the rate of cytochrome *c*-550: its increase with pH is related to the  $\text{pK}_{\text{a}}$  of bound  $\text{H}_2\text{O}_2$ . The drop in activity above pH 10.0 coincides well with the so-called alkaline transition for cytochrome *c*-550 ( $\text{pK}_{\text{a}}$  11.2 [19]), which involves exchange of the Met ligand for a Lys [64,65]. It is to be expected that the activity decreases by such a transition, because Lys is a much stronger ligand for iron than Met and thus is replaced much less easily by the incoming peroxide. The combined and counteracting effects when the pH is increased can thus account for the pH profile of the activity of cytochrome *c*-550 (Fig. 6). This could also be the case with the pH dependence on the activity of horse heart cytochrome *c* (also shown in Fig. 6), although the apparent  $\text{pK}_{\text{a}}$  observed in the activity profile for this protein is slightly lower than its alkaline transition determined by alternative methods ( $\text{pK}_{\text{a}}$  8.9–9.5 [64]). It is

important to note that, as with MP-8 [9], the protein matrix of cytochrome *c*-550 does not participate in lowering the  $\text{pK}_{\text{a}}$  of  $\text{H}_2\text{O}_2$ . Such a function has been shown to be the basis for the high reaction rates for the natural peroxidases [1].

### The activation phase in the peroxidase reaction of cytochrome *c*-550

The steady-state oxidation of guaiacol by cytochrome *c*-550/ $\text{H}_2\text{O}_2$  is preceded by an activation phase, meaning that the protein is activated in the course of the activity assay. As discussed in the Results section, this is due to a process that does not involve changes in the haem environment or iron co-ordination. Increasing the concentration of  $\text{H}_2\text{O}_2$  accelerates this process, indicating an oxidative mechanism. In principle, diffusion of  $\text{H}_2\text{O}_2$  into the protein and binding to the iron could be the reason for the activation phase. However, the activation occurs within seconds but the protein remains activated for at least 30 min after removal of all  $\text{H}_2\text{O}_2$ , suggesting that peroxide diffusion and binding cannot explain the activation phase.

It was pointed out earlier that protein radicals can accumulate on cytochrome *c* when peroxides are present. Moreover, it was suggested that protein radicals might be the true reaction partners with guaiacol. The activation process may thus represent a build up of protein radicals on the cytochrome *c*-550 molecule. Cytochrome *c*-550 contains three tyrosines and two tryptophans, the most common sites of protein radicals [54,66]. Under assay conditions, they thus compete with guaiacol for oxidizing equivalents. A number of studies have shown that, for natural peroxidases, when two substrates are present at the same time, one is preferably oxidized, and the second substrate may remain unreacted until the first substrate is completely consumed [67]. Such a competition might also be the case here between guaiacol and the aromatic residues: once the aromatic residues are 'consumed', oxidation of guaiacol proceeds via these radicals or via the (high-valent) haem.

Attempts to find support for the presence of protein radicals by EPR detection, as applied to horse heart cytochrome *c* [36], failed. Addition of 5- to 10-fold molar excess of  $\text{H}_2\text{O}_2$  to cytochrome *c*-550 leads to reduction of the protein, which is probably related to the high protein concentration (100–500  $\mu\text{M}$ ) used in these experiments, necessary for EPR detection.

An alternative explanation for the observed activation phase may be that the oxidation of one of the aromatic amino acids results in structural changes, which facilitate the peroxidase reaction. In the case of F41W horseradish peroxidase, which also exhibits an activation phase, oxidation of Trp41 was suggested to be responsible for such a mechanism [68].

### Inactivation of the peroxidase activity of cytochrome *c*-550 by hydrogen peroxide

As mentioned in the Results section, up to 25 mM  $\text{H}_2\text{O}_2$  the inactivation of the peroxidase activity follows mono-exponential kinetics and the inactivation rate depends linearly on the concentration of  $\text{H}_2\text{O}_2$ . The rate of inactivation does not depend on the concentration of cytochrome *c*-550. The decrease in activity occurs simultaneously with

bleaching of the UV/visible spectrum (see Fig. 7). Bleaching is common when peroxides are added to haem (proteins), and is ascribed to an oxidative attack at the haem periphery, causing ring opening, loss of iron and further degradation [69–72]. The initial degradation step might be an intramolecular attack on the haem by an activated oxygen species such as oxyferryl [73] or iron-peroxo, as seen with haem oxygenase [74]. In any case, the bleaching is an intramolecular process, in accord with the observation that it is independent of the concentration of cytochrome *c*-550. Formation of the activated oxygen species depends linearly on the concentration of H<sub>2</sub>O<sub>2</sub>. Therefore it is expected that the inactivation rate also depends linearly on the concentration of H<sub>2</sub>O<sub>2</sub>, and as mentioned above, this agrees with the observation that up to 25 mM H<sub>2</sub>O<sub>2</sub>, the rate of inactivation equals  $k_i[\text{H}_2\text{O}_2]$  with  $k_i = 0.51 \text{ M}^{-1}\cdot\text{s}^{-1}$ .

At higher peroxide concentrations (> 50 mM H<sub>2</sub>O<sub>2</sub>) the inactivation characteristics of cytochrome *c*-550 are very different. The higher the concentration of H<sub>2</sub>O<sub>2</sub>, and the lower the concentration of cytochrome *c*-550, the slower the inactivation. Apparently H<sub>2</sub>O<sub>2</sub> can protect the protein when in excess. It is interesting to see that in excess H<sub>2</sub>O<sub>2</sub>, ferrous cytochrome *c*-550 is readily formed (cf [75]), but that when superoxide dismutase is present this is much less (Fig. 9B,C). The latter indicates that superoxide is involved in this process. Superoxide is a potent reductant of cytochrome *c* [33], and the observation that superoxide dismutase inhibits the formation of ferrocycytochrome *c*-550 suggests that O<sub>2</sub><sup>•−</sup> is formed at high concentrations H<sub>2</sub>O<sub>2</sub>, and that this leads to the formation of ferrous cytochrome *c*-550. The appearance of ferrous cytochrome *c*-550 at excess H<sub>2</sub>O<sub>2</sub> is important, because ferrous cytochrome *c* is less susceptible to H<sub>2</sub>O<sub>2</sub>-driven inactivation [76] or protects against it. It can thus be expected that when more cytochrome *c*-550 is in the ferrous state, the inactivation rate is less. Interesting in this respect is, that ferric cytochrome *c* catalyses its own formation from ferrous cytochrome *c* in the presence of H<sub>2</sub>O<sub>2</sub> [76], which explains why cytochrome *c*-550 remains active longer when less protein is present (Fig. 8B).

## CONCLUSION

Despite being a classic electron transport protein, cytochrome *c*-550 clearly possesses peroxidase activity. This activity is characterized by a poor association with H<sub>2</sub>O<sub>2</sub>. The protein matrix of cytochrome *c*-550 does not assist the reaction by lowering the  $pK_a$  of H<sub>2</sub>O<sub>2</sub> unlike the natural peroxidases. Thus the effective concentration of oxidant is much lower for cytochrome *c*-550 than for natural peroxidases, which, in part, is responsible for the relatively low activity of cytochrome *c*-550 versus, for example, horseradish peroxidase. The activity is further lowered by limited accessibility of the haem-iron, as evidenced by the 1000-fold higher peroxidase activity of the highly accessible microperoxidases. Therefore, in order to create an artificial peroxidase out of cytochrome *c*-550, it is necessary to address the following two points: (a) the lack of a properly positioned general base catalyst to deprotonate the incoming H<sub>2</sub>O<sub>2</sub> and (b) the low accessibility of the active site to H<sub>2</sub>O<sub>2</sub>. These topics are currently under investigation.

Interestingly, the studies on the inactivation kinetics of cytochrome *c*-550 at high and low H<sub>2</sub>O<sub>2</sub> concentrations unveiled a complex accumulation of side-reactions in the absence of reducing substrate. Paradoxically, higher peroxide concentrations and lower protein concentrations increasingly protect the haem from degradation. This relative stability of cytochrome *c*-550 under such harsh conditions is remarkable, and this observation helps to identify the major determinants for peroxide-driven inactivation of haem enzymes.

## ACKNOWLEDGEMENT

This work was supported by the Foundation for Chemical Research (SON) and the Foundation for Technical Sciences (STW) with financial aid from the Netherlands Organization for Scientific Research (NWO), and under the auspices of the BIOMAC Graduate Research School of Leiden and Delft.

## REFERENCES

- Dunford, H.B. (1991) Horseradish peroxidase: structure and kinetic properties. In *Peroxidases in Chemistry and Biology* (Everse, J., Everse, K.E. & Grisham, M.B., eds), Vol. II, pp. 1–24. CRC Press, Boca Raton, FL.
- van Deurzen, M.P.J., van Rantwijk, F. & Sheldon, R.A. (1997) Selective oxidations catalyzed by peroxidases. *Tetrahedron* **53**, 13183–13220.
- Cherry, J., Lamsa, M.H., Schneider, P., Vind, J., Svendsen, A., Jones, A. & Pedersen, A.H. (1999) Directed evolution of a fungal peroxidase. *Nat. Biotechnol.* **17**, 379–384.
- Sheldon, R.A. (1999) Enzymes: picking a winner. *Nature* **399**, 636–637.
- Everse, J., Johnson, M.C. & Marini, M.A. (1994) Peroxidative activities of hemoglobin and hemoglobin derivatives. *Methods Enzymol.* **231**, 547–561.
- Giulivi, C. & Cadenas, E. (1994) Ferrylmyoglobin: formation and chemical reactivity toward electron-donating compounds. *Methods Enzymol.* **233**, 189–202.
- Vazquez-Duhalt, R. (1999) Cytochrome *c* as a biocatalyst. *J. Mol. Catal. B: Enzym.* **7**, 241–249.
- Adams, P.A., Baldwin, D.A. & Marques, H.M. (1995) The hemeptides from cytochrome *c*: preparation, physical and chemical properties, and their use as model compounds for the hemoproteins. In *Cytochrome c. A Multidisciplinary Approach* (Scott, R.A. & Mauk, A.G., eds), pp. 635–692. University Science Books, Sausalito, CA.
- Primus, J.-L., Boersma, M.G., Mandon, D., Boeren, S., Veeger, C., Weiss, R. & Rietjens, I.M.C.M. (1999) The effect of iron to manganese substitution on microperoxidase 8 catalysed peroxidase and cytochrome P450 type of catalysis. *J. Biol. Inorg. Chem.* **4**, 274–283.
- Ortiz de Montellano, P.R. (1987) Control of the catalytic activity of prosthetic heme by the structure of hemoproteins. *Acc. Chem. Res.* **20**, 289–294.
- Paléus, S., Ehrenfest, A. & Tuppy, H. (1955) Study of a peptide degradation product of cytochrome *c*. II. Investigation of the linkage between peptide moiety prosthetic group. *Acta Chem. Scand.* **9**, 365–374.
- Ozaki, S.-I., Matsui, S., Roach, M.P. & Watanabe, Y. (2000) Rational molecular design of a catalytic site: engineering of catalytic functions to the myoglobin active site framework. *Coord. Chem. Rev.* **198**, 39–59.
- Wan, L., Twitchett, M.B., Eltis, L.D., Mauk, A.G. & Smith, M. (1998) *In vitro* evolution of horse heart myoglobin to increase peroxidase activity. *Proc. Natl Acad. Sci. USA* **95**, 12825–12831.

14. Hayashi, T., Hitomi, Y., Ando, T., Mizutani, T., Hisaeda, Y., Kitagawa, S., & Ogoshi, H. (1999) Peroxidase activity of myoglobin is enhanced by chemical mutation of heme-propionates. *J. Am. Chem. Soc.* **121**, 7747–7750.
15. Moore, G.R. & Pettigrew, G.W. (1990) *Cytochromes c, Evolutionary, Structural and Physicochemical Aspects*. Springer-Verlag, Berlin, Germany.
16. Ubbink, M., van Beeumen, J. & Canters, G.W. (1992) Cytochrome c<sub>550</sub> from *Thiobacillus versutus*: cloning, expression in *Escherichia coli*, and purification of the heterologous holoprotein. *J. Bacteriol.* **174**, 3707–3714.
17. Benning, M.M., Meyer, T.E. & Holden, H.M. (1994) X-Ray structure of the cytochrome c<sub>2</sub> isolated from *Paracoccus denitrificans* refined to 1.7-Å resolution. *Arch. Biochem. Biophys.* **310**, 460–466.
18. Lommen, A., Ratsma, A., Bijlsma, N., Canters, G.W., van Wielink, J.E., Frank, J. & van Beeumen, J. (1990) Isolation and characterization of cytochrome c<sub>550</sub> from the methylamine-oxidizing electron-transport chain of *Thiobacillus versutus*. *Eur. J. Biochem.* **192**, 653–661.
19. Ubbink, M. & Canters, G.W. (1993) Mutagenesis of the conserved lysine 14 of cytochrome c-550 from *Thiobacillus versutus* affects the protein structure and the electron self-exchange rate. *Biochemistry* **32**, 13893–13901.
20. Ubbink, M., Campos, A.P., Teixeira, M., Hunt, N.I., Hill, H.A.O. & Canters, G.W. (1994) Characterization of mutant Met100Lys of cytochrome c-550 from *Thiobacillus versutus* with lysine-histidine heme ligation. *Biochemistry* **33**, 10051–10059.
21. Ubbink, M., Hunt, N.I., Hill, H.A.O. & Canters, G.W. (1994) Kinetics of the reduction of wild-type and mutant cytochrome c-550 by methylamine dehydrogenase and amicyanin from *Thiobacillus versutus*. *Eur. J. Biochem.* **222**, 561–571.
22. Ubbink, M., Pfuhl, M., van der Oost, J., Berg, A. & Canters, G.W. (1996) NMR assignments and relaxation studies of *Thiobacillus versutus* ferrocyclochrome c-550 indicate the presence of a highly mobile 13-residues long C-terminal tail. *Protein Sci.* **5**, 2494–2505.
23. Nelson, D.P. & Kiesow, L.A. (1972) Enthalpy of decomposition of hydrogen peroxide by catalase at 25 °C (with molar extinction coefficients of H<sub>2</sub>O<sub>2</sub> solutions in the UV). *Anal. Biochem.* **49**, 474–478.
24. Baldwin, D.A., Marques, H.M. & Pratt, J.M. (1987) Hemes and hemoproteins. 5: Kinetics of the peroxidatic activity of microperoxidase-8: model for the peroxidase enzymes. *J. Inorg. Biochem.* **30**, 203–217.
25. Maehly, A.C. & Chance, B. (1954) The assay of catalases and peroxidases. In *Methods of Biochemical Analysis* (Glick, D., ed.), Vol. 1, pp. 357–408. Interscience, New York.
26. Arnao, M.B., Acosta, M., del Río, J.A., Varón, R. & García-Cánovas, F. (1990) A kinetic study on the suicide inactivation of peroxidase by hydrogen peroxide. *Biochim. Biophys. Acta* **1041**, 43–47.
27. Radi, R., Thomson, L., Rubbo, H. & Prodanov, E. (1991) Cytochrome c-catalyzed oxidation of organic molecules by hydrogen peroxide. *Arch. Biochem. Biophys.* **288**, 112–117.
28. Hamachi, I., Fujita, A. & Kunitake, T. (1994) Enhanced N-demethylase activity of cytochrome c bound to a phosphate-bearing synthetic bilayer membrane. *J. Am. Chem. Soc.* **116**, 8811–8812.
29. Rosei, M.A., Blarmino, C., Coccia, R., Foppoli, C., Mosca, L. & Cini, C. (1998) Production of melanin pigments by cytochrome c/H<sub>2</sub>O<sub>2</sub> system. *Int. J. Biochem. Cell Biol.* **30**, 457–463.
30. Hamachi, I., Fujita, A. & Kunitake, T. (1997) Protein engineering using molecular assembly: functional conversion of cytochrome c via noncovalent interactions. *J. Am. Chem. Soc.* **119**, 9096–9102.
31. Cornish-Bowden, A. (1995) *Fundamentals of Enzyme Kinetics*, Portland Press, London, UK.
32. Spee, J.H., Boersma, M.G., Veeger, C., Samyn, B., van Beeumen, J., Warmerdam, G., Canters, G.W., van Dongen, W.M.A.M. & Rietjens, I.M.C.M. (1995) The influence of the peptide chain on the kinetics and stability of microperoxidases. *Eur. J. Biochem.* **241**, 215–220.
33. Bielski, B.H.J. & Richter, H.W. (1977) A study of the superoxide radical chemistry by stopped-flow radiolysis and radiation induced oxygen consumption. *J. Am. Chem. Soc.* **99**, 3019–3023.
34. Davies, M.J. & Puppo, A. (1992) Direct detection of a globin-derived radical in leghaemoglobin treated with peroxides. *Biochem. J.* **281**, 197–201.
35. Puppo, A. & Davies, M.J. (1995) The reactivity of thiol compounds with different redox states of leghaemoglobin: evidence for competing reduction and addition pathways. *Biochim. Biophys. Acta* **1246**, 74–81.
36. Barr, D.P., Gunther, M.R., Deterding, L.J., Tomer, K.B. & Mason, R.P. (1996) ESR spin-trapping of a protein-derived tyrosyl radical from the reaction of cytochrome c with hydrogen peroxide. *J. Biol. Chem.* **271**, 15498–15503.
37. Saleem, M.M.M. & Wilson, M.T. (1988) Ligand binding to cytochrome c and other related haem proteins and peptides. Part I. Equilibrium studies. *Inorg. Chim. Acta* **153**, 93–98.
38. Saleem, M.M.M. & Wilson, M.T. (1988) Ligand binding to cytochrome c and other related haem proteins and peptides. Part II. Kinetic studies. *Inorg. Chim. Acta* **153**, 99–104.
39. Saleem, M.M.M. & Wilson, M.T. (1988) Ligand binding to cytochrome c and other related haem proteins and peptides. Part III. Temperature dependence studies. *Inorg. Chim. Acta* **153**, 105–113.
40. Viola, F., Aime, S., Coletta, M., Desideri, A., Fasano, M., Paoletti, S., Tarricone, C. & Ascenzi, P. (1996) Azide, cyanide, fluoride, imidazole and pyridine binding to ferric and ferrous native horse heart cytochrome c and to its carboxymethylated derivative: a comparative study. *J. Inorg. Biochem.* **62**, 213–222.
41. Gunther, M.R., Tschirret-Guth, R.A., Witkowska, H.E., Fann, Y.C., Barr, D.P., Ortiz de Montellano, P.R. & Mason, R.P. (1998) Site-specific spin trapping of tyrosine radicals in the oxidation of metmyoglobin by hydrogen peroxide. *Biochem. J.* **330**, 1293–1299.
42. Smith, W.L., Garavito, R.M. & DeWitt, D.L. (1996) Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2. *J. Biol. Chem.* **271**, 33157–33160.
43. Doyle, W.A., Blodig, W., Veitch, N.C., Piontek, K. & Smith, A.T. (1998) Two substrate interaction sites in lignin peroxidase revealed by site-directed mutagenesis. *Biochemistry* **37**, 15097–15105.
44. Blodig, W., Smith, A.T., Winterhalter, K. & Piontek, K. (1999) Evidence from spin-trapping for a transient radical on tryptophan residue 171 of lignin peroxidase. *Arch. Biochem. Biophys.* **370**, 86–92.
45. Lardinois, O.M., Medzhradsky, K.F. & Ortiz de Montellano, P.R. (1999) Spin trapping and protein cross-linking of the lactoperoxidase protein radical. *J. Biol. Chem.* **274**, 35441–35448.
46. Deterding, L.J., Barr, D.P., Mason, R.P. & Tomer, K.B. (1998) Characterization of cytochrome c free radical reactions with peptides by mass spectrometry. *J. Biol. Chem.* **273**, 12863–12869.
47. Anni, H. & Israel, Y. (1999) Characterization of adducts of ethanol metabolites with cytochrome c. *Alcohol Clin. Exp. Res.* **23**, 26–37.
48. King, N.K. & Winfield, M.E. (1963) The mechanism of metmyoglobin oxidation. *J. Biol. Chem.* **238**, 1520–1528.
49. Uyeda, M. & Peisach, J. (1981) Ultraviolet difference spectroscopy of myoglobin: assignment of pK values of tyrosyl phenolic groups and the stability of the ferryl derivatives. *Biochemistry* **20**, 2028–2035.
50. Moreau, S., Davies, M.J., Mathieu, C., Hérouart, D. & Puppo, A. (1996) Leghemoglobin-derived radicals – evidence for multiple protein-derived radicals and the initiation of peribacteroid membrane damage. *J. Biol. Chem.* **271**, 32557–32562.

51. Butler, J., Land, E.J., Prütz, W.A. & Swallow, A.J. (1982) Charge transfer between tryptophan and tyrosine in proteins. *Biochim. Biophys. Acta* **705**, 150–162.
52. Faraggi, M., DeFilippis, M.R. & Klapper, M.H. (1989) Long-range electron transfer between tyrosine and tryptophan in peptides. *J. Am. Chem. Soc.* **111**, 5141–5145.
53. Weinstein, M., Alfassi, Z.B., DeFelippis, M.R., Klapper, M.H. & Faraggi, M. (1991) Long range electron transfer between tyrosine and tryptophan in hen egg-white lysozyme. *Biochim. Biophys. Acta* **1076**, 173–178.
54. Irwin, J.A., Østdahl, H. & Davies, M.J. (1999) Myoglobin-induced oxidative damage: Evidence for radical transfer from oxidized myoglobin to other proteins and antioxidants. *Arch. Biochem. Biophys.* **362**, 94–104.
55. Ortiz de Montellano, P.R. & Catalano, C.E. (1985) Epoxidation of styrene by hemoglobin and myoglobin. Transfer of oxidizing equivalents to the protein surface. *J. Biol. Chem.* **260**, 9265–9271.
56. Kelman, D.J., DeGray, J.A. & Mason, R.P. (1994) Reaction of myoglobin with hydrogen peroxide forms a peroxy radical which oxidizes substrates. *J. Biol. Chem.* **269**, 7458–7463.
57. Galaris, D. & Korantzopoulos, P. (1997) On the molecular mechanism of metmyoglobin-catalyzed reduction of hydrogen peroxide by ascorbate. *Free Rad. Biol. Med.* **22**, 657–667.
58. Rao, S.I., Wilks, A. & Ortiz de Montellano, P.R. (1993) The roles of His-64, Tyr-103, Tyr-146, and Tyr-151 in the epoxidation of styrene and beta-methylstyrene by recombinant sperm whale myoglobin. *J. Biol. Chem.* **268**, 803–809.
59. Choe, Y.S., Rao, S.I. & Ortiz de Montellano, P.R. (1994) Requirement of a second oxidation equivalent for ferryl oxygen transfer to styrene in the epoxidation catalyzed by myoglobin-H<sub>2</sub>O<sub>2</sub>. *Arch. Biochem. Biophys.* **314**, 126–131.
60. Giulivi, C., Romero, F.J. & Cadenas, E. (1992) The interaction of Trolox C, a water-soluble vitamin E analog, with ferrylmyoglobin: reduction of the oxoferryl moiety. *Arch. Biochem. Biophys.* **299**, 302–312.
61. Gunther, M.R., Kelman, D.J., Corbett, J.T. & Mason, R.P. (1995) Self-peroxidation of metmyoglobin results in formation of an oxygen-reactive tryptophan-centered radical. *J. Biol. Chem.* **270**, 16075–16081.
62. Wang, J.-S., Baek, H.K. & Van Wart, H.E. (1991) High-valent intermediates in the reaction of *N*<sup>ε</sup>-acetyl microperoxidase-8 with hydrogen peroxide: models for compounds 0, I and II of horseradish peroxidase. *Biochem. Biophys. Res. Comm.* **179**, 1320–1324.
63. Carraway, A.D., Miller, G.T., Pearce, L.L. & Peterson, J. (1998) The alkaline transition of bis (N-acetylated) heme undecapeptide. *Inorg. Chem.* **37**, 4654–4661.
64. Wilson, M.T. & Greenwood, C. (1995) The alkaline transition in ferricytochrome *c*. In *Cytochrome c. A Multidisciplinary Approach* (Scott, R.A. & Mauk, A.G., eds), pp. 611–634. University Science Books, Sausalito, CA.
65. Rosell, F.I., Ferrer, J.C. & Mauk, A.G. (1998) Proton-linked protein conformational switching: definition of the alkaline conformational transition of yeast iso-1-ferricytochrome *c*. *J. Am. Chem. Society* **120**, 11234–11245.
66. Stubbe, J. & van der Donk, W.A. (1998) Protein radicals in enzyme catalysis. *Chem. Rev.* **98**, 705–762.
67. Lebedeva, O.V. & Ugarova, N.N. (1996) Mechanism of peroxidase-catalyzed oxidation. Substrate-substrate activation in horseradish peroxidase-catalyzed reactions. *Russ. Chem. Bull.* **45**, 25–32.
68. Smith, A.T., Sanders, S., Greschik, H., Thorneley, R.N.F., Burke, J.F. & Bray, R.C. (1992) Probing the mechanism of horseradish peroxidase by site-directed mutagenesis. *Biochem. Soc. Trans.* **20**, 340–345.
69. Brown, S.B. & Jones, P. (1967) Reactions between haemin and hydrogen peroxide. Part 2. Destructive oxidation of haemin. *Trans. Faraday Soc.* **64**, 994–998.
70. Brown, S.B., Hatzikostantinou, H. & Herries, D.G. (1978) The role of peroxide in haem degradation. A study of the oxidation of ferrihaems by hydrogen peroxide. *Biochem. J.* **174**, 901–907.
71. Schaefer, W.H., Harris, T.M. & Guengerich, F.P. (1985) Characterization of the enzymatic and nonenzymatic peroxidative degradation of iron porphyrins and cytochrome P-450 heme. *Biochemistry* **24**, 3254–3263.
72. Morishima, I., Fujii, H., Shiro, Y. & Sano, S. (1995) Studies on the iron (II) meso-oxyporphyrin pi-neutral radical as a reaction intermediate in heme catabolism. *Inorg. Chem.* **34**, 1528–1535.
73. Florence, T.M. (1985) The degradation of cytochrome *c* by hydrogen peroxide. *J. Inorg. Biochem.* **23**, 131–141.
74. Ortiz de Montellano, P.R. (1998) Heme oxygenase mechanism: evidence for an electrophilic, ferric peroxide species. *Acc. Chem. Res.* **31**, 543–549.
75. Davison, A.J. & Hulett, L.G. (1971) Consecutive oxidation and reduction of cytochrome *c* in the presence of hydrogen peroxide and copper histidine. *Biochim. Biophys. Acta* **226**, 313–319.
76. Mochan, E. & Degn, H. (1969) Autocatalytic peroxidation of ferrocycytochrome *c*. *Biochim. Biophys. Acta* **189**, 354–359.