



Universiteit
Leiden
The Netherlands

Migraine as a cardiovascular risk factor for women

Linstra, K.M.

Citation


Linstra, K. M. (2023, May 16). *Migraine as a cardiovascular risk factor for women*. Retrieved from <https://hdl.handle.net/1887/3618277>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3618277>

Note: To cite this publication please use the final published version (if applicable).



CHAPTER 3

Cold extremities in migraine: a marker for vascular dysfunction in women

*K.M. Linstra, M.J.L. Perenboom, E.W. van Zwet, F.C. van Welie, R. Fronczek,
M.R. Tannemaat, M. J.H. Wermer, A. Maassenvandenbrink, G.M. Terwindt*

ABSTRACT

Objective: Migraine is recognized as a vascular risk factor, especially in women. Presumably, migraine, stroke and cardiovascular events share pathophysiological mechanisms. We investigated self-reported cold extremities as a marker for vascular dysfunction in migraine. Secondly, we hypothesized that suffering from cold extremities affects sleep quality, possibly exacerbating migraine attack frequency.

Methods: In this case control study, a random sample of 1084 migraine patients and 348 controls (aged 22-65 years) from the LUMINA migraine cohort were asked to complete questionnaires concerning cold extremities, sleep quality and migraine.

Results: A total of 594 migraine patients and 199 controls completed the questionnaires. In women, thermal discomfort and cold extremities (TDCE) was more often reported by migraineurs versus controls (OR 2.3; 95%CI: 1.4-3.7; $p < 0.001$), but not significantly so in men (OR 2.5; 95%CI: 0.9-6.9; $p = 0.09$). There was no difference in TDCE when comparing migraine with or without aura. Female migraineurs who reported TDCE had higher attack frequencies compared to female migraineurs without TDCE (4 versus 3 attacks per month; $p = 0.003$). The association between TDCE and attack frequency was mediated by presence of difficulty initiating sleep (DIS) ($p = 0.02$).

Conclusion: Women with migraine more often reported cold extremities compared with controls, possibly indicating a sex-specific vascular vulnerability. Female migraineurs with cold extremities had higher attack frequencies, partly resulting from sleep disturbances. Future studies need to demonstrate whether cold extremities in female migraineurs are a predictor for cardio- and cerebrovascular events.

INTRODUCTION

Migraine is three times more prevalent in women than in men.¹ Patients with migraine have an increased risk for cerebro- and cardiovascular disease, which is even more pronounced in young women, especially those with migraine with aura.² A systemic vascular vulnerability, possibly influenced by female sex hormones, is hypothesized to underlie this association.³ Comorbidities of migraine include small vessel disease such as coronary artery vasospasm and Raynaud's phenomenon.^{2,3} These syndromes are more often reported in women, similar to increased thermal discomfort and cold extremities (TDCE).⁴ Difficulties in regulating body temperature are associated with a delay in sleep onset, termed difficulty initiating sleep (DIS). Normally, peripheral vasodilation causes redistribution of heat, enabling the reduction of core body temperature that is essential for rapid sleep induction.⁵ Experiencing cold extremities may indicate an autonomic or peripheral microvascular dysfunction and may interfere with falling asleep. For patients with migraine, insufficient sleep may be a trigger factor for attacks.⁶ Based on these observations, the present study aimed to investigate self-reported cold extremities as a marker for vascular dysfunction in migraine. Additionally, we hypothesized that suffering from cold extremities may be associated with difficulties initiating sleep, and therefore, with an increased migraine attack frequency.

METHODS

SUBJECTS

This study was conducted as part of the Leiden University Migraine Neuro-Analysis (LUMINA) programme.⁷ Participants of the LUMINA project are Dutch adults aged 18–80 years who fulfil the criteria for migraine with or without aura according to the International Classification of Headache Disorders (previous ICHD-2, now ICHD-3).⁸ Healthy controls were free of any known neurological or psychiatric disorders and did not have any primary or secondary headaches apart from an occasional episodic tension type headache. Further description of the LUMINA cohort is found as supplement (Supplement 1). The study was approved by the medical ethics committee of Leiden University Medical Centre. All subjects provided written informed consent prior to participation. A random selection of 1084 migraine patients and 348 controls (aged 22–65 years) was made from the LUMINA cohort for this present study.

QUESTIONNAIRES

Participants were invited to fill out questionnaires concerning Thermal Discomfort and Cold Extremities (TDCE) and Difficulties Initiating Sleep (DIS), between December 2016 and January 2017. For TDCE, questions included whether and to what extent participants suffered from cold hands and/or feet over the past month. For DIS, questions concerned to what extent participants experienced problems with falling asleep and/or a sleep onset latency of more than 30 minutes in the last month. The questionnaires and ratings are presented in detail as supplement (Supplement 2). The questionnaires were developed and externally validated with objective finger skin temperature measurements in a large population based study.⁴ Recent information on migraine characteristics and medication use was obtained through

extended questionnaires. Vasoactive medication use was defined as chronic daily use of triptans, ergotamines, beta-blockers, amphetamines and/or SSRI's.

DATA ANALYSIS

In general, non-parametric statistical tests were used. Chi-square statistics were used to calculate prevalence rates of TDCE and DIS in migraineurs and controls. Odds Ratios (OR) were calculated for the association between TDCE and DIS and migraine for each sex separately. The association of migraine subtype and attack frequency to TDCE and DIS was calculated for each sex. Multiple linear regression was performed to investigate association of TDCE and migraine in women, correcting for age, BMI, smoking and vasoactive medication use, followed by correction for multiple comparisons. Sobel test for mediation was applied to investigate whether DIS was a mediator in the association between TDCE and increased migraine attack frequency. All analyses were performed using SPSS 24.0 (SPSS Inc., IBM, USA).

DATA AVAILABILITY

Anonymized data will be shared by request from any qualified investigator for the sole purpose of replicating procedures and results presented in the article and as long as data transfer is in agreement with European Union legislation on the general data protection regulation.

RESULTS

Questionnaires were completed by 594 (55%) migraine patients and 199 (57%) controls. Women were overrepresented in this study and among migraineurs compared to controls (88% vs 61%). Vasoactive medication use was higher among migraineurs compared to controls (15% vs 5%). There were no significant differences in the distribution of age, BMI or smoking (Table 1).

■ **Table 1.** Clinical and headache characteristics^a

Variable	Migraine (n = 594)	Controls (n = 199)
Age, y	45 (13)	46 (10)
Sex		
Female	520 (87.5)	123 (61.8)
Male	74 (12.5)	76 (38.2)
BMI^b	25.0 (6.8)	24.1 (3.6)
Current smoker	77 (13.0)	22 (11.0)
Vasoactive medication^c	90 (15.2)	9 (4.5)
Age at onset, y	19 (10)	n.a.
Migraine with aura	212 (35.7)	n.a.
Attack frequency^d	3.0 (3.4)	n.a.

Abbreviations: BMI=body mass index, n.a.= not applicable.

^a Data is represented as mean ± SD or number of subjects (%)

^b Calculated as weight in kilograms divided by height in meters squared

^c Daily use of triptans, ergotamines, beta-blockers, amphetamines and/or SSRI's.

^d Number of attacks per month


THERMAL DISCOMFORT AND COLD EXTREMITIES (TDCE)

In women, TDCE was more often reported by migraine patients versus controls (OR 2.1; 95%CI: 1.3-3.3) (34% vs 20%; $p<0.01$) (Figure 1). The difference remained statistically significant after adjustment for age, BMI, smoking and vasoactive medication (OR 2.3; 95%CI: 1.4-3.7; $p<0.001$). In men, TDCE was reported more often by migraineurs versus controls (OR 2.5; 95%CI: 0.9-6.9) (18% vs 8%) however this difference was not statistically significant ($p=0.09$). No difference in TDCE was found comparing migraine subtypes (migraine with aura versus migraine without aura) (OR 1.2; 95%CI: 0.8-1.7; $p=0.43$).

DIFFICULTIES INITIATING SLEEP (DIS)

In general, positive outcome of TDCE was associated with DIS (OR: 2.4; 95%CI: 1.7-3.5; $p<0.001$). DIS was reported more often in both women (OR: 5.2; 95%CI: 2.4-11.4; $p<0.0001$) and men (OR: 3.8; 95%CI: 1.2-12.4; $p=0.02$) suffering from migraine compared to healthy controls (Figure 1).


■ **Figure 1.** Thermal discomfort and cold extremities and difficulty initiating sleep in migraine patients



(A) Thermal discomfort and cold extremities (TDCE) and (B) difficulties initiating sleep (DIS). Showing patients with migraine (red) and controls (blue) separately in percentages, within all participants ($n=793$), female participants ($n=520$) and male participants ($n=273$), * $p<0.05$.

Both female and male migraineurs who reported TDCE had significantly higher migraine attack frequencies compared to migraineurs who did not report TDCE, increasing from an average of 3 to 4 attacks per month ($p=0.02$) for women and 6 attacks per month for men ($p=0.002$). Sobel test for mediation indicated that this association between TDCE and migraine attack frequency was mediated by presence of DIS ($p=0.02$) (Figure 2).

■ **Figure 2.** Schematic representation of proposed mechanism and Sobel analysis.



Proposed mechanism of sex-specific vascular vulnerability underlying the association between cerebro- and cardiovascular disease (CVD), Thermal discomfort and cold extremities (TDCE) and migraine (blue) and direction of Sobel analysis for mediation of association between migraine attack frequency as dependent variable and relevant TDCE (yes/no) as independent variable and relevant difficulties initiating sleep (DIS) (yes/no) as mediator (red).

DISCUSSION

We were able to substantiate an association between migraine and cold extremities in our large, well defined cohort of migraine patients. We found that this observation was significant only in women. Previous studies showed decreased peripheral skin temperature in migraineurs. However, these studies had severe limitations with small sample sizes, poor migraine diagnosis, illogical findings (showing an association with right-sided headache only), and all lacked appraisal of sex differences.^{9,10} Our findings coincide with the current views concerning the sex-specificity of the vascular risk for migraine patients, and with suggestions that the underlying vascular vulnerability may be a systemic female type pathophysiology, possibly involving sex hormones.³ It has been debated whether the increased vascular risk for migraineurs is in fact exclusive to women or is less apparent in men due to underpowered analyses for male migraineurs.² This limitation might also have influenced our findings as the majority of our participants were women and therefore the confidence intervals around the OR in men were relatively wide. Larger studies are needed to further clarify the role of TDCE in men with migraine. Based on the higher increased risk of vascular disease for migraine with aura² we expected to find a difference in TDCE between migraine types, but found no evidence for that although the percentage of migraine with aura patients (36%) is what is generally expected in population-based studies.¹ It is noteworthy that our findings are still significant after adjustment for age, since the association of migraine with cardiovascular disease is especially prominent in young women (>45 years).² Although age was present as a co-variate in our regression analysis we did not take pre- or post-menopausal state into account. However, when we divided our groups in ≤45 years (presumably premenopausal) and >45 years (presumably peri- or postmenopausal) we did find that especially the younger female

migraineurs contributed to our findings on TDCE as the OR in this group was 2.3 (95% CI 1.2-4.5; $p=0.012$) comparing migraineurs with controls, whereas for the >45 years groups the OR was 1.9 (95% CI 0.9-3.7; $p=0.067$). Further studies investigating the mechanisms of hormonal contribution could provide interesting insights. Another finding worth mentioning, is the increased attack frequency for migraineurs who reported TDCE. This association was mediated by difficulties falling asleep (DIS). This is comprehensible since the disability to decrease core temperature through peripheral vasodilation may lead to delayed sleep onset and this in turn may be detrimental for migraine sensitivity.⁴⁻⁶ It can be hypothesized that cold extremities are associated with more severe migraine because the underlying vascular vulnerability is likely to be more prominent in these patients. This would concur with suggestions that a higher attack frequency is associated with an even further increased cerebro- and cardiovascular risk in migraine patients.² Using clinical markers such as TDCE might thus provide useful clinical tools to assess vascular vulnerability, but also migraine treatment response in the future.

CONCLUSION

Women with migraine report cold extremities more often than women without migraine, which may indicate the presence of a systemic vascular vulnerability in migraine. Migraineurs with cold extremities experience higher migraine attack frequencies, which is partly due to difficulties initiating sleep. If cold extremities in female migraineurs are proven to be associated with development of vascular complications later in life, this may present an easily assessed marker for vascular vulnerability.

REFERENCES

1. Launer LJ, Terwindt GM, Ferrari MD. The prevalence and characteristics of migraine in a population-based cohort: the GEM study. *Neurology*. 1999;53(3):537–42.

2. Sacco S, Ricci S, Carolei A. Migraine and vascular diseases: a review of the evidence and potential implications for management. *Cephalalgia*. 2012;32(10):785–95.

3. Tietjen GE. Migraine as a systemic vasculopathy. *Cephalalgia*. 2009;29(4):989–96.

4. Kräuchi K, Gasio PF, Vollenweider S, et al. Cold extremities and difficulties initiating sleep: evidence of co-morbidity from a random sample of a Swiss urban population. *J Sleep Res*. 2008;17(4):420–6.

5. Fronczek R, Overeem S, Lammers GJ, et al. Altered skin-temperature regulation in narcolepsy relates to sleep propensity. *Sleep*. 2006;29(11):1444–9.

6. Pavlovic JM, Buse DC, Sollars CM, et al. Trigger factors and premonitory features of migraine attacks: Summary of studies. *Headache*. 2014;54(10):1670–9.

7. van Oosterhout WPJ, Weller CM, Stam AH, et al. Validation of the web-based LUMINA questionnaire for recruiting large cohorts of migraineurs. *Cephalalgia*. 2011;31(13):1359–67.

8. Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. *Cephalalgia*. 2018;38(1):1–211.

9. Zaproudina N, Lipponen JA, Karjalainen PA, et al. Acral coldness in migraineurs. *Auton Neurosci Basic Clin*. 2014;180(1):70–3.

10. McArthur DL, Cohen MJ. Measures of forehead and finger temperature, frontalis EMG, heart rate and finger pulse amplitude during and between migraine headaches. *Headache*. 1980;20(3):134–6.

RESPONSE TO LETTER TO THE EDITOR

Dear Editor,

We would like to take the opportunity to respond to the letter to the Editor of Dr. Celikbilek in response to our recent article *Cold extremities in migraine: a marker for vascular dysfunction in women*, which was recently published in your journal.¹

Dr. Celikbilek points out that temperature regulation and sleep onset are both under the influence of the autonomic nervous system. The onset of sleep follows a cascade of events, including vasodilatory responses in the peripheral microvasculature and a subsequent decrease of body temperature. Dr. Celikbilek reasons that both cold extremities and delayed sleep onset are attributable to an autonomic imbalance and suggests that no sex specific related mechanisms play a role. In response, we would like to clarify that the sex specific vascular vulnerability we propose, can indeed (in part) be dysfunction of the autonomic nervous system. The correspondent suggests that we interpreted the cold extremities and delayed sleep onset to be causative factors of each other, which we did not. What we did observe was that female migraineurs with cold extremities have a higher attack frequency, and that this higher attack frequency is partly explained by the presence of 'difficulty initiating sleep'.

Dysfunction of the autonomic nervous system in migraineurs is widely described.^{2,3} Not only do migraineurs deviate in heart rate variability and autonomic reflex testing, they also more often experience clinical symptoms such as syncope.⁴ Autonomic thermo- and baroregulation are greatly influenced by gonadal hormones, specifically oestradiol, resulting in striking sex differences.⁵ Thus, there is evidence that the autonomic nervous system is influenced by sex.

The correspondent further states that an increased inflammatory state, rather than sleep problems, explain the increased migraine attack frequency. We do not argue that this might also play a role. We do not propose delayed sleep onset as the only pathophysiological mechanism behind an increased attack frequency in migraine, let alone migraine chronification. In the latter, many other important facts are implicated, such as depression, cutaneous allodynia and medication overuse. To conclude, however, we did show a significant role for delayed sleep onset in the association between cold extremities and an increased attack frequency in women with migraine.

REFERENCES

1. Linstra KM, Perenboom MJL, Zwet EW, et al. Cold extremities in migraine: a marker for vascular dysfunction in women. *Eur J Neurol*. May 2020;ene.14289.

2. Mamontov OV, Babayan L, Amelin AV, et al. Autonomous control of cardiovascular reactivity in patients with episodic and chronic forms of migraine. *J Headache Pain*. 2016;17:52.

3. Miglis MG. Migraine and Autonomic Dysfunction: Which Is the Horse and Which Is the Jockey? *Curr Pain Headache Rep*. 2018;22(3):19.

4. Thijs RD, Kruit MC, van Buchem MA, et al. Syncope in migraine: the population-based CAMERA study. *Neurology*. 2006;66(7):1034-1037.

5. Charkoudian N, Hart ECJ, Barnes JN, et al. Autonomic Control of Body Temperature and Blood Pressure: Influences of Female Sex Hormones. *Clin Auton Res*. 2017;27(3).
