



Universiteit
Leiden
The Netherlands

Comparative effectiveness of surgery for traumatic acute subdural hematoma

Essen, T.A. van

Citation

Essen, T. A. van. (2023, June 8). *Comparative effectiveness of surgery for traumatic acute subdural hematoma*. Retrieved from <https://hdl.handle.net/1887/3619981>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3619981>

Note: To cite this publication please use the final published version (if applicable).





Chapter 5

Variation in neurosurgical management of traumatic brain injury: a survey in 68 centers participating in the CENTER-TBI study

Van Essen TA, den Boogert HF, Cnossen MC, de Ruiter GCW, Haitzma I, Polinder S, Steyerberg EW, Menon D, Maas AIR, Lingsma HF, Peul WC; CENTER-TBI Investigators and Participants.

Acta Neurochir (Wien). 2019 Mar;161(3):435-449.

Portions of this work were presented at the 13th Symposium of the International Neurotrauma Society, Toronto, Canada, August 11-16, 2018.

ABSTRACT

BACKGROUND

Neurosurgical management of traumatic brain injury (TBI) is challenging, with only low-quality evidence. We aimed to explore differences in neurosurgical strategies for TBI across Europe.

METHODS

A survey was sent to 68 centers participating in the Collaborative European Neurotrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. The questionnaire contained 21 questions, including the decision when to operate (or not) on traumatic acute subdural hematoma (ASDH) and intracerebral hematoma (ICH), and when to perform a decompressive craniectomy (DC) in raised intracranial pressure (ICP).

RESULTS

The survey was completed by 68 centers (100%). On average, 10 neurosurgeons work in each trauma center. In all centers a neurosurgeon was available within 30 minutes. Forty percent of responders reported a thickness or volume threshold for evacuation of an ASDH. Most responders (78%) decide on a primary DC in evacuating an ASDH during the operation, when swelling is present. For ICH, 3% would perform an evacuation directly to prevent secondary deterioration and 66% only in case of clinical deterioration. Most respondents (91%) reported to consider a DC for refractory high ICP. The reported cut-off ICP for DC in refractory high ICP, however, differed: 60% uses 25 mmHg, 18 % 30 mmHg and 17 % 20 mmHg. Treatment strategies varied substantially between regions, specifically for the threshold for ASDH surgery and DC for refractory raised ICP. Also, within center variation was present: 31% reported variation within the hospital for inserting an ICP monitor and 43% for evacuating mass lesions.

CONCLUSION

Despite a homogeneous organization, considerable practice variation exists of neurosurgical strategies for TBI in Europe. These results provide an incentive for comparative effectiveness research to determine elements of effective neurosurgical care.

INTRODUCTION

Neurosurgical decision-making in patients with traumatic brain injury (TBI) is often challenging for several reasons. First, no two TBI patients are identical – clinical and radiological findings may differ greatly.¹ Second, there is no high-quality evidence to support the range of possible neurosurgical procedures in TBI. Indications for surgical management are summarized in the Brain Trauma Foundation guidelines,² but are merely based on retrospective studies of small groups of selected patients. These guidelines provide general advice on surgical indications for evacuation of acute epidural (EDH), acute subdural (ASDH) and contusions/intracerebral hematomas (ICH) based on the size of the hematoma and midline shift. The guidance for decompressive surgery is even less clear. It is mostly performed to decrease raised intracranial pressure (ICP), either as a primary procedure in an acute setting, or as a secondary procedure to deal with diffuse edema or peri-contusional swelling. The guidelines state that this latter use of secondary decompression can reduce ICP, but does not necessarily improve outcome.³ More fundamentally, the rationale for ICP monitoring has been challenged by the BEST-TRIP randomized controlled trial (RCT), which found no benefit of a management protocol based on intracranial pressure monitoring, compared to one based on serial imaging and clinical examination. These results have generated doubts regarding ICP monitoring.⁴⁻⁸ Overall, there is no clear consensus on the indications, extent and timing of surgery.⁹

This limited high quality evidence for surgical management in TBI arises from a lack of RCTs, which may be difficult to conduct due to pragmatic, ethical and methodological barriers.¹⁰ However, observational studies to determine effectiveness are more prone for bias.¹¹ A promising alternative approach could be comparative effectiveness research (CER).^{12,13} In this design, the heterogeneity and variability, that trouble RCTs in TBI, are accepted and exploited to study effectiveness of treatments as they occur in real-life practice. The current Collaborative European Neurotrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study aims to use CER methodology to study treatment effectiveness of several neurosurgical interventions.¹⁴

The aim of this study was to explore differences in neurosurgical strategies for TBI across Europe to provide a context for CENTER-TBI, an up-to-date insight into European neurosurgical management of TBI, and to identify naturally occurring variation between trauma centers in order to identify substrates for neurosurgical research questions that might be answered using CER in the study.

MATERIALS AND METHODS

This study was conducted within the setting of the international observational study CENTER-TBI.¹⁴ Between 2014 and 2015, all centers participating in the international multicenter observational study CENTER-TBI (www.center-tbi.eu) were asked to complete a questionnaire on neurosurgical management of TBI (Supplementary file 1).¹⁵ The questionnaire was sent to 71 centers (Figure 1), of which 5 centers dropped out and 2 joined in, resulting in 68 eligible centers from Austria (n=2), Belgium (n=4), Bosnia Herzegovina (n=2), Denmark (n=2), Finland (n=2), France (n= 7), Germany (n=4), Hungary (n=3), Israel (n=2), Italy (n=10), Latvia (n=3), Lithuania (n=2), Norway (n=3), Romania (n=1), Serbia (n=1), Spain (n=4), Sweden (n=2), Switzerland (n=1), The Netherlands (n=6) and The United Kingdom (n=7).




Figure 1. Centers and countries included in the Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study

Reprinted and updated from Cnossen et al. (2016) with permission from dr. Cnossen and Maas et al. (2015). Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury: a prospective longitudinal observational study. *Neurosurgery*, 76:67–80, under a CC BY license, with permission from professor A.I. Maas

QUESTIONNAIRE DEVELOPMENT AND ADMINISTRATION

We developed a set of questionnaires based on available literature and experts to measure the structure and processes of TBI care in individual centers. Details regarding this process and the questionnaires used are described in a separate paper.¹⁵ Pilot testing was undertaken in 16 of the participating centers and feedback was incorporated into the final design.

One of the questionnaires was on neurosurgical standard practice. This survey contained 21 questions which could broadly be divided into 3 categories: 1) center characteristics and internal structure; 2) general (neuro)surgical trauma care and processes; 3) site specific neurosurgical management for treating ASDH, EDH, ICH, the use of DC, and policy with regard to orthopedic injuries in the context of patients who had suffered a TBI.

Questions either sought quantitative estimates of key metrics (e.g. annual surgical volume, staff size, ASDH thickness or ICP thresholds for surgery) or attempted to elicit the 'general policy' of the center. To capture the latter these questions were formulated in two ways: respondents were asked to estimate what the management strategy is in more than three quarters of patients in their center in a given context; or respondents were asked to indicate how often they used a particular surgical technique or how often specific factors influence their decision-making (never = 0-10%, rarely = 10-30%, sometimes = 30-70%, frequently = 70-90% and always 90-100%). The options 'frequently' and 'always' were interpreted as 'general policy', in line with a previous report¹⁶ and similar to previous publications on other questionnaires.^{15,17}

The reliability of the surveys were tested by calculation of concordance in a previous publication.¹⁵ Overall, the median concordance rates between duplicate questions, was 0.81 (range 0.44 – 0.97) and specifically for the 'Neurosurgery' survey 0.78 (range 0.68 – 0.86).

ANALYSES

The median and interquartile range (IQR) were calculated for continuous variables, and frequencies were reported along with percentages for categorical variables. Countries were divided into seven geographic regions: Northern Europe (Norway 3, Sweden 2, Finland 2 and Denmark 2 centers), Western Europe (Austria 2, Belgium 4, France 7, Germany 4, Switzerland 1 and The Netherlands 6 centers), The United Kingdom (7 centers), Southern Europe (Italy 10 and Spain 4 centers), Eastern Europe (Hungary 3, Romania 1, Serbia 1 and Bosnia Herzegovina 2 centers), Baltic States (Latvia 3 and Lithuania 2 centers) and Israel (2 centers).

For the following neurosurgical treatment strategies we quantified regional differences: an absolute cutoff of hematoma thickness as an indication for surgery

for ASDH, DC in the primary evacuation of an ASDH, early/pre-emptive surgical evacuation for ICH, and DC as a general policy in case of refractory raised ICP.

To assess the association of region with one of these treatment choices, a logistic regression was performed with treatment choice (general policy or 'yes/no') as a dependent variable and the region (categorical) as independent variable. Nagelkerke R² indicated the variance explained by geographic region. Analyses were done in IBM SPSS Statistics version 20 (IBM, Chicago, IL, USA).

RESULTS

CENTER CHARACTERISTICS

All 68 eligible centers completed the questionnaire on neurosurgery (response rate 100%). Questionnaires were mainly completed by neurosurgeons (n = 53, 78%), followed by local CENTER-TBI investigators (mainly research physicians or nurses: 19%). On average, 10 neurosurgeons (IQR 8-13) and 4 trauma surgeons (IQR 0-12) worked in each center. All centers reported that neurosurgical coverage was available 24 hours a day/7 days a week, either by way of in-house availability of a qualified neurosurgeon (47%), or the availability of such an individual in less than 30 minutes (53%) (Table 1).

GENERAL (NEURO)SURGICAL CARE AND PROCESSES

Treatment decisions regarding cranial surgical interventions in TBI patients within the critical care ER and ICU period are in most centers determined by the neurosurgeon (n= 65, 96%), followed by the orthopedic surgeons and neuro-intensivist in respectively 3% (n=2) and 1% (n=1). Urgent neurosurgical interventions (ICP monitor device insertion not included) for life-threatening traumatic intracranial lesions, are made by the neurosurgeon in 98.5% and trauma surgeons in 1.5% of the centers. Raised ICP will almost always be incorporated in decision making, the time of day almost never (Figure 2).

With regard to extremities fractures, the general policy in 59 (87%) centers was so-called damage control with priority for TBI and delayed definitive treatment of the limb fractures (Table 2). This policy is protocolized in 21 centers (22%).

Of all centers, 58 (85%) estimated the space-occupying effect of traumatic lesions on the surrounding tissue by calculation of the thickness of the hematoma and midline shift on CT. A quarter of centers used actual volume measurement to make surgical decisions (Table 2).

Table 1. Characteristics of centers participating in neurosurgery survey

Characteristic	N completed	No. (%) or median (IQR)
Profession of respondent	68	
Neurologist		3 (4)
Neurosurgeon		53 (78)
Trauma surgeon		3 (4)
ED physician		1 (2)
Intensivist		1 (1)*
Administrative staff member		11 (16)*
CENTER-TBI local investigator		13 (19)*
Volume of surgeries in 2013^a		
ASDH	59	25 (15-49)
ICH/contusion	58	10 (5-21)
EDH	59	10 (5-19)
DC		
Hemicraniectomy	57	10 (5-16)
Bifrontal	57	0 (0-2)
Removal bone flap	55	1 (0-3)
Ventriculostomy	57	7 (2-21)
Cranioplasty	56	10 (6-14)
Depressed skull fracture	57	5 (2-12)
Staffing (FTE)		
Neurosurgeons	66	10 (8-13)
Residents in training	65	5 (3-8)
Residents not in training	61	0 (0-3)
Trauma surgeons	64	4 (0-12)
Organization of care		
Neurosurgical decision making in ICU	68	65 (96)
Neurosurgeon		1 (3)
Trauma surgeon		0
Neurologist		1 (2)
Neurointensivist or general intensivist		
24/7 neurosurgical coverage**	68	32 (47)
Qualified neurosurgeon in-house		30 (44)
Resident neurosurgery in-house		36 (53)
Neurosurgeon within 30 minutes		11 (16)
Neurosurgical resident within 30 minutes		0 (0)
Neurosurgeon more than 30 minutes		

ASDH: acute subdural hematoma, EDH: epidural hematoma, ICH: intracerebral hematoma, DC: decompressive craniectomy, FTE: full time equivalent, ICU: intensive care unit

* Numbers do not add up because the local investigators also depicted their profession and one responder declared to be an intensivist as well as an administrative staff member.

** Multiple options possible

^a Head trauma related surgeries




Figure 2. Factors of influence on neurosurgical decision making

Shown are the percentages of centers that would be never/rarely, sometimes or frequently/always influenced by the described factors in the decision to perform neurosurgical procedures. Question was completed by all 68 centers.

ICP: intracranial pressure; ED: Emergency Department

NEUROSURGICAL MANAGEMENT OF ASDH, EDH, ICH, AND THE USE OF DECOMPRESSIVE CRANIECTOMY

ASDH provided the highest volume of neurosurgical TBI cases, on average 25 cases per year. When performing a DC (for any indication), hemicraniectomy was the preferential technique, and bifrontal craniectomy was rarely performed (Table 1). Less than half of the centers (n=27, 40%) reported an absolute threshold for evacuating an ASDH. Four out of ten centers generally incorporate age in their decision for evacuating an ASDH (Table 2 and Figure 2).

ICH were seldom operated upon pre-emptively, but 67% of centers reported undertaking delayed surgery in the event of deterioration. Almost a third of centers reported within-center variations between individual neurosurgeons in decisions regarding surgical evacuation of contusions or traumatic ICH.

Only a very low proportion of centers would routinely perform a DC at the time of evacuation of either ASDH or ICH (respectively 6% and 1.5% of the centers). For refractory raised ICP, most centers (n= 64, 91%) would consider a decompressive craniectomy, while 32 (47%) see this as a general policy in their center (Figure 3, Table 2 and Figure in supplementary file 2). Ninety-six percent (n=65) reported to have a specific threshold for DC in refractory raised ICP. This was most commonly specified as 25 mmHg (n=39, 58%), followed by 30 mmHg (n=12, 18%) and 20 mmHg (n=11, 17%).

Table 2. Neurosurgical treatment policy of traumatic brain injury

Characteristic	N completed	No. (%) or mean (sd)
Structural estimation of mass lesions on CT*	68	
Visual intuition (e.g. no actual measurement)		27 (40)
Width, diameter and/or amount of MLS of the mass lesion		58 (85)
Volume measurements with imaging software		11 (16)
Volume measurements with direct calculation		17 (25)
Other		1 (2)
ASDH operation determinants		
Age considered important in surgery decision ^A	68	
Size (volume or thickness) threshold for surgery	68	26 (42)
Minimum volume or thickness:	28**	27 (40)
15 mm		
10 mm		2 (3)
10 mm and/or > 5 mm MLS		16 (24)
5 mm		2 (3)
ASDH thickness > width of cranium		3 (4)
Midline shift > thickness ASDH		3 (4)
DC indications	68	2 (3)
Routine		4 (6)
Intra-operative brain swelling		59 (86)
Sometimes as a second procedure in case of uncontrollable ICP		5 (7)
Never		0 (0)
ICH/contusion operation determinants		
General policy	68	
Pre-emptive (to prevent deterioration)		2 (3)
Delayed (after deterioration)		45 (66)
Variable (depends on surgeon)		18 (27)
Other		3 (4)
DC indications	68	
Routine		1 (2)
Intra-operative brain swelling		55 (81)
Sometimes as a delayed procedure in case of uncontrollable ICP		10 (15)
Never		2 (3)
Raised ICP determinants		
DC employed >70 % of refractory high ICP cases	68	32 (46)
Mostly early DC (within 6-12 hrs of refractory ICP)	64	32 (47)
Mostly late DC (as last resort to control ICP)	64	32 (47)
ICP threshold for DC	68	65 (96)
Raised ICP threshold for DC (mmHg):	64***	
30		12 (18)
25		39 (60)
20		11 (17)
15		1 (2)
Not standardized		1 (2)

Table 2. Neurosurgical treatment policy of traumatic brain injury (continued)

Characteristic	N completed	No. (%) or mean (sd)
DC indications considered[*]		7 (10)
Pre-emptive in raised ICP (not last resort)	68	64 (91)
Refractory raised ICP (last resort)		9 (13)
CT evidence of raised ICP		45 (66)
Intra-operative brain swelling		2 (3)
Routine with every ASDH or ICH evacuation		
Policy towards extremity limb fractures^B		59 (87)
Damage control	68	9 (13)
Definitive care		

MLS: midline shift, BTF: Brain Trauma Foundation, ICP: intracranial pressure, hrs: hours

* Multiple options possible. ** One responder did not report a threshold for surgery while answering a specific threshold (10 mm). *** One responder reported to employ a threshold for DC in raised ICP while not giving their specific threshold. ^A The question was whether the responder considers if the decision on surgery in acute SDH is influenced by age (based on a general consensus in their respective center). ^B Damage control is focused on the TBI. All extremity fractures are stabilized, but definitive treatment delayed. Definitive care: the extremity fractures are operated as soon as possible.




Figure 3. Treatment indications for neurosurgical interventions

Shown are the proportions of centers that generally have these specific preferences with regard to operating or not in ASDH, ICH and raised intracranial pressure respectively.

ASDH: acute subdural hematoma; DC: decompressive craniectomy; ICH: intracerebral hematoma; ICP: intracranial pressure

GUIDELINES AND PRACTICE VARIATION

Overall, the reported adherence to the BTF guidelines was high (Figure 4). The use of surgical interventions and specific indications for these interventions varied substantially within and between regions (Table 3). Surgical evacuation of ICH was




Figure 4. Brain Trauma Foundation guideline adherence

Shown are the percentages of centers that reported to never/rarely, sometimes or frequently/always follow the Brain Trauma Foundation guidelines for the management of SDH, EDH or contusions. Question was completed by 68 of the 68 centers.

TBI: traumatic brain injury; SDH: subdural hematoma; EDH: epidural hematoma

Table 3. Within- and between-region variation in surgical management

Decision	Northern Europe	Western Europe	United Kingdom	Southern Europe	Eastern Europe	Baltic States	Israel	Nagelkerke R ² Value
<i>ASDH</i>								
- Size threshold for evacuation	56	29	0	29	71	80	100	0.34
- Routine or intraoperative DC	89	92	100	100	86	80	100	0.17
<i>ICH/contusion</i>								
- Pre-emptive surgery	0	0	0	7	0	20	0	0.35
<i>Refractory raised ICP</i>								
- DC	44	37	29	57	43	80	100	0.15

ASDH: acute subdural hematoma, ICH: intracerebral hematoma, DC: decompressive craniectomy, ICP: intracranial pressure

Table presents the proportion (%) of respondent within each region that indicated that they used the described strategy as their general policy for patients with respectively ASDH, ICH or refractory raised ICP. The Nagelkerke R² value represents the variation in treatment that can be explained by the region.

only performed in the Baltic States and Southern Europe and geographic region explained 35% of the variance in use of the intervention. Having a specific threshold for ASDH surgery and employing a DC for refractory raised ICP showed the largest within-region and also between-region variation. Lastly, when directly asked whether variation in specific management strategies exist, respectively 31% and 43% indicated to have a structural variation within their center staff with regard to ICP sensor insertion and mass lesion evacuation (Table 4).

Table 4. Neurosurgical decision making

Characteristic	N completed	No (%)
Structural variation* ICP monitor insertion	68	
No		47 (69)
Yes		21 (31)
Structural variation* mass lesion evacuation	65	
No		29 (43)
Yes		29 (43)
Depending on lesion type		7 (10)

ED: emergency department, GCS: Glasgow Coma Scale

* Structural variation refers to a situation in which one or more of the clinicians are generally more likely to perform the (diagnostic) intervention than others.

DISCUSSION

The aim of this study was to explore differences in neurosurgical strategies for TBI across Europe. We found substantial variability in practice and thereby provide useful indications regarding potential substrates for CER in CENTER-TBI. The structures and processes of neurosurgical care are generally homogeneous across centers with a comparable number of neurosurgeons, similar organization of neurosurgical coverage and uniform organization of responsibility for most surgical decisions on the ER and ICU. The indications for surgery, however, differ substantially with high within-region and between-region practice variations.

CONTEMPORARY NEUROSURGICAL CARE

There are no recent comparable studies providing an overview of neurosurgical management on this scale. Two recent national surveys, in The United Kingdom and the Republic of Ireland and The Netherlands, have shown a comparable variability among neurosurgeons regarding the decision to evacuate an ASDH or to perform a primary DC.^{18,19}

When comparing our results to existing -much older- surveys, evacuation of a traumatic ICH seems to be less often considered than in the past.^{20,21} Our results are concordant with older surveys in reporting variable use of DC for refractory raised ICP, despite the DECRA trial (the RECUEIcp was not published yet).^{22,23} Interestingly, although the mostly applied cutoff for DC in refractory is reported to be 25 mmHg (60%), a lower value, 20 mmHg, and a higher value, 30 mmHg, are both reported to be used in almost 20 % of centers.

More broadly, our results replicate past data that suggest poor guideline adherence and practice variability. Rayan et al. showed that in only 17% of a random sample of (brain) trauma patients care was delivered according to the BTF guidelines.²⁴ Of

note, in the current study, surveys were sent to the centers between 2014 and 2015, so the more recent, updated BTF guidelines were not published yet, although the update was for medical management mainly (except DC in refractory IC).³ Comparable questionnaires on other aspects of TBI care have recently been published for ER and ICU management that, without exception, show practice variation.^{15,17,25,26} Practice variation has also been reported for other life-threatening or emergency disorders including ruptured abdominal aneurysm²⁷ and the spontaneous intracerebral hemorrhage.²⁸

STRENGTHS AND LIMITATIONS

A strength of the current study is the methodology that we used to investigate practice variation. First, detailed questions were posed to shed light on specific clinical decisions with regard to neurosurgical interventions. Subsequently, (objective) answers on amounts (volume load, mostly from in-hospital registries) were combined with qualitative information (estimations of general policies, using two approaches). When integrated with the high response rate and low amount of missing data in 68 centers, this overview provides a complete picture of reported neurosurgical care across Europe.

This study also had weaknesses. First, responses to the questionnaire may have been biased by the abstract nature of the questions posed, which neglected to provide a more concrete clinical context for judgments about reported practice. Although the respondents were experienced neurosurgeons with a scientific background, the difficulty of weighing individual patient characteristics with potentially fatal consequences can never be fully captured by a theoretical survey. In particular, the rational decision making can obviously be completely different due to the cognitive biases of neurosurgeons in the acute critical care period.

Second, there might be a concern as to how well the individual neurosurgeon respondent can represent the general center neurosurgical policy. Although we urged the respondent to report the general consensus on treatment at their center rather than individual management preferences (see Supplementary file 1), neurosurgical strategies may still be variable within centers between neurosurgeons. However, we did capture a qualitative assessment of this intra-center variability (Table 4). Third, we did not fully account for inherent regional variations such as evidence knowledge, caseload and case-mix due to referral patterns or admission policies, as a potential explanation for differences in neurosurgery policies. Variations in evidence knowledge for some questions, such as those on guidelines, are important. Moreover, while we did assess the center's caseload and case-mix, the caseload and case-mix of the (individual) respondent was not specifically asked. Fourth, the questions dealt with individual decisions in isolation, rather than the more complex

real-life situation where several competing priorities need to be addressed. Fifth, the reports may have been biased (in varying extents) towards how centers would have been liked to be perceived, rather than a faithful report of actual clinical policy and practice. This issue will be addressed by a planned comparison of these Provider Profiling responses with actual treatment strategies employed in patient-level data from these centers in the CENTER-TBI Core study.

Finally, our study sample represents centers participating in TBI-research which are likely specialized neurotrauma centers with a tendency to have practice that is skewed towards up-to-date knowledge. An example is the fact that almost half of all centers stated to have a neurosurgeon in house 24 hours a day. When studying all centers in Europe providing care to TBI patients, variability might be even larger.

IMPLICATIONS

Our results should be interpreted in combination with the current evidence on the effectiveness of different surgical strategies. For the use of DC in refractory raised ICP due to diffuse swelling, two RCTs have provided useful guidance. The DECRA trial showed that early use of DC for modest rises in ICP was associated with worse outcomes.²² More recently however, after the conduct of this survey, the RESCUEicp trial showed that, when used for refractory severe intracranial hypertension, DC can save lives, but results in an excess of severely disabled survivors.²³ It is clear that the intervention is not uniformly beneficial: while some functional improvements occur by 12 months, many survivors remain severely disabled. Rescue-ICP was not published yet at the conduct of this study. In our study the majority of centers indicated that DC is often employed for both indications (pre-emptive and last resort). With regard to focal lesions, a recent study suggested that in patients with an ASDH an aggressive approach towards evacuation is associated with better outcome compared to a conservative approach.²⁹ Similar trends were noted in the STITCH-trauma trial, which suggested better outcome with early surgical management of ICH.³⁰ In our study, a minority of centers considers an early strategy for ICH evacuation.

Lastly, DC in the primary evacuation of an ASDH seems to be associated with more favorable outcomes.³¹ There is no class 1 evidence, although the research question is currently being challenged in an RCT (Rescue-ASDH; ISRCT87370545). In the current survey standard (in some cases preventive) DC in ASDH evacuation is rarely employed but mostly done in case of intraoperative swelling.

There may be several explanations for the practice variation that we observed. Although high practice variation rates can be a sign of poor implementation of evidence based care, in this context it probably reflects the lack of strong evidence to underpin practice. In such a low evidence context, clinical decisions are not driven by careful consideration or penetration of the evidence, but by local customs and surgical

training, handed down over the years from one surgeon to the other in a given center (or country). The professional cultural drivers that underpin such learned treatment preferences are resistant to change, and provide an important hurdle to the design and conduct of randomized studies for neurosurgical interventions in TBI.³²

Additionally, even where the results of RCTs are available, it is possible that many neurosurgeons do not think the RCT results applicable to their (individual) patients, or restrict their focus to short term clinical outcomes such as mortality and complication rates (instead of long term clinical or patient reported outcomes).³³

The results of the questionnaire point out burning clinical questions for neurosurgery in TBI. For ASDH and ICH, important questions include whether to operate or not, the timing of operative evacuation, and whether or not a primary DC should be undertaken. Future studies should address these questions. For DC, the variation should lead to studies exploring the lack of evidence penetration, in addition to studying effectiveness of DC in refractory raised ICP.

While RCTs may provide the security of randomisation as a basis for examining answering these questions, RCTs have no successful history in TBI due to various reasons.¹² The CENTER-TBI Provider Profiling exercise has revealed large practice variation that can be related to variation in patient outcome.³⁴ Such a CER approach may be a pragmatic alternative to RCTs.

Therefore, different steps are required. Firstly, to specify, ideally a-priori, how and where treatment variation occurs. This was one of the goals of this provider profiling. Secondly, the CENTER-TBI Core Study will need to collect patient-level data from a large variety of centers, capturing the range of treatment variation and relate it to outcome. The main challenge is to disentangle the effect of specific surgical strategies in a center from other regional care variation that might affect outcome. To do so we propose random-effect models in which the effect of ‘surgical strategy’ on outcome is estimated with adjustment for other between-hospital differences in a random effect for hospital.^{19,29,35}

CONCLUSIONS

This survey study explored differences in neurosurgical strategies for TBI. Current neurosurgical care differs within Europe (and Israel), while the organization of trauma centers does not. This variation in practice likely reflects the lack of high-quality evidence for these important, potentially life-saving, emergency neurosurgical interventions. In addition, local professional culture may drive practice in ways that are not dependent on the availability or penetration of evidence. The resulting entrenched practice variation does not facilitate equipoise that makes RCTs easy to deliver. CER may provide a pragmatic approach to generate evidence on optimal neurosurgical strategies for TBI patients.

COMPLIANCE WITH ETHICAL STANDARDS

This study was funded by The European Union seventh Framework Program (grant 602150) for Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) and the 'Hersenstichting Nederland' (Dutch Brain Foundation, grant PS2014-06) for The Dutch Neurotraumatology Quality Registry (Net-QuRe).

Author TvE declares that he has no conflict of interest. Author HdB declares that he has no conflict of interest. Author MC declares that she has no conflict of interest. Author GdR declares that he has no conflict of interest. Author IH declares that he has no conflict of interest. Author SP declares that she has no conflict of interest. Author ES declares that he has no conflict of interest. Author DM declares that he has no conflict of interest. Author AM declares that he has no conflict of interest. Author HL declares that she has no conflict of interest. Author WP declares that he has no conflict of interest.

All procedures performed in studies involving human participants were in accordance with ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments.

The CENTER-TBI Investigators and Participants and their affiliations are listed in the Appendix (available online):

<https://link.springer.com/article/10.1007/s00701-018-3761-z#appendices>

Further electronic supplementary material:

<https://link.springer.com/article/10.1007/s00701-018-3761-z#Sec15>

Supplemental material 1. Questionnaire neurosurgery.

Supplemental material 2. Supplementary figure, The use of a decompressive craniectomy.

REFERENCES

1. Maas AIR, Stocchetti N, Bullock R. Moderate and severe traumatic brain injury in adults. *The Lancet Neurology* 2008;7(8):728–41.
2. Bullock MR, Chesnut R, Ghajar J, et al. Introduction. *Neurosurgery* 2006;58(Supplement):S2–1–S2–3.
3. Carney N, Totten AM, O’Reilly C, et al. Guidelines for the Management of Severe Traumatic Brain Injury, Fourth Edition. *Neurosurgery* 2016; 1.
4. Albuquerque FC. Intracranial pressure monitoring after blunt head injuries: conflicting opinions. *World Neurosurg* 2013;79(5-6):598.
5. Chesnut RM, Temkin N, Carney N, et al. A Trial of Intracranial-Pressure Monitoring in Traumatic Brain Injury. *N Engl J Med* 2012; 367(26):2471–81.
6. Ghajar J, Carney N. Intracranial-pressure monitoring in traumatic brain injury. *N Engl J Med* 2013; 368(18):1749.
7. Kahle KT, Duhaime A-C. Intracranial-pressure monitoring in traumatic brain injury. *N Engl J Med* 2013; 368(18):1750.
8. Mattei TA. Intracranial pressure monitoring in severe traumatic brain injury: who is still bold enough to keep sinning against the level I evidence? *World Neurosurg* 2013; 79(5-6):602–4.
9. Servadei F, Compagnone C, Sahuquillo J. The role of surgery in traumatic brain injury. *Curr Opin Crit Care* 2007; 13(2):163–8.
10. Bragge P, Synnot A, Maas AI, et al. A State-of-the-Science Overview of Randomized Controlled Trials Evaluating Acute Management of Moderate-to-Severe Traumatic Brain Injury. *Journal of Neurotrauma* 2016; 33(16):1461–78.
11. Bosco JLF, Silliman RA, Thwin SS, et al. A most stubborn bias: no adjustment method fully resolves confounding by indication in observational studies. *Journal of Clinical Epidemiology* 2010; 63(1):64–74.
12. Maas AIR, Menon DK, Lingsma HF, et al. Re-Orientation of Clinical Research in Traumatic Brain Injury: Report of an International Workshop on Comparative Effectiveness Research. *Journal of Neurotrauma* 2012; 29(1):32–46.
13. Timmons SD, Toms SA. Comparative effectiveness research in neurotrauma. *Neurosurgical FOCUS* 2012;33(1):E3.
14. Maas AIR, Menon DK, Steyerberg EW, et al. Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI): a prospective longitudinal observational study. *Neurosurgery* 2015;76(1):67–80.
15. Cnossen MC, Polinder S, Lingsma HF, et al. Variation in Structure and Process of Care in Traumatic Brain Injury: Provider Profiles of European Neurotrauma Centers Participating in the CENTER-TBI Study. *PLoS ONE* 2016;11(8):e0161367.
16. Hesdorffer DC, Ghajar J. Marked Improvement in Adherence to Traumatic Brain Injury Guidelines in United States Trauma Centers. *The Journal of Trauma: Injury, Infection, and Critical Care* 2007;63(4):841–8.
17. Cnossen MC, Huijben JA, van der Jagt M, et al. Variation in monitoring and treatment policies for intracranial hypertension in traumatic brain injury: a survey in 66 neurotrauma centers participating in the CENTER-TBI study. *Critical Care* 2017;21(1):233.
18. Kolia AG, Scotton WJ, Belli A, et al. Surgical management of acute subdural haematomas: current practice patterns in the United Kingdom and the Republic of Ireland. *Br J Neurosurg* 2013;27(3):330–3.
19. Van Essen TA, De Ruiter GCW, Kho KH, Peul WC. Neurosurgical Treatment Variation of Traumatic Brain Injury: Evaluation of Acute Subdural Hematoma Management in Belgium and The Netherlands. *Journal of Neurotrauma* 2017 Feb 15;34(4):881-889.
20. Compagnone C, Murray GD, Teasdale GM, et al. The Management of Patients with Intracranial Post-Traumatic Mass Lesions: A Multicenter Survey of Current Approaches to Surgical Management in 729 Patients Coordinated by the European Brain Injury Consortium. *Neurosurgery* 2005; 1183–92.
21. Murray GD, Teasdale GM, Braakman R, et al. The European Brain Injury Consortium survey of head injuries. *Acta Neurochir* 1999;141(3):223–36.
22. Cooper DJ, Nichol A, Hodgson C. Craniectomy for Traumatic Intracranial Hypertension. *N Engl J Med* 2016;375(24):2402.
23. Hutchinsonson PJ, Kolia AG, Timofeev IS, et al. Trial of Decompressive Craniectomy for Traumatic Intracranial Hypertension. *N Engl J Med* 2016;375(12):1119–30.
24. Rayan N, Barnes S, Fleming N, et al. Barriers to compliance with evidence-based care in trauma. *J Trauma Acute Care Surg* 2012;72(3):585–92–discussion592–3.
25. Foks KA, Cnossen MC, Dippel DWJ, et al. Management of mild traumatic brain injury at the emergency department and hospital admission in Europe: A survey of 71 neurotrauma centers participating in the

- CENTER-TBI study. *Journal of Neurotrauma* 2017; Sep 1;34(17):2529-2535
26. Huijben JA, van der Jagt M, Cnossen MC, et al. Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. *Journal of Neurotrauma* 2017; neu.2017.5194.
 27. Brattheim BJ, Eikemo TA, Altreuther M, et al. Regional Disparities in Incidence, Handling and Outcomes of Patients with Symptomatic and Ruptured Abdominal Aortic Aneurysms in Norway. *European Journal of Vascular and Endovascular Surgery* 2012; 44(3):267-72.
 28. Gregson BA, Mendelow AD. International Variations in Surgical Practice for Spontaneous Intracerebral Hemorrhage. *Stroke* 2003;34(11):2593-7.
 29. Van Essen TA, Dijkman M, Cnossen MC, et al. Comparative effectiveness of surgery for traumatic acute subdural hematoma in an aging population. *Journal of Neurotrauma* 2019; Apr 1;36(7):1184-1191.
 30. Mendelow AD, Gregson BA, Rowan EN, et al. Early surgery versus initial conservative treatment in patients with traumatic intracerebral haemorrhage [STITCH(Trauma)]: the first randomised trial. *Journal of Neurotrauma* 2015; 32(17):1312-23.
 31. Li LM, Koliass AG, Guilfoyle MR, et al. Outcome following evacuation of acute subdural haematomas: a comparison of craniotomy with decompressive craniectomy. *Acta Neurochir* 2012; 154(9):1555-61.
 32. Macefield RC, Boulind CE, Blazeby JM. Selecting and measuring optimal outcomes for randomised controlled trials in surgery. *Langenbeck's Archives of Surgery* 2014; 399(3):263-72.
 33. Ergina PL, Cook JA, Blazeby JM, et al. Challenges in evaluating surgical innovation. *The Lancet* 2009; 374(9695):1097-104.
 34. Lingsma HF, Roozenbeek B, Li B, et al. Large Between-Center Differences in Outcome After Moderate and Severe Traumatic Brain Injury in the International Mission on Prognosis and Clinical Trial Design in Traumatic Brain Injury (IMPACT) Study. *Neurosurgery* 2011; 68(3):601-8.
 35. Cnossen MC, van Essen TA, Ceyisakar IE, et al. Adjusting for confounding by indication in observational studies: a case study in traumatic brain injury. *Clinical Epidemiology* 2018; 10:841-52.

