



Universiteit
Leiden
The Netherlands

Comparative effectiveness of surgery for traumatic acute subdural hematoma

Essen, T.A. van

Citation

Essen, T. A. van. (2023, June 8). *Comparative effectiveness of surgery for traumatic acute subdural hematoma*. Retrieved from <https://hdl.handle.net/1887/3619981>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)


Downloaded from: <https://hdl.handle.net/1887/3619981>

Note: To cite this publication please use the final published version (if applicable).

PART IV

**THE EFFECTIVENESS OF
NEUROSURGICAL INTERVENTIONS
IN ACUTE SUBDURAL HEMATOMA**





Chapter 9

Comparative effectiveness of surgery for traumatic acute subdural hematoma in an aging population

Van Essen TA, Dijkman MD, Cnossen MC, Moudrous W, Ardon H, Schoonman
GG, Steyerberg EW, Peul WC, Lingsma HF, de Ruiter GCW.

J Neurotrauma. 2019 Apr 1;36(7):1184-1191.

Portions of this work were presented at the 12th Symposium of the International Neurotrauma
Society, Capetown, South-Africa, February 1-4, 2016.

ABSTRACT

Uncertainty remains on the optimal initial management of patients with traumatic acute subdural hematoma, leading to regional variation in surgical policy. This can be exploited to compare the effect of various management strategies and determine best practices. This article reports such a comparative effectiveness analysis of a retrospective observational cohort of traumatic acute subdural hematoma patients in two geographically distinct neurosurgical departments chosen for their – a-priori defined – diverging treatment preferences. Region A favored a strategy focused on surgical hematoma evacuation, while region B employed a more conservative approach, performing primary surgery less often. Region was used as a proxy for preferred treatment strategy to compare outcomes between groups, adjusted for potential confounders using multivariable logistic regression with imputation of missing data. In total, 190 patients were included: 108 from region A and 82 from region B. There were 104 males (54.7%). Matching current epidemiological developments, the median age was relatively high at 68 years (IQR, 54-76). Baseline characteristics were comparable between regions. Primary evacuation was carried out in 84% of patients in region A and 65% in region B ($p < .01$). Mortality was lower in region A (37% vs. 45%, $p = .29$), as well as unfavorable outcome (53% vs. 62%, $p = .23$). The strategy favoring surgical evacuation was associated with significantly lower odds of mortality (OR: 0.43; 95% CI: 0.21 to 0.88) and unfavorable outcome (OR: 0.53; 95% CI: 0.27 to 1.02) 3-9 months post-injury. Therefore, in the aging population of patients with acute subdural hematoma a treatment strategy favoring emergency hematoma evacuation might be associated with lower odds of mortality and unfavorable outcome.

INTRODUCTION

In 1983, Gelpke and colleagues compared survival after severe traumatic brain injury (TBI) between two neurosurgical departments in the Netherlands. They concluded that of the 18% difference in survival rate between centers, 10.5% was accounted for by differences in baseline characteristics, and the remaining 7.5% could reflect differences in management strategies.¹ This study can be regarded as comparative effectiveness research (CER) *avant la lettre* in the field of TBI. CER aims to relate naturally occurring treatment variation to differences in patient-relevant outcomes, measured in day-to-day clinical settings and broad populations.² This is increasingly recognized as a powerful alternative to randomized controlled trials (RCTs).³ Many surgical trials have been hampered by protracted recruitment periods^{4,5} and financial constraints,⁶ while providing limited benefit to clinical management.^{5,7-9} Moreover, CER could be worthwhile because, partly due to current treatment guidelines featuring low class evidence, considerable between-center differences in the clinical management of TBI are seen.¹⁰⁻¹⁶

“To operate or not?” is a burning clinical question every surgeon frequently has to answer. In the case of traumatic acute subdural hematoma (ASDH), whether the patient will benefit from direct surgical evacuation of the hematoma remains a contested issue. Though prompt surgery has been associated with improved survival,¹⁷⁻²⁰ reported mortality rates remain high, ranging from 40 to 60% in surgically treated patients of all trauma severities.²¹ Conversely, it has been suggested that certain patients with a poor prognosis may not be treated at all,^{22,23} and patients with minimal symptomatology or comatose ASDH patient with a minimal hematoma without extracranial explanations for his/her coma can successfully be treated conservatively.²⁴⁻²⁷ Conservative treatment comprises of observation, serial neurological examinations and sometimes placement of an intracranial pressure (ICP) monitor with non-operative/medical management of intracranial hypertension when appropriate.

Thus, the decision to operate or not poses a clinical challenge for the neurosurgeon and should preferably be aided by guidelines based on comparison of treatment regimes. However, RCTs have never been done and comparative observational cohort studies with balanced treatment groups are unavailable. The most widely used surgical guidelines published by the Brain Trauma Foundation are consequently made up of weak evidence.²¹ Therefore clinical decision-making is presently influenced by neurosurgeons' individual preferences for, or familiarity with treatment of traumatic ASDH. As a result, management strategies vary considerably between neurosurgical departments, even between those within dedicated level 1 trauma centers with protocolled emergency medicine schemes.¹⁰⁻¹² Clearly, there is a need for comparative studies to elucidate the optimal treatment of traumatic

ASDH. Since randomized surgical trials are not practically feasible because of moral boundaries of treating physicians, the comparison of cohorts between homogenous regions managing ASDH in different ways may be the best available alternative to assess the effectiveness of surgical therapy.

The objective of this study was to compare mortality and functional outcome between an immediate surgical and a conservative, less surgery driven approach for the treatment of traumatic ASDH.

MATERIALS AND METHODS

This study was conducted and reported according to the criteria of the ‘Strengthening the Reporting of Observational Based Studies’ (STROBE) statement.²⁸

THE DUTCH HEALTHCARE SYSTEM AND REGIONAL TREATMENT VARIATION

Healthcare in the Netherlands is uniformly accessible to all patients, with equal distribution of resources among hospitals. Neurosurgical care for patients with TBI is provided at 11 level 1 trauma centers, serving separate areas according to regional referral policies. Training and licensing is equal among these centers. However, neurosurgical practices differ across these centers, probably due to local surgical customs handed down over the years: a recent survey assessing whether neurosurgeons would perform an operation in various patients with ASDH showed considerable between-center variation among Dutch neurosurgeons’ attitudes towards patients with traumatic ASDH.¹² Based on this questionnaire, the centers that showed the most divergent view on whether to operate acutely or not in cases reflecting the whole clinical TBI severity spectrum (90.0% vs 14.3% for moderate TBI, 90.0% and 42.9% for severe TBI) were chosen before the start of the current study. In both centers the same Brain Trauma Foundation guideline-based ICU protocol for (refractory) raised ICP is employed while the neurosurgical departments do not employ a surgical protocol (e.g. the Brain Trauma Foundation guideline).

STUDY DESIGN, SETTING AND POPULATION

Two patient cohorts were retrospectively identified at two neurosurgical departments, in geographically separate regions. Neurosurgeons from region A treat TBI patients at two neurosurgically collaborating hospitals within level 1 trauma centers and advocate primary surgical treatment of ASDH (through hematoma evacuation with or without decompression). Neurosurgeons from region B operate in a single level 1 trauma center and opt for surgery less often, suggesting a more conservative approach.¹² Both regions serve a homogeneous population of about 2 million people. Regions A and B employed 16 and 18 neurosurgeons respectively.

ELIGIBILITY CRITERIA

The Dutch registry system for hospital funding appoints diagnosis and treatment codes to all patients visiting a hospital. Using this system as a screening tool, we consecutively identified all patients treated by a neurosurgeon for TBI between 2008 and 2012 in both regions. Thereafter, the national trauma registry was checked for any missed inclusions.²⁹ A broad set of inclusion criteria was applied: patients were included if they had sustained head injury with direct presentation to the emergency room, were over 16 years of age, and showed a hyperdense, crescent shaped lesion – indicative of an ASDH – on the CT-scan. Exclusion criteria were penetrating injury, non-traumatic ASDH, ASDH secondary to an earlier procedure, and patients presenting with concomitant intracranial focal lesions (i.e. intracerebral hematoma or epidural hematoma) that required emergency surgery. Patients withheld from treatment due to severe comorbidity or because they were deemed unsalvageable, were also excluded on the premises the outcome would have been the same regardless of treatment.

VARIABLES

Data from electronic patient files were gathered on demographics, medical history, use of anticoagulants or antiplatelet agents, injury related variables, radiological variables, treatment variables, complications and outcome variables.

Injury related variables included trauma mechanism, first emergency room Glasgow Coma Scale (GCS) score, focal neurological symptoms (paresis, aphasia or cranial nerve deficit), pupillary light reflex, clinical deterioration (i.e. a decrease of more than 1 point on the GCS, new abnormal pupillary light reflex, or new focal neurological symptoms, from the time of first assessment), presence of significant extra-cranial injury, and primary presentation to the study hospitals.

Radiological variables were assessed from the first CT-scan. They included clot thickness, midline shift, patency of the basal cisterns, presence of cranial fractures, and presence of concomitant intracranial hemorrhage (subarachnoid hemorrhage, epidural hematoma, contralateral subdural hematoma, intraventricular hemorrhage, or intracerebral hematoma/contusion). We also noted if a second preoperative CT-scan was made and whether it showed radiological deterioration (i.e. presence of new focal lesion or >5mm increase in hematoma thickness).

Therapy related variables included type of management (conservative or surgical), type of surgery (craniotomy, decompressive craniectomy (DC), or other), use of an ICP monitor, and any delayed surgical procedures performed after the initial treatment was received. The maximum diameter of the DC was measured from postoperative CT-scans. Patients were considered surgically treated when a report was made on the indication and the surgical procedure was started after the last CT made on the emergency room. Patients were considered conservatively managed when the

neurosurgeon on call reported not to operate after the CT. Patients that were primarily conservatively managed could require surgery later on, after secondary deterioration. All complications requiring medical attention during admission (e.g. antibiotic treatment for infection) were noted. Complications were defined as intracranial (e.g. seizure, hydrocephalus), cardiovascular (e.g. arrhythmia, ischemia), respiratory (e.g. respiratory insufficiency, hypoxia), metabolic (e.g. electrolyte disturbances, renal failure), infections (e.g. pneumonia, wound infection), or other.

Outcome measures were mortality at discharge and functional outcome according to the Glasgow Outcome Score (GOS). Functional outcome was judged from outpatient follow-up letters at 3-9 months post trauma, and dichotomized into favorable (GOS 4-5) or unfavorable (GOS 1-3).³⁰

Two authors from region A collected all data. To ensure uniform and unbiased collection, both contributors independently gathered data according to a standardized collection sheet. In case of uncertainty, variables were coded after consensus was reached through discussion.

This study was approved by the ethics committees of the two participating hospitals.

STATISTICAL ANALYSIS

The median and interquartile range (IQR) were calculated for continuous variables and frequencies were reported along with percentages for categorical variables. To test for differences in patient characteristics, treatments and outcomes between regions, we used Pearson's chi-square test for categorical, and the Mann-Whitney *U* test for continuous variables. In order to provide a summary baseline prognostic score for both regions, a multivariable logistic regression model, based on all available variables featured in the CRASH-CT head injury prognosis model (age, GCS score, pupil reactivity to light, major extracranial injury, midline shift >5mm, traumatic subarachnoid hemorrhage, and obliteration of the basal cisterns), was used to calculate predicted probabilities for mortality and unfavorable outcome.³¹

To assess the effect of treatment strategy on outcome, we used region as proxy for treatment strategy. That strategy correlates with more surgeries (and possibly other aspects, see discussion). Assuming regional variation was the only determinant of treatment strategy, the region in which patients were treated was used as grouping variable to compare outcomes. A multivariable logistic regression model was used to estimate the effect of treatment strategy on outcome, adjusted for the confounders age, GCS score, pupil reactivity and hematoma thickness. In addition, subgroup analysis restricted to patients with mild to moderate TBI (GCS 9 to 15) was done. The regression analyses were done using a multiple imputed dataset created with the automatic imputation method of the *impute* function of SPSS. Missing data were reported for all variables at baseline. Results were considered statistically significant if $p < .05$. All analyses were performed using IBM's SPSS version 23.

RESULTS

PARTICIPANTS AND PRESENTATION

A total of 612 cases were screened (N, region A=294; N, region B=318) of which 195 cases met the eligibility criteria: 109 from region A and 86 from region B. Five cases were excluded due to severe comorbidity (N, region A=1), or were deemed unsalvageable (N, region B=4, figure 1).

The study population included 104 males (54.7%) and the median age was 68 years (IQR, 54-76). Regions did not differ significantly on age, sex, use of anticoagulants, injury mechanism, GCS score, occurrence of focal neurological symptoms, pupillary exam, or frequency of major extracranial injury (table 1). Region A received more primary referrals than region B (63% vs. 21%, $p<.01$). Patients in region B more often experienced clinical deterioration (34% vs. 51%, $p<.01$). Region A had larger median hematoma thickness (14mm vs. 10mm, $p<.01$), larger median midline shift (12mm vs. 9mm, $p<.01$), and less cranial fractures (28% vs. 44%, $p<.03$). The baseline CRASH-CT predicted risk of unfavorable outcome was similar: 56% in region A and 53% in region B ($p=.18$). Missing values per variable used in both univariable and multivariable regression analyses did not exceed 5.8% (table 1).




Figure 1. Patient inclusion flow chart showing the number of patients included in each region (region A: liberal surgical policy, region B: more conservative policy) and the definitive treatment received (either primary surgical or conservative).

Table 1. Clinical and radiological variables for the entire study population

Variable, % missing	Region A (N=108)	Region B (N=82)	p-value
Age, median (IQR), years, 0%	70 (54 – 78)	65 (53 – 74)	.07
Male sex, No. (%), 0%	57/108 (53%)	47/82 (57%)	.56
History of diabetes, No. (%), 1.6%	14/105 (13%)	10/82 (12%)	1.00
Vascular history, No. (%), 1.6%	60/105 (57%)	45/82 (55%)	.769
Anticoagulant, No. (%), 1.6%			
Anticoagulants	34/107 (32%)	18/80 (23%)	.19
Platelet inhibitors	20/107 (20%)	16/80 (19%)	1.00
Mechanism of injury, No. (%), 2.5%			
Fall	58/104 (56%)	45/81 (56%)	1.00
Assault	5/104 (5%)	1/81 (1%)	.23
Motor vehicle accident	12/104 (12%)	13/81 (16%)	.39
Fall from bike	12/104 (12%)	11/81 (14%)	.82
Other	17/104 (16%)	11/81 (14%)	.68
Primary presentation, No. (%), 0%	67/106 (63%)	17/82 (21%)	<.01
Focal neurological symptoms, No. (%), 24.7%	42/79 (53%)	28/64 (44%)	.31
Abnormal pupils, No. (%), 5.3%			
One	13/101 (13%)	12/79 (15%)	.67
Two	14/101 (14%)	9/79 (12%)	.66
First GCS score, median (IQR), 2.6%	9 (6 – 14)	12 (7 – 15)	.48
Initial motor (M) score, median (IQR), 29.5%	6 (5 – 6)	5 (4 – 6)	.04
TBI severity, No. (%), 1.6%			
Mild (GCS 13 to 15)	40/107 (37%)	29/80 (36%)	1.00
Moderate (GCS 9 to 12)	18/107 (17%)	20/80 (25%)	.20
Severe (GCS 3 to 8)	49/107 (46%)	31/80 (39%)	.37
Clinical deterioration, No. (%), 0%	37/108 (34%)	44/82 (54%)	<.01
GCS score after deterioration, median (IQR), 5.2%	8 (4 – 13)	8 (5 – 13)	.45
Major extracranial injury, No. (%), 3.2%	12/102 (12%)	11/82 (13%)	.82
CT characteristics			
Thickness, median (IQR), mm, 5.8%	14 (9 – 18)	10 (7 – 14)	<.01
Midline shift, median (IQR), mm, 4.2%	12 (7 – 16)	8 (3 – 13)	<.01
Basal cisterns obliterated, No. (%), 4.2%	39/101 (39%)	24/81 (30%)	.22
Concomitant SAH, No. (%), 2.1%	24/104 (23%)	23/82 (28%)	.50
Concomitant contusion, No. (%), 2.6%	20/104 (19%)	25/81 (31%)	.08
Concomitant EDH, No. (%), 2.1%	7/104 (7%)	4/82 (5%)	.76
Concomitant SDH, No. (%), 2.1%	8/104 (8%)	11/82 (13%)	.23
Concomitant IVH, No. (%), 2.1%	2/104 (2%)	2/82 (2%)	1.00
Cranial fracture, No. (%), 2.1%	30/106 (28%)	35/80 (44%)	.03
Second CT scan, No. (%), 1.6%	23/105 (22%)	19/82 (23%)	.86
Radiologic deterioration, 2.4%	14/22 (64%) ^a	10/19 (53%) ^a	.54
Predicted unfavorable outcome based on CRASH-CT variables (GOS ≤ 3, %), mean (IQR), 12.6%	56% (42% – 75%)	53% (32% – 70%)	.18

Table 1. Clinical and radiological variables for the entire study population (continued)

Variable, % <i>missing</i>	Region A (N=108)	Region B (N=82)	p-value
^a percentage of all second CT scans			
IQR, interquartile range; No., Number; GCS, Glasgow Coma Scale; TBI, Traumatic Brain Injury; CT, Computer Tomography; SAH, subarachnoid hemorrhage; EDH, epidural hematoma; SDH, subdural hematoma; IVH, intraventricular hemorrhage; GOS, Glasgow Outcome Score			

INTERVENTIONS

Patients in region A more often received emergency surgery (84% vs. 65%, $p < .01$) or an ICP sensor (38% vs. 15%, $p < .01$). When undergoing surgery, patients were more likely to undergo DC in region B (32% vs. 55%, $p < .01$). The median size of DC in region B was somewhat larger (110mm vs. 116mm, $p = .15$). Both regions had a similar rate of delayed procedures (15% vs 16%, $p = .57$) and complications (table 2).

Table 2. Hospital course variables for the entire study population

Variable, % <i>missing</i>	Region A (N=108)	Region B (N=82)	p-values
ICP monitor placed, No. (%), 1.1%	40/106 (38%)	12/82 (15%)	<.01
Emergency surgery, 0%			
<i>Total</i> , No. (%)	91/108 (84%)	53/82 (65%)	<.01
Craniotomy, No. (%)	60/91 (66%) ^a	24/53 (45%) ^a	.02
DC, No. (%)	29/91 (32%) ^a	29/53 (55%) ^a	<.01
Other, No. (%)	2/91 (2%) ^a	0/53 (0%) ^a	.53
Size of DC, mdn (IQR)	110 (97 – 115)	116 (103.5 – 127)	.15
Delayed surgery, No. (%), 2.6%			
<i>Total</i>	16/103 (15%)	16/82 (19%)	.57
Craniotomy	4/16 (25%) ^a	7/15 (44%) ^a	.46
DC	6/16 (38%) ^a	4/15 (25%) ^a	.70
Burrhole	4/16 (25%) ^a	3/15 (19%) ^a	1.00
Other	2/16 (13%) ^a	2/15 (13%) ^a	1.00
Complications, No. (%), 7.4%			
<i>Total</i>	56/95 (59%)	50/81 (62%)	.76
Intracranial	30/95 (32%)	29/81 (36%)	.63
Cardiovascular	13/95 (14%)	17/81 (21%)	.23
Respiratory	4/95 (4%)	9/81 (11%)	.15
Metabolic	3/95 (3%)	9/81 (11%)	.07
Infection	30/95 (32%)	18/81 (22%)	.18
Other	1/100 (1%)	7/81 (9%)	.02
^a Percentage within group of emergency or delayed procedures			
ICP, intracranial pressure; mdn, median; No., Number; IQR, interquartile range; DC, decompressive craniectomy			

Table 3. Output of logistic regression analyses. Numbers are odds ratios and 95% confidence intervals for the effect of treatment strategy on mortality and unfavorable outcome (Glasgow Outcome Score ≤ 3)

Analysis type	Mortality at discharge	Unfavorable outcome at 3-9 months
All patients		
Univariable (N=190), OR (95% CI)	0.75 (0.42 to 1.10)	0.74 (0.41 to 1.02)
Multivariable (N=190), OR (95% CI) ^a	0.43 (0.21 to 0.88)	0.53 (0.27 to 1.02)
Subgroup of mild-moderate TBI patients		
Univariable (N=110), OR (95% CI)	0.51 (0.22 to 1.18)	0.45 (0.21 to 0.96)
Multivariable (N=110), OR (95% CI) ^b	0.34 (0.13 to 0.91)	0.33 (0.14 to 0.77)

^a controlling for age, GCS score, pupillary reactivity and hematoma thickness
^b controlling for age, GCS score and hematoma thickness
OR, odds ratio; CI, confidence interval; TBI, traumatic brain injury; GCS, Glasgow coma scale

THE EFFECT OF TREATMENT STRATEGY

Hospital mortality in region A was 37%, compared to 45% in region B ($p=.29$). Also, in region A 53% had an unfavorable outcome after 3-9 months, compared to 62% in region B ($p=.23$). Region showed a favorable effect of a more aggressive treatment strategy (region A) in both unadjusted (mortality OR, 0.75; 95% CI, 0.42 to 1.10; unfavorable outcome OR, 0.74; 95% CI, 0.41 to 1.02) and adjusted analysis (mortality OR, 0.43; 95% CI, 0.21 to 0.88; unfavorable outcome OR, 0.53; 95% CI, 0.27 to 1.02). Subgroup analysis restricted to patients with mild to moderate TBI (N=110) showed a similar effect for both mortality (adjusted OR, 0.34; 95% CI, 0.13 to 0.91) and unfavorable outcome (adjusted OR, 0.33; 95% CI 0.14 to 0.77) (table 3). A post-hoc analysis with adjustment for decompressive craniectomy and primary referral confirmed the main results (*supplementary tables 1 and 2*).

THE EFFECT OF SURGERY ON PATIENT LEVEL

In multivariable analysis, surgery was associated with higher odds of in-hospital mortality (OR, 6.6; 95% CI, 1.43 to 30.8) and unfavorable outcome (OR, 1.56, 95% CI: 0.64 to 3.82, table 4).

Table 4. Output of logistic regression analyses. Numbers are odds ratio's and 95% confidence intervals for the effect of surgery on mortality and unfavorable outcome (Glasgow Outcome Score ≤ 3)

Analysis type	Mortality at discharge	Unfavorable outcome at 3-9 months
Effect of surgery on patient level		
Univariable, OR (95% CI)	9.7 (3.3 to 28.6)	3.5 (1.7 to 7.3)
Multivariable, OR (95% CI) ^a	6.6 (1.4 to 30.8)	1.6 (0.6 to 3.8)

^a controlling for age, GCS score, pupillary reactivity and hematoma thickness

DISCUSSION

This study compared an aggressive surgical policy to a more conservative management strategy for traumatic ASDH. Though the study groups were comparable in terms of baseline prognosis, patients were significantly more likely to undergo emergency surgery in the expected region. A primarily surgically focused strategy was associated with a lower in-hospital mortality and unfavorable outcome.

Estimating causal treatment effects in non-randomized data is often impossible due to confounding by indication. Some of the baseline imbalance can be corrected through stratification or multivariable modeling (amongst other selection or analysis techniques). However, when we compared a surgical with a non-operative treatment on patient level in this study (table 4), in contrast to comparing treatment strategies on a center level, emergency surgery was associated with worse outcome even after adjusting for confounders. This suggests unmeasured confounding by indication, causing misleading effect estimates. The region-based comparison is a major strength of this study. It allowed for a 'natural experiment' where patients were 'allocated' to one of both policies based on where the accident occurred and not based on their baseline characteristics. This provided an opportunity for an analysis that likely has a considerable reduction of unmeasured confounding compared to an analysis comparing the actual treatment received. Based on the balance between groups with regard to measured confounders, we may speculate that there is also balance in unmeasured confounders. This assumption, however, cannot be statistically proven (as would neither be possible or rational in an RCT).

A weakness is that we compared surgical strategy in only two centers. It cannot be excluded that differences between regions in other aspects of care, e.g. other treatments or organization of care, caused the difference in outcome in addition to the surgical approach. To account for these differences in such a hospital-level approach more centers are generally needed to disentangle the care aspects that correlate with the treatment decision of interest. Therefore, we need to interpret our results with caution. The wide confidence interval of the effect estimates clearly underwrites this statement even more. A study with more hospitals is needed to allow adjustment for differences between the hospitals other than in the treatment of interest.

An obvious limitation in this context, although inherent to the observational nature, is that the region-based approach did not completely eliminate all differences in baseline confounders. Patients from region A presented directly to the study hospital more often (63% vs. 21%, $p < .01$). Primary presentation to a neurosurgical center improves patient outcome and has a close relationship with time to surgery,³² both of which may have contributed to the lower rates of mortality and unfavorable outcome observed in region A. The higher incidence of clinical deterioration in region B may

be consequent to the larger number of secondary referrals. On the other hand, region B is characterized by relatively smaller hematoma thickness and midline shift (on the same time point CT), while having a higher proportion of accompanying contusions. However, the possible resulting confounding is adjusted for appropriately. Moreover, apparently the imbalances are evenly spread out because the cohorts of both regions have a similar prognosis in the validated CRASH model. So they were similar for the most important predictors of outcome on arrival to the study hospital.

Another limitation may be that the data collectors could not be blinded for hospital of admission, which might give rise to information bias. To minimize potential bias, a standardized and controlled data collection protocol was followed. Nevertheless, we need to consider that the GOS was estimated from clinic discharge and visit letters, instead of a structured interview.

Furthermore, this study needs an important note on generalizability. This is a relatively old population, with a high proportion of fall-related ASDH (56%). Recent comparable series of patients with ASDH of varying severities report mean ages of 55 to 58 years.^{19,33,34} The high age in this study should come as no surprise considering the aging population in developed countries, but should be kept in mind when applying its results. Also, for a clinician to determine to which patients these results are relevant she/he needs to consider for what kind of ASDH patients the neurosurgery is consulted in the Netherlands. An unknown number of mild cases reviewed by the neurologist or neurology resident without consultation of a neurosurgeon were not included. This explains the relatively small number of conservatively treated patients (24% overall) in our cohort compared to recent studies, reporting 74-83% of ASDHs being treated nonoperatively.^{33,34} And, though much less common, this also applies to very severe cases, not operated due to an extremely poor prognosis. Because the mild cases constitute the majority of TBI, and this patient category undoubtedly has an above-average prognosis, applying the favorable results of surgery to all ASDH patients would not be reasonable. While both the age and consultation patterns might influence the generalizability, we nonetheless consider our selection useful in research on surgical decision making, because our cohort is representative of the ASDH population currently presented to neurosurgeons.

THE ROLE OF SURGERY FOR ASDH

Our results are in line with other recent studies reporting a positive effect of early/aggressive surgical management, mainly for severe TBI on mortality and functional outcome.^{15,35} A recent report showed this could also be true for routine use of invasive ICP measurement for severe TBI.¹⁴ On the other hand, when restricting to surgery, and the complete patient domain, i.e. ASDH patients with mild to severe TBI, previous studies have reported worse or similar outcomes in patients undergoing surgical

treatment.^{24-26,32,36,37} However, all studies compared outcomes between surgically and non-operatively treated patients at the patient level rather than a regional level, potentially leading to confounding by indication, because more severely injured patients are more likely to be operated. As mentioned above, when performing a conventional analysis on our cohort, surgery was associated with increased mortality and unfavorable outcome in our data. Hence, the distorting effect of confounding by indication in TBI supports the use of a comparative effectiveness approach when an RCT is not feasible for whatever reason.

When discussing the role of surgery in TBI, the use of decompressive craniectomy cannot be left unmentioned. Region B used DC significantly more often, despite equal distribution of prognostic factors. This could reflect another variation in surgical strategy or be consequent to the higher number of secondary referrals, resulting in presentation at a later stage, when brain swelling has started to occur. There is limited evidence on the use of primary DC in the treatment of ASDH, and studies comparing craniotomy to DC are likely confounded by indication as well.³⁸⁻⁴⁰ Consequently, the effect of more frequent use of DC in region B on our outcome measures is uncertain. The planned RESCUE-ASDH trial will use an experimental comparative effectiveness design to clarify the value of DC as a primary treatment in severe TBI.¹¹

This study reports an example of the practice variation in surgical treatment of traumatic ASDH. Advantage can be taken of this variation for CER. Larger cohorts with more hospitals are required to perform robust analyses to explore the ability of this method to infer causality, and with enough statistical power to study specific substrata according to age and trauma severity. The availability of such cohorts, with prospectively gathered, trial quality data from real-world settings, will allow initiatives such as Net-QuRe and CENTER-TBI to answer urgent clinical questions to provide the much needed guidance for the treatment of TBI.⁴¹

CONCLUSION

In this comparative study of surgery for ASDH, an aggressive management strategy favoring emergency evacuation of the hematoma was associated with better outcome. This conclusion derives from a contemporary cohort of relatively old traumatic ASDH patients that reflects the current population presented to a neurosurgeon. Larger, prospective, comparative studies with more hospitals are needed to confirm this effect of surgery and to explore generalizability.

ACKNOWLEDGEMENTS

FUNDING/SUPPORT

The European Union seventh Framework Program (grant 602150) for Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI). Hersenstichting Nederland (Dutch Brain Foundation, project PS2014-06) for Neurotraumatology Quality Registry (Net-QuRe). No industry affiliation.

Portions of this work were presented in an oral presentation as a plenary lecture at the 12th Symposium of the International Neurotrauma Society, Capetown, South Africa, February 1-4, 2016.

AUTHOR DISCLOSURE STATEMENT

No competing financial interests exist.

Supplemental material available online: <https://www.liebertpub.com/doi/suppl/10.1089/neu.2018.5869>

Tables S1 and S2

REFERENCES

1. Gelpke GJ, Braakman R, Habbema, JD, Hilden J. Comparison of outcome in two series of patients with severe head injuries. *J Neurosurg* 1982; 59, 745–750.
2. Timmons SD, Toms SA. Comparative effectiveness research in neurotrauma. *Neurosurg FOCUS* 2012 33, E3.
3. Maas AI, Menon DK, Lingsma HF, et al. Re-orientation of clinical research in traumatic brain injury: report of an international workshop on comparative effectiveness research. *J Neurotrauma* 29, 32–46.
4. Hutchinson, P.J., Cordeen, E., Czosnyka, M., et al. Decompressive craniectomy in traumatic brain injury: the randomized multicenter RESCUEicp study. *Acta Neurochir Suppl* 96, 17–20.
5. Chesnut RM, Temkin N, Carney N, et al. A trial of intracranial-pressure monitoring in traumatic brain injury. *N Engl J Med* 2012; 367, 2471–2481.
6. Gregson BA, Rowan EN, Francis R, et al. Surgical Trial In Traumatic intraCerebral Haemorrhage (STITCH): a randomised controlled trial of Early Surgery compared with Initial Conservative Treatment. *Heal Technol Assess* 19, 1–138.
7. Cooper DJ, Rosenfeld JV, Murray L, et al. Decompressive craniectomy in diffuse traumatic brain injury. *N Engl J Med* 2011; 364, 1493–1502.
8. Maas AIR, Roozenbeek B, Manley GT. Clinical trials in traumatic brain injury: past experience and current developments. *Neurotherapeutics* 2010; 7, 115–126.
9. Toliaas CM, Bullock MR. Critical appraisal of neuroprotection trials in head injury: what have we learned? *NeuroRx* 2004. 1, 71–79.
10. Koliass AG, Belli A, Li LM, Timofeev I, et al. Primary decompressive craniectomy for acute subdural haematomas: results of an international survey. *Acta Neurochir* 2012; 154, 1563–1565.
11. Koliass AG, Scotton WJ, Belli A, et al. Surgical management of acute subdural haematomas: current practice patterns in the United Kingdom and the Republic of Ireland. *Br J Neurosurg* 2013; 27, 330–333.
12. Van Essen TA, De Ruiter GCW, et al. Neurosurgical Treatment Variation of Traumatic Brain Injury: Evaluation of Acute Subdural Hematoma Management in Belgium and The Netherlands. *J Neurotrauma* 2017; 34, 881–889.
13. Dawes AJ, Sacks GD, Cryer HG, et al Compliance With Evidence-Based Guidelines and Interhospital Variation in Mortality for Patients With Severe Traumatic Brain Injury. *JAMA Surg* 2005; 150, 965–972.
14. Alali AS, Fowler RA, Mainprize TG., et al. Intracranial pressure monitoring in severe traumatic brain injury: results from the American College of Surgeons Trauma Quality Improvement Program. *J Neurotrauma* 2013; 30, 1737–1746.
15. Hartings JA, Vidgeon S, Strong AJ, et al. Surgical management of traumatic brain injury: a comparative-effectiveness study of 2 centers. *J Neurosurg* 2014; 120, 434–446.
16. Compagnone C, Murray GD, Teasdale, et al. The management of patients with intradural post-traumatic mass lesions: a multicenter survey of current approaches to surgical management in 729 patients coordinated by the European Brain Injury Consortium. *Neurosurgery* 2005; 57, 1183–1192.
17. Seelig JM, Becker DP, Miller JD, et al. Traumatic acute subdural hematoma: major mortality reduction in comatose patients treated within four hours. *N Engl J Med* 1981; 304, 1511–1518.
18. Wilberger Jr JE, Harris M, Diamond DL. Acute subdural hematoma: morbidity, mortality, and operative timing. *J Neurosurg* 1991 74, 212–218.
19. Leitgeb J, Mauritz W, Brazinova A, et al. (2012). Outcome after severe brain trauma due to acute subdural hematoma. *J Neurosurg* 2012; 117, 324–333.
20. Matsushima K, Inaba K, Siboni S, et al. Emergent operation for isolated severe traumatic brain injury: Does time matter? *J Trauma Acute Care Surg* 2015; 79, 838–842.
21. Bullock MR, Chesnut R, Ghajar J, et al. Surgical Management of Traumatic Brain Injury Surgical management of acute subdural hematomas. *Neurosurgery* 2006 58, S16–24; discussion Si–iv.
22. Scotter J, Hendrickson S, Marcus HJ, Wilson MH. Prognosis of patients with bilateral fixed dilated pupils secondary to traumatic extradural or subdural haematoma who undergo surgery: a systematic review and meta-analysis. *Emerg Med J* 2015; 32, 654–659.
23. Nijboer JMM, Van der Naalt J, Ten Duis HJ. Patients beyond salvation?. Various categories of trauma patients with a minimal Glasgow Coma Score. *Injury* 2010 41, 52–57.

24. Croce MA, Dent DL, Menke PG, et al. Acute subdural hematoma: nonsurgical management of selected patients. *J Trauma* 1994 36, 820–827.
25. Servadei F, Nasi MT, Cremonini AM, et al. Importance of a reliable admission Glasgow Coma Scale score for determining the need for evacuation of posttraumatic subdural hematomas: a prospective study of 65 patients. *J Trauma* 1998 44, 868–873.
26. Feliciano CE, De Jesus O. Conservative management outcomes of traumatic acute subdural hematomas. *P R Heal. Sci J* 2008 27, 220–223.
27. Mathew P, Oluoch-Olunya DL, Condon BR, Bullock R. Acute subdural haematoma in the conscious patient: outcome with initial non-operative management. *Acta Neurochir* 1993 121, 100–108.
28. Von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *Int J Surg* 2014; 12, 1495–1499.
29. Goslings JC, Gouma DJ. [Trauma registration. A promising, but difficult process]. *Ned Tijdschr Geneesk* 2009 153, 744–746.
30. Jennett B, Bond M. Assessment of outcome after severe brain damage. *The Lancet* 1975; 1, 480–484.
31. Perel P, Arango M, Clayton T, et al. Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. *BMJ* 336, 425–429.
32. Stone JL, Lowe RJ, Jonasson O, et al. Acute subdural hematoma: direct admission to a trauma center yields improved results. *J Trauma* 1986 26, 445–450.
33. Bajsarowicz P, Prakash I, Lamoureux J, et al. Nonsurgical acute traumatic subdural hematoma: what is the risk? *J Neurosurg* 2015 123, 1–8.
34. Ryan CG, Thompson RE, Temkin NR, et al. Acute traumatic subdural hematoma: current mortality and functional outcomes in adult patients at a Level I trauma center. *J Trauma Acute Care Surg* 2012 73, 1348–1354.
35. Hedges JR, Newgard CD, Veum-Stone J, et al. Early neurosurgical procedures enhance survival in blunt head injury: propensity score analysis. *J Emerg Med* 2009 37, 115–123.
36. Dent DL, Croce MA, Menke PG, et al. Prognostic factors after acute subdural hematoma. *J Trauma* 1995 39, 33–36.
37. Tierney KJ, Nayak NV, Prestigiacomo CJ, Sifri ZC. Neurosurgical intervention in patients with mild traumatic brain injury and its effect on neurological outcomes. *J Neurosurg* 2015 124, 538–545.
38. Li LM, Koliass AG, Guilfoyle MR, et al. Outcome following evacuation of acute subdural haematomas: a comparison of craniotomy with decompressive craniectomy. *Acta Neurochir* 2012 154, 1555–1561.
39. Chen SH., Chen Y, Fang WK, et al. Comparison of craniotomy and decompressive craniectomy in severely head-injured patients with acute subdural hematoma. *J Trauma* 2011; 71, 1632–1636.
40. Woertgen C, Rothoerl RD, Schebesch KM, Albert R. Comparison of craniotomy and craniectomy in patients with acute subdural haematoma. *J Clin Neurosci* 2006 13, 718–721.
41. Maas AIR, Menon DK, Steyerberg EW, et al. Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI): a prospective longitudinal observational study. *Neurosurgery* 2015; 76, 67–80.

