



Universiteit  
Leiden  
The Netherlands

## On topological properties of massless fermions in a magnetic field

Lemut, G.

### Citation

Lemut, G. (2023, June 13). *On topological properties of massless fermions in a magnetic field*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/3620153>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3620153>

**Note:** To cite this publication please use the final published version (if applicable).

# Bibliography

- [1] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, *The electronic properties of graphene*, Rev. Mod. Phys. **81**, 109 (2009).
- [2] M. I. Katsnelson, *Graphene: carbon in two dimensions*, materialstoday **10**, 20 (2007).
- [3] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, *Theory of Superconductivity*, Phys. Rev. **108**, 1175 (1957).
- [4] E. Majorana, *Teoria simmetrica dell'elettrone e del positrone*, Il Nuovo Cimento **14**, 171-184 (2010).
- [5] J. K. Asbóth, L. Oroszlány and A. Palyi, *A Short Course on Topological Insulators*, Lect. Notes Phys **919**, (2016).
- [6] T. Meng and L. Balents, *Weyl superconductors*, Phys. Rev. B **86**, 054504 (2012). *Erratum*: Phys. Rev. B **96**, 019901 (2017).
- [7] G. Bednik, A. A. Zyuzin, and A. A. Burkov, *Superconductivity in Weyl metals*, Phys. Rev. B **92**, 035153 (2015).
- [8] Liang Fu and C. L. Kane, *Superconducting proximity effect and Majorana fermions at the surface of a topological insulator*, Phys. Rev. Lett. **100**, 096407 (2008).
- [9] G. Chang, B. J. Wieder, F. Schindler, D. S. Sanchez, I. Belopolski, S.-M. Huang, B. Singh, D. Wu, T. Neupert, T.-R. Chang, S.-Y. Xu, H. Lin, and M. Z. Hasan, *Topological quantum properties of chiral crystals*, Nature Mat. **17**, 978 (2018).
- [10] C. Shekhar, *Chirality meets topology*, Nature Mat. **17**, 953 (2018).
- [11] M. Z. Hasan and C. L. Kane, *Colloquium: Topological insulators*, Rev. Mod. Phys. **82**, 3045 (2010).
- [12] Xiao-Liang Qi and Shou-Cheng Zhang, *Topological insulators and superconductors*, Rev. Mod. Phys. **83**, 1057 (2011).

## Bibliography

- [13] B. Yan and C. Felser, *Topological Materials: Weyl Semimetals*, Annu. Rev. Condens. Matter Phys. **8**, 337 (2017).
- [14] L. D. Landau and E. M. Lifshitz *Quantum Mechanics: Non-Relativistic Theory*, Course of Theoretical Physics **3**, (1965).
- [15] A.A. Abrikosov, *The magnetic properties of superconducting alloys*, Journal of Physics and Chemistry of Solids **3**, 199 (1957).
- [16] A. S. Mel'nikov, *Quantization of the quasiparticle spectrum in the mixed state of d-wave superconductors*, J. Phys. Condens. Matter **11**, 4219 (1999).
- [17] L. Alvarez-Gaumé, *Supersymmetry and the Atiyah-Singer index theorem*, Communications in Mathematical Physics **90**, 161 (1983).
- [18] M. Filoche and S. Mayboroda, *Universal mechanism for Anderson and weak localization*, PNAS **109**, 14761 (2012).
- [19] M. Filoche and S. Mayboroda, *The landscape of Anderson localization in a disordered medium*, Contemp. Math. **601**, 113 (2013).
- [20] D. N. Arnold, G. David, D. Jerison, S. Mayboroda, and M. Filoche, *Effective confining potential of quantum states in disordered media*, Phys. Rev. Lett. **116**, 056602 (2016).
- [21] S. Steinerberger, *Localization of quantum states and landscape functions*, Proc. Amer. Math. Soc. **145**, 2895 (2017).
- [22] M. Filoche, M. Piccardo, Y.-R. Wu, C.-K. Li, C. Weisbuch, and S. Mayboroda, *Localization landscape theory of disorder in semiconductors. I. Theory and modeling*, Phys. Rev. B **95**, 144204 (2017).
- [23] M. Piccardo, C.-K. Li, Y.-R. Wu, J. S. Speck, B. Bonef, R. M. Farrell, M. Filoche, L. Martinelli, J. Peretti, and C. Weisbuch, *Localization landscape theory of disorder in semiconductors. II. Urbach tails of disordered quantum well layers*, Phys. Rev. B **95**, 144205 (2017).
- [24] C.-K. Li, M. Piccardo, L.-S. Lu, S. Mayboroda, L. Martinelli, J. Peretti, J. S. Speck, C. Weisbuch, M. Filoche, and Y.-R. Wu, *Localization landscape theory of disorder in semiconductors. III. Application to carrier transport and recombination in light emitting diodes*, Phys. Rev. B **95**, 144206 (2017).

- [25] Y. Chalopin, F. Piazza, S. Mayboroda, C. Weisbuch, and M. Filoche *Universality of fold-encoded localized vibrations in enzymes*, arXiv:1902.09939
- [26] D. Arnold, D. Guy, M. Filoche, D. Jerison, and S. Mayboroda, *Computing spectra without solving eigenvalue problems*, SIAM J. Scientific Comput. **41**, B69 (2019).
- [27] E. M. Harrell II and A. V. Maltsev, *Localization and landscape functions on quantum graphs*, arXiv:1803.01186.
- [28] C. W. J. Beenakker, *Hidden landscape of an Anderson insulator*, J. Club Cond. Matt. (August, 2019,
- [29] A. Ostrowski, *Über die Determinanten mit überwiegender Hauptdiagonale*, Comment. Mathemat. Helvet. **10**, 69 (1937). The comparison inequality is on page 71.
- [30] A. Ostrowski, *Determinanten mit überwiegender Hauptdiagonale und die absolute Konvergenz von linearen Iterationsprozessen*, Comment. Mathemat. Helvet. **30**, 175 (1956).
- [31] Ostrowski originally used the name “companion matrix” (*Begleitmatrix*), which now refers to a different construction. The notation  $\langle H \rangle$  for the comparison matrix of  $H$  is common in the mathematical literature, but to avoid confusion with the physics notation for expectation value we use  $\overline{H}$  instead. One more piece of nomenclature: If  $\overline{H}$  is positive-definite, then it is called an M-matrix while  $H$  is called an H-matrix.
- [32] For background on comparison matrices, see A. Berham and R. J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences* (SIAM, 1994).
- [33] The discretization (2.9) is appropriate near the bottom of the tight-binding band at  $p = 0$ . Near the top of the band at  $p = \pi/a$  a different discretization produces a different landscape function, as discussed by M. L. Lyra, S. Mayboroda, and M. Filoche, *Dual landscapes in Anderson localization on discrete lattices*, EPL **109**, 47001 (2015).
- [34] G. Giovannetti, P. A. Khomyakov, G. Brocks, P. J. Kelly, and J. van den Brink, *Substrate-induced band gap in graphene on hexagonal boron nitride*, Phys. Rev. B **76**, 073103 (2007).

## Bibliography

- [35] R. Stacey, *Eliminating lattice fermion doubling*, Phys. Rev. D **26**, 468 (1982).
- [36] J. Tworzydło, C. W. Groth, and C. W. J. Beenakker, *Finite difference method for transport properties of massless Dirac fermions*, Phys. Rev. B **78**, 235438 (2008).
- [37] L. Susskind, *Lattice fermions*, Phys. Rev. D **16**, 3031 (1977).
- [38] A. R. Hernández and C. H. Lewenkopf, *Finite-difference method for transport of two-dimensional massless Dirac fermions in a ribbon geometry*, Phys. Rev. B **86**, 155439 (2012).
- [39] C. W. J. Beenakker and L. P. Kouwenhoven, *A road to reality with topological superconductors*, Nature Physics **12**, 618 (2016).
- [40] N. Read and D. Green, *Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect*, Phys. Rev. B **61**, 10267 (2000).
- [41] M. Wimmer, A. R. Akhmerov, M. V. Medvedyeva, J. Tworzydło, and C. W. J. Beenakker, *Majorana bound states without vortices in topological superconductors with electrostatic defects*, Phys. Rev. Lett. **105**, 046803 (2010).
- [42] A. F. Tzortzakakis, K. G. Makris, and E. N. Economou, *Non-Hermitian disorder in two-dimensional optical lattices*, Phys. Rev. B **101**, 014202 (2020).
- [43] Yi Huang and B. I. Shklovskii, *Anderson transition in three-dimensional non-Hermitian disorder*, arXiv:1911.00562.
- [44] D. S. Wiersma, *The physics and applications of random lasers*, Nature Phys. **4**, 359 (2008).
- [45] F. Evers and A. D. Mirlin, *Anderson transitions*, Rev. Mod. Phys. **80**, 1355 (2008).
- [46] S. Balasubramanian, Y. Liao, and V. Galitski, *Many-body localization landscape*, Phys. Rev. B **101**, 014201 (2020).
- [47] C. W. Groth, M. Wimmer, A. R. Akhmerov, and X. Waintal, *Kwant: A software package for quantum transport*, New J. Phys. **16**, 063065 (2014).

- [48] Zhicheng Rao, Hang Li, Tiantian Zhang, Shangjie Tian, Chenghe Li, Binbin Fu, Cenyao Tang, Le Wang, Zhilin Li, Wenhui Fan, Jiajun Li, Yaobo Huang, Zhehong Liu, Youwen Long, Chen Fang, Hongming Weng, Youguo Shi, Hechang Lei, Yujie Sun, Tian Qian, and Hong Ding, *Observation of unconventional chiral fermions with long Fermi arcs in CoSi*, Nature **567**, 496 (2019).
- [49] D. S. Sanchez, I. Belopolski, T. A. Cochran, X. Xu, J.-X. Yin, G. Chang, W. Xie, K. Manna, V. Süß, C.-Y. Huang, N. Alidoust, D. Multer, S. S. Zhang, N. Shumiya, X. Wang, G.-Q. Wang, T.-R. Chang, C. Felser, S.-Y. Xu, S. Jia, H. Lin, M. Z. Hasan, *Topological chiral crystals with helicoid-arc quantum states*, Nature **567**, 500 (2019).
- [50] D. Takane, Z. Wang, S. Souma, K. Nakayama, T. Nakamura, H. Oinuma, Y. Nakata, H. Iwasawa, C. Cacho, T. Kim, K. Horiba, H. Kumigashira, T. Takahashi, Y. Ando, T. Sato, *Observation of chiral fermions with a large topological charge and associated Fermi-arc surface states in CoSi*, Phys. Rev. Lett. **122**, 076402 (2019).
- [51] Qian-Qian Yuan, Liqin Zhou, Zhi-Cheng Rao, Shangjie Tian, Wei-Min Zhao, Cheng-Long Xue, Yixuan Liu, Tiantian Zhang, Cen-Yao Tang, Zhi-Qiang Shi, Zhen-Yu Jia1, Hongming Weng, Hong Ding, Yu-Jie Sun, Hechang Lei, and Shao-Chun Li, *Quasiparticle interference evidence of the topological Fermi arc states in chiral fermionic semimetal CoSi*, Science Adv. **5**, eaaw9485 (2019).
- [52] N. B. M. Schröter, D. Pei, M. G. Vergniory, Y. Sun, K. Manna, F. de Juan, J. A. Krieger, V. Süß, M. Schmidt, P. Dudin, B. Bradlyn, T. K. Kim, T. Schmitt, C. Cacho, C. Felser, V. N. Strocov, and Y. Chen, *Chiral topological semimetal with multifold band crossings and long Fermi arcs*, Nature Phys. **15**, 759 (2019).
- [53] C.-L. Zhang, F. Schindler, H. Liu, T.-R. Chang, S.-Y. Xu, G. Chang, W. Hua, H. Jiang, Z. Yuan, J. Sun, H.-T. Jeng, H.-Z. Lu, H. Lin, M. Z. Hasan, X. C. Xie, T. Neupert, and S. Jia, *Ultraquantum magnetoresistance in Kramers Weyl semimetal candidate  $\beta$ -Ag<sub>2</sub>Se*, Phys. Rev. B **96**, 165148 (2017).
- [54] B. Wan, F. Schindler, K. Wang, K. Wu, X. Wan, T. Neupert, and H.-Z. Lu, *Theory for the negative longitudinal magnetoresistance in the quantum limit of Kramers Weyl semimetals*, J. Phys. Condens. Matter **30**, 505501 (2018).

## Bibliography

- [55] Wen-Yu He, Xiao Yan Xu, K. T. Law, *Kramers Weyl semimetals as quantum solenoids and their applications in spin-orbit torque devices*, arXiv:1905.12575.
- [56] R. R. Gerhardts, D. Weiss, and K. von Klitzing, *Novel magnetoresistance oscillations in a periodically modulated two-dimensional electron gas*, Phys. Rev. Lett. **62**, 1173 (1989).
- [57] R. W. Winkler, J. P. Kotthaus, and K. Ploog, *Landau band conductivity in a two-dimensional electron system modulated by an artificial one-dimensional superlattice potential*, Phys. Rev. Lett. **62**, 1177 (1989).
- [58] C. W. J. Beenakker, *Guiding-center-drift resonance in a periodically modulated two-dimensional electron gas*, Phys. Rev. Lett. **62**, 2020 (1989).
- [59] P. Štředa and A. H. MacDonald, *Magnetic breakdown and magnetoresistance oscillations in a periodically modulated two-dimensional electron gas*, Phys. Rev. B **41**, 11892 (1990).
- [60] V. M. Gvozdkov, *Magnetoresistance oscillations in a periodically modulated two-dimensional electron gas: The magnetic-breakdown approach*, Phys. Rev. B **75**, 115106 (2007).
- [61] A. C. Potter, I. Kimchi, and A. Vishwanath, *Quantum oscillations from surface Fermi-arcs in Weyl and Dirac semi-metals*, Nature Comm. **5**, 5161 (2014).
- [62] Y. Zhang, D. Bulmash, P. Hosur, A. C. Potter, and A. Vishwanath, *Quantum oscillations from generic surface Fermi arcs and bulk chiral modes in Weyl semimetals*, Sci. Rep. **6**, 23741 (2016).
- [63] A. R. Akhmerov and C. W. J. Beenakker, *Boundary conditions for Dirac fermions on a terminated honeycomb lattice*, Phys. Rev. B **77**, 085423 (2008).
- [64] M. Z. Hasan, S.-Y. Xu, I. Belopolski, and S.-M. Huang, *Discovery of Weyl fermion semimetals and topological Fermi arc states*, Annu. Rev. Condens. Matter Phys. **8**, 289(2017).
- [65] A. A. Burkov, *Weyl Metals*, Annu. Rev. Condens. Matter Phys. **9**, 359 (2018).

- [66] N. P. Armitage, E. J. Mele, and A. Vishwanath, *Weyl and Dirac semimetals in three-dimensional solids*, Rev. Mod. Phys. **90**, 15001 (2018).
- [67] N. Bovenzi, M. Breitzkreiz, T. E. O'Brien, J. Tworzydło, and C. W. J. Beenakker, *Twisted Fermi surface of a thin-film Weyl semimetal*, New J. Phys. **20**, 023023 (2018).
- [68] V. Barsan and V. Kuncser, *Exact and approximate analytical solutions of Weiss equation of ferromagnetism and their experimental relevance*, Phil. Mag. Lett. **97**, 359 (2017).
- [69] A. B. Pippard, *Magnetic breakdown*, in: *Physics of Solids in Intense Magnetic Fields* (Springer, Boston, 1969).
- [70] M. I. Kaganov and A. A. Slutskin, *Coherent magnetic breakdown*, Phys. Rep. **98**, 189 (1983).
- [71] R. W. Stark and L. M. Falicov, *Magnetic breakdown in metals*, Prog. Low Temp. Phys **5**, 235 (1967).
- [72] V. M. Gvozdkov, *Thermodynamic oscillations in periodic magnetic breakdown structures*, Fiz. Nizk. Temp. **12**, 705 (1986).
- [73] A. Alexandradinata and L. Glazman, *Geometric phase and orbital moment in quantization rules for magnetic breakdown*, Phys. Rev. Lett. **119**, 256601 (2017).
- [74] M. Breitzkreiz, N. Bovenzi, and J. Tworzydło, *Phase shift of cyclotron orbits at type-I and type-II multi-Weyl nodes*, Phys. Rev. B **98**, 121403 (2018).
- [75] E. P. Wigner, *Random matrices in physics*, SIAM Rev. **9**, 1 (1967).
- [76] F. Haake, S. Gnutzmann, and M. Kuś, *Quantum Signatures of Chaos* (Springer, Berlin, 2018).
- [77] F. J. Dyson, *The threefold way: Algebraic structure of symmetry groups and ensembles in quantum mechanics*, J. Math. Phys. **3**, 1199 (1962).
- [78] T. Guhr, A. Mueller-Groeling, and H. A. Weidenmüller, *Random matrix theories in quantum physics: common concepts*, Phys. Rep. **299**, 189 (1998).



## Bibliography

- [79] Ying-Cheng Lai, Hong-Ya Xu, Liang Huang, Celso Grebogi, *Relativistic quantum chaos — An emergent interdisciplinary field*, *Chaos* **28**, 052101 (2018).
- [80] M. Kieburg, J. J. M. Verbaarschot, and S. Zafeiropoulos, *Dirac spectra of two-dimensional QCD-like theories*, *Phys. Rev. D* **90**, 085013 (2014).
- [81] M. Kieburg and T. R. Würfel, *Shift of symmetries of naive and staggered fermions in QCD-like lattice theories*, *Phys. Rev. D* **96**, 034502 (2017).
- [82] C. W. J. Beenakker, *Random-matrix theory of quantum transport*, *Rev. Mod. Phys.* **69**, 731 (1997).
- [83] B. L. Altshuler, *Fluctuations in the extrinsic conductivity of disordered conductors*, *JETP Lett.* **41**, 648 (1985).
- [84] P. A. Lee and A. D. Stone, *Universal conductance fluctuations in metals*, *Phys. Rev. Lett.* **55**, 1622 (1985).
- [85] Our computer codes are available at a repository: [10.5281/zenodo.6394211](https://zenodo.org/record/6394211)
- [86] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, *Distribution of the ratio of consecutive level spacings in random matrix ensembles*, *Phys. Rev. Lett.* **110**, 084101 (2013).
- [87] R. Modak and S. Mukerjee, *Finite size scaling in crossover among different random matrix ensembles in microscopic lattice models*, *New J. Phys.* **16**, 093016 (2014).
- [88] H. B. Nielsen and M. Ninomiya, *A no-go theorem for regularizing chiral fermions*, *Phys. Lett. B* **105**, 219 (1981).
- [89] D. B. Kaplan, *Chiral symmetry and lattice fermions*, in: *Modern Perspectives in Lattice QCD: Quantum Field Theory and High Performance Computing*, Lecture Notes of the Les Houches Summer School (Oxford University Press, 2009).
- [90] N. P. Ong and Sihang Liang, *Review of experiments on the chiral anomaly in Dirac-Weyl semimetals*, *Nature Rev. Phys.* **3** 394 (2021).

- [91] M. Zahid Hasan, Guoqing Chang, Ilya Belopolski, Guang Bian, Su-Yang Xu, and Jia-Xin Yin, *Weyl, Dirac and high-fold chiral fermions in topological quantum matter*, arXiv:2103.01714, to appear in Nature Rev. Mat. (2021).
- [92] H. B. Nielsen and M. Ninomiya, *The Adler-Bell-Jackiw anomaly and Weyl fermions in a crystal*, Nucl. Phys. B **130**, 389 (1983).
- [93] D. Kharzeev, *The Chiral Magnetic Effect and anomaly-induced transport*, Progr. Part. Nucl. Phys. **75**, 133 (2014).
- [94] A. A. Burkov, *Chiral anomaly and transport in Weyl metals*, J. Phys. Condens. Matter **27**, 113201 (2015).
- [95] T. E. O'Brien, C. W. J. Beenakker, and I. Adagideli, *Superconductivity provides access to the chiral magnetic effect of an unpaired Weyl cone*, Phys. Rev. Lett. **118**, 207701 (2017).
- [96] M. J. Pacholski, C. W. J. Beenakker, and I. Adagideli, *Topologically protected Landau level in the vortex lattice of a Weyl superconductor*, Phys. Rev. Lett. **121**, 037701 (2018).
- [97] G. Lemut, M. J. Pacholski, I. Adagideli, and C. W. J. Beenakker, *Effect of charge renormalization on the electric and thermoelectric transport along the vortex lattice of a Weyl superconductor*, Phys. Rev. B **100**, 035417 (2019).
- [98] A. Donís Vela, G. Lemut, M. J. Pacholski, and C. W. J. Beenakker, *Chirality inversion of Majorana edge modes in a Fu-Kane heterostructure*, arXiv:2105.04433.
- [99] P. W. Anderson, *Anomalous magnetothermal resistance of high- $T_c$  superconductors: Anomalous cyclotron orbits at a Dirac point*, arXiv:cond-mat/9812063.
- [100] Z. Faraei and S. A. Jafari, *Superconducting proximity in three dimensional Dirac materials: odd-frequency, pseudoscalar, pseudovector and tensor-valued superconducting orders*, Phys. Rev. B **96**, 134516 (2017).
- [101] G. E. Volovik, *Quantum phase transitions from topology in momentum space*, Lect. Notes Phys. **718**, 31 (2007).

## Bibliography

- [102] D. F. Agterberg, P. M. R. Brydon, and C. Timm, *Bogoliubov Fermi surfaces in superconductors with broken time-reversal symmetry*, Phys. Rev. Lett. **118**, 127001 (2017).
- [103] Noah F. Q. Yuan and Liang Fu, *Zeeman-induced gapless superconductivity with a partial Fermi surface*, Phys. Rev. B **97**, 115139 (2018).
- [104] S. Autti, J. T. Mäkinen, J. Rysti, G. E. Volovik, V. V. Zavjalov, and V. B. Eltsov, *Exceeding the Landau speed limit with topological Bogoliubov Fermi surfaces*, Phys. Rev. Res. **2**, 033013 (2020).
- [105] J. M. Link and I. F. Herbut, *Bogoliubov-Fermi surfaces in non-centrosymmetric multi-component superconductors*, Phys. Rev. Lett. **125**, 237004 (2020).
- [106] Zhen Zhu, Michal Papa, Xiao-Ang Nie, Hao-Ke Xu, Yi-Sheng Gu, Xu Yang, Dandan Guan, Shiyong Wang, Yaoyi Li, Canhua Liu, Jianlin Luo, Zhu-An Xu, Hao Zheng, Liang Fu, and Jin-Feng Jia, *Discovery of segmented Fermi surface induced by Cooper pair momentum*, arXiv:2010.02216. For a commentary on this experiment, see DOI: 10.36471/JCCM\_October\_2020\_01
- [107] S. Rachel, L. Fritz, and M. Vojta, *Landau levels of Majorana fermions in a spin liquid*, Phys. Rev. Lett. **116**, 167201 (2016).
- [108] B. Perreault, S. Rachel, F. J. Burnell, and J. Knolle, *Majorana Landau-level Raman spectroscopy*, Phys. Rev. B **95**, 184429 (2017).
- [109] R. Jackiw and P. Rossi, *Zero modes of the vortex-fermion system*, Nucl. Phys. B **190**, 681 (1981).
- [110] The data in Fig. 6.2 is obtained from the tight-binding Hamiltonian (6.14) of the topological insulator layer. The parameters are  $\Delta_0 = 20 \hbar v / d_0$ ,  $d_0 = 302 a_0$ ,  $B = \hbar / e d_0^2$ ,  $\mu = 0$ ,  $M_0 = 0$ ,  $M_1 = 0.2 a_0$ . The vortex pair in a unit cell is at the positions  $(x, y) = (d_0/4)(1, 1)$  and  $(d_0/4)(3, 3)$ . The superflow momentum  $K$  equals  $0.8 \Delta_0 / v$  in the left panel and  $2 \Delta_0 / v$  in the right panel.
- [111] M. I. Katsnelson and M. F. Prokhorova, *Zero-energy states in corrugated bilayer graphene*, Phys. Rev. B **77**, 205424 (2008).
- [112] J. Kailasvuori, *Pedestrian index theorem à la Aharonov-Casher for bulk threshold modes in corrugated multilayer graphene*, EPL **87**, 47008 (2009).

- [113] Y. Aharonov and A. Casher, *Ground state of a spin-1/2 charged particle in a two-dimensional magnetic field*, Phys. Rev. A **19**, 2461 (1979).
- [114] M. Franz and Z. Tešanović, *Quasiparticles in the vortex lattice of unconventional superconductors: Bloch waves or Landau levels?*, Phys. Rev. Lett. **84**, 554 (2000).
- [115] Details of the solution of the Helmholtz equation are given in Apps. 6.B and 6.C of the Supplemental Material.
- [116] B. Sutherland, *Localization of electronic wave functions due to local topology*, Phys. Rev. B **34**, 5208 (1986).
- [117] V. M. Pereira, F. Guinea, J. M. B. Lopes-dos Santos, N. M. R. Peres, and A. H. Castro-Neto, *Disorder induced localized states in graphene*, Phys. Rev. Lett. **96**, 036801 (2006).
- [118] Wen-Yu Shan, Hai-Zhou Lu, and Shun-Qing Shen, *Effective continuous model for surface states and thin films of three-dimensional topological insulators*, New J. Phys. **12**, 043048 (2010).
- [119] Song-Bo Zhang, Hai-Zhou Lu, and Shun-Qing Shen, *Edge states and integer quantum Hall effect in topological insulator thin films*, Scientif. Rep. **5**, 13277 (2015).
- [120] Details of the method of numerical simulation, with supporting data, are given in App. 6.A of the Supplemental Material.
- [121] The renormalized charge  $q_{\text{eff}}$  in the Majorana Landau level is calculated in App. 6.D of the Supplemental Material. That calculation also gives the renormalized Fermi velocity  $v_{\text{eff}} = \sqrt{v_x v_y} = \sqrt{q_{\text{eff}}} v$  that appears in the Landau level energy  $E_L$ .
- [122] The chiral symmetry at  $\mu = 0$  is broken by the mass term  $M(k)$  in the Hamiltonian (6.14). This residual chiral symmetry breaking is visible in Fig. 6.3 as a very small splitting of the  $\mu = 0$  Landau levels (green flat bands).
- [123] The comparison between numerics and analytics in Fig. 6.4 involves no adjustable parameters. To compare the same state in the degenerate zeroth Landau level we choose the state with left-right reflection symmetry. There are two of these, the other is compared in App. 6.E of the Supplemental Material.

## Bibliography

- [124] Hao-Hua Sun and Jin-Feng Jia, *Detection of Majorana zero mode in the vortex*, npj Quantum Mat. **2**, 34 (2017).
- [125] S. Cho, B. Dellabetta, A. Yang, J. Schneeloch, Z. Xu, T. Valla, G. Gu, M. J. Gilbert, and N. Mason, *Symmetry protected Josephson supercurrents in three-dimensional topological insulators*, Nature Comm. **4**, 1689 (2013).
- [126] D. I. Pikulin and M. Franz, *Black hole on a chip: proposal for a physical realization of the SYK model in a solid-state system*, Phys. Rev. X **7**, 031006 (2017).  
 $K_n(r) = \frac{1}{2}(r/2)^n \int_0^\infty t^{-n-1} \exp(-t - \frac{1}{4}r^2/t) dt$ , upon the substitution  $p = \frac{1}{2}(t - 1/t)$ . <https://mathoverflow.net/q/379157/11260>.
- [127] N. B. Kopnin and M. M. Salomaa, *Mutual friction in superfluid  $^3\text{He}$ : Effects of bound states in the vortex core*, Phys. Rev. B **44**, 9667 (1991).
- [128] G. Volovik, *Fermion zero modes on vortices in chiral superconductors*, JETP Lett. **70**, 609 (1999).
- [129] S. Frolov, <https://espressospin.org/2012/04/18/zen-particle/>
- [130] C. W. J. Beenakker, *Search for Majorana fermions in superconductors*, Annu. Rev. Con. Mat. Phys. **4**, 113 (2013).
- [131] S. Das Sarma, M. Freedman, and C. Nayak *Majorana zero modes and topological quantum computation*, npj Quantum Inf. **1**, 15001 (2015) .
- [132] R. M. Lutchyn, E. P. A. M. Bakkers, L. P. Kouwenhoven, P. Krogstrup, C. M. Marcus, and Y. Oreg, *Majorana zero modes in superconductor–semiconductor heterostructures*, Nature Rev. Mat. **3**, 52 (2018).
- [133] P. Nozières and W. F. Vinen, *The motion of flux lines in type II superconductors*, Phil. Mag. **14**, 667 (1966) 10.1080/14786436608211964.
- [134] Yu. G. Makhlin and G. E. Volovik, *Spectral flow in Josephson junctions and effective Magnus force*, JETP Lett. **62**, 941 (1995) .
- [135] M. L. Stone, *Spectral flow, Magnus force, and mutual friction via the geometric optics limit of Andreev reflection*, Phys. Rev. B **54**, 13222 (1996).

- [136] E. B. Sonin *Magnus force in superfluids and superconductors*, Phys. Rev. B **55**, 485 (1997) 10.1103/PhysRevB.55.485.
- [137] M. J. Pacholski, G. Lemut, O. Ovdad, I. Adagideli, and C. W. J. Beenakker, *Deconfinement of Majorana vortex modes produces a superconducting Landau level*, Phys. Rev. Lett. **126**, 226801 (2021).
- [138] Zhen Zhu, Michał Papaż, Xiao-Ang Nie, Hao-Ke Xu, Yi-Sheng Gu, Xu Yang, Dandan Guan, Shiyong Wang, Yaoyi Li, Canhua Liu, Jianlin Luo, Zhu-An Xu, Hao Zheng, Liang Fu, and Jin-Feng Jia, *Discovery of segmented Fermi surface induced by Cooper pair momentum*, Science **374**, 1381 (2021).
- [139] M. Tinkham, *Introduction to Superconductivity* (Dover, New York, 2004).
- [140] D. Xiao, M.-C. Chang, and Q. Niu, *Berry phase effects on electronic properties*, Rev. Mod. Phys. **82**, 1959 (2010) 10.1103/RevModPhys.82.1959.
- [141] Zhi Wang, Liang Dong, Cong Xiao, and Qian Niu, *Berry curvature effects on quasiparticle dynamics in superconductors*, Phys. Rev. Lett. **126**, 187001 (2021).
- [142] J. R. Clem, *Simple model for the vortex core in a type II superconductor*, J. Low Temp. Phys. **18**, 427 (1975)
- [143] T. Kloss, J. Weston, B. Gaury, B. Rossignol, C. Groth, and X. Waintal, *Tkwant: a software package for time-dependent quantum transport*, New J. Phys. **23**, 023025 (2021).
- [144] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *Non-Abelian anyons and topological quantum computation*, Rev. Mod. Phys. **80**, 1083 (2008).
- [145] Hong-Hao Tu, Yi Zhang, and Xiao-Liang Qi, *Momentum polarization: An entanglement measure of topological spin and chiral central charge*, Phys. Rev. B **88**, 195412 (2013).
- [146] D. Ariad and E. Grosfeld, *Signatures of the topological spin of Josephson vortices in topological superconductors*, Phys. Rev. B **95**, 161401(R) (2017).