



Universiteit
Leiden
The Netherlands

Targets for improving patient outcomes after major gastrointestinal cancer surgery: the value of perioperative care

Kooten, R.T. van

Citation

Kooten, R. T. van. (2023, June 15). *Targets for improving patient outcomes after major gastrointestinal cancer surgery: the value of perioperative care*. Retrieved from <https://hdl.handle.net/1887/3620458>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3620458>

Note: To cite this publication please use the final published version (if applicable).

Part I: Identification of Prognostic Factors for Postoperative Complications

Chapter 2: Preoperative Risk Factors for Major Postoperative Complications After Complex Gastrointestinal Cancer Surgery: A Systematic Review

Chapter 3: Patient-Related Prognostic Factors for Anastomotic Leakage, Major Complications and Short-term Mortality Following Esophagectomy for Cancer: A Systematic Review and Meta-analysis

Chapter 4: Conventional Regression Analysis and Machine Learning in Prediction of Anastomotic Leakage and Pulmonary Complications after Esophagogastric Cancer Surgery

Chapter 5: Computer Tomography-based Preoperative Muscle Measurements as Prognostic Factors for Anastomotic Leakage Following Oncological Sigmoid and Rectal Resections



Chapter 2

Preoperative Risk Factors for Major Postoperative Complications After Complex Gastrointestinal Cancer Surgery: A Systematic Review

Robert T. van Kooten MD¹, Renu R. Bahadoer MD¹, Koen C.M.J. Peeters MD PhD¹, Jetty H.L. Hoeksema PhD¹, Ewout W. Steyerberg PhD², Henk H. Hartgrink MD PhD¹, Cornelis J.H. van de Velde MD PhD FACS¹, Michel W.J.M. Wouters MD PhD^{2,3} and Rob A.E.M. Tollenaar MD PhD¹

¹ Department of Surgery, Leiden University Medical Center, Leiden, The Netherlands

² Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

³ Department of Surgical Oncology, Netherlands Cancer Institute-Antoni van Leeuwenhoek, Amsterdam, The Netherlands

List of abbreviations

ACS NSQIP; American College of Surgeons National Surgical Quality Improvement Program, CD; Clavien-Dindo, CRP; C-Reactive Protein, DICA; Dutch Institute for Clinical Auditing, ERAS; Enhanced Recovery After Surgery

Abstract

Patients undergoing complex gastrointestinal surgery are at high risk of major postoperative complications (e.g., anastomotic leakage, sepsis), classified as Clavien-Dindo (CD) $\geq$ IIIa. Identification of preoperative risk factors can lead to the identification of high-risk patients. These risk factors can also be used to design personalized perioperative care. This systematic review focuses on the identification of these factors. The Medline and Embase databases were searched for prospective, retrospective cohort studies and randomized controlled trials investigating the effect of risk factors on the occurrence of major postoperative complications and/or mortality after complex gastrointestinal cancer surgery. Risk of bias was assessed using the Quality in Prognostic Studies tool. The level of evidence was graded based on the number of studies reporting a significant association between risk factors and major complications. A total of 207 eligible studies were retrieved, identifying 33 risk factors for major postoperative complications and 13 preoperative laboratory results associated with postoperative complications. The present systematic review provides a comprehensive overview of preoperative risk factors associated with major postoperative complications. A wide range of risk factors are amenable to actions in perioperative care and prehabilitation programs, which may lead to improved outcomes for high-risk patients. Additionally, the knowledge of this study is important for benchmarking surgical outcomes.

Introduction

Postoperative complications can occur after every type of surgery, and can lead to increased morbidity and mortality, as well as increased hospital length of stay and healthcare costs [1]. Complex gastrointestinal surgery (e.g., colorectal, gastric and esophagus resections) is associated with high complication rates [2, 3]. A large number of studies have focused on reducing complications by improving surgical techniques; however, relatively few have addressed improving perioperative care. The latter contributes largely to the avoidance of complications and is responsible for shorter recovery time after surgery, together with less morbidity and increased survival. Some studies have suggested that perioperative care more accurately dictates outcomes and

postoperative complications than surgery itself [4]. Perioperative care is currently being standardized in the form of enhanced recovery after surgery (ERAS) protocols, which provide guidelines for improved perioperative care. A meta-analysis by Vardhan et al. showed that the use of ERAS protocols reduces the rate of complications following major abdominal surgery by up to 50% [5]. The period of time before admission is used for screening for medical conditions that can negatively alter the surgical outcome (e.g., smoking and malnutrition). This can be particularly beneficial when the screening focuses on modifiable risk factors, which subsequently can be (partially) reversed (e.g., physical therapy, nutritional support).

Reduction of postoperative complications is also important in relation to long-term outcomes, especially in patients with cancer. The severity of complications is often graded using the Clavien-Dindo (CD) classification, a therapy-based complication severity classification [6]. It has been demonstrated that major complications (CD $\geq$ IIIa) are associated with postponement of adjuvant therapy and worse oncological outcomes, like local recurrences and shortened recurrence-free survival [7, 8]. The majority of studies addressing the prevention of postoperative complications have concentrated on operation-specific risk factors (e.g., anastomosis technique). However, for complex surgeries, the standard perioperative care protocols may not be adequate to reduce major complications for every individual patient.

Additionally, identifying risk factors for adverse outcomes is important for case-mix correction in benchmarking quality of care in nation-wide clinical auditing and surgical improvement programs, such as the Dutch Institute of Clinical Auditing (DICA) and American College of Surgeons National Surgical Quality Improvement Program (ACS NSQIP) [9, 10].

This review focuses on the identification of preoperative risk factors for major postoperative complications (CD $\geq$ IIIa) after major abdominal surgery with the construction of an intestinal anastomosis, which includes esophagectomy, gastrectomy, and colorectal surgery. Since, these types of surgery have technical similarities and are all high risk procedures. Furthermore, this study aims to identify the strengths and possible improvements in ERAS protocols.

Methods

The study protocol for this systematic review was registered with the PROSPERO database (CRD42020198812). This review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA).

Criteria for study eligibility

To evaluate the effect of preoperative factors on the incidence of major postoperative complications, studies were selected based on the type of surgery. Only studies addressing complex gastrointestinal cancer surgery (e.g., esophagectomy, gastrectomy, and colectomy), including the construction of an intestinal anastomosis, were selected. As an outcome, a study was required to report on the associations between major complications and an independent preoperative factor. Major complications were classified as CD ≥ IIIa or severe complications that were classified as such (e.g., anastomotic leakage, endoscopic intervention) [6]. Retrospective, prospective cohort studies, and randomized-controlled trials with full-text articles published in English or Dutch were assessed for eligibility. Case reports and case series (< 40 patients) were excluded. Only studies including adult patients (≥ 18 years of age) were selected, and animal studies were excluded.

Search method

The Medline and Embase electronic databases were searched to identify all relevant publications. Search terms included those from MeSH in PubMed and EMtree in Embase, as well as free text terms. Reference lists of identified studies will be checked for additional relevant studies. Included studies were restricted to those that were published between January 2005 and July 2021. Authors were contacted in case of full-text unavailability.

Study selection

Assessment of eligibility was performed independently by RB and RvK. Any disagreement regarding eligibility was resolved by discussion with MW as an arbitrator when necessary. The initial screening was based on title and abstract. Full texts were independently screened by two authors (RB and RvK). Again, disagreement was resolved by discussion with MW, who acted as an arbitrator. Study selection was performed using Endnote X9 (Clarivate Analytics, Philadelphia, PA, USA) and Rayyan QCRI (a mobile web app for systematic reviews).

Assessment of risk of bias

All eligible studies were independently assessed for potential risk of bias by RB and RvK, using the Quality in Prognostic Studies (QUIPS) tool for classification of prognostic factor studies [11]. Discrepancies were resolved by discussion, with MW as an arbitrator when necessary. The risk of bias in clinical trials was assessed in the following domains: study participation, study attrition, prognostic factor measurement, outcome measurement, adjustment bias, and statistical analysis bias. Each domain was graded as high, low, or unclear. The results of risk of bias screening are summarized in Supplementary File A.

Data extraction and management

Data extraction was performed by RvK, and subsequently verified by RB using a predefined, standardized form designed by RB and RvK. Any discrepancies were resolved through discussion.

Grading the level of evidence

The level of evidence regarding the association between a risk factor and major complications (CD ≥ IIIa) was scored using a grading system (Table 1) [12]. The score resulted from the number of studies conducting a multivariable analysis of the association and percentage of statistically significant results of these analyses.

Table 1 - Grading the level of evidence adapted from the grading score used by Lagarde et al. [12].

Level of evidence	Criteria
None	No significant evidence
Minor	Evidence significant from multivariable analysis form one article
Considerable	Evidence significant from multivariable analysis in three or less articles and/or in less than 50% of the articles describing this risk factor
Strong	Evidence significant from multivariable analysis in more than three articles and in more than 50% of all articles describing this risk factor
Very strong	Evidence significant from multivariable analysis in ten or more articles and in more than 70% of all articles describing this risk factor

Results

The literature search retrieved 207 eligible studies (Figure 1), all of which used an observational study design. An overview of the results reported in these studies on preoperative risk factors associated with major complications (i.e., CD ≥ IIIa) after major gastrointestinal cancer surgery is shown in Table 2, together with the type of surgery (lower or upper gastrointestinal surgery), and the level of evidence graded according to Table 1. The fourth column reports the number of studies, including the risk factor, in multivariable analysis and the percentage of significant results. This section is divided into six subsections: patient characteristics, comorbidities, intoxication, nutritional indicators, disease-related factors, and neoadjuvant therapy-related factors.






Figure 1 - PRISMA flowchart of study selection

Patient characteristics

Age and frailty

Age is an important risk factor. Many studies reported an independent association between older age and major complications and mortality (Table 2). The elderly are believed to exhibit less healing capacity, which leads to more postoperative complications [13, 14]. Another term reported in studies is “frailty”, which is a physiological state of cumulative deficits (e.g., advanced age, poor physical performance), which render patients more susceptible to major complications [15]. In a large population-based retrospective cohort, Sparreboom et al. reported an association between frailty and anastomotic leakage [2]. Along with frailty, functional status, and activities of daily living dependency have demonstrated an association with postoperative complications and mortality (Table 2).

Table 2- This table represents all pre-operative risk factors for major complications (Clavien-Dindo (CD) ≥ IIIa) and mortality described in literature References used in this table are listed in Supplementary Information B. BMI= body-mass index; CD= Clavien-Dindo; HbA1c= glycated hemoglobin; Low= lower GI surgery; Up= upper GI surgery. *30-day mortality or in-hospital mortality.

Preoperative risk factors	Type of complications	Type of	Number of articles favoring (%)	Level of evidence	Reference(s)		
Patient characteristics							
Age	Anastomotic leakage	Up & Low	10/22 (45)	Considerable	[2, 26, 66, 88-157]		
	Intra-abdominal infection	Up	2/2 (100)	Considerable			
	Reoperation	Up & Low	2/2 (100)	Considerable			
	Venous thrombo-embolism	Up & Low	2/3 (67)	Considerable			
	Mortality*	Up & Low	17/20 (85)	Very strong			
	CD≥ IIIa	Up & Low	18/29 (62)	Strong			
Male gender	Anastomotic leakage	Up & Low	27/34 (79)	Very Strong	[2, 14, 26, 88, 89, 92, 95, 97, 102, 104, 105, 107, 111, 112, 114, 119-124, 126, 128, 129, 133, 137-140, 146, 149, 152, 156, 158-190]		
	Pancreatic fistula	Up	1/2 (50)	Minor			
		Low	1/1 (100)	Minor			
	Intra-abdominal infection	Up & Low	3/5 (60)	Considerable			
	Reoperation	Up	1/1 (100)	Minor			
	Venous thrombo-embolism	Up	1/1 (100)	Minor			
	Mortality*	Low	5/6 (83)	Strong			
	CD ≥ IIIa	Up & Low	9/17 (53)	Strong			
	American Society of Anesthesiologists (ASA) score	Anastomotic leakage	Up & Low	10/19 (53)		Strong	[2, 26, 88, 90, 91, 98, 99, 101, 102, 105, 110, 114, 120, 124, 126, 128-130, 133, 137-140, 144, 146, 148, 151-153, 162, 166, 170, 189, 191-196]
		Reintervention	Low	1/1 (100)		Minor	
Mortality*		Up & Low	8/10 (80)	Strong			
CD ≥ IIIa		Up & Low	6/15 (40)	Considerable			
Mortality*		Up & Low	1/2 (50)	Minor			
Physical fitness	Anastomotic leakage	Low	1/2 (50)	Minor	[138, 160, 197, 198]		
	CD ≥ IIIa	Up & Low	2/2 (100)	Considerable			
	Frailty	Mortality	Up	2/2 (100)		Considerable	[92, 138, 153, 199]
CD ≥ IIIa		Up & Low	2/3 (67)	Considerable			
Comorbidity							
Comorbidity	Anastomotic leakage	Up & Low	6/8 (75)	Strong	[2, 105-107, 129, 133, 146, 147, 149, 152, 162, 168, 200-203]		
	Reoperation	Up & Low	2/2 (100)	Considerable			
	Respiratory failure	Up	1/1 (100)	Minor			
	Mortality*	Up & Low	3/3 (100)	Considerable			
	CD ≥ IIIa	Up	3/4 (75)	Considerable			

Table 2- Continued

Preoperative risk factors	Type of complications	Type of	Number of articles favoring (%)	Level of evidence	Reference(s)
Hypertension	Anastomotic leakage	Up & Low	6/6 (100)	Strong	[136, 192, 193, 204-206]
	Anastomotic leakage	Up & Low	2/3 (67)	Considerable	[135,
	CD ≥ IIIa	Up	3/3 (100)	Considerable	150, 155, 184, 207-209]
	Anastomotic leakage	Up & Low	4/8 (50)	Strong	[100-102,
	Acute respiratory distress syndrome (ARDS)	Up	1/1 (100)	Minor	104, 114, 137, 140, 141, 154, 155, 170, 189, 190, 194,
	Respiratory failure	Up	1/2 (50)	Minor	202, 210-214]
	Mortality*	Up & Low	5/5 (100)	Strong	
	CD ≥ III	Up & Low	4/6 (67)	Strong	
	Anastomotic leakage	Up & Low	8/12 (75)	Strong	[100, 102,
	Duodenal stump fistula	Up	1/1 (100)	Minor	104, 114, 117, 121, 131,
	Respiratory failure	Up	1/1 (100)	Minor	135-137, 154, 155, 158, 159, 170, 185, 194,
	Venous thrombo-embolism	Up	1/1 (100)	Minor	209, 210, 212, 213, 215-219]
	Mortality*	Up & Low	4/5 (80)	Strong	
	CD ≥ IIIa	Up & Low	8/11 (73)	Strong	
Chronic hepatic disease	Anastomotic leakage	Up	1/1 (100)	Minor	[102, 135,
	Duodenal stump fistula	Up	1/1 (100)	Minor	140, 181, 215, 220, 221]
	Intra-abdominal infection	Up	1/1 (100)	Minor	
	Mortality*	Up & Low	2/2 (100)	Considerable	
	CD ≥ IIIa	Up	1/1 (100)	Minor	
Chronic kidney failure	Anastomotic leakage	Up	2/3 (67)	Considerable	[98, 102, 193, 209]
	Mortality*	Low	1/1 (100)	Minor	
Diabetes	Anastomotic leakage	Up & Low	9/18 (50)	Considerable	[26, 98, 104, 140, 158, 176, 182, 186, 194, 207, 208, 210, 212, 221-226]
	CD IV-V	Up	1/2 (50)	Minor	[100, 145]
Steroid use (chronically)	Anastomotic leakage	Up & Low	3/5 (60)	Strong	[98, 114, 129, 140, 165, 227, 228]
	Mortality*	Up & Low	2/2 (100)	Considerable	
Anti-coagulant therapy	Anastomotic leakage	Low	1/1 (100)	Minor	[189]
Prior abdominal surgery	Anastomotic leakage	Up & Low	3/5 (60)	Considerable	[2, 95, 142, 148, 167, 200]

Table 2- Continued

Preoperative risk factors	Type of complications	Type of	Number of articles favoring (%)	Level of evidence	Reference(s)
Intoxications					
Smoking	Anastomotic leakage	Up & Low	15/20 (75)	Very strong	[14, 25, 26,
	Mortality*	Low	2/2 (100)	Considerable	98, 102, 114, 124, 129, 138, 142, 158, 160, 165, 166, 200, 201, 210, 220, 221, 229-232]
	CD ≥ IIIa	Up	2/2 (100)	Considerable	
	Anastomotic leakage	Low	4/6 (67)	Considerable	[124, 129, 138, 158, 166, 210, 232]
Nutritional indicators					
Overweight (BMI >25)	Anastomotic leakage	Up & Low	4/9 (44)	Considerable	[2, 49, 88,
	Pancreatic fistula	Up	2/2 (100)	Considerable	94, 95, 104, 105, 111,
	Intra-abdominal infection	Up & Low	3/5 (60)	Considerable	113-115, 122, 123, 133, 138, 146, 160, 163, 174, 180, 188, 195, 233-240]
	Reoperation	Up	1/1 (100)	Minor	
	Venous thrombo-embolism	Up & Low	1/3 (33)	Minor	
	Mortality*	Up & Low	2/4 (50)	Considerable	
	CD ≥ IIIa	Up	6/11 (55)	Strong	
	Anastomotic leakage	Up & Low	6/7 (86)	Strong	[14, 114, 124,
Obesity (BMI >30)	Venous thrombotic-embolism	Up & Low	1/3 (33)	Minor	140, 148, 165, 189, 207, 210, 233, 235, 238, 239, 241]
	CD ≥ IIIa	Up & Low	1/3 (33)	Minor	
	Venous thrombo-embolism	Up & Low	1/3 (33)	Minor	
	CD ≥ IIIa	Low	1/1 (100)	Minor	
	CD ≥ IIIa	Up	1/1 (100)	Minor	[130]
	Anastomotic leakage	Up & Low	1/2 (50)	Minor	[128, 184, 227, 235, 239]
Sarcopenic obesity	Anastomotic leakage	Up & Low	2/3 (67)	Considerable	[242-247]
	Prolonged intubation	Up	1/1 (100)	Minor	
	CD ≥ IIIa	Up & Low	3/4 (75)	Strong	
Malnutrition/ preoperative weight loss	Anastomotic leakage	Up & Low	5/8 (63)	Strong	[106, 108,
	Duodenal stump fistula	Up	1/1 (100)	Minor	114, 115, 130, 138, 140, 179, 210, 215, 248, 249]
	Mortality*	Up	2/2 (100)	Considerable	
	CD ≥ IIIa	Up	4/5 (80)	Strong	[144, 153, 198, 250]
High visceral fat area (VFA)	Anastomotic leakage	Up & Low	2/3 (100)	Considerable	[103,
	Intra-abdominal infection	Up	2/2 (100)	Considerable	234, 251-253]
	Mortality*	Up	1/1 (100)	Minor	

Table 2- Continued

Preoperative risk factors	Type of complications	Type of	Number of articles favoring (%)	Level of evidence	Reference(s)
Perineal Fat Surface area (PRF) $\geq 40 \text{ cm}^2$	CD $\geq$ III	Low	1/1 (100)	Minor	[124]
Disease related risk factors					
Tumor stage/ tumor size	Anastomotic leakage	Up & Low	9/16 (56)	Considerable	[2, 49, 90, 91, 95, 104, 105, 110, 113, 119, 120, 123, 131, 133, 137, 139, 144-146, 148, 152, 153, 159-161, 164, 172, 173, 176-178, 187, 191, 216-219, 225, 254-260]
	Postoperative hemorrhage	Up	2/2 (100)	Considerable	
	Intra-abdominal infection	Up & Low	2/3 (67)	Considerable	
	Major adverse cardiac event (MACE)	Up	1/1 (100)	Minor	
	Mortality*	Up & Low	2/4 (50)	Minor	
	CD $\geq$ IIIa	Up	13/22 (59)	Strong	
Preoperative	Anastomotic leakage	Low	1/3 (33)	Minor	[2, 126, 139, 147, 184, 261, 262]
	Mortality*	Up & Low	2/3 (67)	Considerable	
	CD $\geq$ IIIa	Up	1/1 (100)	Minor	
Neoadjuvant therapy-related risk factors					
	Anastomotic leakage	Up	1/1 (100)	Minor	[53, 149, 207, 263, 264]
	CD $\geq$ IIIa	Up	1/2 (50)	Minor	
	Mortality*	Up & Low	1/2 (50)	Minor	
	Anastomotic leakage	Low	4/7 (57)	Strong	[2, 105, 166, 175, 196, 256, 265, 266]
	Chylothorax	Up	1/1 (100)	Minor	
	Intra-abdominal complication (CD $\geq$ IIIa)	Low	1/1 (100)	Minor	
	Anastomotic leakage	Up & Low	2/3 (67)	Considerable	[113, 114, 133, 153, 164, 195, 259, 267]
	CD $\geq$ IIIa	Up & Low	2/4 (50)	Considerable	
	Mortality*	Up & Low	1/2 (50)	Minor	
	Anastomotic leakage	Up & Low	4/7 (57)	Strong	[2, 105, 138, 184, 237, 268, 269]
Preoperative laboratory tests					
	Anastomotic leakage	Up & Low	2/4 (50)	Considerable	[104, 108, 128, 184, 210]
Platelet count increased	Anastomotic leakage	Low	1/1 (100)	Minor	[165]
Platelet count decreased		Low	1/1 (100)	Minor	[171]
White blood cell count (WBC) increased	Anastomotic leakage	Up & Low	1/2 (50)	Minor	[143, 210, 224]
	Venous thrombo-embolism	Low	1/1 (100)	Minor	
Neutrophil-to-lymphocyte Ratio (NLR)	Anastomotic leakage	Low	1/2 (50)	Minor	[160, 270]

Table 2- Continued

Preoperative risk factors	Type of complications	Type of	Number of articles favoring (%)	Level of evidence	Reference(s)
C-reactive protein (CRP) increased	Anastomotic leakage	Up & Low	2/2 (100)	Considerable	[121, 128, 158, 178, 186]
	CD $\geq$ IIIa	Up	2/3 (67)	Considerable	
CRP/Albumin ratio (CAR)	Anastomotic leakage	Low	1/1 (100)	Minor	[142]
Increased creatinine	Anastomotic leakage	Up & Low	1/2 (50)	Minor	[104, 117, 139, 140, 208, 216]
	Mortality*	Up	1/2 (50)	Minor	
	CD $\geq$ IIIa	Up	1/2 (50)	Minor	
Decreased estimated glomerular filtration rate (eGFR)	CD $\geq$ IIIa	Up	1/1 (100)	Minor	[159]
Serum albumin	Anastomotic leakage	Up & Low	5/8 (63)	Strong	[26, 104, 123, 128, 130, 137, 139, 140, 151, 164, 168, 178, 183, 189, 190, 216, 232, 250, 263, 266]
	Mortality	Up	2/3 (67)	Considerable	
	CD $\geq$ IIIa	Up	5/10 (50)	Considerable	
Total protein decreased	Anastomotic leakage	Low	3/3 (100)	Considerable	[14, 121, 189, 190, 210]
	CD $\geq$ IIIa	Up & Low	2/2 (100)	Considerable	
	CD > IIIa	Up	1/1 (100)	Minor	[186]
Albumin-to-fibrinogen ratio (AFR)					
Increased HbA1c	Anastomotic leakage	Up	1/1 (100)	Minor	[220]

Male sex

A wide variety of studies have confirmed that male sex is a risk factor for major postoperative complications. Several theories have been proposed to address this issue. Historically, the incidence of smoking and alcohol consumption in the male population has been higher. However, these confounding variables have not been measured in many studies and, therefore, their effect on males may be overestimated [16]. Another theory is that differences in cortisol-induced sex hormones change after surgically induced stress, which could render males more susceptible to postoperative complications [17]. A third theory is that the narrower pelvis of male patients can make surgery for tumors located in this region technically more difficult [2, 14].

American Society of Anesthesiologists score

One of the most studied risk indicators in the context of predicting postoperative complications is the American Society of Anesthesiologists (ASA) score. Multiple studies found an independent association between ASA score and a higher incidence of anastomotic leakage and major complications [2, 18, 19]. Furthermore, the ASA score is a reliable predictor of 30-day mortality (Table 2).

Preoperative inflammatory biomarkers

Several studies described an association between elevated levels of preoperative inflammatory biomarkers (e.g., white blood cell count, C-reactive protein [CRP]) and postoperative complications (Table 2). Similarly, the neutrophil-to-lymphocyte ratio, a proxy measure of inflammation status in the body, is independently associated with an increased risk for major complications (Table 2). The association between preoperative inflammation and complications, however, is not yet fully understood.

Serum albumin is a negative acute-phase protein. It decreases during inflammation due to increased capillary leakage [20]. It is also known as a nutritional biomarker reflecting malnutrition (Section 3.4.1). In the Glasgow Prognostic Score, an inflammation-based prognostic score for cancer prognosis, albumin and CRP are combined to predict perioperative complications [21]. Similarly, You et al. proposed the albumin/fibrinogen ratio as a predictor of major complications (Table 2). Fibrinogen is an essential protein in the coagulation cascade as well as an acute-phase reaction protein in the response to systemic inflammation [22].

Comorbidities

Patients with ≥ 1 comorbidities and those using ≥ 5 drugs per day are more susceptible to complications [23]. Several studies have demonstrated that heart failure, hypertension, and renal insufficiency are independently associated with major complications and anastomotic leakage (Table 2). Vascular disease, particularly arterial calcifications, is an important risk factor for major complications, especially anastomotic leakage (Table 2). Furthermore, the relationship between major complications and diabetes is well understood, whereas hyperglycemia induces microvascular damage, yielding a reduced capacity for anastomotic healing [24] (Table 2).

Intoxication

Smoking

A history of smoking is a risk factor for the occurrence of postoperative complications after major abdominal surgery (Table 2). In a large retrospective cohort study, Sharma et al. estimated the increased risk for major postoperative complications and mortality after smoking to be 30% [16]. Quan et al. reported that the number of pack-years

significantly influenced the risk for major complications [25]. Smoking is believed to induce microvascular damage, leading to decreased healing ability of the anastomosis, thereby leading to an increased rate of anastomotic leakage [26].

Alcohol consumption

Several studies have shown that habitual use of alcohol (≥ 3 units per day) increases the risk for postoperative complications (Table 2). Alcohol causes alcohol-induced liver and pancreatic disorders, as well as impaired immune capacity, hemostasis, and surgical stress response [27, 28]. Alcohol cessation before elective surgery significantly decreased postoperative complications [27, 28].

Nutrition-related risk factors

Malnutrition/preoperative weight loss

Among cancer patients, 63% experience weight loss before treatment. In those with gastric and esophageal cancers, this figure has been reported to be as high as 79% to 83% [29, 30]. Absolute weight loss can be an indication of malnutrition, which can also be measured according to nutritional indexes (e.g., Prognostic Nutritional Index, Nutritional Risk Screening). A more advanced stage of malnutrition leads to cancer anorexia-cachexia syndrome—a hypercatabolic state characterized by weight loss and sarcopenia—which occurs in 15% to 40% of cancer patients [31, 32]. Malnutrition and preoperative weight loss were significantly associated with major complications and mortality (Table 2). Lack of nutrients has been implicated in decreased function of the immune, respiratory and cardiac systems, as well as decreased healing function [33, 34] and further deterioration due to a more catabolic metabolic state [13]. Collectively, this leads to an increased incidence of infectious complications as well as anastomotic leakage (Table 2). Low preoperative serum albumin levels are independently associated with an increased risk for major complications (Section 3.1.4).

Sarcopenia

Sarcopenia refers to the loss of skeletal muscle volume and/or strength, which have a close relationship, and primarily originates from malnutrition (Section 3.4.1). Sarcopenia is especially prevalent in patients with esophageal and gastric cancers (up to 56%), but also in elderly patients [35-37]. As shown in Table 2, sarcopenia was independently associated with worse surgical outcomes. The relationship between sarcopenia and major postoperative complications and mortality is due to reduced healing capacity resulting from a lack of nutrients and, therefore, a catabolic state.

Overweight and obesity

Obese and overweight patients are at higher risk for postoperative complications and mortality after major gastrointestinal surgery (Table 2). There are multiple theories addressing the association between overweight and major complications. First, obese patients often exhibit a significantly increased number of comorbidities, including diabetes, hyperlipidemia, coronary artery disease, and hypertension [38, 39]. Second, overweight and obesity are associated with an increased incidence of anastomotic leakage believed to be caused by a preoperative inflammatory state and increased insulin resistance, leading to decreased healing capacity [40, 41]. Third, increased visceral fat in those undergoing abdominal surgery may lead to more complications due to more technical difficulties (e.g., thicker mesocolon, increased abdominal wall pressure leading to decreased intraoperative visibility) [39, 42], which in turn leads to longer operation time and greater transfusion requirements [19]. Some retrospective studies have explored the relationship between visceral fat area, body mass index, and the impact of excessive abdominal fat tissue on surgical outcomes. However, whether visceral fat area is a better parameter than body mass index remains controversial [43].

Disease-related risk factors

Preoperative tumor complications

Of all preoperative tumor complications, anemia and iron deficiency are the most common. The prevalence of any degree of anemia has been suggested to be 50% to 75% in patients with colorectal cancer [44, 45]. Anemia leads to decreased healing capacity. Therewithal, patients receiving preoperative transfusion exhibited an increased rate of postoperative complications [46]. Blood transfusions appear to induce an immunosuppressive effect; therefore, a policy restricting transfusion is recommended [47]. Local preoperative tumor complications (e.g., bowel obstruction and tumor perforation) are independently associated with major complications (Table 2), theoretically, due to greater technical difficulty caused by an inflammatory response of the abdominal cavity and by the frailty of the tissue used for anastomosis and/or the spill of gastrointestinal fluids.

Advanced tumor stage

Advanced tumor stage, including those from poorly differentiated cancer types, lead to more extensive resections and technically more demanding surgery, followed by more intraoperative organ damage and postoperative complications [42, 48]. Second, extensive lymph node dissections and additional splenic resection, especially in gastric and esophageal resections, are high-risk procedures [49, 50]. Additionally, larger tumors and more extensive resections lead to more non-radical resections [51]. Furthermore, patients with a higher tumor grade or TNM stage are more likely to exhibit a form of systemic immune-inflammation, which is also associated with major complications [52] (Section 3.1.4).

Neoadjuvant therapy-related factors

Neoadjuvant therapy aims to reduce tumor volume to achieve R0 resections and mitigate—if not eliminate—micrometastases and, eventually, cancer recurrence. However, the use of neoadjuvant therapy is also associated with an increase in postoperative complications caused by a diminished healing capacity of damaged tissue (Table 2). Additionally, a possible decrease in psychological performance after neoadjuvant therapy may lead to impaired postoperative recovery [53]. Preoperative radiotherapy in those treated for rectal cancer has a high prevalence of postoperative complications and anastomotic leakage (Table 2). After neoadjuvant therapy, patients also experience postoperative cardiopulmonary complications more frequently [54]. Patients unable to complete neoadjuvant therapy often experience increased postoperative complications, which may be a confounder due to poor underlying health conditions [55].

Discussion

Results of the present study provide a comprehensive and structured overview of the associations between preoperative risk factors and major complications and mortality following complex gastrointestinal cancer surgery. Our findings provide unambiguous evidence supporting the association between age and major postoperative complications, as well as for the association between anastomotic leakage and male sex and smoking. Furthermore, substantial evidence has been provided regarding the association between major postoperative complications and age, male sex, comorbidities, malnutrition, sarcopenia and overweight/obesity. This study also provides strong evidence supporting an association between different comorbidities, obesity, malnutrition, decreased serum albumin, more advanced tumor stages, neoadjuvant radiotherapy, neoadjuvant chemotherapy and the occurrence of anastomotic leakage. Furthermore, strong evidence exists for an association between 30-day mortality and male sex, higher ASA score, and cardiac comorbidity. This systematic review also shows that risk factors for postoperative major complications in lower – and upper gastrointestinal cancer surgery show a substantial overlap.

The identification of risk factors may afford opportunities to optimize perioperative care by managing preoperative risk factors, thereby decreasing the risk for postoperative complications and mortality. This may reduce healthcare costs, in contrast to major complications, which lead to an increase in healthcare expenditures [1]. The described associations may contribute to focused and personalized preoperative care by enrolling patients with certain risk factors (e.g., frailty and malnutrition) into prehabilitation programs. Subsequently, identification of high-risk patients may prompt closer

postoperative surveillance. Additionally, the identification of high-risk patients may also influence decision making regarding treatment options, for example, a ‘watch and wait’ strategy after clinical complete response to neoadjuvant therapy [56, 57].

Preoperative care

In literature, several prehabilitation programs have been described for modifiable risk factors, acting on the associations between preoperative factors and postoperative complications (Table 3). Preoperative control/management of these factors could improve postoperative outcomes. For example, adequate preoperative glycemic control in diabetic patients should lead to less postoperative hyperglycemia, which is associated with postoperative infectious complications and, could therefore, decrease the complication rate [58]. Furthermore, several prehabilitation programs incorporating for instance physical resistance training and nutritional support have been described in the literature (Table 3). Theoretically, these prehabilitation programs should lead to a reduction in postoperative complications, although there is limited evidence to support this [59, 60]. A limitation—present in the majority of research investigating preoperative interventions—could be that prehabilitation is not specifically targeted at patient-specific risk factors. Physical endurance training in a population >70 years of age with ASA III-IV, led to a 20% reduction in complications [61], indicating that preoperative care should be tailored to and specified for patients targeting their risk factors. Smoking cessation, which leads to a significant reduction in postoperative complications, is such an example [25, 62]. Currently, growing interest of perioperative research is focused on the implementation and further improvement of ERAS protocols, which may lead to a reduction in overall complications by up to 50%, as shown in a meta-analysis [5]. However, studies included in this systematic review have been published during the period in which ERAS protocols have been gaining interest and were widely implemented. This means that perioperative care has been improved and optimization of risk factors (e.g., malnutrition, smoking cessation) is standard in daily practice [63]. Also standard in ERAS protocols for gastrointestinal surgery is nutritional support, this is important for patients to cope with the metabolic and physiological stress inflicted by gastrointestinal cancer surgery and increased protein requirements [64]. In addition to nutritional support the so-called “Immunonutrition” which entails nutritional supplements (e.g., arginine, omega-3 fatty acids) is being studied, this is thought to lead to a reduction of surgical stress [65](Table 3). In the light of ERAS protocols studies have shown that an abbreviated period (2 h versus 12 h) of fasting leads to significantly reduced time-to-first-stool and complete oral intake [66, 67]. In the ERAS protocol for lower gastrointestinal surgery, bowel preparation is an important point of discussion because this could lead to changes in electrolyte levels, metabolic imbalance, and dehydration, especially in elderly and/or frail patients [68]. The suggestion to omit this from the protocol, if possible, especially in frail patients, is supported by a meta-

analysis that revealed an advantage to no-bowel preparation with regard to anastomotic leakage, intra-abdominal infections, and wound infections [69]. In this context the role of perioperative prophylactic antibiotics usage is studied, which may have a preventive effect on surgical site infections, anastomotic leakage and mortality [70].

Table 3 – Table includes actable or improvable risk factors and subsequent in literature described prehabilitation options to reduce the risk of postoperative morbidity. References used in this table are listed in Supplementary Information B.

Risk factors	Prehabilitation	Reference
Physical performance	Resistance training	[47, 59, 61, 271-275]
	Endurance training	
	Physical therapy	
	Breathing exercises	
	Nutritional support	
	Immunonutrition	
Pulmonary comorbidity	Preoperative inspiratory muscle training	[276-279]
Malnutrition	Nutritional support	[63-65, 280-283]
	Oral nutritional supplements	
	Immunonutrition	
Sarcopenia	Nutritional support	[275, 284, 285]
	Resistance training	
	Nutritional supplements	
Smoking	Smoking cessation	[16, 25, 62, 286]
Alcohol consumption	Alcohol cessation	[28]
Iron deficiency anemia	Intravenous iron supplementation	[287]
Dental plaque	Preoperative oral management by dentist	[288, 289]

Intraoperative techniques and care

Furthermore, ERAS protocols have been further improved intraoperative care in terms of: minimally invasive surgery, pain management, temperature management and fluid administration [71]. During the publishing of the included studies minimally invasive surgery has become more standard procedure. Other intra-operative ERAS-principles that have been studied and implemented such as goal-directed fluid administration and use of fewer use of intra-operative vasopressors have been independently associated with decreased postoperative complications [71-73]. Also intra-operative normothermia has been shown to have a positive effect on prevention of postoperative infections [74].



The use of opioid-sparing analgesia has been shown to increase recovery time, but no reduction in postoperative complications [75].

Postoperative care

With the current increase in data-driven approaches in healthcare, the risk factors reported in Table 2 could be assessed in analysis of large datasets, in which the development of artificial intelligence may play an important role. Machine learning models usually demonstrate similar performance for predicting medical outcomes compared with logistic regression [76]. With increasingly larger datasets, machine learning holds the potential to unravel subtle associations that are not—or cannot—be identified using classic regression approaches. For suspected low-risk patients, machine learning has been suggested to support early discharge decisions [77]. Suspected high-risk patients may benefit from closer postoperative surveillance. Earlier detection of deterioration in patients may reduce the severity of complications and lessen the incidence of failure-to-rescue [78]. A proposed method for augmented postoperative surveillance involves wearable devices for constant postoperative monitoring [79]. These devices continuously transmit vital signs that alert healthcare personnel in case of deterioration.

Benchmarking surgical outcomes

Reduction of postoperative complications can also be established by clinical auditing and benchmarking of surgical outcomes [80, 81]. Auditing is used to measure quality of care using structure, process, and outcome indicators [82, 83]. The information provided by this review can be used for fair comparison of outcomes between different hospitals and institutions, which can only be established when using robust casemix models.

Limitations

The present study had some limitations. First, it provided only an overview of the associations between preoperative risk factors and major complications. As such, additional evidence is needed to confirm that these risk factors are causally related to poor surgical outcomes. Second, heterogeneous patient populations and study designs may have hindered adequate interpretation of the study results. The included studies were all conducted in an observational manner, and most of them were designed retrospectively. There was a wide variety between risk factor reporting between studies, not all risk factors (e.g., renal disease, pulmonary comorbidity) were defined within the studies therefore making interpretation difficult. A similar reporting absence was seen in the implementation and usage of ERAS protocols within the included patient population. ERAS protocols have been widely implemented in surgery in recent years, that's why we limited our study period to 2005. This type of study is subjected to bias, although we suspect that due to the large number of studies, this bias was limited.

However, all patients included in this study were preoperatively selected to be fit for surgery by expert opinion undergoing surgery, leading to allocation bias. This is a limiting factor for generalization of risk factor research in general. Although the present study provides an overview of all known risk factors, not all factors are described or necessarily applicable to every patient. Additionally, this study provides a theoretical overview; therefore, no quantitative effect of the specific risk factors is reported. An additional meta-analysis should be conducted to calculate the quantitative effects of each risk factor. Moreover, the inclusion of risk factors described in Table 2 was based on the significant outcomes in multivariable analysis. This selection was performed to minimize the risk of including confounding factors. However, this may have excluded risk factors studied in low-powered studies, which could also have led to the lack of research investigating risk factors. In the present study, both upper gastrointestinal and lower gastrointestinal cancer surgery were considered by examining esophageal, gastric, and colorectal resections in a large subset of patients undergoing these operations.

Conclusions

In conclusion, identification of improvable/modifiable risk factors exposes possibilities for augmentation of perioperative care, which may lead to improved surgical outcomes. Furthermore, the identified risk factors can lead to alteration and additions to already existing ERAS protocols, which have already resulted in improved perioperative care and reduction in complications [5, 63]. In addition, the identification of preoperative risk factors could lead to further improved and personalized perioperative care, thereby reducing major postoperative complications (e.g., risk factor-targeted prehabilitation). This study also contains important information to improve benchmarking of surgical outcomes in nation-wide clinical audits. The reduction of postoperative complications may prolong (recurrence-free) survival and lead to improved quality of life [84-87].

References

1. Selby, L.V., et al., *Association of Hospital Costs With Complications Following Total Gastrectomy for Gastric Adenocarcinoma*. JAMA Surg, 2017. **152**(10): p. 953-958.
2. Sparreboom, C.L., et al., *Different Risk Factors for Early and Late Colorectal Anastomotic Leakage in a Nationwide Audit*. 2018. **61**(11): p. 1258-1266.
3. Busweiler, L.A., et al., *Failure-to-rescue in patients undergoing surgery for esophageal or gastric cancer*. Eur J Surg Oncol, 2017. **43**(10): p. 1962-1969.
4. Nygren, J., et al., *A comparison in five European Centres of case mix, clinical management and outcomes following either conventional or fast-track perioperative care in colorectal surgery*. Clin Nutr, 2005. **24**(3): p. 455-61.
5. Varadhan, K.K., et al., *The enhanced recovery after surgery (ERAS) pathway for patients undergoing major elective open colorectal surgery: a meta-analysis of randomized controlled trials*. Clin Nutr, 2010. **29**(4): p. 434-40.
6. Clavien, P.A., et al., *The Clavien-Dindo classification of surgical complications: five-year experience*. Ann Surg, 2009. **250**(2): p. 187-96.
7. Aurello, P., et al., *Recurrence Following Anastomotic Leakage After Surgery for Carcinoma of the Distal Esophagus and Gastroesophageal Junction: A Systematic Review*. Anticancer Res, 2019. **39**(4): p. 1651-1660.
8. Ramphal, W., et al., *Oncologic outcome and recurrence rate following anastomotic leakage after curative resection for colorectal cancer*. Surg Oncol, 2018. **27**(4): p. 730-736.
9. van Groningen, J.T., et al., *Identifying best performing hospitals in colorectal cancer care; is it possible?* Eur J Surg Oncol, 2020. **46**(6): p. 1144-1150.
10. Eisenstein, S., S. Stringfield, and S.D. Holubar, *Using the National Surgical Quality Improvement Project (NSQIP) to Perform Clinical Research in Colon and Rectal Surgery*. Clin Colon Rectal Surg, 2019. **32**(1): p. 41-53.
11. Hayden, J.A., et al., *Assessing bias in studies of prognostic factors*. Ann Intern Med, 2013. **158**(4): p. 280-6.
12. Lagarde, S.M., et al., *Prognostic factors in adenocarcinoma of the esophagus or gastroesophageal junction*. J Clin Oncol, 2006. **24**(26): p. 4347-55.
13. Lago, V., et al., *Risk factors for anastomotic leakage after colorectal resection in ovarian cancer surgery: A multi-centre study*. 2019. **153**(3): p. 549-554.
14. Klose, J., et al., *A Nomogram to Predict Anastomotic Leakage in Open Rectal Surgery- Hope or Hype?* 2018. **22**(9): p. 1619-1630.
15. Rockwood, K. and A. Mitnitski, *Frailty defined by deficit accumulation and geriatric medicine defined by frailty*. Clin Geriatr Med, 2011. **27**(1): p. 17-26.
16. Sharma, A., et al., *Tobacco smoking and postoperative outcomes after colorectal surgery*. Vol. 258. 2013. 296-300.
17. Scheingraber, S., et al., *Gender-specific differences in sex hormones and cytokines in patients undergoing major abdominal surgery*. Surg Today, 2005. **35**(10): p. 846-54.
18. Smit, L.C., et al., *Infectious Complications after Major Abdominal Cancer Surgery: In Search of Improvable Risk Factors*. 2016. **17**(6): p. 683-693.
19. Benoist, S., et al., *Impact of obesity on surgical outcomes after colorectal resection*. Am J Surg, 2000. **179**(4): p. 275-81.
20. Fleck A, Raines G, Hawker F, Trotter J, Wallace PI, et al., *Increased vascular permeability: a major cause of hypoalbuminaemia in disease and injury*. Lancet, 1985. **1**(8432): p. 781-4.
21. Proctor MJ, Horgan PG, Talwar D, Fletcher CD, Morrison DS, McMillan DC., *Optimization of the systemic inflammation-based Glasgow prognostic score: a Glasgow Inflammation Outcome Study*. Cancer, 2013. **119**(12): p. 2325-32.
22. Kijima T, Arigami T, Uckikado Y, et al., *Combined fibrinogen and neutrophil-lymphocyte ratio as a prognostic marker of advanced esophageal squamous cell carcinoma*. Cancer Sci, 2017. **108**(2): p. 193-199.
23. Xue, D.-D., et al., *Comprehensive geriatric assessment prediction of postoperative complications in gastrointestinal cancer patients: a meta-analysis*. 2018. **13**: p. 723-736.
24. Li, S.-J., et al., *Diabetes mellitus and risk of anastomotic leakage after esophagectomy: a systematic review and meta-analysis*. 2017. **30**(6): p. 1-12.
25. Quan, H., et al., *The effect of preoperative smoking cessation and smoking dose on postoperative complications following radical gastrectomy for gastric cancer: a retrospective study of 2469 patients*. 2019. **17**(1): p. 61.
26. Parthasarathy, M., et al., *Risk factors for anastomotic leakage after colorectal resection: a retrospective analysis of 17 518 patients*. 2017. **19**(3): p. 288-298.
27. Mantziari S, Hübner M, Demartines N, Schäfer M, *Impact of preoperative risk factors on morbidity after esophagectomy: is there room for improvement?* World J Surg, 2014. **38**(11): p. 2882-90.
28. Oppedal K, Moller AM, Pedersen B, Tonnesen H., *Preoperative alcohol cessation prior to elective surgery*. Cochrane Database Syst Rev, 2012(7): p. Cd008343.
29. Laviano A, Meguid MM, Inui A, Muscaritoli M, et al., *Therapy insight: Cancer anorexia-cachexia syndrome--when all you can eat is yourself*. Nat Clin Pract Oncol, 2005. **2**(3): p. 158-65.
30. Laviano A, Meguid MM., *Nutritional issues in cancer management*. Nutrition, 1996. **12**(5): p. 358-71.
31. Muliawati, Y., H. Haroen, and L.W. Rotty, *Cancer anorexia - cachexia syndrome*. Acta Med Indones, 2012. **44**(2): p. 154-62.
32. Ruiz Garcia, V., et al., *Megestrol acetate for treatment of anorexia-cachexia syndrome*. Cochrane Database Syst Rev, 2013(3): p. Cd004310.

33. Schneider, S.M., et al., *Malnutrition is an independent factor associated with nosocomial infections*. Br J Nutr, 2004. **92**(1): p. 105-11.
34. Jayarajan, S. and J.M. Daly, *The relationships of nutrients, routes of delivery, and immunocompetence*. Surg Clin North Am, 2011. **91**(4): p. 737-53, vii.
35. Nakashima, Y., et al., *Assessment of Sarcopenia as a Predictor of Poor Outcomes After Esophagectomy in Elderly Patients With Esophageal Cancer*. Ann Surg, 2018. **267**(6): p. 1100-1104.
36. Shachar, S.S., et al., *Prognostic value of sarcopenia in adults with solid tumours: A meta-analysis and systematic review*. Eur J Cancer, 2016. **57**: p. 58-67.
37. Nishigori, T., et al., *Impact of Sarcopenic Obesity on Surgical Site Infection after Laparoscopic Total Gastrectomy*. Ann Surg Oncol, 2016. **23**(Suppl 4): p. 524-531.
38. Bachmann, R., et al., *Comparison between abdominal fat measured by CT and anthropometric indices as prediction factors for mortality and morbidity after colorectal surgery*. Vol. 81. 2018. 477-483.
39. Hotouras, A., et al., *The influence of obesity and body mass index on the outcome of laparoscopic colorectal surgery: a systematic literature review*. 2016. **18**(10): p. O337-O366.
40. Zhang, X., et al., *The effect of increased body mass index values on surgical outcomes after radical resection for low rectal cancer*. 2019. **49**(5): p. 401-409.
41. De Magistris, L., et al., *Preoperative inflammation increases the risk of infection after elective colorectal surgery: results from a prospective cohort*. Int J Colorectal Dis, 2016. **31**(9): p. 1611-7.
42. Sparreboom, C.L., et al., *Different Risk Factors for Early and Late Colorectal Anastomotic Leakage in a Nationwide Audit*. Dis Colon Rectum, 2018. **61**(11): p. 1258-1266.
43. Yang, S.J., et al., *Visceral Fat Area (VFA) Superior to BMI for Predicting Postoperative Complications After Radical Gastrectomy: a Prospective Cohort Study*. J Gastrointest Surg, 2019.
44. Shander A, Knight K, Thurer R, Adamson J, Spence R., *Prevalence and outcomes of anemia in surgery: a systematic review of the literature*. Am J Med, 2004. **5**(116): p. 013.
45. Leightle SW, Mouawad NJ, Lampman R, Signal B, Cleary RK., *Does preoperative anemia adversely affect colon and rectal surgery outcomes?* J Am Coll Surg, 2011. **212**(2): p. 187-94.
46. Papageorge, C.M., G.D. Kennedy, and E.H. Carchman, *Preoperative blood transfusion is a predictor of worse short-term postoperative outcomes after colectomy*. Vol. 161. 2017. 1067-1075.
47. Ejaz, A., Schmidt, C., Johnston, F.M., Frank, S.M., Pawlik, T.M., *Risk factors and prediction model for inpatient surgical site infection after major abdominal surgery*. J Surg Res, 2017. **217**: p. 153-159.
48. Bertelsen CA, Neuenschwander AU, Jansen JE, et al., *Short-term outcomes after complete mesocolic excision compared with 'conventional' colonic cancer surgery*. Br J Surg, 2016. **103**(5): p. 581-9.
49. Lu, J., et al., *Major perioperative complications in laparoscopic spleen-preserving total gastrectomy for gastric cancer: perspectives from a high-volume center*. 2016. **30**(3): p. 1034-42.
50. Kulig P, Sierzega M, Pach R, et al., *Differences in prognosis of Siewert II and III oesophagogastric junction cancers are determined by the baseline tumour staging but not its anatomical location*. Eur J Surg Oncol, 2016. **42**(8): p. 1215-21.
51. Vogelsang RP, Klein MF, Gögenur I., *Risk Factors for Compromised Surgical Resection: A Nationwide Propensity Score-Matched Study on Laparoscopic and Open Resection for Colonic Cancer*. Dis Colon Rectum, 2019. **62**(4): p. 438-446.
52. Shi, H., et al., *Nomogram Based on Systemic Immune-Inflammation Index to Predict Overall Survival in Gastric Cancer Patients*. Dis Markers, 2018. **2018**: p. 1787424.
53. Mungo, B., et al., *Does neoadjuvant therapy for esophageal cancer increase postoperative morbidity or mortality?* Diseases of the esophagus : official journal of the International Society for Diseases of the Esophagus, 2015. **28**(7): p. 644-651.
54. Bosch, D.J., et al., *Impact of neoadjuvant chemoradiotherapy on postoperative course after curative-intent transthoracic esophagectomy in esophageal cancer patients*. Ann Surg Oncol, 2014. **21**(2): p. 605-11.
55. Claassen, Y.H.M., et al., *Surgical morbidity and mortality after neoadjuvant chemotherapy in the CRITICS gastric cancer trial*. Eur J Surg Oncol, 2018. **44**(5): p. 613-619.
56. van der Valk, M.J.M., et al., *Long-term outcomes of clinical complete responders after neoadjuvant treatment for rectal cancer in the International Watch & Wait Database (IWWDD): an international multicentre registry study*. Lancet, 2018. **391**(10139): p. 2537-2545.
57. van der Valk, M.J.M., et al., *Importance of patient reported and clinical outcomes for patients with locally advanced rectal cancer and their treating physicians. Do clinicians know what patients want?* Eur J Surg Oncol, 2020. **46**(9): p. 1634-1641.
58. McConnell, Y.J., P.M. Johnson, and G.A. Porter, *Surgical site infections following colorectal surgery in patients with diabetes: association with postoperative hyperglycemia*. Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract, 2009. **13**(3): p. 508-515.
59. Bruns, E.R., et al., *The effects of physical prehabilitation in elderly patients undergoing colorectal surgery: a systematic review*. Colorectal Dis, 2016. **18**(8): p. O267-77.
60. Carli, F., et al., *Effect of Multimodal Prehabilitation vs Postoperative Rehabilitation on 30-Day Postoperative Complications for Frail Patients Undergoing Resection of Colorectal Cancer: A Randomized Clinical Trial*. JAMA Surg, 2020.
61. Barberan-Garcia, A., et al., *Personalised Prehabilitation in High-risk Patients Undergoing Elective Major Abdominal Surgery: A Randomized Blinded Controlled Trial*. Ann Surg, 2018. **267**(1): p. 50-56.
62. Lindstrom, D., et al., *Effects of a perioperative smoking cessation intervention on postoperative complications: a randomized trial*. Ann Surg, 2008. **248**(5): p. 739-45.

63. Ljungqvist, O., *Jonathan E. Rhoads lecture 2011: Insulin resistance and enhanced recovery after surgery*. JPEN J Parenter Enteral Nutr, 2012. **36**(4): p. 389-98.
64. Wischmeyer, P.E., et al., *American Society for Enhanced Recovery and Perioperative Quality Initiative Joint Consensus Statement on Nutrition Screening and Therapy Within a Surgical Enhanced Recovery Pathway*. Anesth Analg, 2018. **126**(6): p. 1883-1895.
65. Moya, P., et al., *Perioperative Standard Oral Nutrition Supplements Versus Immunonutrition in Patients Undergoing Colorectal Resection in an Enhanced Recovery (ERAS) Protocol: A Multicenter Randomized Clinical Trial (SONVI Study)*. Medicine (Baltimore), 2016. **95**(21): p. e3704.
66. Reis, P.G.A., et al., *Abbreviated preoperative fasting favours postoperative oral intake at lower hospital admission costs for cancer patients*. Rev Col Bras Cir, 2019. **46**(3): p. e20192175.
67. Svanfeldt, M., et al., *Randomized clinical trial of the effect of preoperative oral carbohydrate treatment on postoperative whole-body protein and glucose kinetics*. Br J Surg, 2007. **94**(11): p. 1342-50.
68. Beloosesky, Y., et al., *Electrolyte disorders following oral sodium phosphate administration for bowel cleansing in elderly patients*. Arch Intern Med, 2003. **163**(7): p. 803-8.
69. Bucher, P., et al., *Mechanical bowel preparation for elective colorectal surgery: a meta-analysis*. Arch Surg, 2004. **139**(12): p. 1359-64; discussion 1365.
70. Rollins, K.E. and D.N. Lobo, *The Controversies of Mechanical Bowel and Oral Antibiotic Preparation in Elective Colorectal Surgery*. Ann Surg, 2021. **273**(1): p. e13-e15.
71. Hikasa, Y., et al., *Intraoperative fluid therapy and postoperative complications during minimally invasive esophagectomy for esophageal cancer: a single-center retrospective study*. J Anesth, 2020. **34**(3): p. 404-412.
72. Mukai, A., et al., *Impact of intraoperative goal-directed fluid therapy on major morbidity and mortality after transthoracic oesophagectomy: a multicentre, randomised controlled trial*. Br J Anaesth, 2020. **125**(6): p. 953-961.
73. Van Dessel, E., et al., *Perioperative fluid management in esophagectomy for cancer and its relation to postoperative respiratory complications*. Dis Esophagus, 2020.
74. Fahim, M., et al., *Effect of Intra-Operative Hypothermia on Post-Operative Morbidity in Patients with Colorectal Cancer*. Surg Infect (Larchmt), 2021.
75. Levy, B.F., et al., *Randomized clinical trial of epidural, spinal or patient-controlled analgesia for patients undergoing laparoscopic colorectal surgery*. Br J Surg, 2011. **98**(8): p. 1068-78.
76. Christodoulou, E., et al., *A systematic review shows no performance benefit of machine learning over logistic regression for clinical prediction models*. J Clin Epidemiol, 2019. **110**: p. 12-22.
77. Grass, F., et al., *Identification of patients eligible for discharge within 48 h of colorectal resection*. Br J Surg, 2020. **107**(5): p. 546-551.
78. Nguyen, H.B., et al., *Early goal-directed therapy in severe sepsis and septic shock: insights and comparisons to ProCESS, ProMISe, and ARISE*. Crit Care, 2016. **20**(1): p. 160.
79. Weenk, M., et al., *Wireless and continuous monitoring of vital signs in patients at the general ward*. Resuscitation, 2019. **136**: p. 47-53.
80. Wouters, M.W., M.L. Jansen-Landheer, and C.J. van de Velde, *The Quality of Cancer Care initiative in the Netherlands*. Eur J Surg Oncol, 2010. **36 Suppl 1**: p. S3-S13.
81. Marang-van de Mheen, P.J., et al., *The relative importance of quality of care information when choosing a hospital for surgical treatment: a hospital choice experiment*. Med Decis Making, 2011. **31**(6): p. 816-27.
82. Waljee, J., et al., *Patient expectations and patient-reported outcomes in surgery: a systematic review*. Surgery, 2014. **155**(5): p. 799-808.
83. Dimick, J.B., H.G. Welch, and J.D. Birkmeyer, *Surgical mortality as an indicator of hospital quality: the problem with small sample size*. Jama, 2004. **292**(7): p. 847-51.
84. Beck, C., et al., *The influence of postoperative complications on long-term prognosis in patients with colorectal carcinoma*. Int J Colorectal Dis, 2020.
85. Brown, S.R., et al., *The impact of postoperative complications on long-term quality of life after curative colorectal cancer surgery*. Ann Surg, 2014. **259**(5): p. 916-23.
86. Hayashi, T., et al., *Impact of infectious complications on gastric cancer recurrence*. Gastric Cancer, 2015. **18**(2): p. 368-74.
87. Kasahara, N., et al., *A lack of postoperative complications after pancreatectomy contributes to the long-term survival of patients with pancreatic cancer*. Pancreatology, 2019. **19**(5): p. 686-694. Tables and Figures

Supplementary information A

Table S1 – Table showing the classification of study quality of all included studies using the QUIPS tool [1]. On the horizontal axis show the different assessed domains per study. The vertical axis shows all included studies. The green bullet = little risk of bias, yellow = unclear risk of bias, red = high risk of bias.

	Study participation	Study attrition	Prognostic factor measurement	Outcome measurement	Adjustment for other prognostic factors	Statistical analysis and reporting
Abunnsra 2005	●	●	●	●	●	●
Akiyoshi 2011(1)	●	●	●	●	●	●
Akiyoshi 2011(2)	●	●	●	●	●	●
Aminian 2011	●	●	●	●	●	●
Amri 2013	●	●	●	●	●	●
Aoyama 2020	●	●	●	●	●	●
Baucom 2015	●	●	●	●	●	●
Bausys 2018	●	●	●	●	●	●
Beckmann 2017	●	●	●	●	●	●
Bertelsen 2010	●	●	●	●	●	●
Beuran 2018	●	●	●	●	●	●
Bickenbach 2013	●	●	●	●	●	●
Bisgard 2013	●	●	●	●	●	●
Blom 2012	●	●	●	●	●	●
Bludau 2015	●	●	●	●	●	●
Boccola 2011	●	●	●	●	●	●
Borggreve 2018	●	●	●	●	●	●
Brinkmann 2019	●	●	●	●	●	●
Chang 2014	●	●	●	●	●	●
Coimbra 2019	●	●	●	●	●	●
Cong 2009	●	●	●	●	●	●
Cooke 2009	●	●	●	●	●	●
Deguchi 2012	●	●	●	●	●	●
Der Hagopian 2018	●	●	●	●	●	●
Elliott 2017	●	●	●	●	●	●
Eom 2014	●	●	●	●	●	●
Eto 2016	●	●	●	●	●	●
Flynn 2020	●	●	●	●	●	●
Frasson 2015	●	●	●	●	●	●
Frasson 2016	●	●	●	●	●	●
Frouws 2017	●	●	●	●	●	●
Gessler 2016	●	●	●	●	●	●
Goense 2016	●	●	●	●	●	●
Gong 2014	●	●	●	●	●	●
Gooszen 2018	●	●	●	●	●	●
Ooto 2016	●	●	●	●	●	●
Ouner 2018	●	●	●	●	●	●
Hall 2019	●	●	●	●	●	●
Hamai 2015	●	●	●	●	●	●
Harustiak 2015	●	●	●	●	●	●
Hayashi 2019	●	●	●	●	●	●
Herrod 2019	●	●	●	●	●	●
Horne 2020	●	●	●	●	●	●
Hu 2015	●	●	●	●	●	●
Huang 2015	●	●	●	●	●	●
Huang 2020	●	●	●	●	●	●
Iancu 2008	●	●	●	●	●	●
Ishino 2014	●	●	●	●	●	●
Iversen 2008	●	●	●	●	●	●
Jeong 2011	●	●	●	●	●	●
Jeong 2014	●	●	●	●	●	●
Jiang 2012	●	●	●	●	●	●
Jiang 2014	●	●	●	●	●	●
Jones 2015	●	●	●	●	●	●
Josse 2016	●	●	●	●	●	●
Julooi 2014	●	●	●	●	●	●
Jung 2012	●	●	●	●	●	●
Jung 2014	●	●	●	●	●	●
Kanaji 2016	●	●	●	●	●	●
Kang 2011	●	●	●	●	●	●
Kang 2016	●	●	●	●	●	●
Karup 2012	●	●	●	●	●	●
Kassir 2013	●	●	●	●	●	●
Katai 2005	●	●	●	●	●	●
Katsuno 2016	●	●	●	●	●	●
Kenig 2018	●	●	●	●	●	●
Kim 2003	●	●	●	●	●	●
Kim 2009	●	●	●	●	●	●
Kim 2015	●	●	●	●	●	●
Kim 2016	●	●	●	●	●	●
Kim 2017	●	●	●	●	●	●
Kievelro 2019	●	●	●	●	●	●
Klose 2018	●	●	●	●	●	●
Kobayashi 2015	●	●	●	●	●	●
Koeter 2015	●	●	●	●	●	●
Konishi 2006	●	●	●	●	●	●
Kosuga 2017	●	●	●	●	●	●
Kouyoumdjian 2021	●	●	●	●	●	●
Kruthikava 2017	●	●	●	●	●	●
Kruschewski 2007	●	●	●	●	●	●
Kumagai 2011	●	●	●	●	●	●
Kumagai 2014	●	●	●	●	●	●
Kumar 2011	●	●	●	●	●	●
Krasnovsky 2016	●	●	●	●	●	●
Lagarde 2008	●	●	●	●	●	●
Landi 2016	●	●	●	●	●	●
Leake 2013	●	●	●	●	●	●
Lee 2012	●	●	●	●	●	●
Lee 2016 (1)	●	●	●	●	●	●
Lee 2016 (2)	●	●	●	●	●	●
Lee 2016 (3)	●	●	●	●	●	●
Lee 2017	●	●	●	●	●	●
Lee 2018	●	●	●	●	●	●
Li 2016	●	●	●	●	●	●
Li 2017	●	●	●	●	●	●
Li 2021	●	●	●	●	●	●
Lim 2016	●	●	●	●	●	●
Lindner 2017	●	●	●	●	●	●
Liou 2018	●	●	●	●	●	●
Lipska 2006	●	●	●	●	●	●
Lo 2008	●	●	●	●	●	●
Lu 2018	●	●	●	●	●	●
Manfredi 2017	●	●	●	●	●	●
Marinello 2016	●	●	●	●	●	●
Marlin 2015	●	●	●	●	●	●
Marusch 2005	●	●	●	●	●	●
Matsui 2021	●	●	●	●	●	●
Matsumoto 2016	●	●	●	●	●	●
Merikow 2008	●	●	●	●	●	●
Miki 2014	●	●	●	●	●	●
Milgrom 2014	●	●	●	●	●	●
Miltzman 2018	●	●	●	●	●	●
Miyakita 2017	●	●	●	●	●	●
Monta 2011	●	●	●	●	●	●
Mungo 2015	●	●	●	●	●	●
Mungo 2017	●	●	●	●	●	●
Nakagawa 2017	●	●	●	●	●	●
Nevo 2018	●	●	●	●	●	●
Nikolian 2017	●	●	●	●	●	●
Nishikawa 2018	●	●	●	●	●	●
Nobuoka 2008	●	●	●	●	●	●
Nordholm-Carlsten 2019	●	●	●	●	●	●
Norero 2019	●	●	●	●	●	●
Ohkura 2018	●	●	●	●	●	●
Ojima 2009	●	●	●	●	●	●
Okabe 2019	●	●	●	●	●	●
Okada 2019	●	●	●	●	●	●
Okamura 2016	●	●	●	●	●	●
Okamura 2017	●	●	●	●	●	●
Omerup 2019	●	●	●	●	●	●
Orsenigo 2014	●	●	●	●	●	●
Oshi 2018	●	●	●	●	●	●
Ostenfeld 2015	●	●	●	●	●	●
Ozer 2010	●	●	●	●	●	●
Park 2013	●	●	●	●	●	●
Parthasarathy 2017	●	●	●	●	●	●
Pellino 2018	●	●	●	●	●	●
Poelmeijer 2018	●	●	●	●	●	●
Quan 2019	●	●	●	●	●	●
Ragg 2009	●	●	●	●	●	●
Reilly 2014	●	●	●	●	●	●
Rencuzogullari 2017	●	●	●	●	●	●
Saito 2019	●	●	●	●	●	●
Salem 2017	●	●	●	●	●	●
Sato 2020	●	●	●	●	●	●
Schlotmann 2018	●	●	●	●	●	●
Schreckenbach 2018	●	●	●	●	●	●
Shen 2019	●	●	●	●	●	●
Shichonohe 2019	●	●	●	●	●	●
Shinji 2018	●	●	●	●	●	●
Smit 2016	●	●	●	●	●	●
Song 2014	●	●	●	●	●	●
Souwer 2020	●	●	●	●	●	●
Sparreboom 2018	●	●	●	●	●	●
Stromark 2016	●	●	●	●	●	●
Sugisawa 2012	●	●	●	●	●	●
Sunpawerawong 2012	●	●	●	●	●	●
Tackauchi 2016	●	●	●	●	●	●
Takeuchi 2015	●	●	●	●	●	●
Tan 2013	●	●	●	●	●	●
Tanaka 2015	●	●	●	●	●	●
Tanaka 2016	●	●	●	●	●	●
Tegels 2014	●	●	●	●	●	●
Tian 2017	●	●	●	●	●	●
Tokunaga 2009	●	●	●	●	●	●
Tran 2015	●	●	●	●	●	●
Tu 2016	●	●	●	●	●	●
Tu 2017 (1)	●	●	●	●	●	●
Tu 2017 (2)	●	●	●	●	●	●
Ushiku 2015	●	●	●	●	●	●
Van Daele 2016	●	●	●	●	●	●
Van Leersum 2014	●	●	●	●	●	●
Van Rossum 2015	●	●	●	●	●	●
Verweij 2016	●	●	●	●	●	●
Veyrie 2007	●	●	●	●	●	●
Wang 2017	●	●	●	●	●	●
Wang 2019 (1)	●	●	●	●	●	●
Wang 2019 (2)	●	●	●	●	●	●
Wang 2020	●	●	●	●	●	●
Wantanbe 2014	●	●	●	●	●	●
Wrightmann 2017	●	●	●	●	●	●
Xiao 2019 (1)	●	●	●	●	●	●
Xiao 2019 (2)	●	●	●	●	●	●
Xie 2017	●	●	●	●	●	●
Xu 2020	●	●	●	●	●	●
Yang 2017 (1)	●	●	●	●	●	●
Yang 2017 (2)	●	●	●	●	●	●
Yasui 2017	●	●	●	●	●	●
Yeh 2005	●	●	●	●	●	●
Yoshida 2014	●	●	●	●	●	●
Yoshida 2016	●	●	●	●	●	●
Yoshida 2018	●	●	●	●	●	●
You 2019	●	●	●	●	●	●
Yu 2019 (1)	●	●	●	●	●	●
Yu 2019 (2)	●	●	●	●	●	●
Yun 2017	●	●	●	●	●	●
Zaimi 2018	●	●	●	●	●	●
Zhang 2016	●	●	●	●	●	●
Zhang 2018	●	●	●	●	●	●
Zhao 2016	●	●	●	●	●	●
Zheng 2018	●	●	●	●	●	●
Zheng 2019	●	●	●	●	●	●
Zhou 2015	●	●	●	●	●	●
Zingg 2011	●	●	●	●	●	●

Table S1 –Continued

	Study participation	Study attrition	Prognostic factor measurement	Outcome measurement	Adjustment for other prognostic factors	Statistical analysis and reporting
Marin 2015	●	●	●	●	●	●
Marusch 2005	●	●	●	●	●	●
Matsui 2021	●	●	●	●	●	●
Matsumoto 2016	●	●	●	●	●	●
Merikow 2008	●	●	●	●	●	●
Miki 2014	●	●	●	●	●	●
Milgrom 2014	●	●	●	●	●	●
Miltzman 2018	●	●	●	●	●	●
Miyakita 2017	●	●	●	●	●	●
Monta 2011	●	●	●	●	●	●
Mungo 2015	●	●	●	●	●	●
Mungo 2017	●	●	●	●	●	●
Nakagawa 2017	●	●	●	●	●	●
Nevo 2018	●	●	●	●	●	●
Nikolian 2017	●	●	●	●	●	●
Nishikawa 2018	●	●	●	●	●	●
Nobuoka 2008	●	●	●	●	●	●
Nordholm-Carlsten 2019	●	●	●	●	●	●
Norero 2019	●	●	●	●	●	●
Ohkura 2018	●	●	●	●	●	●
Ojima 2009	●	●	●	●	●	●
Okabe 2019	●	●	●	●	●	●
Okada 2019	●	●	●	●	●	●
Okamura 2016	●	●	●	●	●	●
Okamura 2017	●	●	●	●	●	●
Omerup 2019	●	●	●	●	●	●
Ong 2014	●	●	●	●	●	●
Oishi 2018	●	●	●	●	●	●
Ofeld 2015	●	●	●	●	●	●
Ozer 2010	●	●	●	●	●	●
Park 2013	●	●	●	●	●	●
Pathy 2017	●	●	●	●	●	●
Plino 2018	●	●	●	●	●	●
Pijer 2018	●	●	●	●	●	●
Puan 2019	●	●	●	●	●	●
Ragg 2009	●	●	●	●	●	●
Reilly 2014	●	●	●	●	●	●
Rencuzogullari 2017	●	●	●	●	●	●
Saito 2019	●	●	●	●	●	●
Salem 2017	●	●	●	●	●	●
Sato 2020	●	●	●	●	●	●
Schlottmann 2018	●	●	●	●	●	●
Schreckenbach 2018	●	●	●	●	●	●
Shen 2019	●	●	●	●	●	●
Shichonohne 2019	●	●	●	●	●	●
Shinji 2018	●	●	●	●	●	●
Smit 2016	●	●	●	●	●	●
Song 2014	●	●	●	●	●	●
Souwer 2020	●	●	●	●	●	●
Sparreboom 2018	●	●	●	●	●	●
Stromark 2016	●	●	●	●	●	●
Sugisawa 2012	●	●	●	●	●	●
Sunpawerawong 2012	●	●	●	●	●	●
Tackeuchi 2016	●	●	●	●	●	●
Takeuchi 2015	●	●	●	●	●	●
Tan 2013	●	●	●	●	●	●
Tanaka 2015	●	●	●	●	●	●
Tanaka 2016	●	●	●	●	●	●
Tegels 2014	●	●	●	●	●	●
Tian 2017	●	●	●	●	●	●
Tokunaga 2009	●	●	●	●	●	●
Tran 2015	●	●	●	●	●	●
Tu 2016	●	●	●	●	●	●
Tu 2017 (1)	●	●	●	●	●	●
Tu 2017 (2)	●	●	●	●	●	●
Ushiku 2015	●	●	●	●	●	●
Van Daele 2016	●	●	●	●	●	●
Van Leersum 2014	●	●	●	●	●	●
Van Rossum 2015	●	●	●	●	●	●
Verweij 2016	●	●	●	●	●	●
Veyrie 2007	●	●	●	●	●	●
Wang 2017	●	●	●	●	●	●
Wang 2019 (1)	●	●	●	●	●	●
Wang 2019 (2)	●	●	●	●	●	●
Wang 2020	●	●	●	●	●	●
Wantanbe 2014	●	●	●	●	●	●
Wightmann 2017	●	●	●	●	●	●
Xiao 2019 (1)	●	●	●	●	●	●
Xiao 2019 (2)	●	●	●	●	●	●
Xie 2017	●	●	●	●	●	●
Xu 2020	●	●	●	●	●	●
Yang 2017 (1)	●	●	●	●	●	●
Yang 2017 (2)	●	●	●	●	●	●
Yasui 2017	●	●	●	●	●	●
Yeh 2005	●	●	●	●	●	●
Yoshida 2014	●	●	●	●	●	●
Yoshida 2016	●	●	●	●	●	●
Yoshida 2018	●	●	●	●	●	●
You 2019	●	●	●	●	●	●
Yu 2019 (1)	●	●	●	●	●	●
Yu 2019 (2)	●	●	●	●	●	●
Yun 2017	●	●	●	●	●	●
Zaimi 2018	●	●	●	●	●	●
Zhang 2016	●	●	●	●	●	●
Zhang 2018	●	●	●	●	●	●
Zhao 2016	●	●	●	●	●	●
Zheng 2018	●	●	●	●	●	●
Zheng 2019	●	●	●	●	●	●
Zhou 2015	●	●	●	●	●	●
Zingg 2011	●	●	●	●	●	●

Supplementary Information B

2. Sparreboom, C.L., et al., *Different Risk Factors for Early and Late Colorectal Anastomotic Leakage in a Nationwide Audit*. 2018. **61**(11): p. 1258-1266.
25. Quan, H., et al., *The effect of preoperative smoking cessation and smoking dose on postoperative complications following radical gastrectomy for gastric cancer: a retrospective study of 2469 patients*. 2019. **17**(1): p. 61.
26. Parthasarathy, M., et al., *Risk factors for anastomotic leakage after colorectal resection: a retrospective analysis of 17 518 patients*. 2017. **19**(3): p. 288-298.
49. Lu, J., et al., *Major perioperative complications in laparoscopic spleen-preserving total gastrectomy for gastric cancer: perspectives from a high-volume center*. 2016. **30**(3): p. 1034-42.
53. Mungo, B., et al., *Does neoadjuvant therapy for esophageal cancer increase postoperative morbidity or mortality?* Diseases of the esophagus : official journal of the International Society for Diseases of the Esophagus, 2015. **28**(7): p. 644-651.
88. Tanaka, K., et al., *Risk factors for anastomotic leakage after laparoscopic surgery with the double stapling technique for stage 0/I rectal carcinoma: a subgroup analysis of a multicenter, single-arm phase II trial*. Surgery today, 2017. **47**(10): p. 1215-1222.
89. Manfredi, S., et al., *Time trends in colorectal cancer early postoperative mortality. A French 25-year population-based study*. International journal of colorectal disease, 2017. **32**(12): p. 1725-1731.
90. Wang, W.-J., et al., *Systematic assessment of complications after robotic-assisted total versus distal gastrectomy for advanced gastric cancer: A retrospective propensity score-matched study using Clavien-Dindo classification*. International journal of surgery (London, England), 2019. **71**: p. 140-148.
91. Yang, S.Y., et al., *Surgical Outcomes After Open, Laparoscopic, and Robotic Gastrectomy for Gastric Cancer*. Annals of surgical oncology, 2017. **24**(7): p. 1770-1777.
92. Kenig, J., et al., *The Surgical Apgar Score predicts outcomes of emergency abdominal surgeries both in fit and frail older patients*. 2018. **76**: p. 54-59.
93. Yang, J.-Y., et al., *Short- and Long-Term Outcomes After Gastrectomy in Elderly Gastric Cancer Patients*. 2017. **24**(2): p. 469-477.
94. Katai, H., et al., *Risk factors for pancreas-related abscess after total gastrectomy*. 2005. **8**(3): p. 137-41.
95. Kim, J.S., et al., *Risk factors for anastomotic leakage after laparoscopic intracorporeal colorectal anastomosis with a double stapling technique*. Journal of the American College of Surgeons, 2009. **209**(6): p. 694-701.
96. Lo, C.-H., et al., *Risk factors and management of intra-abdominal infection after extended radical gastrectomy*. 2008. **196**(5): p. 741-5.
97. Jeong, O., et al., *Risk factors and learning curve associated with postoperative morbidity of laparoscopic total gastrectomy for gastric carcinoma*. 2014. **21**(9): p. 2994-3001.
98. Van Daele, E., et al., *Risk factors and consequences of anastomotic leakage after Ivor Lewis oesophagectomy†*. 2016. **22**(1): p. 32-7.
99. Ragg, J.L., D.A. Watters, and G.D. Guest, *Preoperative risk stratification for mortality and major morbidity in major colorectal surgery*. Diseases of the colon and rectum, 2009. **52**(7): p. 1296-1303.
100. Lagarde, S.M., et al., *Preoperative prediction of the occurrence and severity of complications after esophagectomy for cancer with use of a nomogram*. The Annals of thoracic surgery, 2008. **85**(6): p. 1938-1945.
101. Liou, D.Z., et al., *Predictors of Failure to Rescue After Esophagectomy*. 2018. **105**(3): p. 871-878.
102. Marusch, F., et al., *The impact of the risk factor "age" on the early postoperative results of surgery for colorectal carcinoma and its significance for perioperative management*. World J Surg, 2005. **29**(8): p. 1013-21; discussion 1021-2.
103. Takeuchi, M., et al., *Excessive visceral fat area as a risk factor for early postoperative complications of total gastrectomy for gastric cancer: a retrospective cohort study*. 2016. **16**(1): p. 54.
104. Eom, B.W., et al., *A body shape index has a good correlation with postoperative complications in gastric cancer surgery*. 2014. **21**(4): p. 1115-22.
105. Frouws, M.A., et al., *Clinical Relevance of a Grading System for Anastomotic Leakage After Low Anterior Resection: Analysis From a National Cohort Database*. Diseases of the colon and rectum, 2017. **60**(7): p. 706-713.
106. Lee, J.Y., et al., *Clinical Significance of the Prognostic Nutritional Index for Predicting Short- and Long-Term Surgical Outcomes After Gastrectomy: A Retrospective Analysis of 7781 Gastric Cancer Patients*. 2016. **95**(18): p. e3539.
107. Tian, Y., et al., *Comorbidity and the risk of anastomotic leak in Chinese patients with colorectal cancer undergoing colorectal surgery*. International journal of colorectal disease, 2017. **32**(7): p. 947-953.
108. Tu, R.H., et al., *Development of a nomogram for predicting the risk of anastomotic leakage after a gastrectomy for gastric cancer*. European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology, 2017. **43**(2): p. 485-492.
109. Zaimi, I., et al., *The effect of age on anastomotic leakage in colorectal cancer surgery: A population-based study*. Journal of surgical oncology, 2018. **118**(1): p. 113-120.
110. Zhou, J., et al., *Evaluation of Clavien-Dindo classification in patients undergoing total gastrectomy for gastric cancer*. 2015. **32**(4): p. 120.
111. Kodera, Y., et al., *Identification of risk factors for the development of complications following extended and superextended lymphadenectomies for gastric cancer*. The British journal of surgery, 2005. **92**(9): p. 1103-1109.

112. Martin, J.T., et al., *Identifying Esophagectomy Patients at Risk for PredischARGE Versus Postdischarge Venous Thromboembolism*. The Annals of thoracic surgery, 2015. **100**(3): p. 932-938.
113. Bickenbach, K.A., et al., *Impact of obesity on perioperative complications and long-term survival of patients with gastric cancer*. Annals of surgical oncology, 2013. **20**(3): p. 780-787.
114. Mungo, B., et al., *The Impact of Operative Approach on Postoperative Complications Following Colectomy for Colon Cancer*. World journal of surgery, 2017. **41**(8): p. 2143-2152.
115. Shichinohe, T., et al., *Impact of Preoperative Skeletal Muscle Mass and Nutritional Status on Short-and Long-Term Outcomes After Esophagectomy for Esophageal Cancer: A Retrospective Observational Study: Impact of Psoas Muscle Mass and Body Mass on Esophagectomy*. 2019. **26**(5): p. 1301-1310.
116. Morita, M., et al., *In-hospital mortality after a surgical resection for esophageal cancer: analyses of the associated factors and historical changes*. Annals of surgical oncology, 2011. **18**(6): p. 1757-1765.
117. Shinji, S., et al., *Male sex and history of ischemic heart disease are major risk factors for anastomotic leakage after laparoscopic anterior resection in patients with rectal cancer*. BMC gastroenterology, 2018. **18**(1): p. 117-117.
118. Landi, F., et al., *Morbidity after laparoscopic and open rectal cancer surgery: a comparative analysis of morbidity in octogenarians and younger patients*. Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland, 2016. **18**(5): p. 459-467.
119. Park, J.S., et al., *Multicenter analysis of risk factors for anastomotic leakage after laparoscopic rectal cancer excision: the Korean laparoscopic colorectal surgery study group*. Annals of surgery, 2013. **257**(4): p. 665-671.
120. Krarup, P.-M., et al., *A nationwide study on anastomotic leakage after colonic cancer surgery*. Vol. 14. 2012. e661-7.
121. Saito, T., et al., *Novel prognostic score of postoperative complications after transthoracic minimally invasive esophagectomy for esophageal cancer: a retrospective cohort study of 90 consecutive patients*. 2019. **16**(2): p. 155-161.
122. Gong, J.-P., et al., *Outcomes based on risk assessment of anastomotic leakage after rectal cancer surgery*. Asian Pacific journal of cancer prevention : APJCP, 2014. **15**(2): p. 707-712.
123. Guner, A., et al., *Parameters for Predicting Surgical Outcomes for Gastric Cancer Patients: Simple Is Better Than Complex*. 2018. **25**(11): p. 3239-3247.
124. Bisgård, A.S., et al., *Perioperative statin therapy is not associated with reduced risk of anastomotic leakage after colorectal resection*. Diseases of the colon and rectum, 2013. **56**(8): p. 980-986.
125. der Hagopian, O., et al., *Perirenal fat surface area as a risk factor for perioperative difficulties and 30-day postoperative complications in elective colon cancer surgery*. Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland, 2018. **20**(12): p. 1078-1087.
126. Iversen, L.H., et al., *Postoperative medical complications are the main cause of early death after emergency surgery for colonic cancer*. The British journal of surgery, 2008. **95**(8): p. 1012-1019.
127. Schlottmann, F., et al., *Postoperative outcomes of esophagectomy for cancer in elderly patients*. 2018. **229**: p. 9-14.
128. Kumagai, K., et al., *Potentially fatal complications for elderly patients after laparoscopy-assisted distal gastrectomy*. 2014. **17**(3): p. 548-55.
129. Ostensfeld, E.B., et al., *Preadmission glucocorticoid use and anastomotic leakage after colon and rectal cancer resections: a Danish cohort study*. BMJ open, 2015. **5**(9): p. e008045-e008045.
130. Zhang, W.-T., et al., *Sarcopenic Obesity Is Associated with Severe Postoperative Complications in Gastric Cancer Patients Undergoing Gastrectomy: a Prospective Study*. 2018. **22**(11): p. 1861-1869.
131. Hamai, Y., et al., *Effects of neoadjuvant chemoradiotherapy on postoperative morbidity and mortality associated with esophageal cancer*. Vol. 28. 2015. 358-64.
132. Kumar, A., et al., *Anterior resection for rectal carcinoma - risk factors for anastomotic leaks and strictures*. 2011. **17**(11): p. 1475-9.
133. Zhang, K., et al., *Ability of Serum C-Reactive Protein Concentrations to Predict Complications After Laparoscopy-Assisted Gastrectomy: A Prospective Cohort Study*. Medicine, 2016. **95**(21): p. e3798-e3798.
134. Kanaji, S., et al., *Can the intraoperative leak test prevent postoperative leakage of esophagojejunal anastomosis after total gastrectomy? Surgery today*, 2016. **46**(7): p. 815-820.
135. Coimbra, F.J.F., et al., *Predicting overall and major postoperative morbidity in gastric cancer patients*. Journal of surgical oncology, 2019. **120**(8): p. 1371-1378.
136. Goense, L., et al., *Aortic Calcification Increases the Risk of Anastomotic Leakage After Ivor-Lewis Esophagectomy*. 2016. **102**(1): p. 247-52.
137. Jung, M.R., et al., *Definition and classification of complications of gastrectomy for gastric cancer based on the accordion severity grading system*. 2012. **36**(10): p. 2400-11.
138. Bertelsen, C.A., et al., *Anastomotic leakage after anterior resection for rectal cancer: risk factors*. 2010. **12**(1): p. 37-43.
139. Tran, T.B., et al., *Outcomes of Gastric Cancer Resection in Octogenarians: A Multi-institutional Study of the U.S. Gastric Cancer Collaborative*. 2015. **22**(13): p. 4371-9.
140. Rencuzogullari, A., et al., *Predictors of Anastomotic Leak in Elderly Patients After Colectomy: Nomogram-Based Assessment From the American College of Surgeons National Surgical Quality Program Procedure-Targeted Cohort*. 2017. **60**(5): p. 527-536.



141. Abunasra, H., et al., *Predictors of operative death after oesophagectomy for carcinoma*. 2005. **92**(8): p. 1029-33.
142. Yu, Y., et al., *Preoperative C-reactive protein-to-albumin ratio predicts anastomotic leak in elderly patients after curative colorectal surgery*. *Cancer biomarkers : section A of Disease markers*, 2019: p. 10.3233/CBM-190470.
143. Lee, E., et al., *Prospective Study on the Incidence of Postoperative Venous Thromboembolism in Korean Patients with Colorectal Cancer*. *Cancer research and treatment : official journal of Korean Cancer Association*, 2016. **48**(3): p. 978-989.
144. Lee, K.-G., et al., *Risk factors associated with complication following gastrectomy for gastric cancer: retrospective analysis of prospectively collected data based on the Clavien-Dindo system*. 2014. **18**(7): p. 1269-77.
145. Nevo, Y., et al., *Risk Factors for Complications of Total/Subtotal Gastrectomy for Gastric Cancer: Prospectively Collected, Based on the Clavien-Dindo Classification System*. 2018. **5**(20): p. 277-280.
146. Okumura, N., et al., *Robotic gastrectomy for elderly gastric cancer patients: comparisons with robotic gastrectomy in younger patients and laparoscopic gastrectomy in the elderly*. *Gastric cancer : official journal of the International Gastric Cancer Association and the Japanese Gastric Cancer Association*, 2016. **19**(4): p. 1125-1134.
147. Huang, C.-M., et al., *A scoring system to predict the risk of postoperative complications after laparoscopic gastrectomy for gastric cancer based on a large-scale retrospective study*. 2015. **94**(17): p. e812.
148. Wang, W.-J., et al., *Severity and incidence of complications assessed by the Clavien-Dindo classification following robotic and laparoscopic gastrectomy for advanced gastric cancer: a retrospective and propensity score-matched study*. *Surgical endoscopy*, 2019. **33**(10): p. 3341-3354.
149. Beckmann, K., et al., *Short-term outcomes after surgical resection for colorectal cancer in South Australia*. 2017. **23**(2): p. 316-324.
150. Nakagawa, A., et al., *The Surgical Apgar Score Predicts Not Only Short-Term Complications But Also Long-Term Prognosis After Esophagectomy*. *Annals of surgical oncology*, 2017. **24**(13): p. 3934-3946.
151. Ozer, I., et al., *Surgical treatment for gastric cancer in Turkish patients over age 70: early postoperative results and risk factors for mortality*. 2010. **395**(8): p. 1101-6.
152. van Leersum, N.J., et al., *Synchronous colorectal carcinoma: a risk factor in colorectal cancer surgery*. Vol. 57. 2014. 460-6.
153. Tegels, J.J.W., et al., *Value of geriatric frailty and nutritional status assessment in predicting postoperative mortality in gastric cancer surgery*. 2014. **18**(3): p. 439-45; discussion 445-6.
154. Flynn, D.E., Mao, D., Yerkovich, S.T., et al., *The impact of comorbidities on post-operative complications following colorectal cancer surgery*. 2020. **15**(12): p. e0243995.
155. Coimbra, F.J.F., de Jesus V.H.F., Franco, C.P., et al., *Predicting overall and major postoperative morbidity in gastric cancer patients*. 2019. **120**(8): p. 1371-1378.
156. Horne, Z.D., Wegner, R.E., Colonias, A., et al., *Drivers of 30- and 90-day Postoperative Death After Neoadjuvant Chemoradiation for Esophageal Cancer*. 2020. **109**(3): p. 921-926.
157. Li, Z.-Y., Zhao, Y.-L., Qian, F., et al., *Incidence and risk factors of postoperative complications after robotic gastrectomy for gastric cancer: an analysis of 817 cases based on 10-year experience in a large-scale center*. 2021.
158. Huang, C., Yao, H., Huang, Q.J., et al., *A novel nomogram to predict the risk of anastomotic leakage in patients after oesophagectomy*. 2020. **20**(1): p. 64.
159. Tanaka, Y., et al., *Usefulness of preoperative estimated glomerular filtration rate to predict complications after curative gastrectomy in patients with clinical T2-4 gastric cancer*. 2017. **20**(4): p. 736-743.
160. Miyakita, H., et al., *Risk scores as useful predictors of perioperative complications in patients with rectal cancer who received radical surgery*. 2017. **22**(2): p. 324-331.
161. Ushiku, H., et al., *A Risk Model for Surgical Site Infection in the Gastric Cancer Surgery Using Data of 790 Patients*. 2015. **32**(6): p. 472-9.
162. Gessler, B., et al., *Risk factors for anastomotic dehiscence in colon cancer surgery--a population-based registry study*. *International journal of colorectal disease*, 2016. **31**(4): p. 895-902.
163. Nobuoka, D., et al., *Prevention of postoperative pancreatic fistula after total gastrectomy*. 2008. **32**(10): p. 2261-6.
164. Miki, Y., et al., *Perioperative risk assessment for gastrectomy by surgical apgar score*. *Annals of surgical oncology*, 2014. **21**(8): p. 2601-2607.
165. Nikolian, V.C., et al., *Anastomotic leak after colorectal resection: A population-based study of risk factors and hospital variation*. *Surgery*, 2017. **161**(6): p. 1619-1627.
166. Kim, C.W., et al., *Anastomotic Leakage After Low Anterior Resection for Rectal Cancer Is Different Between Minimally Invasive Surgery and Open Surgery*. *Annals of surgery*, 2016. **263**(1): p. 130-137.
167. Lipska, M.A., et al., *Anastomotic leakage after lower gastrointestinal anastomosis: men are at a higher risk*. *ANZ journal of surgery*, 2006. **76**(7): p. 579-585.
168. Yu, B., Park, K.B., Park, J.Y., et al., *Bioelectrical Impedance Analysis for Prediction of Early Complications after Gastrectomy in Elderly Patients with Gastric Cancer: the Phase Angle Measured Using Bioelectrical Impedance Analysis*. *Journal of gastric cancer*, 2019. **19**(3): p. 278-289.
169. Leake, P.A., et al., *Colorectal anastomotic leakage at the university hospital of the west indies: an analysis of risk factors*. *The West Indian medical journal*, 2013. **62**(8): p. 711-715.
170. Verweij, N.M., et al., *Colorectal Cancer Resections in the Oldest Old Between 2011 and 2012 in The Netherlands*. 2016. **23**(6): p. 1875-82.

171. Yasui, M., et al., *Comparison of bleeding risks related to venous thromboembolism prophylaxis in laparoscopic vs open colorectal cancer surgery: a multicenter study in Japanese patients*. American journal of surgery, 2017. **213**(1): p. 43-49.
172. Katsuno, H., et al., *Comparison of symptomatic anastomotic leakage following laparoscopic and open low anterior resection for rectal cancer: a propensity score matching analysis of 1014 consecutive patients*. Surgical endoscopy, 2016. **30**(7): p. 2848-2856.
173. Goto, S., et al., *Differences in surgical site infection between laparoscopic colon and rectal surgeries: sub-analysis of a multicenter randomized controlled trial (Japan-Multinational Trial Organization PREV 07-01)*. International journal of colorectal disease, 2016. **31**(11): p. 1775-1784.
174. Kobayashi, D., et al., *Factors related to occurrence and aggravation of pancreatic fistula after radical gastrectomy for gastric cancer*. Journal of surgical oncology, 2015. **112**(4): p. 381-386.
175. Reilly, F., et al., *Incidence, risks and outcome of radiological leak following early contrast enema after anterior resection*. International journal of colorectal disease, 2014. **29**(4): p. 453-458.
176. Zheng, H., et al., *Laparoscopic surgery may decrease the risk of clinical anastomotic leakage and a nomogram to predict anastomotic leakage after anterior resection for rectal cancer*. 2019. **34**(2): p. 319-328.
177. Stormark, K., et al., *Nationwide implementation of laparoscopic surgery for colon cancer: short-term outcomes and long-term survival in a population-based cohort*. Surgical endoscopy, 2016. **30**(11): p. 4853-4864.
178. Hoshino, N., et al., *Nomogram for predicting anastomotic leakage after low anterior resection for rectal cancer*. 2018. **33**(4): p. 411-418.
179. Lee, S.Y., et al., *Nutritional risk screening score is an independent predictive factor of anastomotic leakage after rectal cancer surgery*. 2018. **72**(4): p. 489-495.
180. Wang, L.J., et al., *[Ridge regression analysis of pulmonary infection rate after transthoracic esophagectomy for esophageal cancer]*. Zhonghua Yi Xue Za Zhi, 2012. **92**(19): p. 1310-3.
181. Kosuga, T., et al., *Clinical and surgical factors associated with organ/space surgical site infection after laparoscopic gastrectomy for gastric cancer*. 2017. **31**(4): p. 1667-1674.
182. Hu, X. and Y. Cheng, *A Clinical Parameters-Based Model Predicts Anastomotic Leakage After a Laparoscopic Total Mesorectal Excision: A Large Study With Data From China*. 2015. **94**(26): p. e1003.
183. Kang, S.C., H.I. Kim, and M.G. Kim, *Low Serum Albumin Level, Male Sex, and Total Gastrectomy Are Risk Factors of Severe Postoperative Complications in Elderly Gastric Cancer Patients*. 2016. **16**(1): p. 43-50.
184. Yun, J.-A., et al., *Clinical manifestations and risk factors of anastomotic leakage after low anterior resection for rectal cancer*. 2017. **87**(11): p. 908-914.
185. Pellino, G., et al., *Predictors of complications and mortality following left colectomy with primary stapled anastomosis for cancer: results of a multicentric study with 1111 patients*. Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland, 2018. **20**(11): p. 986-995.
186. You, X., et al., *Preoperative albumin-to-fibrinogen ratio predicts severe postoperative complications in elderly gastric cancer subjects after radical laparoscopic gastrectomy*. 2019. **19**(1): p. 931.
187. Chang, J.S., et al., *Preoperative chemoradiotherapy effects on anastomotic leakage after rectal cancer resection: a propensity score matching analysis*. Annals of surgery, 2014. **259**(3): p. 516-521.
188. Li, P., et al., *Risk factors affecting unplanned reoperation after laparoscopic gastrectomy for gastric cancer: experience from a high-volume center*. 2017. **31**(10): p. 3922-3931.
189. Frasson, M., et al., *Risk Factors for Anastomotic Leak After Colon Resection for Cancer: Multivariate Analysis and Nomogram From a Multicentric, Prospective, National Study With 3193 Patients*. Annals of surgery, 2015. **262**(2): p. 321-330.
190. Frasson, M., et al., *Risk factors for anastomotic leak and postoperative morbidity and mortality after elective right colectomy for cancer: results from a prospective, multicentric study of 1102 patients*. International journal of colorectal disease, 2016. **31**(1): p. 105-114.
191. Schreckenbach, T., et al., *Identification of factors predictive of postoperative morbidity and short-term mortality in older patients after colorectal carcinoma resection: A single-center retrospective study*. J Geriatr Oncol, 2018. **9**(6): p. 649-658.
192. Tan, W.P., et al., *American Society of Anesthesiologists class and Charlson's comorbidity index as predictors of postoperative colorectal anastomotic leak: a single-institution experience*. 2013. **184**(1): p. 115-9.
193. Zhao, L., et al., *Calcification of arteries supplying the gastric tube increases the risk of anastomotic leakage after esophagectomy with cervical anastomosis*. Journal of thoracic disease, 2016. **8**(12): p. 3551-3562.
194. Gooszen, J.A.H., et al., *Intrathoracic versus cervical anastomosis and predictors of anastomotic leakage after oesophagectomy for cancer*. The British journal of surgery, 2018. **105**(5): p. 552-560.
195. Hayashi, M., et al., *A low surgical Apgar score is a predictor of anastomotic leakage after transthoracic esophagectomy, but not a prognostic factor*. Esophagus : official journal of the Japan Esophageal Society, 2019. **16**(4): p. 386-394.
196. Lee, K., et al., *Surgical Outcome of Colon Interposition in Esophageal Cancer Surgery: Analysis of Risk Factors for Conduit-Related Morbidity*. The Thoracic and cardiovascular surgeon, 2018. **66**(5): p. 384-389.

197. Onerup, A., et al., *Self-assessed preoperative level of habitual physical activity predicted postoperative complications after colorectal cancer surgery: A prospective observational cohort study*. European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology, 2019. **45**(11): p. 2045-2051.
198. Yoshida, N., et al., *Preoperative Nutritional Assessment by Controlling Nutritional Status (CONUT) is Useful to estimate Postoperative Morbidity After Esophagectomy for Esophageal Cancer*. 2016. **40**(8): p. 1910-7.
199. Okabe, H., et al., *Frailty predicts severe postoperative complications after elective colorectal surgery*. American journal of surgery, 2019. **217**(4): p. 677-681.
200. Cooke, D.T., et al., *Analysis of cervical esophagogastric anastomotic leaks after transhiatal esophagectomy: risk factors, presentation, and detection*. Ann Thorac Surg, 2009. **88**(1): p. 177-84; discussion 184-5.
201. Nordholm-Carstensen, A., M. Schnack Rasmussen, and P.-M. Krarup, *Increased Leak Rates Following Stapled Versus Handsewn Ileocolic Anastomosis in Patients with Right-Sided Colon Cancer: A Nationwide Cohort Study*. Diseases of the colon and rectum, 2019. **62**(5): p. 542-548.
202. Zingg, U., et al., *Factors associated with postoperative pulmonary morbidity after esophagectomy for cancer*. Annals of surgical oncology, 2011. **18**(5): p. 1460-1468.
203. Kim, M.C., et al., *Risk factors associated with complication following laparoscopy-assisted gastrectomy for gastric cancer: a large-scale korean multicenter study*. Annals of surgical oncology, 2008. **15**(10): p. 2692-2700.
204. van Rossum, P.S.N., et al., *Calcification of arteries supplying the gastric tube: a new risk factor for anastomotic leakage after esophageal surgery*. 2015. **274**(1): p. 124-32.
205. Borggreve, A.S., et al., *Generalized cardiovascular disease on a preoperative CT scan is predictive for anastomotic leakage after esophagectomy*. 2018. **44**(5): p. 587-593.
206. Brinkmann, S., et al., *Stenosis of the celiac trunk is associated with anastomotic leak after Ivor-Lewis esophagectomy*. Diseases of the esophagus : official journal of the International Society for Diseases of the Esophagus, 2019. **32**(7): p. doy107.
207. Harustiak, T., et al., *Anastomotic leak and stricture after hand-sewn versus linear-stapled intrathoracic oesophagogastric anastomosis: single-centre analysis of 415 oesophagectomies*. European journal of cardio-thoracic surgery : official journal of the European Association for Cardio-thoracic Surgery, 2016. **49**(6): p. 1650-1659.
208. Aminian, A., et al., *Predictors and outcome of cervical anastomotic leakage after esophageal cancer surgery*. 2011. **7**(4): p. 448-53.
209. Kassis, E.S., et al., *Predictors of anastomotic leak after esophagectomy: an analysis of the society of thoracic surgeons general thoracic database*. The Annals of thoracic surgery, 2013. **96**(6): p. 1919-1926.
210. Iancu, C., et al., *Host-related predictive factors for anastomotic leakage following large bowel resections for colorectal cancer*. Journal of gastrointestinal and liver diseases : JGLD, 2008. **17**(3): p. 299-303.
211. Deguchi, Y., et al., *Identification of risk factors for esophagojejunal anastomotic leakage after gastric surgery*. 2012. **36**(7): p. 1617-22.
212. Beuran, M., et al., *Nonelective Left-Sided Colon Cancer Resections are Associated with Worse Postoperative and Oncological Outcomes: A Propensity-Matched Study*. Chirurgia (Bucharest, Romania : 1990), 2018. **113**(2): p. 218-226.
213. Klevebro, F., et al., *Cardiorespiratory Comorbidity and Postoperative Complications following Esophagectomy: a European Multicenter Cohort Study*. 2019. **26**(9): p. 2864-2873.
214. Bludau, M., et al., *Preoperative airway colonization prior to transthoracic esophagectomy predicts postoperative pulmonary complications*. 2015. **400**(6): p. 707-14.
215. Orsenigo, E., et al., *Duodenal stump fistula after gastric surgery for malignancies: a retrospective analysis of risk factors in a single centre experience*. 2014. **17**(4): p. 733-44.
216. Yoshida, N., et al., *Influence of preoperative docetaxel, cisplatin, and 5-fluorouracil on the incidence of complications after esophagectomy for resectable advanced esophageal cancer*. 2014. **27**(4): p. 374-9.
217. Matsumoto, S., et al., *Preoperative Cardiac Risk Assessment and Surgical Outcomes of Patients with Gastric Cancer*. 2016. **23**: p. S222-9.
218. Matsui, K., Sakuramoto, S., Sugita, H., et al., *C-reactive protein adjusted for body mass index as a predictor of postoperative complications following laparoscopic gastrectomy for gastric cancer*. 2021.
219. Shen, Z., An, Y., Shi, Y., et al., *The Aortic Calcification Index is a risk factor associated with anastomotic leakage after anterior resection of rectal cancer*. Vol. 21. 2019, England. 1397-1404.
220. Okamura, A., et al., *Preoperative Glycosylated Hemoglobin Levels Predict Anastomotic Leak After Esophagectomy with Cervical Esophagogastric Anastomosis*. 2017. **41**(1): p. 200-207.
221. Kruschewski, M., et al., *Risk factors for clinical anastomotic leakage and postoperative mortality in elective surgery for rectal cancer*. 2007. **22**(8): p. 919-27.
222. Cong, Z.-j., et al., *Influencing factors of symptomatic anastomotic leakage after anterior resection of the rectum for cancer*. World journal of surgery, 2009. **33**(6): p. 1292-1297.
223. Nishikawa, K., et al., *Association of level of anastomosis and anastomotic leak after esophagectomy in anterior mediastinal reconstruction*. Esophagus : official journal of the Japan Esophageal Society, 2018. **15**(4): p. 231-238.
224. Hall, B.R., et al., *Risk factors for anastomotic leak after esophagectomy for cancer: A NSQIP procedure-targeted analysis*. 2019. **120**(4): p. 661-669.

225. Xu, H., and Kong, K., *Malnutrition-Related Factors Increased the Risk of Anastomotic Leak for Rectal Cancer Patients Undergoing Surgery*. 2020. **2020**: p. 5059670.
226. Sato, S., Nakatani, E., Higashizono, K., et al., *Size of the thoracic inlet predicts cervical anastomotic leak after retrosternal reconstruction after esophagectomy for esophageal cancer*. 2020. **168**(3): p. 558-566.
227. Sunpaweravong, S., et al., *Prediction of major postoperative complications and survival for locally advanced esophageal carcinoma patients*. Asian journal of surgery, 2012. **35**(3): p. 104-109.
228. Konishi, T., et al., *Risk factors for anastomotic leakage after surgery for colorectal cancer: results of prospective surveillance*. Journal of the American College of Surgeons, 2006. **202**(3): p. 439-444.
229. Yoshida, N., et al., *Clinical utility of exhaled carbon monoxide in assessing preoperative smoking status and risks of postoperative morbidity after esophagectomy*. 2018. **31**(9).
230. Jung, K.H., et al., *Preoperative smoking cessation can reduce postoperative complications in gastric cancer surgery*. Gastric cancer : official journal of the International Gastric Cancer Association and the Japanese Gastric Cancer Association, 2015. **18**(4): p. 683-690.
231. Baucom, R.B., et al., *Smoking as dominant risk factor for anastomotic leak after left colon resection*. 2015. **210**(1): p. 1-5.
232. Aoyama, T., Atsumi, Y., Hara, K., et al., *Risk Factors for Postoperative Anastomosis Leak After Esophagectomy for Esophageal Cancer*. 2020. **34**(2): p. 857-862.
233. Merkow, R.P., et al., *Effect of body mass index on short-term outcomes after colectomy for cancer*. 2009. **208**(1): p. 53-61.
234. Tokunaga, M., et al., *Effect of individual fat areas on early surgical outcomes after open gastrectomy for gastric cancer*. 2009. **96**(5): p. 496-500.
235. Wightman, S.C., et al., *Extremes of body mass index and postoperative complications after esophagectomy*. Diseases of the esophagus : official journal of the International Society for Diseases of the Esophagus, 2017. **30**(5): p. 1-6.
236. Ojima, T., et al., *Influence of overweight on patients with gastric cancer after undergoing curative gastrectomy: an analysis of 689 consecutive cases managed by a single center*. 2009. **144**(4): p. 351-8; discussion 358.
237. Juloori, A., et al., *Influence of preoperative radiation field on postoperative leak rates in esophageal cancer patients after trimodality therapy*. 2014. **9**(4): p. 534-40.
238. Poelemeijer, Y.Q.M., et al., *Obesity as a determinant of perioperative and postoperative outcome in patients following colorectal cancer surgery: A population-based study (2009-2016)*. European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology, 2018. **44**(12): p. 1849-1857.
239. Mitzman, B., et al., *Complications After Esophagectomy Are Associated With Extremes of Body Mass Index*. 2018. **106**(4): p. 973-980.
240. Tu, R.-H., et al., *A scoring system to predict the risk of organ/space surgical site infections after laparoscopic gastrectomy for gastric cancer based on a large-scale retrospective study*. 2016. **30**(7): p. 3026-34.
241. Akiyoshi, T., et al., *Effect of body mass index on short-term outcomes of patients undergoing laparoscopic resection for colorectal cancer: a single institution experience in Japan*. Surgical laparoscopy, endoscopy & percutaneous techniques, 2011. **21**(6): p. 409-414.
242. Herrod, P.J.J., et al., *Quick and simple; psoas density measurement is an independent predictor of anastomotic leak and other complications after colorectal resection*. Techniques in coloproctology, 2019. **23**(2): p. 129-134.
243. Elliott, J.A., et al., *Sarcopenia: Prevalence, and Impact on Operative and Oncologic Outcomes in the Multimodal Management of Locally Advanced Esophageal Cancer*. Annals of surgery, 2017. **266**(5): p. 822-830.
244. Jones, K.I., et al., *Simple psoas cross-sectional area measurement is a quick and easy method to assess sarcopenia and predicts major surgical complications*. 2015. **17**(1): p. O20-6.
245. Lu, J., et al., *A Novel Preoperative Skeletal Muscle Measure as a Predictor of Postoperative Complications, Long-Term Survival and Tumor Recurrence for Patients with Gastric Cancer After Radical Gastrectomy*. 2018. **25**(2): p. 439-448.
246. Wang, P.-Y., Chen, X.-K., Liu, Q., et al., *Highlighting sarcopenia management for promoting surgical outcomes in esophageal cancers: Evidence from a prospective cohort study*. 2020. **83**: p. 206-215.
247. Souwer, E.T.D., Moos, S.L., van Rooden, C.J., et al., *Physical performance has a strong association with poor surgical outcome in older patients with colorectal cancer*. 2020. **46**(3): p. 462-469.
248. Oshi, M., et al., *Risk Factors for Anastomotic Leakage of Esophagojejunostomy after Laparoscopy-Assisted Total Gastrectomy for Gastric Cancer*. 2018. **35**(1): p. 28-34.
249. Veyrie, N., et al., *Anastomotic leakage after elective right versus left colectomy for cancer: prevalence and independent risk factors*. Journal of the American College of Surgeons, 2007. **205**(6): p. 785-793.
250. Zheng, H.-L., et al., *Short- and Long-Term Outcomes in Malnourished Patients After Laparoscopic or Open Radical Gastrectomy*. World journal of surgery, 2018. **42**(1): p. 195-203.
251. Sugisawa, N., et al., *Intra-abdominal infectious complications following gastrectomy in patients with excessive visceral fat*. 2012. **15**(2): p. 206-12.
252. Watanabe, J., et al., *The impact of visceral obesity on surgical outcomes of laparoscopic surgery for colon cancer*. 2014. **29**(3): p. 343-51.
253. Okada, K., Nishigori, T., Obama, K., et al., *The Incidence of Postoperative Complications after Gastrectomy Increases in Proportion to the Amount of Preoperative Visceral Fat*. 2019. **2019**: p. 8404383.

254. Norero, E., Quezada, J.L., Cerda, J., et al., *Risk factors for severe postoperative complications after gastrectomy for gastric and esophagogastric junction cancers*. 2019. **32**(4): p. e1473.
255. Boccola, M.A., et al., *Risk factors and outcomes for anastomotic leakage in colorectal surgery: a single-institution analysis of 1576 patients*. World journal of surgery, 2011. **35**(1): p. 186-195.
256. Kang, J., et al., *Risk factor analysis of postoperative complications after robotic rectal cancer surgery*. World journal of surgery, 2011. **35**(11): p. 2555-2562.
257. Song, W., et al., *The delayed massive hemorrhage after gastrectomy in patients with gastric cancer: characteristics, management opinions and risk factors*. 2014. **40**(10): p. 1299-306.
258. Kim, D.J., J.H. Lee, and W. Kim, *Comparison of the major postoperative complications between laparoscopic distal and total gastrectomies for gastric cancer using Clavien-Dindo classification*. Surgical endoscopy, 2015. **29**(11): p. 3196-3204.
259. Jiang, N., et al., *Effect of complication grade on survival following curative gastrectomy for carcinoma*. World journal of gastroenterology, 2014. **20**(25): p. 8244-8252.
260. Xie, X.-S., et al., *A risk prediction system of postoperative hemorrhage following laparoscopy-assisted radical gastrectomy with D2 lymphadenectomy for primary gastric cancer*. 2017. **8**(46): p. 81511-81519.
261. Marinello, F.G., et al., *Anastomotic leakage after colon cancer resection: does the individual surgeon matter?* Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland, 2016. **18**(6): p. 562-569.
262. Kouyoumdjian, A., Trepanier, M., Al Shehhi, R., et al., *The Effect of Preoperative Anemia and Perioperative Transfusion on Surgical Outcomes After Gastrectomy for Gastric Cancer*. 2021. **259**: p. 523-531.
263. Xiao, C., et al., *Novel nomogram with microvascular density in the surgical margins can accurately predict the risk for anastomotic leakage after anterior resection for rectal cancer*. Journal of surgical oncology, 2019. **120**(8): p. 1412-1419.
264. Eto, K., et al., *Surgical Apgar Score Predicted Postoperative Morbidity After Esophagectomy for Esophageal Cancer*. 2016. **40**(5): p. 1145-51.
265. Ohkura, Y., et al., *Risk Factors for Postoperative Chylothorax After Radical Subtotal Esophagectomy*. Annals of surgical oncology, 2018. **25**(9): p. 2739-2746.
266. Xiao, C., Zhou, M., Yang, X., et al., *Novel nomogram with microvascular density in the surgical margins can accurately predict the risk for anastomotic leakage after anterior resection for rectal cancer*. 2019. **120**(8): p. 1412-1419.
267. Milgrom, S.A., et al., *Neoadjuvant radiation therapy prior to total mesorectal excision for rectal cancer is not associated with postoperative complications using current techniques*. Annals of surgical oncology, 2014. **21**(7): p. 2295-2302.
268. Lim, S.-B., et al., *Late anastomotic leakage after low anterior resection in rectal cancer patients: clinical characteristics and predisposing factors*. 2016. **18**(4): p. O135-40.
269. Koëter, M., et al., *Radiation dose does not influence anastomotic complications in patients with esophageal cancer treated with neoadjuvant chemoradiation and transhiatal esophagectomy*. 2015. **10**: p. 59.
270. Josse, J.M., et al., *The neutrophil-to-lymphocyte ratio predicts major perioperative complications in patients undergoing colorectal surgery*. Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland, 2016. **18**(7): p. O236-O242.
271. Achilli, P., et al., *Preoperative immunonutrition in frail patients with colorectal cancer: an intervention to improve postoperative outcomes*. Int J Colorectal Dis, 2020. **35**(1): p. 19-27.
272. Minnella, E.M., et al., *Effect of Exercise and Nutrition Prehabilitation on Functional Capacity in Esophagogastric Cancer Surgery: A Randomized Clinical Trial*. JAMA Surg, 2018. **153**(12): p. 1081-1089.
273. Pouwels, S., et al., *Preoperative exercise therapy for elective major abdominal surgery: a systematic review*. Int J Surg, 2014. **12**(2): p. 134-40.
274. Souwer, E.T.D., et al., *Comprehensive multidisciplinary care program for elderly colorectal cancer patients: "From prehabilitation to independence"*. Eur J Surg Oncol, 2018. **44**(12): p. 1894-1900.
275. Yamamoto, K., et al., *Effectiveness of a preoperative exercise and nutritional support program for elderly sarcopenic patients with gastric cancer*. Gastric Cancer, 2017. **20**(5): p. 913-918.
276. Inoue, J., et al., *Prevention of postoperative pulmonary complications through intensive preoperative respiratory rehabilitation in patients with esophageal cancer*. Dis Esophagus, 2013. **26**(1): p. 68-74.
277. Valkenet, K., et al., *Preoperative inspiratory muscle training to prevent postoperative pulmonary complications in patients undergoing esophageal resection (PREPARE study): study protocol for a randomized controlled trial*. Trials, 2014. **15**: p. 144.
278. Yamana, I., et al., *Randomized Controlled Study to Evaluate the Efficacy of a Preoperative Respiratory Rehabilitation Program to Prevent Postoperative Pulmonary Complications after Esophagectomy*. Dig Surg, 2015. **32**(5): p. 331-7.
279. Katsura, M., et al., *Preoperative inspiratory muscle training for postoperative pulmonary complications in adults undergoing cardiac and major abdominal surgery*. Cochrane Database Syst Rev, 2015(10): p. Cd010356.
280. Ligthart-Melis, G.C., et al., *Dietician-delivered intensive nutritional support is associated with a decrease in severe postoperative complications after surgery in patients with esophageal cancer*. Dis Esophagus, 2013. **26**(6): p. 587-93.
281. Williams, D.G.A., J. Molinger, and P.E. Wischmeyer, *The malnourished surgery patient: a silent epidemic in perioperative outcomes?* Curr Opin Anaesthesiol, 2019. **32**(3): p. 405-411.

282. Previtali, P., et al., *Malnutrition and Perioperative Nutritional Support in Retroperitoneal Sarcoma Patients: Results from a Prospective Study*. Ann Surg Oncol, 2019.
283. Drover, J.W., et al., *Perioperative use of arginine-supplemented diets: a systematic review of the evidence*. J Am Coll Surg, 2011. **212**(3): p. 385-99, 399.e1.
284. Phillips, S.M., *Nutritional supplements in support of resistance exercise to counter age-related sarcopenia*. Adv Nutr, 2015. **6**(4): p. 452-60.
285. Di Girolamo, F.G., et al., *Omega-3 fatty acids and protein metabolism: enhancement of anabolic interventions for sarcopenia*. Curr Opin Clin Nutr Metab Care, 2014. **17**(2): p. 145-50.
286. Moller, A.M., et al., *Effect of preoperative smoking intervention on postoperative complications: a randomised clinical trial*. Lancet, 2002. **359**(9301): p. 114-7.
287. Froessler, B., et al., *The Important Role for Intravenous Iron in Perioperative Patient Blood Management in Major Abdominal Surgery: A Randomized Controlled Trial*. Ann Surg, 2016. **264**(1): p. 41-6.
288. Shin, J., et al., *Effects of preoperative oral management by dentists on postoperative outcomes following esophagectomy: Multilevel propensity score matching and weighting analyses using the Japanese inpatient database*. Medicine (Baltimore), 2019. **98**(17): p. e15376.
289. Soutome, S., et al., *Preventive Effect on Post-Operative Pneumonia of Oral Health Care among Patients Who Undergo Esophageal Resection: A Multi-Center Retrospective Study*. Surg Infect (Larchmt), 2016. **17**(4): p. 479-84.