



Universiteit
Leiden
The Netherlands

Targets for improving patient outcomes after major gastrointestinal cancer surgery: the value of perioperative care

Kooten, R.T. van

Citation

Kooten, R. T. van. (2023, June 15). *Targets for improving patient outcomes after major gastrointestinal cancer surgery: the value of perioperative care*. Retrieved from <https://hdl.handle.net/1887/3620458>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3620458>

Note: To cite this publication please use the final published version (if applicable).



Chapter 4

Conventional Regression Analysis and Machine Learning in Prediction of Anastomotic Leakage and Pulmonary Complications after Esophagogastric Cancer Surgery

Robert T. van Kooten MD¹, Renu R. Bahadoer MD¹, Boudewijn ter Burkes de Vries MSc², Michel W.J.M. Wouters, MD PhD^{2,3}, Rob A.E.M. Tollenaar MD PhD¹, Henk H. Hartgrink MD PhD¹, Hein Putter PhD², Johan L. Dikken MD PhD¹

¹ Department of Surgery, Leiden University Medical Center, Leiden, The Netherlands

² Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

³ Department of Surgical Oncology, Netherlands Cancer Institute-Antoni van Leeuwenhoek, Amsterdam, The Netherlands

List of abbreviations

ARDS; Acute Respiratory Distress Syndrome, ASA; American Society of Anesthesiologists, AUC; Area Under the Curve, CI; Confidence Interval, DUCA; Dutch Upper Gastrointestinal Cancer Audit, ERAS; Enhanced Recovery After Surgery, GLM; Generalized Linear Model, KNN; k-Nearest Neighbors, OR; Odds Ratio, ROC; Receiver Operating Characteristics, SVM; Support Vector Machine.

Abstract

Background and Objectives: With the current advanced data-driven approach to health care, machine learning is gaining more interest. The current study investigates the added value of machine learning to linear regression in predicting anastomotic leakage and pulmonary complications after upper gastrointestinal cancer surgery.

Methods: All patients in the Dutch Upper Gastrointestinal Cancer Audit (DUCA) undergoing curatively intended esophageal- or gastric cancer surgery in 2011-2017 were included. Anastomotic leakage was defined as any clinically or radiologically proven anastomotic leakage. Pulmonary complications entailed: pneumonia, pleural effusion, respiratory failure, pneumothorax and/or acute respiratory distress syndrome. Different machine learning models were tested. Nomograms were constructed using logistic regression.

Results: Between 2011-2017, 4,228 patients underwent surgical resection for esophageal cancer of which 18% developed anastomotic leakage and 30% a pulmonary complication. Of the 2,199 patients with surgical resection for gastric cancer, 7% developed anastomotic leakage and 15% a pulmonary complication. In all cases, linear regression had the highest predictive value with area under the curves (AUCs) varying between 61.9-68.0, but the difference with machine learning models did not reach statistical significance.

Conclusion: Machine learning models can predict postoperative complications in upper gastrointestinal cancer surgery, but they do not outperform the current gold standard, linear regression.

Introduction

The incidence of esophageal cancer in the western world has increased over the past decades and is currently the seventh most common malignancy worldwide and accounts for 5% of the cancer-related mortality in 2018. Although the incidence of gastric cancer decreased over the last years, it is still the fifth most common malignancy worldwide and was responsible for 8% of the cancer-related mortality in 2018. [1] Curative treatment of these upper gastrointestinal cancers consists in most cases of (neo)adjuvant therapy and surgical resection. These resections are complex procedures. Present-day, the 5-year survival rates of resectable esophageal- and gastric carcinoma lie around 28-42%. [2]. Although in centers of excellence, postoperative mortality is around 2%, the overall-complication rate of around 60-65% after esophagectomy is high compared to most procedures for gastrointestinal malignancies. [3] Of all severe postoperative complications, anastomotic leakage and pulmonary complications are the most common. [2-5] The incidence of major complications (Clavien-Dindo \geq IIIa) is 20-31%, with a failure-to-rescue rate of 13-25%. [6, 7] Postoperative complications are associated with higher tumor recurrence and lower overall (cancer-related) survival. [8] Reduction of postoperative complications will enhance recovery, lead to fewer readmissions and may increase long-term quality of life.

With the present increase of data-driven approaches in healthcare, preoperative risk factors can be appraised by analyzing large datasets. Machine learning holds the potential to unravel subtle associations that are not—or cannot—be identified using conventional regression analyses. In the current literature, no consensus exists on the added value of machine learning in predicting postoperative outcomes. [9, 10]

The aim of this study is twofold. First, to investigate the added value of machine learning methods in predicting postoperative outcomes after esophageal- and gastric carcinoma surgery and compare it to conventional regression analyses. Second, to use the best performing method to develop a predictive model for anastomotic leakage and cardiopulmonary complications after esophagectomy and gastrectomy.

Methods

Data source and study population

Data were retrieved from the Dutch Upper GI Cancer Audit (DUCA). A prospective nationwide audit, initiated in 2011, containing all patients undergoing surgery with the intention of resection for esophageal- or gastric cancer in the Netherlands. [11] Participation in the DUCA has been incorporated as a mandatory quality standard,

leading to data completeness of 99.8% and accuracy of 94%-100%. Validation of completeness and accuracy of this data registration has been performed by external data verification. [3] All Dutch hospitals register detailed patient, tumor, and treatment characteristics, pathology, 30-day morbidity, and 30-day/in-hospital mortality. [12]

Patient selection

Patients that underwent elective surgery with curative intent for primary esophageal- or gastric cancer were selected. Only patients with histologically proven adenocarcinoma or squamous cell carcinoma, a known surgery date between 2011 and 2017 and a recorded surgical technique were included. Patients with surgery with palliative or prophylactic intent and patients with non-epithelial tumors were excluded. Additionally, patients with missing essential values (age, sex, length, weight, surgical approach, American Society of Anesthesiologists (ASA)-score, preoperative therapy and TMN-stage) were excluded.

Definitions of complications

The studied postoperative outcomes were anastomotic leakage and pulmonary complications in patients with esophageal carcinoma, and anastomotic leakage in patients with gastric carcinoma. Anastomotic leakage was defined as any clinically or radiologically proven anastomotic leakage. Pulmonary complications entailed: pneumonia, pleural effusion, respiratory failure, pneumothorax and/or acute respiratory distress syndrome (ARDS).

Statistical analysis For each outcome, the dataset was randomly divided in training (75%) and testing (25%) data. All models used variables documented in the Dutch Upper GI Cancer Audit (DUCA), which covers patient characteristics, comorbidity, treatment characteristics and outcome [13]; a total of 28 prognostic variables were included. The following machine learning models, which are frequently described in literature, were used: k-Nearest Neighbors (KNN), support vector machine (SVM), Neural Networks, Random Forest, AdaBoost and SuperLearner. [14-16] These models were compared with linear regression, for which a generalized linear model (GLM) was used. Background information on the models used can be found in table 1. Afterwards, nomograms were constructed using a regression model fit. The predictive strength of the models was measured by the Area Under the Receiver Operating Characteristics (ROC) Curve (AUC). Odds ratios (OR) with 95% confidence intervals and P-values were reported for each variable to assess the impact on the risk of all patient characteristics. All analyses were done using R version 3.6.1 in RStudio. The Caret packages were used for pipelining and data splitting. ROC curves and AUC scores were calculated using the pROC package, plots were made using the ggplot2 package. The rms package was used to make the nomograms.

Table 1 - Explanation of the models used

| |
|---|
| Logistic Regression |
| Describes the relationship between a discrete binary outcome and one or several predictor variables. The outcome is expressed as the log odds of one class over the other. This can be transformed to odds or probabilities. |
| Lasso Regression |
| The difference between the logistic regression model and the lasso model is that the lasso model can exclude coefficients that have little weight in the solution. This may increase interpretability. |
| k-Nearest Neighbour (kNN) |
| Predicts new instances of a class by looking at k other instances in the neighborhood. The predictor variables are transformed by centering and scaling to improve numerical stability. For each outcome a separate kNN model is fit. |
| Neural Networks (NN) |
| The inspiration for NN comes from the architecture of the human brain. The idea is that artificial neurons send the next neuron a signal based on the input they are receiving. A network of artificial neurons is called a neural network. A NN consists of layers. The first being an input layer (the predictor variables), followed by one or more hidden layers (the artificial neurons) and finally resulting in an output layer (the prediction). For each outcome in the data a NN is fit. |
| Support Vector Machine (SVM) |
| A classification (and regression) algorithm that can classify non linearly separable data by constructing a hyperplane (or a set of hyperplanes) in high dimensional space. A SVM tries to find a hyperplane that best separates two groups. This is the hyperplane whose distance to the nearest element of each class is the largest. For data that is not linearly separable the kernel trick is used. This is a method of adding dimensions to the data while at the same time keep the calculations feasible. For each outcome a <i>polynomial</i> (kernel) SVM and a <i>radial</i> (kernel) SVM is fit. |
| Random Forest |
| A random forest is an ensemble of decision trees. The model is trained with a technique called bootstrap aggregation (bagging). Bagging reduces variance and avoids overfitting in ensemble methods. With this technique many bootstrap samples are taken and a decision tree is trained on each sample. The outcome of all trees together is averaged, which leads to the final outcome. For each outcome a random forest is trained. |
| Adaboost |
| Boosting is similar to a random forest. The main differences are that the trees are now built sequentially and the results are averaged along the way. Boosting is an ensemble method that combines weak classifiers to output a single strong predicted response. The technique is considered to be an improvement over random forests in some occasions. For each outcome an Adaboost.m1 model is trained. |
| Super Learner |
| The super learner finds an optimal weighted combination of candidate learners. The candidate learners can be any prediction algorithm. The super learner itself is a prediction algorithm as well. The performance of the candidate learners is assessed by cross-validation. For each outcome a super learner model is trained. The candidate learners consist of all models mentioned above. With the exception of Adaboost.m1, which is replaced by XGBoost (an alternative boosting algorithm). |

Results

Study population

Between 2011 and 2017, 8,173 patients were included in the DUCA. Of these, 6,427 were included in the final dataset (figure 1). Of the excluded patients, 403 were a result of missing essential values; the outcomes of these patients were not significantly different from those included. In total, 4,228 patients underwent a surgical resection for esophageal carcinoma, of which 2,540 had a postoperative complication (60%). Of the 2199 patients with a resection for gastric carcinoma, 883 patients had a postoperative complication (40%). Patient characteristics are described in table 2, and figure 2 presents an overview of the type of postoperative complications.

Esophageal carcinoma

Anastomotic leakage occurred in 31% (799 of 4,228) patients following esophagectomy and pulmonary complications in 54% (1380 of 4228), figure 2. From all prediction models, the generalized linear model had the highest AUC, both for anastomotic leakage (61.9; 95%CI 57.9-65.9) and for pulmonary complications (64.4; 95%CI 60.9-67.9), figures 3a and 3b. Closely followed by the machine learning models: Neural Networks (AUC 61.7; 95%CI 57.7-65.6), LASSO (AUC 61.7; 95%CI 57.7-65.7) and SuperLearner (AUC 61.7; 95%CI 57.7-65.8) for anastomotic leakage. And the machine learning model LASSO (AUC 64.3; 95%CI 60.9-67.8) for pulmonary complications. For preoperative prediction, nomograms, based on a generalized linear model, have been constructed for anastomotic leakage (figure 5a) and pulmonary complications (figure 5b). For anastomotic leakage: steroid use, advanced tumor stage, distant metastasis, surgical approach and preoperative weight loss factors with the most prognostic value. For pulmonary complications, these are weight loss, ASA III/IV, advanced tumor stage, type of resection and location of anastomosis.

Gastric carcinoma

After gastrectomy, anastomotic leakage was reported in 18% (156 of 2,199) patients. Generalized linear model had the highest AUC (68.0; 95%CI 60.2-75.8) (Figure 4), followed by the machine learning model Neural Networks (AUC 67.9; 95%CI 60.4-75.5). A nomogram for the preoperative prediction of anastomotic leakage after gastric resection is displayed in figure 5c. Tumor histology and lymph node involvement are factors with the most prognostic value.




Figure 1 - Patient selection




Table 2 - Patient characteristics

| | Esophageal cancer resection | | Gastric cancer resection | |
|---------------------------------|-----------------------------|---------|--------------------------|---------|
| | (n = 4,228) | | (n = 2,199) | |
| Age, median (IQR) | 66 | (59-71) | 70 | (62-77) |
| Gender | | | | |
| Male | 3,272 | (77%) | 1,366 | (62%) |
| Female | 956 | (23%) | 833 | (38%) |
| BMI | | | | |
| < 20 | 276 | (7%) | 170 | (8%) |
| 20-24 | 1,614 | (38%) | 949 | (43%) |
| 25-29 | 1,663 | (39%) | 789 | (36%) |
| ≥ 30 | 675 | (16%) | 291 | (13%) |
| Comorbidity | | | | |
| None | 998 | (24%) | 433 | (20%) |
| Yes | 3,229 | (76%) | 1,764 | (80%) |
| Of which | | | | |
| cardiac | 978 | (30%) | 676 | (31%) |
| diabetes | 639 | (20%) | 375 | (17%) |
| pulmonary | 769 | (18%) | 352 | (16%) |
| thrombotic | 173 | (5%) | 154 | (7%) |
| Unknown | 1 | (<1%) | 2 | (<1%) |
| Preoperative weight loss | | | | |
| None | 1,138 | (27%) | 639 | (29%) |
| 1-5 kg | 1,184 | (28%) | 543 | (25%) |
| 6-10 kg | 891 | (21%) | 473 | (22%) |
| 11-15 kg | 279 | (7%) | 146 | (7%) |
| 16-20 kg | 106 | (3%) | 55 | (3%) |
| 21-35 kg | 56 | (1%) | 24 | (1%) |
| Unknown | 574 | (14%) | 319 | (15%) |
| Previous surgery* | | | | |
| No | 2,943 | (70%) | 1,313 | (60%) |
| Yes | 1,276 | (30%) | 882 | (40%) |
| Unknown | 9 | | 4 | |
| Histology | | | | |
| Adenocarcinoma | 3,383 | (80%) | 2195 | (>99%) |
| Squamous cell carcinoma | 845 | (20%) | 4 | (<1%) |
| Type of surgery | | | | |
| Transhiatal | 1,395 | (33%) | - | |
| Transthoracic | 2,833 | (67%) | - | |
| McKeown | 1,353 | (48%) | | |

Table 2 - Continued

| | Esophageal cancer resection | | Gastric cancer resection | |
|------------------------------|-----------------------------|-------|--------------------------|-------|
| | (n = 4,228) | | (n = 2,199) | |
| Total gastrectomy | - | | 924 | (42%) |
| cTNM-7 stage | | | | |
| Stage 0 | 6 | (<1%) | 16 | (1%) |
| Stage I | 566 | (13%) | 465 | (21%) |
| Stage II | 1,116 | (26%) | 842 | (38%) |
| Stage III | 2,155 | (51%) | 185 | (8%) |
| Stage IV | 40 | (1%) | 39 | (2%) |
| Stage X | 345 | (8%) | 652 | (30%) |
| Neoadjuvant treatment | | | | |
| None | 314 | (7%) | 848 | (39%) |
| Chemotherapy | 286 | (7%) | 1,316 | (60%) |
| Chemoradiotherapy | 3,628 | (86%) | 35 | (2%) |
| ASA-score | | | | |
| I | 712 | (17%) | 305 | (14%) |
| II | 2,592 | (61%) | 1,237 | (56%) |
| III | 908 | (22%) | 639 | (29%) |
| IV | 16 | (<1%) | 18 | (1%) |
| Steroid use | | | | |
| No | 4,093 | (97%) | 2,118 | (96%) |
| Yes | 107 | (3%) | 46 | (2%) |
| Unknown | 28 | (1%) | 35 | (2%) |


*Thoracic- and/or abdominal surgery



| | AUC% (95% CI) |
|--------------------------|------------------|
| Generalized Linear Model | 61.9 (57.9-65.9) |
| Lasso | 61.7 (57.7-65.7) |
| kNN | 56.9 (52.6-61.2) |
| Neural Networks | 61.7 (57.7-65.6) |
| SvmPoly | 59.3 (55.2-63.4) |
| SvmRadial | 54.0 (49.6-58.4) |
| Random Forest | 59.0 (54.7-63.2) |
| Adaboost | 61.3 (57.4-65.2) |
| SuperLearner | 61.7 (57.7-65.8) |


Figure 3a - Anastomotic leakage after esophageal cancer resection

Figure 2 - Type of complications after surgery after (A) esophagectomy and (B) gastrectomy



| | AUC% (95% CI) |
|--------------------------|------------------|
| Generalized Linear Model | 64.4 (60.9-67.8) |
| Lasso | 64.3 (60.9-67.8) |
| kNN | 61.1 (57.6-64.6) |
| Neural Net | 63.6 (60.2-67.1) |
| SvmPoly | 61.1 (57.5-64.6) |
| SvmRadial | 58.4 (54.7-62.1) |
| Random Forest | 61.1 (57.5-64.7) |
| Adaboost M1 | 63.1 (59.7-66.6) |
| SuperLearner | 63.9 (60.5-67.4) |

Figure 3b - Pulmonary complications after esophageal cancer resection



| | AUC% (95% CI) |
|---------------------|------------------|
| Logistic Regression | 68.0 (60.2-75.8) |
| Lasso | 67.7 (59.4-76.0) |
| kNN | 59.4 (50.8-68.1) |
| Neural Net | 67.9 (60.4-75.5) |
| SvmPoly | 63.0 (55.3-70.8) |
| SvmRadial | 56.4 (47.7-65.2) |
| Random Forest | 62.0 (53.7-70.3) |
| Adaboost M1 | 64.9 (56.7-73.1) |
| SuperLearner | 67.8 (59.5-76.1) |

Figure 4 - Anastomotic leakage after gastric cancer resection




Figure 5a - Nomogram for the prediction of anastomotic leakage after esophageal cancer resection. AUC: 61.9%




Figure 5b - Nomogram for the prediction of pulmonary complications after esophageal cancer resection. AUC: 64.4%






Figure 5c- Nomogram for the prediction of anastomotic leakage after gastric cancer resection. AUC: 68.1%

Discussion

This study presents the development of various (machine learning) models for the prediction of anastomotic leakage and pulmonary complications in a population-based cohort of 4,228 patients that underwent esophagectomy and 2,199 patients that underwent gastrectomy in the Netherlands. Linear regression had the highest accuracy of all models for the prediction of anastomotic leakage after esophagectomy and gastrectomy, as well as for pulmonary complications after esophagectomy. Of the machine learning models, the Neural Networks had the highest accuracy for predicting anastomotic leakage after esophageal cancer surgery and after gastric cancer surgery. LASSO had the highest accuracy of the machine learning models for the prediction of pulmonary complications following esophageal cancer surgery. Furthermore, highest accuracy of all studied models was 68.0%, suggesting that none of the models had superior predictive ability for postoperative complications in this patient cohort.

It is thought that for the development of machine models, a large population is necessary to adequately train the models. For example, the study of Nudel et al. included over 436,000 patients and 40 different variables. In their study, artificial neural networks and gradient boosting machines outperformed the traditional linear regression in predicting anastomotic leakage after weight loss surgery. [10] However, a study that included merely 321 patients successfully designed a support vector classification model to predict postoperative complications in patients undergoing gastrectomy, using 23 clinical features. Their model had an accuracy of 78% in external validation. Like in the current study, age and tumor stage were the most predictive for the development of complications. [17]. With a broad array of machine learning models available, it is difficult to decide which model to use for each particular outcome. The systematic review of Elfanagely et al. reviewed 45 papers published between 2015 and 2020. [18] They concluded that machine learning in surgical research is still in its infancy, but these early-published papers show potential. However, they found great heterogeneity exists between the different studies; various models are being used, and different variables and outcomes are being investigated.[18] They have also shown large variation in sample sizes, ranging from 71 to 130,945, implying that sample size is not the only factor for successful machine learning models. However, it might be possible that a certain publication and confirmation bias is present in the current literature, which could be deceiving.

In line with current literature, our study has demonstrated that ASA-score \geq III is associated with anastomotic leakage and pulmonary complications. [17, 19] Both patients with advanced age and high ASA-classification are thought to have lower healing capacity



causing higher susceptibility to postoperative complications. Patients with a more advanced tumor stage may require more extensive resections and technically more demanding surgery to reach an R0 resection. This may lead to more intraoperative organ damage and subsequent postoperative complications. [20] Furthermore, lymph node involvement and, therefore, extensive lymph node dissections and possibly additional splenic resection are high-risk procedures. [21] As shown in this study, chronic use of steroids preoperatively is associated with postoperative complications, which is thought to be due to a reduced healing capacity. [22, 23]

In daily practice, it is difficult to estimate the surgical risk and make treatment decisions based on individual predictive factors. Therefore incorporating multiple factors into prediction models can be used to combine information in a simple and more useful manner. [24] However, the use of these models is often limited since they are often created in a selected patient population or specialized centers, making generalization hard; hence this study used a nationwide population-based cohort. [25, 26] Furthermore, clinical judgment and expertise are still needed for correct interpretation and usage of clinical prediction models. With the current more data-driven approach to health care and the availability of nationwide clinical audits, big data becomes available, eliminating this limitation of generalization of models. In addition, with big data, the interest in machine learning for prediction models has increased. In our study, linear regression was superior to the machine learning models. Another study, which used a more extensive amount of variables, did show favor for machine learning models. [10] However, one could question the use in daily clinical practice when using more extensive models, which subsequently leads to more administrative burden to include all variables unless being automated. Furthermore, in some machine learning models (e.g., neural networks) individual prognostic factors are not known, in contrast to linear regression. As demonstrated in this study, linear regression can be used to construct nomograms, which are easy to use in daily practice and might expose improvable prognostic factors (e.g., weight loss, steroid use) for postoperative complications. Subsequently, nomograms can easily be formed into web-based models of mobile phone applications, which might increase usability in daily practice.

As part of the currently ongoing implementation and standardization of perioperative care into enhanced recovery after surgery (ERAS) protocols, preoperative optimization of patients has gained interest. [27] Research towards improving perioperative care for upper gastrointestinal surgery also focuses on identifying preoperative high-risk patients and developing prehabilitation programs for these patients. Upper gastrointestinal cancer patients are at high risk for malnutrition due to the anatomical localization of the tumor. Therefore, nutritional interventions are important in preoperative prehabilitation. [28] Prehabilitation programs for patients with esophageal

carcinoma have been shown to improve objective measures of physical fitness but are less clear on postoperative outcomes. [29] However, good physical fitness and nutritional status are widely recognized as a protective factor against postoperative complications. It has shown to lead to sooner return to bowel function, oral feeding and restored metabolic equilibrium and is therefore currently being standardized and implemented into ERAS protocols. [27, 30] This may indicate that prehabilitation programs might have to be more specific towards certain risk factors.

Although our study provides insight on a different aspect of the clinical applicability of machine learning models, it has some limitations. The use of Dutch national audit data, DUCA, might lead to less generalizability to other countries. However, in The Netherlands, participation in the DUCA has been incorporated as a mandatory quality standard, leading to an exceptionally complete and reliable database. Voluntary participation of some other audits and registries could give a distorted view if their participation did not concern all patients. External validation of the models did not occur in this study. However, the accuracy of the models was tested using a random internal sample of 25% of patients. Another limitation of the study is that the experience and expertise of the individual centers and/or surgeons could not be included. Hospital volume is thought to be a predictor of mortality after high-risk surgery. [31] Patients treated in high-volume hospitals benefit from more experience and more advanced expertise. However, according to the DUCA research regulations, no data is provided that can be used to derive individual hospitals. If these restrictions are lifted in the future, this variable could be implemented in the model to improve accuracy. Additionally, accuracy may be improved by adding more variables, which are currently not in the the DUCA registry, such as preoperative laboratory results (e.g., C-reactive protein (CRP), albumin) [32] and other predictors such as smoking and alcohol usage [33, 34] were not included in our models, whereas these variables may serve as strong preoperative predictors. The use of intraoperative variables such as intraoperative hypotension or blood transfusion may improve predictive accuracy of the model. However, these factors cannot be used during patient selection for prehabilitation programs or for surgery. [35-37] Additionally, the anastomotic leakage rate of 18% following esophagectomy in the Netherlands is relatively high compared to other countries, an explanation is the learning curve for new techniques (e.g., minimally invasive) in recent years. Around 2010 minimally invasive surgery was introduced and many surgeons changed from a McKeown to an Ivor Lewis technique. [38] Furthermore, both clinically and radiologically proven anastomotic leakages were included. Finally, all patients included in this study were selected to be fit for surgery by expert opinion preoperatively, leading to allocation bias. However, occurrence of this type of bias is unavoidable in this type of study.

Conclusion

This study demonstrates that the studied machine-learning models are able to predict postoperative complications in upper gastrointestinal cancer surgery, but they are not superior to the current gold standard, logistic regression. However, the accuracy of all studied models was relatively low. Furthermore, the use of prediction models does serve a purpose for preoperative risk estimation and treatment decisions, but clinical expertise is still needed. Additionally, identifying predictive individual factors within prediction models (e.g., malnutrition) may improve perioperative care and might lead to improved preoperative physical fitness of patients, which can improve ERAS protocols and therewith surgical outcomes.

References

1. Bray, F., et al., *Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries*. CA Cancer J Clin, 2018. **68**(6): p. 394-424.
2. Ruol, A., et al., *Trends in management and prognosis for esophageal cancer surgery: twenty-five years of experience at a single institution*. Arch Surg, 2009. **144**(3): p. 247-54; discussion 254.
3. van der Werf, L.R., et al., *Reporting National Outcomes After Esophagectomy and Gastrectomy According to the Esophageal Complications Consensus Group (ECCG)*. Ann Surg, 2020. **271**(6): p. 1095-1101.
4. Lagarde, S.M., et al., *Preoperative prediction of the occurrence and severity of complications after esophagectomy for cancer with use of a nomogram*. Ann Thorac Surg, 2008. **85**(6): p. 1938-45.
5. Sunpaweravong, S., et al., *Prediction of major postoperative complications and survival for locally advanced esophageal carcinoma patients*. Asian J Surg, 2012. **35**(3): p. 104-9.
6. Busweiler, L.A., et al., *Failure-to-rescue in patients undergoing surgery for esophageal or gastric cancer*. Eur J Surg Oncol, 2017. **43**(10): p. 1962-1969.
7. Abdelsattar, Z.M., et al., *Understanding Failure to Rescue After Esophagectomy in the United States*. Ann Thorac Surg, 2020. **109**(3): p. 865-871.
8. Aurello, P., et al., *Recurrence Following Anastomotic Leakage After Surgery for Carcinoma of the Distal Esophagus and Gastroesophageal Junction: A Systematic Review*. Anticancer Res, 2019. **39**(4): p. 1651-1660.
9. Christodoulou, E., et al., *A systematic review shows no performance benefit of machine learning over logistic regression for clinical prediction models*. J Clin Epidemiol, 2019. **110**: p. 12-22.
10. Nudel, J., et al., *Development and validation of machine learning models to predict gastrointestinal leak and venous thromboembolism after weight loss surgery: an analysis of the MBSAQIP database*. Surg Endosc, 2021. **35**(1): p. 182-191.
11. Busweiler, L.A., et al., *Early outcomes from the Dutch Upper Gastrointestinal Cancer Audit*. Br J Surg, 2016. **103**(13): p. 1855-1863.
12. van der Werf, L.R., et al., *A National Cohort Study Evaluating the Association Between Short-term Outcomes and Long-term Survival After Esophageal and Gastric Cancer Surgery*. Ann Surg, 2019. **270**(5): p. 868-876.
13. Wouters, M.W., M.L. Jansen-Landheer, and C.J. van de Velde, *The Quality of Cancer Care initiative in the Netherlands*. Eur J Surg Oncol, 2010. **36 Suppl 1**: p. S3-S13.
14. Christie, S.A., et al., *Machine learning without borders? An adaptable tool to optimize mortality prediction in diverse clinical settings*. J Trauma Acute Care Surg, 2018. **85**(5): p. 921-927.

15. Kim, J.S., et al., *Examining the Ability of Artificial Neural Networks Machine Learning Models to Accurately Predict Complications Following Posterior Lumbar Spine Fusion*. Spine (Phila Pa 1976), 2018. **43**(12): p. 853-860.
16. Hofer, I.S., et al., *Development and validation of a deep neural network model to predict postoperative mortality, acute kidney injury, and reintubation using a single feature set*. NPJ Digit Med, 2020. **3**: p. 58.
17. Lu, S., et al., *Machine-learning-assisted prediction of surgical outcomes in patients undergoing gastrectomy*. Chin J Cancer Res, 2019. **31**(5): p. 797-805.
18. Elfanagely, O., et al., *Machine Learning and Surgical Outcomes Prediction: A Systematic Review*. J Surg Res, 2021. **264**: p. 346-361.
19. van Kooten, R.T., et al., *Patient-Related Prognostic Factors for Anastomotic Leakage, Major Complications, and Short-Term Mortality Following Esophagectomy for Cancer: A Systematic Review and Meta-Analyses*. Ann Surg Oncol, 2021.
20. Kulig, P., et al., *Differences in prognosis of Siewert II and III oesophagogastric junction cancers are determined by the baseline tumour staging but not its anatomical location*. Eur J Surg Oncol, 2016. **42**(8): p. 1215-21.
21. Lu, J., et al., *Major perioperative complications in laparoscopic spleen-preserving total gastrectomy for gastric cancer: perspectives from a high-volume center*. Surg Endosc, 2016. **30**(3): p. 1034-42.
22. van Rossum, P.S.N., et al., *Calcification of arteries supplying the gastric tube: a new risk factor for anastomotic leakage after esophageal surgery*. Radiology, 2015. **274**(1): p. 124-32.
23. Anstead, G.M., *Steroids, retinoids, and wound healing*. Adv Wound Care, 1998. **11**(6): p. 277-85.
24. Steyerberg, E.W., et al., *Surgical mortality in patients with esophageal cancer: development and validation of a simple risk score*. J Clin Oncol, 2006. **24**(26): p. 4277-84.
25. Bartels, H., H.J. Stein, and J.R. Siewert, *Preoperative risk analysis and postoperative mortality of oesophagectomy for resectable oesophageal cancer*. Br J Surg, 1998. **85**(6): p. 840-4.
26. Klose, J., et al., *A Nomogram to Predict Anastomotic Leakage in Open Rectal Surgery- Hope or Hype?* J Gastrointest Surg, 2018. **22**(9): p. 1619-1630.
27. Low, D.E., et al., *Guidelines for Perioperative Care in Esophagectomy: Enhanced Recovery After Surgery (ERAS((R))) Society Recommendations*. World J Surg, 2019. **43**(2): p. 299-330.
28. Steenhagen, E., et al., *Nutrition in peri-operative esophageal cancer management*. Expert Rev Gastroenterol Hepatol, 2017. **11**(7): p. 663-672.
29. Bolger, J.C., et al., *Perioperative prehabilitation and rehabilitation in esophagogastric malignancies: a systematic review*. Dis Esophagus, 2019. **32**(9).
30. Gianotti, L., et al., *Enhanced recovery programs in gastrointestinal surgery: Actions to promote optimal perioperative nutritional and metabolic care*. Clin Nutr, 2020. **39**(7): p. 2014-2024.
31. Ra, J., et al., *Postoperative mortality after esophagectomy for cancer: development of a preoperative risk prediction model*. Ann Surg Oncol, 2008. **15**(6): p. 1577-84.
32. You, X., et al., *Preoperative albumin-to-fibrinogen ratio predicts severe postoperative complications in elderly gastric cancer subjects after radical laparoscopic gastrectomy*. BMC Cancer, 2019. **19**(1): p. 931.
33. Oppedal, K., et al., *Preoperative alcohol cessation prior to elective surgery*. Cochrane Database Syst Rev, 2012(7): p. CD008343.
34. Quan, H., et al., *The effect of preoperative smoking cessation and smoking dose on postoperative complications following radical gastrectomy for gastric cancer: a retrospective study of 2469 patients*. World J Surg Oncol, 2019. **17**(1): p. 61.
35. Bernard, A.C., et al., *Intraoperative transfusion of 1 U to 2 U packed red blood cells is associated with increased 30-day mortality, surgical-site infection, pneumonia, and sepsis in general surgery patients*. J Am Coll Surg, 2009. **208**(5): p. 931-7, 937.e1-2; discussion 938-9.
36. Fumagalli, U., et al., *Intra-operative hypotensive episodes may be associated with post-operative esophageal anastomotic leak*. Updates Surg, 2016. **68**(2): p. 185-90.
37. Kim, S.H., et al., *Risk Factors for Anastomotic Leakage: A Retrospective Cohort Study in a Single Gastric Surgical Unit*. J Gastric Cancer, 2015. **15**(3): p. 167-75.
38. Voeten, D.M., et al., *Outcomes of Esophagogastric Cancer Surgery During Eight Years of Surgical Auditing by the Dutch Upper Gastrointestinal Cancer Audit (DUCA)*. Ann Surg, 2021. **274**(5): p. 866-873.