

## Promoting early recognition of persistent somatic symptoms in primary care

Kitselaar, W.M.

### Citation

Kitselaar, W. M. (2023, June 27). *Promoting early recognition of persistent somatic symptoms in primary care*. Retrieved from https://hdl.handle.net/1887/3628068

Version: Publisher's Version

Licence agreement concerning inclusion of doctoral

License: thesis in the Institutional Repository of the University

of Leiden

Downloaded from: <a href="https://hdl.handle.net/1887/3628068">https://hdl.handle.net/1887/3628068</a>

**Note:** To cite this publication please use the final published version (if applicable).

## Chapter 3

The General Practitioners' Perspective Regarding Registration of Persistent Somatic Symptoms in Primary Care: a survey

Willeke M. Kitselaar, Rosalie van der Vaart, Madelon van Tilborg-den Boeft, Hedwig M.M. Vos, Mattijs E. Numans, Andrea W.M. Evers

Based on: Kitselaar WM, van der Vaart R, van Tilborg-den Boeft M, Vos HMM, Numans ME, and Evers AWM. The general practitioners perspective regarding registration of persistent somatic symptoms in primary care: a survey. BMC Fam Pract 22, 182 (2021).

DOI: 10.1186/s12875-021-01525-6

### Abstract

**Background:** Persistent somatic symptoms (PSS) are common in primary care and often accompanied by an increasing disease burden for both the patient and healthcare. In medical practice, PSS is historically considered a diagnosis by exclusion or primarily seen as psychological. Besides, registration of PSS in electronic health records (EHR) is ambiguous and possibly does not reflect classification adequately. The present study explores how general practitioners (GPs) currently register PSS, and their view regarding the need for improvements in classification, registration, and consultations.

**Method:** Dutch GPs were invited by email to participate in a national cross-sectional online survey. The survey addressed ICPC-codes used by GPs to register PSS, PSS-related terminology added to free text areas, usage of PSS-related syndrome codes, and GPs' need for improvement of PSS classification, registration and care.

**Results:** GPs (n = 259) were most likely to use codes specific to the symptom presented (89.3%). PSS-related terminology in free-text areas was used sparsely. PSS-related syndrome codes were reportedly used by 91.5% of GPs, but this was primarily the case for the code for irritable bowel syndrome. The ambiguous registration of PSS is reported as problematic by 47.9% of GPs. Over 56.7% of GPs reported needing additional training, tools or other support for PSS classification and consultation. GPs also reported needing other referral options and better guidelines.

**Conclusions:** Registration of PSS in primary care is currently ambiguous. Approximately half of GPs felt a need for more options for registration of PSS and reported a need for further support. In order to improve classification, registration and care for patients with PSS, there is a need for a more appropriate coding scheme and additional training.

### Introduction

Up to 50% of primary care visits in Western societies are related to symptoms that cannot be fully explained by well-known biomedical pathology. 1-4 While most of these symptoms are self-limiting, 2.5-10% of cases persist without clear medical explanation. 5-9 These persistent somatic symptoms (PSS) are accompanied by increasing disease burden for both the patient and the healthcare system. 10 Differentiation from well-known chronic medical conditions and classification of symptoms as PSS is challenging. 11 Challenges arise from similarities between symptoms of PSS and other conditions, possible co-existence with a well-documented medical disorder, the heterogeneity of symptoms, lack of universal guidelines and the lack of biomarkers. 3, 12-14 Delayed identification of PSS impedes early management of symptoms, which in turn can result in inappropriate healthcare utilization and high costs. 15-17 Additionally, it may hinder reusability of electronic health records (EHR) for research, quality monitoring and proactive population health management. 17-20

Across medical and psychological specialties, a variety of terminology and aetiology is reflected in different concepts of PSS. While some countries have specific guidelines for PSS, widely accepted guidelines for classifying (and treating) PSS are missing.<sup>3</sup> PSS is currently diagnosed as either a somatic disease or a mental disorder, since diagnostic classifications are inherently dualistic in nature. <sup>21</sup> In the medical field, patients may be classified under umbrella terms such as 'medically unexplained physical symptoms' (MUPS), 'functional somatic symptoms', and 'somatically fixed', 9, 22 which indicate a negative symptomology – i.e. a lack of medical pathology.<sup>23</sup> PSS may also be classified as syndromes such as irritable bowel syndrome (IBS), chronic fatigue syndrome (CFS) or fibromyalgia (FM). Ongoing debate about terminology has redirected the most recent versions of the diagnostic and statistical manual of mental disorders (DSM-5)<sup>24</sup> and the international classification of disease (ICD-11)<sup>25</sup> towards no longer requiring the explicit exclusion of any underlying medical condition (this only applies to PSS-related classifications in the mental health chapter). Both focus on positive symptomology, such as maladaptive cognitions, emotions and/or behaviours related to the somatic symptoms, 3, 13 as described in the DSM-5 as the so-called B-criteria of somatic symptom

disorder (SSD).<sup>24, 26</sup> Still, consensus on labelling and addressing these symptoms is limited. In this paper, the term 'persistent somatic symptoms' (PSS) is used since the descriptive nature of the term transcends the problem of dualism. Moreover, recent research has found that this term is generally preferred over other terms.<sup>27</sup>

In the Dutch health care system, as well as in many (Western) countries, the GP serves as a gatekeeper for health care in general. The classification of symptoms and illnesses in EHRs by Dutch GPs is based on the International Classification of Primary Care (ICPC) system.<sup>28</sup> Since medical practice historically operates according to mind-body dualism, physicians are required to locate complaints either in the body or the mind.<sup>21</sup> Accordingly, most symptoms and disorders – physical and psychological – have a domain specific diagnostic code in the ICPC. Nonetheless, the current ICPC lacks a specific and clearly defined code for PSS and the ICPC system instructs to register symptoms not fulfilling the criteria for a diagnosis on a symptom level.<sup>29, 30</sup> Arguably, registration of cases with PSS is less straight forward due to the multi-domain nature of PSS even though the ICPC does contain a chapter with multi-domain codes (A-chapter).

Nonetheless, there are international codes available for some PSS-related syndromes (such as, IBS), and the Dutch ICPC also contains codes for FM and CFS.<sup>30</sup>

While several studies have documented ample diagnostic variation regarding patients with PSS in general practice, <sup>31, 32</sup> it is not well documented which codes or other methods GPs use for registration of PSS and if they find their current approach to registration satisfactory. The primary aim of the present study was, therefore, to explore how GPs currently register PSS. The secondary aim was to gauge GPs' perspective on their needs to improve classification, registration and care for PSS.

### Methods

### Study design

A cross-sectional online survey was developed to reach our aims. The survey questions were developed in collaboration with experts in general practice, medical psychology and PSS. The survey was set up in Qualtrics.<sup>33</sup> This paper focuses on GPs' registration behaviour and needs, using the STROBE cross sectional reporting guidelines.<sup>34</sup> Prior to distribution, the survey was pilot tested among four GPs and modified based on their feedback. Informed consent was included at the start of the survey. The ethics committee of Leiden University Medical Centre supplied a waiver of ethical approval (C1O8.045/DJ/gk).

### Procedure

The survey was sent out via e-mail between June and September 2018 to mailing lists of Dutch GPs who consented to be approached anonymously for research purposes, and to email addresses obtained through an overarching Dutch healthcare website (www.zorgkaartnederland.nl). This method ensured optimal distribution over all regions of the Netherlands in order to obtain a representative and generalizable sample. Reminders were sent two weeks after first distribution. Ten gift cards of 25 euro were allotted to GPs who participated and provided us with their email address. The email addresses were not linked to the survey responses.

### Measures

To adhere to the term currently used in guidelines for PSS-related complaints, the Dutch term for MUPS ('SOLK') was used to indicate PSS in the survey. Somatisch onvoldoende verklaarde lichamelijke klachten (SOLK) is literally translated as somatic insufficiently explained physical complaints. In the introduction of the survey, a description of the definition of SOLK was presented: 'We speak of SOLK when regular medical care cannot find an adequate explanation for the complaints with which the patient presents him/herself. Patients with a well-known somatic condition can also have SOLK, either presenting with a totally different complaint or presenting with more severe complaints than is expected.' Distinction and explanation about self-limiting and persistent symptoms were provided. To address conceptual differences between GPs regarding PSS

and to ensure that both the medical and psychological domain of PSS was captured, separate questions were added which specifically addressed PSS in patients with a(n explained) chronic medical condition (i.e., 'patients presenting with more or more severe symptoms than you would expect') and/or the B-criteria of SSD (i.e., 'patients who have maladaptive cognitions, emotions and/or behaviours related to the somatic symptoms'). At the start of the survey, GPs were asked to fill in non-identifying demographic questions. All questions required at least one response to continue to the next question, except comment sections. Below you find a description of the survey questions (for an exact outline of the survey, see appendix A).

Primary aim ('registration of PSS') – The following four items were constructed to reach the primary aim: (1) First a description of a hypothetical patient was given as follows: 'Imagine a patient visiting your office who has consulted you frequently in the previous 6 months with the same or differing complaints. Extensive research has excluded a medical explanation for the complaint(s). For each complaint presented, choose the ICPC-code which you would use most often. You can choose a maximum of three ICPC-codes per complaint.' Then followed 4 complaints on different pages: bowel problems, fatigue, neck and back pain, and shortness of breath. A drop-down menu contained all codes related to the complaint in the thesaurus menu from ICPC-online, 30 which reflects the presentation in GPs' EHR. This list was supplemented with suitable codes based on a PSSexpert panel of GPs (see appendix B for the full list of ICPC-codes from which GPs could choose). The four separate complaints were offered in random order to minimize bias due to presentation order. Next, (2) GPs were asked whether they use the PSS-related syndrome codes A04.01 (CFS), D93 (IBS) and/or L18.01 (FM). Respondents selected one or more of five options: 'Yes, I diagnose the syndromes myself sometimes', 'Yes, I use this code when the syndrome is diagnosed by a medical specialist', 'No, I think these complaints should be reported on a symptom level', 'No, I am not convinced these are distinguishable syndromes', and 'Other, namely...' (with an additional comment section). This question was added to the survey in a later stage and was therefore only presented to 73% (n=189) of the GPs.

Subsequently, (3) GPs were asked whether they mention PSS-related terminology in the (3a) episode name or (3b) free text area (two 4-point scale items (ranging from 'never' to 'always')). Lastly, (4) a description of a hypothetical consultation with a patient with a diagnosed medical condition was given, whereby the patient presented with specific cognitive, emotional and behavioural problems (conforming to the B-criteria of SSD).<sup>24</sup> GPs were asked if they would mention this in their EHR (yes/no, and a comment section).

Secondary aim ('GPs needs') – The following four items were constructed to reach the secondary aim: First, (1) GPs were asked if the lack of an unambiguous way of classifying or coding PSS was problematic for them (yes/no, and a comment section). Next, (2) GPs were asked if they had a need for a code which captures the specific cognitions, emotions and behaviour conforming to the B-criteria of SSD (yes/no, and a comment section). Subsequently, an open-ended question was presented where GPs were asked (3) what they need to be able to improve registration and classification of PSS; and, in order to ensure not missing any needs, this was followed by three specific semi-open-ended questions – (4) if they have needs regarding training, (online) tools or other support, to improve consultations and classification of PSS (response options were: 'no' and 'yes, namely...').

### Data analysis

All results are based on descriptive statistics. Survey responses were summarized as is, using sample sizes and percentages, unless otherwise specified above. For the first hypothetical consultation, codes were first categorized into four groups: symptom-specific codes (e.g., A04-fatigue), general codes – i.e., non-specific codes (e.g., P28-limited function/disability(p)), somatization (P75-somatization disorder) and syndromes (A04.01-CFS, D93-IBS and L18.01-FM) (see appendix B). The responses on the four single complaints were analysed both combined and as separate complaints. For the question regarding the use of PSS-related syndrome codes, the two 'Yes, ...' answering options were combined and the two 'No, ...' answering options were combined to construct total scores.

### Results

Of the approximately 12,000 active GPs in the Netherlands, an estimated 2,000 GPs were reached through our distribution method. In total, 259 GPs (13%) fully completed the survey, with exception to the fourth item (4) which was completed by 189 GPs. Table 1 displays the characteristics of the total sample. Of the GPs who filled out the survey, 60.2% were female, which reflects the current trend towards increasing numbers of female GPs in the Netherlands.<sup>35</sup> GPs from all regions in the Netherlands completed the survey. GPs years since graduation is reasonably evenly distributed over 5-year periods, varying between the smallest group of GPs graduating between 26 to 30 years since participating in the survey (8.5%) and the largest group of GPs graduating between 6 to 10 years before participation (17.8%).

Table 1. Characteristics of the 259 Dutch GPs participating in the study

General practitioners	n=259 (%)				
Gender (female)	156 (60.2)				
Years since graduation					
< 5	41 (15.8)				
6 - 10	46 (17.8)				
11 - 15	44 (14.0)				
16 - 20	39 (15.1)				
21 - 25	31 (12.0)				
26 - 30	22 (8.5)				
> 30	36 (13.9)				
Location of practice					
Urban (Randstad)	81 (31.0)				
North	62 (23.8)				
Middle	102 (35.6)				
South	23 (8.8)				


### Registration of PSS

As shown in Figure 1, GPs vary in their way of reporting PSS. Combining the preferred first choices of code for the four PSS case examples, the general trend indicates that GPs were most likely to register PSS on a symptom-specific level (89.3%). The frequency of choosing general codes increased from 6.9% as a first choice to 31.1% for the second and

45.5% for the third choice. The choice for ICPC code P75 (somatization disorder) increased from 1% as a first choice, to 5.1% and 8.0% for the second and third choice respectively. When presented with fatigue complaints, more than 35.7% chose to report the complaint with P75 (as a second or third choice). Looking at the number of times a syndrome code (IBS, CFS or FM) was generally chosen either as a first, second or third choice, 144 chose D93 (IBS) in case of bowel complaints, 69 GPs chose A04.01 (CFS) in case of fatigue complaints, and 6 chose L18.01 (FM) in case of neck and back pain. For a more detailed description of the choices of ICPC codes per presented symptom, see appendix C.

### Chapter 3

*Figure 1.* Visualizations of general practitioners' order of choosing ICPC-codes for specified persistent somatic symptoms (PSS).



General


Third choice

# Order of ICPC-code preference for PSS-related Bowel Problems Syndrome No answer Syndrome Symptom-specific Symptom-specific Symptom-specific Symptom-specific

First choice

General

Second choice






Table 2 shows the reported likelihood that GPs mention PSS-related terminology and cognitions, emotions or behaviour related to PSS in their EHR. Some GPs mentioned in the comment section that a fear of stigmatization was the reason for avoiding PSS-related terms.

Table 2. GPs (n=259) mentioning PSS-related terminology in their EHR

Do you mention PSS in the episode name?	n (%)		
never	79 (30.5)		
occasionally	146 (56.4)		
often	28 (10.8)		
always	6 (2.3)		
Do you mention PSS in the free-text area?			
never	29 (11.2)		
occasionally	156 (60.2)		
often	64 (24.7)		
always	10 (3.9)		
Do you mention components <sup>a</sup> of PSS in the free-text area?			
Yes	204 (78.8)		

<sup>&</sup>lt;sup>a</sup> Specific thoughts, feelings and behaviour conforming to the B-criteria of SSD

Of all GPs who answered the question regarding ICPC-codes for recognized PSS-syndromes (n=189), 91.5% indicated that they use the codes for IBS (D93), CVS (A04.01) and FM (L18.01) (not shown in Table). The answering options given in the survey are depicted in Table 3. While 68.3% of GPs reported diagnosing the syndrome themselves, several GPs commented that this was only the case for IBS (which was also the case in most GPs who selected the answer 'Other, namely...').

Table 3. GPs (n=189) use of PSS-related syndrome codes

Do you use the PSS-syndrome ICPC codes? a,b	n = 189
Yes, I diagnose the syndromes myself sometimes	129 (68.3)
Yes, I use this code when the syndrome is diagnosed by a medical specialist	74 (39.2)
No, I think these complaints should be reported on a symptom level	25 (12.2)
No, I am not convinced these are distinguishable syndromes	9 (4.8)
Other, namely	19 (10.1)

<sup>&</sup>lt;sup>a</sup> A04.01 – CFS; D93 – IBS; L18.01 - FM

<sup>&</sup>lt;sup>b</sup> Multiple answers possible per GP

### GPs needs

Table 4 shows the results relating to the second aim of this study. Approximately half of GPs (47.9%) reported that the lack of an unambiguous way of classifying or coding PSS is a problem for them. Many GPs commented that a specific code for PSS would be helpful, and some suggested that PSS-codes per tract or a specific code with different severity levels would be helpful. GPs commented requiring widely accepted guidelines in combination with a new PSS-code. Of those who did not see the lack of a specific PSS code as a problem (52.1%), many commented they sometimes describe PSS in the available free text area when registering the patient's somatic complaint. Others felt there is still too much uncertainty regarding PSS to code it, felt unwilling to apply "that label" to a patient, or commented that registration at a symptom level was sufficient. Of all GPs, 32.8% reported that they would like to be able to express PSS-related components – i.e., specific thoughts, feelings and behaviour conforming to the B-criteria of SSD – in a code. Although some of these GPs commented that they found it difficult to specify the components. Additionally, GPs indicated a need for training (56.7%) and/or an – (online) classification and/or risk assessment – tool (58.3%) and/or other support (69.7%). Other PSS-related needs mentioned by GPs in the elective comment sections regarded clearer or more referral options, more time and financial compensation for consultations and better guidelines (although others explicitly mentioned that they found the current guidelines adequate).

**Table 4.** GPs' needs for improving registration and classification of persistent somatic symptoms in EHR

There is no ICPC code for PSS; is this a problem for you?	n=259 (%)
Yes	127 (47.9)
Would you like to express components <sup>a</sup> of PSS in an ICPC code?	n=259 (%)
Yes	85 (32.8)
Do you have a need for to improve consultations/classification for PSS?	n=254 (%)
training (yes)	144 (56.7)
an (online) tool (yes)	148 (58.3)
other support (yes)	177 (69.7)

<sup>&</sup>lt;sup>a</sup> Specific thoughts, feelings and behaviour conforming to the B-criteria of SSD

### Discussion and conclusion

The results of this survey indicate that codes used for registration of PSS in primary care varies widely among GPs. PSS are primarily coded at a specific somatic symptom level and GPs often avoid using terminology related to PSS in their EHR. In addition, GPs indicate they us the codes for well-known PSS-syndromes as IBS, CFS, or FM, although IBS is coded more often than CFS and FM. Besides, the cognitive, emotional, or behavioural components of PSS are sparsely reported in EHRs. Some GPs indicated that they have difficulties in specifying these components. Overall, half of GPs are unsatisfied with current registration options for PSS. Many GPs have a need for additional tools, training or support regarding PSS registration and classification. Still, while GPs provide several suggestions for improvements of the classification system, there is little consensus on this matter.

Looking more specifically at the first aim of this study, many GPs are struggling with registration of PSS and are hesitant to use codes beyond the somatic complaints they objectively observe. This is in line with instructions of the ICPC <sup>13, 30</sup> and previous research findings, reporting that GPs' fear of stigmatization may lead them to avoid codes related to social and psychological problems. <sup>29, 37, 39</sup> On the other hand, respondents did indicate more frequently diagnosing the PSS-related syndrome IBS, compared to CFS and FM, which is in line with previous research indicating that GPs are more proficient in diagnosing IBS. <sup>11</sup> This suggests that registration behaviour may be more depended upon the GP's confidence in classifying PSS than upon fear of stigmatization.

Regarding the second aim of this study, our results show that the current registration and classification options for PSS are insufficient for a substantial number of GPs. These GPs reportedly require a specific code for PSS, in combination with training, tools, a widely accepted guideline, and referral options. In contrast, the literature shows that there are a variety of training options, 40 concise and validated screening questionnaires, 26, 38, 41 and referral options 3, 13, 42 available to GPs. Besides, the Dutch GP association has an elaborate PSS guideline. 13 In line with this guideline, some GPs suggested coding of PSS should be done by severity, which is also in line with studies

which propose the introduction of codes that specify severity to improve documentation of mild PSS.<sup>8, 9, 29</sup> Interestingly, research demonstrated that the GPs' use of subcategories directed at classifying severity is challenged by the GPs' conceptual understanding of PSS.<sup>8</sup> It is therefore conceivable that GPs do indeed need training, and knowledge of the availability of training, to increase their understanding of PSS.

### Strengths and limitations

To the best of our knowledge, this study is the first to capture an overview of GPs' perspectives regarding registration and classification of PSS through exploration of their specific ICPC-registration behaviour. Our data sheds light on GPs' reasoning regarding PSS, confirms the lack of consensus on registration and classification and offers guidance for improvements in registration and classification based on the GPs' reported needs. Nonetheless, some limitations should be noted. First, in order to distribute the survey as broadly as we have, we involved third parties (i.e. regional GP-networks) to promote distribution. This resulted in a limited overview of the number of GPs reached, leading to a rough estimate of the response rate. Second, responses may have been biased by elective participation. Still, although adequate reference data is limited, responses appear fairly representative for the population of GPs in the Netherlands.<sup>36, 43</sup> Regarding the content of the survey, a strength is that face valid answers were facilitated for the choice in ICPC codes by presenting GPs with codes in a drop-down menu, similar to their EHRs' set-up. Nonetheless, this came with the limitation that it is unclear if the more frequently coded 'P75-somatization disorder' in case of fatigue compared to other complaints is a true finding, or whether it demonstrates the limitations of the ICPC coding system itself to facilitate classification of PSS, or if it is related to a lack of potential alternative codes (see appendix B). Lastly, generalisation of the findings should be done with caution, since many questions were based on hypothetical situations.

### **Practical implications**

The great variance in responses and methods for registration of PSS found in our research suggests that clinical practice may be improved by better registration of PSS. Improving classification and providing adequate registration options may support GPs in the overall care for PSS. To improve registration, a clear definition with a specific code for

PSS should be implemented in the ICPC system. Introduction of such a code should be combined with (communication on) training options for GPs, which also broadens the GPs' knowledge on currently available diagnostic tools, guidelines, and referral options. Besides providing more accessible coding and training options, research could support the GP further by developing a data-based screening tool for early identification of patients at risk for PSS. This could be another way to support the GP with their challenges in conceptualizing PSS. Besides, this may promote timely treatment of the cognitive, emotional, and behavioural components of PSS, which, in turn, may decrease the burden of PSS and reduce the risk of iatrogenic harm.

### Conclusion

Registration of PSS in primary care is currently ambiguous. Specific complaints presented by patients with PSS are primarily coded on a symptom-specific level. Approximately half of GPs expressed a need for more coding options for PSS and over half of GPs reported a need for further training, tools or other support regarding PSS. Since many of the latter already exist, improvements should be directed at new options for registration, specifically coding, and increasing and spreading knowledge about PSS, guidelines, available tools, and referral options.

### Acknowledgements

The authors gratefully thank the Dutch GP associations, Huisartsenkringen and Academisch Netwerk Huisartsgeneeskunde (Extramural Leiden Academic Network, Amsterdam UMC, Groningen UMC and Erasmus MC/PrimEUR-network) for distributing the survey and we are very grateful to the many GPs that spent time filling out the survey. Additionally, we appreciate the suggestion of the Amsterdam UMC GP association for adding the question regarding PSS-related syndrome ICPC-codes.

### References

- 1 Khan AA, Khan A, Harezlak J, Tu W, Kroenke K. Somatic Symptoms in Primary Care: Etiology and Outcome. Psychosomatics. 2003;44(6):471–8.
- 2 Kroenke K. Patients presenting with somatic complaints: Epidemiology, psychiatric co-morbidity and management. Int J Methods Psychiatr Res. 2003;12(1):34–43.
- 3 Henningsen P, Zipfel S, Herzog W. Management of functional somatic syndromes. Lancet. 2007;369:946-55.
- 4 Eikelboom EM, Tak LM, Roest AM, Rosmalen JGM. A systematic review and meta-analysis of the percentage of revised diagnoses in functional somatic symptoms. J Psychosom Res. 2016;88(C):60–7.
- De Waal MWM, Arnold IA, Eekhof JAH, Van Hemert AM. Somatoform disorders in general practice: Prevalence, functional impairment and comorbidity with anxiety and depressive disorders. British Journal of Psychiatry. 2004;184:470–6.
- Toft T, Fink P, Oernboel E, Christensen K, Frostholm L, Olesen F. Mental disorders in primary care: prevalence and co-morbidity among disorders. results from the functional illness in primary care (FIP) study. Psychol Med. 2005;35(8):1175–84
- 7 Verhaak PF, Meijer SA, Visser AP, Wolters G. Persistent presentation of medically unexplained symptoms in general practice. Fam Pract. 2006;23(4):414–20.
- 8 Rask MT, Andersen RS, Bro F, Fink P, Rosendal M. Towards a clinically useful diagnosis for mild-to-moderate conditions of medically unexplained symptoms in general practice: A mixed methods study.

  BMC Fam Pract. 2014;15(1).
- 9 Rosendal M, Olde Hartman TC, Aamland A, Van der Horst H, Lucassen P, Budtz-Lilly A, et al. "Medically unexplained" symptoms and symptom disorders in primary care: prognosis-based recognition and classification. BMC Fam Pract. 2017;18(1):1–9.
- Sirri L, Grandi S, Tossani E. Medically unexplained symptoms and general practitioners: a comprehensive survey about their attitudes, experiences and management strategies. Fam Pract. 2017;34(2):201–5.
- 11 Warren JW, Clauw DJ. Functional somatic syndromes: Sensitivities and specificities of self-reports of physician diagnosis. Psychosom Med. 2012;74(9):891–5.
- Spiegel BMR, Farid M, Esrailian E, Talley J, Chang L. Is irritable bowel syndrome a diagnosis of exclusion?: A survey of primary care providers, gastroenterologists, and ibs experts. Am J Gastroenterol. 2010;105(4):848–58.
- Hartman TO, Blankenstein N, Molenaar B, Bentz Van Den Berg D, Van Der Horst H, Arnold I, et al. NHG-Standaard Somatisch Onvoldoende verklaarde Lichamelijke Klachten (SOLK). Huisarts Wet. 2013;56(5):1–18.

- 14 Claassen van Dessel N, van der Wouden JC, Hoekstra T, Dekker J, van der Horst HE. The 2-year course of Medically Unexplained Physical Symptoms (MUPS) in terms of symptom severity and functional status: results of the PROSPECTS cohort study. J Psychosom Res. 2018;104:76–87.
- 15 Konnopka A, Schaefert R, Heinrich S, Kaufmann C, Luppa M, Herzog W, et al. Economics of medically unexplained symptoms: A systematic review of the literature. Psychother Psychosom. 2012;81:265–75.
- Zonneveld LN, Sprangers MA, Kooiman CG, van 't Spijker A, Busschbach JJ. Patients with unexplained physical symptoms have poorer quality of life and higher costs than other patient groups: a cross-sectional study on burden. BMC Health Serv Res. 2013;13:520.
- 17 Pohontsch NJ, Zimmermann T, Jonas C, Lehmann M, Löwe B, Scherer M. Coding of medically unexplained symptoms and somatoform disorders by general practitioners an exploratory focus group study. BMC Fam Pract. 2018;19(1):129.
- Jensen-Doss A, Weisz JR. Diagnostic Agreement Predicts Treatment Process and Outcomes in Youth Mental Health Clinics. 2008;76(5):711-22.
- 19 Majeed A, Car J, Sheikh A. Accuracy and completeness of electronic patient records in primary care. Fam Pract. 2008;25:213–4.
- 20 Swinglehurst D, Greenhalgh T. Caring for the patient, caring for the record: An ethnographic study of "back office" work in upholding quality of care in general practice. BMC Health Serv Res. 2015;15(1):177.
- 21 White PD, Rickards H, Zeman AZJ. Time to end the distinction between mental and neurological illnesses. BMJ. 2012;344:e3454.
- Page L, Wessely S. Medically Unexplained Symptoms: Exacerbating Factors in the Doctor-Patient Encounter. J R Soc Med . 2003;96(5):223–7.
- 23 Creed F, Guthrie E, Fink P, Henningsen P, Rief W, Sharpe M, et al. Is there a better term than "Medically unexplained symptoms"? J Psychosom Res .2010;68:5–8.
- 24 American Psychiatric Association,, Force DSMT. Diagnostic and statistical manual of mental disorders: DSM-5, 2013.
- 25 Fink P, Schröder A. One single diagnosis, bodily distress syndrome, succeeded to capture 10 diagnostic categories of functional somatic syndromes and somatoform disorders. J Psychosom Res . 2010;68(5):415–26.
- 26 Kop WJ, Toussaint A, Mols F, Löwe B. Somatic symptom disorder in the general population: Associations with medical status and health care utilization using the SSD-12. Gen Hosp Psychiatry. 2019;56:36–41.
- 27 Marks EM, Hunter MS. Medically Unexplained Symptoms: An acceptable term? Br J Pain. 2015;9(2):109–14.
- Het Nederlands Huisartsen Genootschap is de wetenschappelijke vereniging van huisartsen: ICPC. https://www.nhg.org/themas/artikelen/icpc. Accessed 08 Dec 2017.

- Schaefert R, Laux G, Kaufmann C, Schellberg D, Bölter R, Szecsenyi J, et al. Diagnosing somatisation disorder (P75) in routine general practice using the International Classification of Primary Care. J Psychosom Res . 2010;69(3):267–77.
- 30 WHO | International Classification of Primary Care, Second edition (ICPC-2). Available from: https://www.who.int/classifications/icd/adaptations/icpc2/en/. Accessed 12 Apr 2021.
- 31 Rosendal M, Bro F, Fink P, Christensen KS, Olesen F. Diagnosis of somatisation: effect of an educational intervention in a cluster randomised controlled trial. BrJ GenPract. 2003;53(497):917-22.
- 32 Rosendal M, Carlsen AH, Rask MT, Moth G. Symptoms as the main problem in primary care: A cross-sectional study of frequency and characteristics. Scand J Prim Health Care. 2015;33(2):91-9.
- 33 Qualtrics. Provo, Utah, USA: Qualtrics, LLC 2019; 2005.
- 34 The EQUATOR Network | Enhancing the QUAlity and Transparency Of Health Research. Available from: https://www.equator-network.org/. Accessed 10 Sept 2020.
- 35 Het Nederlands Huisartsen Genootschap is de wetenschappelijke vereniging van huisartsen: ICPC-online. Available from: https://www.nhg.org/themas/artikelen/icpc-online. Accessed 20 Dec 2018.
- 36 Van Hassel DTP, Kasteleijn A, Kenens RJ. Cijfers uit de registratie van huisartsen peiling 2015 [Internet]. 2016.
- 37 Freidl M, Piralic-Spitzl S, Grohe N, Aigner M. Association between fear of stigma, depressive and anxiety symptoms in patients with somatoform pain disorder. Psychiatr Prax. 2012;39(6):263–6.
- den Boeft M, van der Wouden JC, Rydell-Lexmond TR, de Wit NJ, van der Horst HE, Numans ME.
  Identifying patients with medically unexplained physical symptoms in electronic medical records in primary care: a validation study. BMC Fam Pract. 2014;15:109.
- 39 Nilsson G, Åhlfeldt H, Strender LE. Computerisation, coding, data retrieval and related attitudes among Swedish general practitioners A survey of necessary conditions for a database of diseases and health problems.Int J Med Inform. 2002;65(2):135–43.
- 40 Weiland A, Blankenstein AH, Van Saase JLCM, Van der Molen HT, Jacobs ME, Abels DC, et al. Training Medical Specialists to Communicate Better with Patients with Medically Unexplained Physical Symptoms (MUPS). A Randomized, Controlled Trial. PLoS One. 2015;10(9):e0138342.
- 41 Terluin B, van Marwijk HWJ, Adèr HJ, de Vet HCW, Penninx BWJH, Hermens MLM, et al. The Four-Dimensional Symptom Questionnaire (4DSQ): A validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety and somatization. BMC Psychiatry. 2006;6.
- 42 van Dessel N, den Boeft M, van der Wouden JC, Kleinstäuber M, Leone SS, Terluin B, et al. Nonpharmacological interventions for somatoform disorders and medically unexplained physical symptoms (MUPS) in adults. Cochrane Database of Syst Rev. 2014;(11):CD011142.
- 43 Velden LFJ, van der Hingstman L, Kenens RJ, Batenburg RS. Regionale spreiding van huisartsen: over mogelijke regionale tekorten aan huisartsen anno 2009 en in de nabije toekomst. NIVEL. Utrecht; 2011.

### **Appendixes**

### **Appendix A.** Survey translation: SOLK in general practice, a general practitioners' perspective

This file contains SOLK-related definitions and questions outlined in the measures section of the article are presented below.

### Definition of 'somatisch onvoldoende verklaarde lichamelijke klachten' (SOLK):

We speak of SOLK when regular medical care cannot find an adequate explanation for the complaints with which the patient presents him/herself. Patients with a well-known somatic condition can also have SOLK, either presenting with a totally different complaint or presenting with more severe complaints regarding their diagnosed condition than would be expected.

Part 1: Imagine a patient visiting your office who has consulted you frequently in the precious 6 months with the same or differing complaints. Extensive research has excluded a medical explanation for the complaint(s). For each complaint presented, choose the ICPC-code which you would use most often. You can choose a maximum of three ICPC-codes per complaint. a) Bowel problems Three drop-down menus (see appendix B for list of ICPC codes) b) Fatigue Three drop-down menus (see appendix B for list of ICPC codes) c) Neck and back pain Three drop-down menus (see appendix B for list of ICPC codes) d) Shortness of breath Three drop-down menus (see appendix B for list of ICPC codes) (Each complaint was presented on a new page with the description of the hypothetical patients at the top of the page) Part 2: Do you use the ICPC-codes for fibromyalgia (L18.01), chronic fatigue syndrome (A04.01) and/or irritable bowel syndrome (D93)? (multiple answers possible) Yes, I diagnose these syndromes myself sometimes Yes, I use these codes when the diagnosis is made by a medical specialist No, I prefer reporting these complaints at a symptom level No, I am not convinced that these are discernable syndromes Other, namely... Comments: \_\_\_\_\_\_ Part 3: The next questions are about if or how you report SOLK in your electronic health record (EHR) when you have enough reasons to assume that the complaints are SOLK. now and then often always a) Do you write SOLK in the episode name?

### Chapter 3

b) Do you write SOLK in the free text area of the EHR?	$\bigcirc$	$\circ$	$\bigcirc$	$\circ$
Comments:		 the EHR (there is no	ICPC-code for SO	LK).
O No				
Yes				
Comments:				
Part 4: The questions below are about reg Imagine a patient with a well-known med office. This patient experiences the disease disproportion thoughts, feelings or behavi compared to other patients who have the a) Would you report this in your EHR?	ically explained diso e differently than mo iors related to the di	rder (such as, Crohn ost people with this o sorder or displays se	's disease or asthr disorder; this patio	ma) visiting your ent may have
O No				
Yes				
Comments:b) Would you like to have the option to re	port this (behavior)	 in a specific ICPC-co	de?	
O No				
O Yes				
Comments:				
Part 6: Research indicates that (primary confidence of patients with SOLK. The following questa) Do you have a need for an (online) tool	tions are addressing		ties with consulta	tions and treatment
○ No				
Yes, namely				
Comments:				
b) Do you have a need for training regardi	ing SOLK?			
O No				
Yes, namely				
Comments:				
c) Do you have a need other support rega	rding care for patier	nts with SOLK?		
○ No				
Yes, namely				
Comments:				

Appendix B. Overview of items, ICPC codes and categorization

	Type of complaint					
ICPC	Name	Neck & back	Bowel	Shortness of breath	Fatigue	Category
A01	Pain general/multiple sites	Х	Х			General
A04	Weakness/tiredness general			X	Χ	Symptom-specific
A04.01	Chronic fatigue syndrome			X	Χ	Syndrome
A28	Limited function/disability NOS	Χ	Χ	X	Χ	General
A97	No disease	Χ	Χ	X	Χ	General
A99	General disease NOS	Χ	Χ	X	Χ	General
D01	Abdominal pain/cramps general		Χ			Symptom-specific
D02	Abdominal pain epigastric		Χ			Symptom-specific
D06	Abdominal pain localized other		Χ			Symptom-specific
D20	Mouth/tongue/lip symptom/complt.			X		Symptom-specific
D28	Limited function/disability (d)		Χ			General
D29	Digestive symptom/complaint other		Χ			General
D93	Irritable bowel syndrome		Χ			Syndrome
K02	Pressure/tightness of heart			X		Symptom-specific
K03	Cardiovascular pain NOS			X		General
K04	Palpitations/awareness of heart			Х		Symptom-specific
K05	Irregular heartbeat other			Х		Symptom-specific
L01	Neck symptom/complain	Х				Symptom-specific
L02	Back symptom/complaint	Х				Symptom-specific
L03	Low back symptom/complaint	Χ				Symptom-specific
L18	Muscle pain	Х				Symptom-specific
L18.01	Fibromyalgia	Х				Syndrome
L27	Fear musculoskeletal disease other	Х				General
L28	Limited function/disability (I)	Х				General
L29	Sympt/com. musculoskeletal other	X				General
L79	Sprain/strain of joint NOS	Х				Symptom-specific
L79.01	Whiplashtrauma cervical spine	Х				Symptom-specific
L83	Neck syndrome	Х				Symptom-specific
L86	Back syndrome with radiating pain	X				Symptom-specific
L86.01	Spinal disc hernia (thoracic/lumbar)	X				Symptom-specific
N01	Headache	X				Symptom-specific
N28	Limited function/diability (n)	X	Х			General
N29	Neurological symptom/com other	X	X	Х		General
P28	Limited function/disability (p)	X	X	X	Χ	General
P29	Psychological symptom/com other	X	X	X	X	General
P75	Somatization disorder	X	X	X	X	Somatization
P99	Psychological disorders, other	X	X	X	X	General
R01	Pain respiratory system	**		X		Symptom-specific
R02	Shortness of breath/dyspnoea			X		Symptom-specific
R04	Breathing problem, other			X		General
R28	Limited function/disability (r)			X		General
U13	Bladder symptom/complaint other		Х	^		General
U28	Limited function/disability urinary		X			General

Appendix C. Registration of PSS-related complaints using ICPC (n = 259)

Registration method	First choice <sup>a</sup>	Second choice <sup>a</sup>	Third choice <sup>a</sup>
Combination <sup>b</sup>	n = 1035	n = 791	n = 500
Symptom-specific <sup>c</sup>	924 (89.3)	377 (47.7)	172 (34.4)
General <sup>d</sup>	71 (6.9)	246 (31.1)	227 (45.5)
Somatization <sup>e</sup>	10 (1.0)	40 (5.1)	40 (8.0)
Syndrome <sup>f</sup>	30 (2.9)	128 (16.1)	61 (12.2)
Fatigue	n = 259	n = 154	n = 88
Symptom-specific <sup>c</sup>	252 (97.3)	14 (9.1)	11 (12.5)
General <sup>d</sup>	2 (0.8)	61 (39.6)	47 (53.4)
Somatization <sup>e</sup>	3 (1.2)	24 (15.6)	18 (20.5)
Syndrome <sup>f</sup>	2 (0.8)	55 (35.7)	12 (13.6)
Bowel problems	n = 259	n = 216	n = 139
Symptom-specific <sup>c</sup>	218 (84.2)	104 (48.1)	46 (33.1)
General <sup>d</sup>	11 (4.2)	37 (17.1)	38 (27.3)
Somatization <sup>e</sup>	2 (0.8)	4 (1.9)	10 (7.2)
Syndrome <sup>f</sup>	28 (10.8)	71 (32.9)	45 (32.4)
Shortness of Breath	n = 259	n = 190	n = 107
Symptom-specific <sup>c</sup>	210 (81.1)	67 (35.3)	30 (28.0)
General <sup>d</sup>	45 (17.4)	114 (60.0)	70 (65.4)
Somatization <sup>e</sup>	4 (1.5)	9 (4.7)	7 (6.5)
Syndrome <sup>f</sup>	0 (0)	0 (0)	0 (0)
Neck and back pain	n = 258	n = 231	n = 166
Symptom-specific <sup>c</sup>	244 (94.6)	192 (83.1)	85 (51.2)
General <sup>d</sup>	13 (5.0)	34 (14.7)	72 (43.4)
Somatization <sup>e</sup>	1 (0.4)	3 (1.3)	5 (3.0)
Syndrome <sup>f</sup>	0 (0)	2 (0.9)	4 (2.4)

<sup>&</sup>lt;sup>a</sup> Top three ICPC codes (by category) GPs most likely choose for PSS-related complaints (second and third choice elective).

<sup>&</sup>lt;sup>b</sup> Combination of all responses for the four complaints (fatigue, bowel problems, shortness of breath, and neck and back pain).

<sup>&</sup>lt;sup>c</sup> Includes range of (mainly symptomatic) ICPC codes.

 $<sup>^{\</sup>rm d}$  Includes range of more general ICPC codes without specified location or diagnosis.

<sup>&</sup>lt;sup>e</sup> P75 – somatization disorder.

<sup>&</sup>lt;sup>f</sup> Fatigue: A04.01 – CFS; bowel problems: D93 – IBS; shortness of breath: no code available; neck and back pain: L18.01

<sup>-</sup> FM.