



Universiteit  
Leiden  
The Netherlands

## **Concise review: the endothelial cell extracellular matrix regulates tissue homeostasis and repair**

Witjas, F.M.R.; Berg, B.M. van den; Berg, C.W. van den; Engelse, M.A.; Rabelink, T.J.

### **Citation**

Witjas, F. M. R., Berg, B. M. van den, Berg, C. W. van den, Engelse, M. A., & Rabelink, T. J. (2018). Concise review: the endothelial cell extracellular matrix regulates tissue homeostasis and repair. *Stem Cells Translational Medicine*, 8(4), 375-382.  
doi:10.1002/sctm.18-0155

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3630718>

**Note:** To cite this publication please use the final published version (if applicable).

## Concise Review: The Endothelial Cell Extracellular Matrix Regulates Tissue Homeostasis and Repair

FRANCA M. R. WITJAS , BERNARD M. VAN DEN BERG, CATHELIJNE W. VAN DEN BERG, MARTEN A. ENGELSE, TON J. RABELINK

**Key Words.** Endothelial cell • Induced pluripotent stem cells • Microvasculature • Angiogenesis • Cell adhesion molecules • Cell interactions • Glycosaminoglycan • Tissue regeneration

### ABSTRACT

All tissues are surrounded by a mixture of noncellular matrix components, that not only provide physical and mechanical support to cells, but also mediate biochemical signaling between cells. The extracellular matrix (ECM) of endothelial cells, also known as the perivascular matrix, forms an organ specific vascular niche that orchestrates mechano-, growth factor, and angiocrine signaling required for tissue homeostasis and organ repair. This concise review describes how this perivascular ECM functions as a signaling platform and how this knowledge can impact the field of regenerative medicine, for example, when designing artificial matrices or using decellularized scaffolds from organs. *STEM CELLS TRANSLATIONAL MEDICINE* 2019;8:375–382

### SIGNIFICANCE STATEMENT

In general, all tissues are surrounded by a mixture of noncellular matrix components, that not only provide physical and mechanical support to cells, but also mediate cross-talk between cells. The extracellular matrix of endothelial cells, also known as the perivascular matrix, forms a vascular niche that orchestrates processes required for tissue homeostasis and organ repair. This concise review describes which messages are sent by this perivascular matrix and how this knowledge can be used in the field of regenerative medicine.

### INTRODUCTION

The extracellular matrix (ECM) has central roles in tissue integrity and remodeling. While collagens, laminins, and proteoglycans are the most abundant structural components of the ECM in most tissues, tissue-specific molecular complexity during repair is determined by ECM glycoproteins and the capacity to secrete specific matrix proteins. In mammals, this comprises approximately 300 proteins, often referred to as the matrisome [1]. In particular, remodeling of ECM at the level of capillaries, in a very fundamental way, enables the interaction between endothelial cells, circulating immunocytes, and tissue that is required for homeostasis and regeneration. A lot of insight in this adaptive vascular ECM remodeling has been derived from research into developmental biology and cancer metastasis [2–7]. However, more recently these same processes have also been implicated in tissue repair [8]. From this perspective, understanding the biology and regulation of the ECM is key to the field of regenerative medicine, where the use of artificial matrices to drive stem

cell maturation is an important theme. In this concise review, we will in particular discuss the microvascular ECM structure as an essential signaling platform for tissue repair and how understanding its biology may help advance the field of regenerative medicine.

### THE CONCEPT OF BIDIRECTIONAL ANGIOCRINE SIGNALING IN TISSUE HOMEOSTASIS AND REPAIR

#### Vascularization Induces Maturation of Stem Cell-Derived Tissue and Tissue Regeneration

In addition to delivering oxygen and nutrients to cells as passive and permissive conduits, and clearing cellular waste products simultaneously, blood vessels also provide instructive signals for organogenesis. During embryogenesis, endothelial progenitor cells invade into nascent organs, develop into different endothelial phenotypes, and establish organ-specific vascular niches which release endothelium-derived growth factors and trophogens (angiocrine factors) to

The Eindhoven Laboratory for Vascular and Regenerative Medicine, Department of Internal Medicine, Division of Nephrology, Leiden University Medical Center, Leiden, The Netherlands

Correspondence: Ton J. Rabelink, M.D. Ph.D., Division of Nephrology & Internal Medicine at Leiden University Medical Center, Postbus 9600, Postzone C7-Q, Albinusdreef 2 2300 RC Leiden, The Netherlands. Telephone: +31 71 526 2148; e-mail: a.j.rabelink@lumc.nl

Received July 17, 2018; accepted for publication October 26, 2018; first published December 11, 2018.

<http://dx.doi.org/10.1002/sctm.18-0155>

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

promote stem cell differentiation (extensively reviewed by Raffi et al. [7]).

The relevance of this biology for regenerative medicine is illustrated by recent experiments where we demonstrated that prolonged culture of induced pluripotent stem cells (iPSCs)-derived kidney organoids failed to induce maturation of kidney beyond early embryonic tissue. However, maturation and segmental cell specification could be induced when vascularization of the organoids occurred upon transplantation *in vivo*, illustrating the requirement of vascular signaling for differentiation [9, 10]. Functional differentiation could also be observed in stem cell-derived beta cells upon vascularization after transplantation under the kidney capsule [9–11]. Similarly, it was recently shown that transplantation of human brain organoids into adult mouse brains resulted in progressive neuronal differentiation and maturation, gliogenesis and axonal growth of engrafted organoids. After the organoids became vascularized, neuronal activity could be demonstrated [12].

### Endothelial Cells Are Crucial for Tissue Regeneration Through Angiocrine Signaling

The angiocrine function of EC is preserved beyond embryogenesis in specialized microvascular areas such as sinusoids where ECs are in close apposition with progenitor cells or diploid epithelial cells that have preserved proliferative potential [13]. For example, capillary sinusoidal endothelial cells (SECs) within liver and bone marrow comprise phenotypically and functionally discrete populations of endothelial cells. After partial hepatectomy, liver SECs (LSECs) stimulate hepatocyte proliferation through a process of angiocrine production of hepatocyte growth factor and Wnt2 [14]. Subsequently, LSECs undergo angiogenesis to meet the demand in blood supply to regenerate liver tissue. Similarly, after chemotherapy and irradiation, activated bone marrow SECs reconstitute hematopoiesis by angiocrine expression of Notch ligands and insulin-like growth factor-binding proteins [7, 15, 16]. Pulmonary capillary endothelial cells also produce angiocrine growth factors that induce proliferation of epithelial progenitor cells thereby supporting alveologenesis [17]. There is a large variety in such angiocrine factors that can be produced by the endothelium, while the secretion is very tissue specific and still largely unknown [7].

### Organotypic EC Phenotype Depends upon Instruction from Neighboring Cells and Vice Versa


Endothelial cells in turn depend for their survival and organotypic function upon instructive signaling from neighboring cells as well. For example, the highly specialized fenestrated capillary EC phenotype in the glomerulus is dependent upon vascular endothelial growth factor (VEGF) signaling from the podocytes [18], as illustrated by experiments where podocyte specific deletion of VEGF secretion results in loss of fenestrae and glomerular endothelioses. Angiopoietin 1 is another growth factor that is secreted by perivascular stromal cells such as pericytes, to maintain a stable endothelial cell phenotype [19]. Loss of angiopoietin 1 mediated Tie2 kinase signaling in the endothelium leads to endothelial cell proliferation and loss of organotypic characteristics such as fenestration. Conditional deletion of VEGF-A receptor-2 in liver ECs or bone marrow ECs [20], thus disrupting the stable EC phenotype, inhibits liver and bone marrow regeneration underscoring the necessity of these angiocrine endothelial beds to respond to

signaling molecules such as VEGF to exert their angiocrine regenerative properties.

### Endothelial Cell Extracellular Matrix Serves as an Important Platform for Bidirectional Signaling of Angiocrine EC to Its Neighboring Cells

The bidirectional exchange takes place at the EC extracellular matrix, where the matrix should allow for the formation of specific gradients of growth factors and subsequent engagement with surface receptors (see Figs. 1, 2). The ultrastructure of ECM that enables endothelial cells to engage with their direct surroundings through angiocrine signaling consists first of all of a very dynamically regulated subendothelial basement membrane (BM), containing collagen type IV isoforms, laminin isoforms, heparan sulfate proteoglycans (perlecan or agrin), and nidogen-1 and/or nidogen-2. Collagen type IV and laminins are the two major BM components that self-assemble to form independent networks that confer structural stability and biological activity, respectively. These two networks are connected by perlecan and nidogens [21]. Vascular endothelial cells express laminin  $\alpha 4$  and  $\alpha 5$  chains that combine with laminin  $\beta 1$  and  $\gamma 1$  chains to form laminins 411 and 511 (isotypes 8 and 10), respectively. Laminins are considered to be the major BM components responsible for the biological functions of BMs; that is, for transducing signals that control cell migration, survival, proliferation, and differentiation. Laminin  $\alpha$ -chain distribution and expression depends on endothelial cell type, state of vessel growth, and activation state [22, 23]. Laminin  $\alpha 5$  chain, for example, is strongly expressed in most capillary BMs and inhibits leukocyte transmigration [24]. Laminin isoforms also bind to growth factors (VEGF, platelet-derived growth factor [PDGF], fibroblast growth factor [FGF], and epidermal growth factor [EGF]) with high affinity, through their heparin-binding domains located in the  $\alpha$  chain laminin-type G domain [25]. In addition, membrane-bound proteoglycans and glycosaminoglycans such as hyaluronan, chondroitin sulfate, and heparan sulfate, with terminal sialic acids, contribute to binding and sequestering of growth factors and morphogens with their surface receptors. In embryology altering the proteoglycan composition of this layer results in abnormal vascular development and loss of angiocrine signaling molecules [26]. Using endothelial specific deletion of hyaluronan synthesis, we recently found that also in adult life this layer is essential for endothelial stabilization through specific binding of vascular growth factors [27].

Finally, a fibrillar interstitial matrix underlies the endothelial BM and serves to further interconnect the endothelium with its neighboring tissue. In general, it is composed largely of the fibrillar collagen types I and III together with chondroitin sulfate and dermatan sulfate proteoglycans such as decorin and biglycan, and again multiadhesive glycoproteins [28], although its composition may change from one capillary bed to another. The composition of this latter layer is primarily controlled by resident tissue fibroblasts, that act as a signaling hub for tissue homeostasis and repair. Like the endothelium, they have an organ-specific phenotype where anatomical localization is related to the specific transcriptome profiles, such as embryonic patterning genes, as well as function [29]. While these cells intimately communicate with tissue cells and the immune system to preserve an anti-inflammatory and trophic milieu, they can respond to injury by differentiating into proliferating myofibroblasts and eventually converting into a



**Figure 1.** The endothelial basement membrane, composed of laminin and collagen IV strands which self-assemble to form an independent network crosslinked by perlecan and nidogen, is bound by the endothelial cell through integrins (right lower panel). The glycocalyx, formed by membrane-bound proteoglycans (GAGs), is able to sequester growth factors and constitutes a part of the interstitial matrix functioning as a signaling platform. Long polymeric hyaluronan (orange) crosslinks proteoglycans that carry heparan sulfate (gray) and sialated surface proteins (red), together constituting a gel like layer. Abbreviation: EC, endothelial cell.


nonproliferating, matrix-producing phenotype. Where the role of the latter in scar formation is undisputed, the way by which tissue fibroblast use and regulate their extracellular microenvironment for homeostasis is not well understood.

#### THE PERIVASCULAR EXTRACELLULAR MATRIX AS A SIGNALING PLATFORM FOR REPAIR AND REGENERATION

During tissue repair and regeneration, where downstream signaling pathways such as Akt (protein kinase B) and mammalian target of rapamycin regulate endothelial cellular events, multilayered perivascular ECM remodeling can be discerned. Matrix metalloproteinases (MMPs), which normally reside in a quiescent state within the ECM can be activated upon injury, in particular inflammation, through proteolytic cleavage by proteases such as MT1-MMP on leukocytes, or by direct redox activation. The subsequent ECM degradation leads to the production and release of

ECM components that further modulate ECM remodeling and set the stage to resynthesize the ECM. These include:

- a. During EC injury rapid activation of latent MMPs and subsequent generation of matrix fragments occurs through activation of endothelial cells and invading immunocytes (Fig. 3). The resulting protease-mediated fragmentation of matrix proteins results in generation of matrikines. Matrikines are ECM fragments that are generated through proteolysis in a highly regulated manner and which, for example, through exposition of binding domains, can elicit specific local biological responses [30]. Collagen and fibronectin-derived peptides are well-known matrikines that have been implicated in further activation of immunocytes and fibroblasts and modulating endothelial function. For example, exposition of the terminal domain of fibronectin during ECM degradation can stimulate MMP1 formation by fibroblasts and hence further stimulate ECM remodeling [30].




**Figure 2.** Matricellular proteins and glycosaminoglycans that contain heparan sulfate bind with growth factors and orchestrate receptor interaction at the endothelial cell membrane. This will elicit growth factor-induced downstream cellular events such as Akt (protein kinase B) and mTOR signaling, which define endothelial adaptive responses, such as the secretion of angiocrine factors. The transmission of angiocrine factors to neighboring cells again is governed in turn by these same endothelial extracellular matrix components. Abbreviations: EC, endothelial cell; FGFR, fibroblast growth factor receptor; HAS/hyaluronan, hyaluronan acid synthase with its product hyaluronan; mTOR, mammalian target of rapamycin; TIE2, tyrosine kinase with Immunoglobulin-like and EGF-like domains 2; VEGF, vascular endothelial growth factor.

Another example is the proteolytic cleavage of laminin and collagen which has been shown to result in fragments that contain EGF-like repeats capable of EGF receptor transactivation and reinitiating endothelial ECM synthesis (reviewed in [31]). Together this leads to the initial remodeling of the structural ECM network in the BM and allows for resynthesis of a new provisional EC matrix. This initial step in remodeling of the perivascular matrix is important to set the stage for tissue regeneration.

- b. Matricellular proteins, such as CCN (acronym of CYR61, connective tissue growth factor [CTGF], and NOV)-family of proteins, TSP-1 (thrombospondin-1), tenascin-C and SPARC (secreted protein acidic and rich in cysteine) have been identified as components of the ECM during developmental processes and organogenesis (for review see Adams [32]). The term matricellular has been applied to this group of extracellular proteins that do not contribute directly to the formation of structural ECM elements but serve to modulate

cell–matrix interactions and cell function. While they usually have limited expression in adult life, they can become upregulated and released from the ECM during injury and repair [33–36]. The production and release of matricellular proteins upon injury allows for binding of these proteins to a variety of receptors, growth factors and proteases in the capillary pericellular space, generally reducing cell adhesion and proliferation, and thus tuning ECM remodeling down (Fig. 3). These effects are very specific for the various matricellular proteins [37]; for example the matricellular protein TSP-1 has been shown to act as a strong inhibitor of angiogenesis. It binds to the multiligand CD36 that is expressed on activated endothelium and that induces multiple proapoptotic antiproliferative and anti-inflammatory downstream signaling events [38, 39]. Through these effects TSP-1 can modulate response-to-injury processes. For example, during the proliferative phase of infarct healing, localized induction of TSP-1 in the ischemic zone of the myocardium prevents expansion



**Figure 3.** Activated leukocytes, through the release of Cathepsin L can activate heparanase, leading to the degradation of the luminal endothelial surface glycoalyx [43]. This allows the leukocyte to engage with endothelial surface adhesion receptors (such as ICAM-1) and form a podosome that can invade into the subendothelial matrix. The podosome delivers MT-1 leading to subsequent binding and local activation of MMPs and matrix degradation. The release of matrikines and matricellular proteins such as TSP-1 from the matrix counter-regulate matrix degradation through their respective activation of EGF and CD36 receptors on the subendothelial surface. Abbreviations: EGF, epidermal growth factor; ICAM-1, intracellular adhesion molecule 1; MMPs, matrix metalloproteinases; MT-1, membrane type I-metalloproteinase; TSP-1, thrombospondin-1.

of inflammation into viable myocardial areas through local activation of anti-inflammatory signals [40]. Another example is that TSP-1 through its modulation of endothelial function can drive lineage-specific differentiation of bronchoalveolar stem cells [41]. CCN2/CTGF, expressed by endothelial cells, plays an important role in pericyte recruitment through PDGF signaling and subsequent vascular stabilization [42]. Through integrin binding, CCN2 mediates vascular cell migration and proliferation and regulates endothelial BM formation. The general association of matricellular proteins with stem cell niches across species further suggests that matricellular proteins should be considered more closely for their potential role tissue regeneration [5, 32].

c. Finally, dynamic remodeling and chemical editing of the gel-like glycoalyx envelope of endothelial cells occurs during injury and repair (Fig. 3). The changes in this glycoalyx are probably among the most critical in controlling angiogenesis and angiocrine signaling. In particular, the ability of heparan sulfate domains to mediate binding of specific proteins is fundamental to the organization of protein–receptor interactions at the cell surface and for the creation of chemotactic gradients of growth factors and chemokines [43, 44]. For example, heparan sulfate facilitates the interaction of growth factors, such as FGF-2, VEGF and heparin-binding epidermal-like

growth factor, with their respective receptors [45–49]. Upon EC injury resynthesis of membrane bound glycosaminoglycans occurs where sulfation patterns, and consequently also growth factor and morphogen binding, are modified to adapt to the altered microenvironment (for review see Rabelink et al. [43]). Several lines of evidence support the concept that such perivascular ECM remodeling favors tissue regeneration. For example, in newts, cardiac regeneration following injury was preceded by formation of a matrix network comprising tenascin-C, fibronectin, and hyaluronan. This specialized matrix network may serve as a path for progenitor cells as well as play an active role in activation of a regenerative program [50].

## IMPLICATIONS FOR REGENERATIVE MEDICINE

### Mimicking the Regenerative Endothelial ECM

Understanding the biology of the provisional perivascular EC matrix during repair has implications for biomaterials that are currently being developed for regenerative medicine and tissue engineering in particular. Today, many micro/nanofabrication technologies have been developed for tissue engineering applications. These technologies can be used to create geometrically

defined matrix structures that enable interactions with a specific matrix topography. However, still little knowledge is available as to how to engineer the qualitative characteristics of these materials, such that they mimic the regenerative properties of the perivascular EC matrix. Recently, synthetic neoproteoglycans (neoPGs) with affinity for FGF2 were synthesized, thus mimicking an essential component of the EC glycocalyx envelope. These neoPGs were introduced into plasma membranes of ESCs deficient in HS biosynthesis. Here, they assumed the function of native heparan sulfate proteoglycans, rescued FGF2-mediated kinase activity, and promoted neural stem cell specification [51]. A 2018 study described the covalent incorporation of laminin heparin binding domains into fibrin matrices, thus recreating components of the BM in a hydrogel [24]. This addition improved retention of growth factors and significantly enhanced the efficacy of VEGF and PDGF to promote wound healing *in vivo*, under conditions where the growth factors alone in fibrin are inefficient [25]. Another example, where insight in the biology of the EC matrix was used was a biomolecular engineering approach, which introduced cell-instructive peptide motifs of the matricellular protein CNN within a recombinant elastin-like protein polymer. This polymer elicits integrin-specific endothelial cell survival and function, and would therefore serve as a promising biomaterial for tissue regeneration [52].

Understanding how ECM components induce cell differentiation is also critical to the field of stem cell and organoid culture. Stem cell-derived organoids recapitulate multiple aspects of real organs making them promising models to study organ development, function, and disease. However, the full potential of organoids has remained unrealized, owing to the poorly defined animal-derived matrices in which they are grown. Typically, such organoids are grown in Matrigel, which is a solubilized BM preparation extracted from the Engelbreth-Holm-Swarm mouse sarcoma, a tumor rich in such ECM proteins as laminin (a major component), collagen IV, heparin sulfate proteoglycans, entactin/nidogen, and growth factors. A 2014 study showed that such a Matrigel environment can be recreated by using enzymatically crosslinked polyethylene glycol hydrogels that include the RGD (Arg-Gly-Asp) peptide of fibronectin [53]. It appeared that in addition to the presence of ECM components mechanical properties of the hydrogel were of relevance as well: matrix stiffness significantly enhanced stem cell expansion through a yes-associated protein 1-dependent mechanism [54]. Stem cell differentiation and organoid formation, on the other hand, required a soft matrix and laminin-based adhesion [55, 56].

### The Use of Decellularized Matrices as Regenerative Matrix

Another approach to capitalize on the regenerative and instructive potential of the ECM in general and that of the vasculature in particular, is to use decellularized ECM obtained from organs. This field was pioneered by Ott and coworkers who developed a standard process for pressure-controlled perfusion decellularization of whole organs for generating acellular 3D scaffolds with preserved ECM protein content, architecture, and perfusable vascular conduits [57]. By applying antegrade perfusion of detergents and subsequent washes to arterial vasculature at low physiological pressures, successful decellularization of complex

organs (i.e., hearts, lungs, and kidneys) can be performed [57–59]. The resulting scaffolds can be used in *ex vivo* culture models where the ECM may provide important information to repopulating cells through retention of growth factor or niche-specific cues. For example, it was shown that hepatic SECs maintained their differentiated state longer when cultured on ECM- from the liver than on ECM-derived from the urinary bladder or the small intestine, emphasizing the crosstalk and organ specificity of the ECM and SECs [60]. The instructive capacity of decellularized matrix on differentiation of early progenitor or stem cells into endothelial cells has proven successful. For instance, embryonic stem cells or human nephrosphere-derived PAX2 positive cells differentiate into endothelial cells when cultured on decellularized kidney scaffolds [61–63]. The challenge here is to unravel the culture conditions during which the interaction between early progenitor cells and decellularized matrix scaffolds are optimal. For example, we could recently demonstrate that decellularized scaffolds of human kidney's still contain endothelial glycosaminoglycans that can be reloaded with EC growth factors, and which subsequently resulted in improved adherence of iPSC-derived ECs and functional recellularized vascular conduits [64].

In addition, it was recently demonstrated [65] that the recellularization of decellularized rat gut using both epithelial and endothelial iPSCs showed functionality after 4 weeks, indicating the therapeutic potential of this approach.

### CONCLUSION

Further insight into the matrixome, at the very critical interface between the perivascular capillary EC and niches of tissue with proliferative capacity (progenitor cells) may prove to be key in developing successful biomaterials for tissue engineering. At the same time, the complexity of the biology of glycosaminoglycans and matricellular proteins is daunting, incompletely understood and currently hard to resynthesize and recreate *in vitro*. Given the central role of the endothelial cell matrix in coordinating tissue homeostasis, we argue that focusing on understanding the dynamic regulation and composition of this matrix is of particular relevance.

### ACKNOWLEDGMENTS

The work of the authors is supported by RECellularizing ORgan Donors for KIDney bioengineering (RECORD KID, Dutch Kidney Foundation, 15RN02). We thank M. Zuurmond for preparing the illustrations.

### AUTHOR CONTRIBUTIONS

F.M.R.W and T.J.R. wrote and finalized the manuscript. C.W.v.d.B, B.M.v.d.B, M.A.E, commented and improved the manuscript writing.

### DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

The authors indicated no potential conflicts of interest.

## REFERENCES

- 1 Hynes RO, Naba A. Overview of the matrisome—An inventory of extracellular matrix constituents and functions. *Cold Spring Harb Perspect Biol* 2012;4:a004903.
- 2 Pettersson A, Nagy JA, Brown LF et al. Heterogeneity of the angiogenic response induced in different normal adult tissues by vascular permeability factor/vascular endothelial growth factor. *Lab Invest* 2000;80:99–115.
- 3 Davis GE, Senger DR. Endothelial extracellular matrix: Biosynthesis, remodeling, and functions during vascular morphogenesis and neovessel stabilization. *Circ Res* 2005;97:1093–1107.
- 4 Jarvelainen H, Sainio A, Koulu M et al. Extracellular matrix molecules: Potential targets in pharmacotherapy. *Pharmacol Rev* 2009;61:198–223.
- 5 Vorotnikova E, McIntosh D, Dewilde A et al. Extracellular matrix-derived products modulate endothelial and progenitor cell migration and proliferation in vitro and stimulate regenerative healing in vivo. *Matrix Biol* 2010;29:690–700.
- 6 Carmeliet P, Jain RK. Molecular mechanisms and clinical applications of angiogenesis. *Nature* 2011;473:298–307.
- 7 Rafii S, Butler JM, Ding BS. Angiocrine functions of organ-specific endothelial cells. *Nature* 2016;529:316–325.
- 8 Keane TJ, Horejs CM, Stevens MM. Scarring vs. functional healing: Matrix-based strategies to regulate tissue repair. *Adv Drug Deliv Rev* 2018;129:407–419.
- 9 van den Berg CW, Ritsma L, Avramut MC et al. Renal subcapsular transplantation of PSC-derived kidney organoids induces neo-vasculogenesis and significant glomerular and tubular maturation in vivo. *Stem Cell Reports* 2018;10:751–765.
- 10 Takebe T, Enomura M, Yoshizawa E et al. Vascularized and complex organ buds from diverse tissues via mesenchymal cell-driven condensation. *Cell Stem Cell* 2015;16:556–565.
- 11 Raikwar SP, Kim EM, Sivitz WI et al. Human iPSC cell-derived insulin producing cells form vascularized organoids under the kidney capsules of diabetic mice. *PLoS One* 2015;10:e0116582.
- 12 Mansour AA, Goncalves JT, Bloyd CW et al. An in vivo model of functional and vascularized human brain organoids. *Nat Biotechnol* 2018;36:432–441.
- 13 Wang B, Zhao L, Fish M et al. Self-renewing diploid Axin2(+) cells fuel homeostatic renewal of the liver. *Nature* 2015;524:180–185.
- 14 Ding BS, Nolan DJ, Butler JM et al. Inductive angiocrine signals from sinusoidal endothelium are required for liver regeneration. *Nature* 2010;468:310–315.
- 15 Kobayashi H, Butler JM, O'Donnell R et al. Angiocrine factors from Akt-activated endothelial cells balance self-renewal and differentiation of haematopoietic stem cells. *Nat Cell Biol* 2010;12:1046–1056.
- 16 Butler JM, Kobayashi H, Rafii S. Instructive role of the vascular niche in promoting tumour growth and tissue repair by angiocrine factors. *Nat Rev Cancer* 2010;10:138–146.
- 17 Ding BS, Nolan DJ, Guo P et al. Endothelial-derived angiocrine signals induce and sustain regenerative lung alveolarization. *Cell* 2011;147:539–553.
- 18 Bartlett CS, Jeansson M, Quaggin SE. Vascular growth factors and glomerular disease. *Annu Rev Physiol* 2016;78:437–461.
- 19 Jeansson M, Gawlik A, Anderson G et al. Angiopoietin-1 is essential in mouse vasculature during development and in response to injury. *J Clin Invest* 2011;121:2278–2289.
- 20 Butler JM, Nolan DJ, Vertes EL et al. Endothelial cells are essential for the self-renewal and repopulation of Notch-dependent hematopoietic stem cells. *Cell Stem Cell* 2010;6:251–264.
- 21 Miner JH. Laminins and their roles in mammals. *Microsc Res Tech* 2008;71:349–356.
- 22 Frieser M, Nockel H, Pausch F et al. Cloning of the mouse laminin alpha 4 cDNA. Expression in a subset of endothelium. *Eur J Biochem* 1997;246:727–735.
- 23 Sorokin LM, Pausch F, Durbeek M et al. Differential expression of five laminin alpha (1-5) chains in developing and adult mouse kidney. *Dev Dyn* 1997;210:446–462.
- 24 Wu C, Ivars F, Anderson P et al. Endothelial basement membrane laminin alpha5 selectively inhibits T lymphocyte extravasation into the brain. *Nat Med* 2009;15:519–527.
- 25 Ishihara J, Ishihara A, Fukunaga K et al. Laminin heparin-binding peptides bind to several growth factors and enhance diabetic wound healing. *Nat Commun* 2018;9:2163.
- 26 Henderson-Toth CE, Jahnsen ED, Jamarani R et al. The glycocalyx is present as soon as blood flow is initiated and is required for normal vascular development. *Dev Biol* 2012;369:330–339.
- 27 van den Berg BM, Wang G, Boels MGS et al. Endothelial glucose metabolism regulates glycocalyx hyaluronan synthesis and thereby vessel stability. *International Vascular Biology Meeting*; 2018.
- 28 Yousif LF, Di Russo J, Sorokin L. Laminin isoforms in endothelial and perivascular basement membranes. *Cell Adh Migr* 2013;7:101–110.
- 29 Leuning DG, Reinders ME, Li J et al. Clinical-grade isolated human kidney perivascular stromal cells as an organotypic cell source for kidney regenerative medicine. *STEM CELLS TRANSLATIONAL MEDICINE* 2017;6:405–418.
- 30 Bonnans C, Chou J, Werb Z. Remodelling the extracellular matrix in development and disease. *Nat Rev Mol Cell Biol* 2014;15:786–801.
- 31 Tran KT, Lamb P, Deng JS. Matrikines and matricryptins: Implications for cutaneous cancers and skin repair. *J Dermatol Sci* 2005;40:11–20.
- 32 Adams JC. Matricellular proteins: Functional insights from non-mammalian animal models. *Curr Top Dev Biol* 2018;130:39–105.
- 33 Pi L, Fu C, Lu Y et al. Vascular endothelial cell-specific connective tissue growth factor (CTGF) is necessary for development of chronic hypoxia-induced pulmonary hypertension. *Front Physiol* 2018;9:138.
- 34 Paulitti A, Andreuzzi E, Bizzotto D et al. The ablation of the matricellular protein EMILIN2 causes defective vascularization due to impaired EGFR-dependent IL-8 production affecting tumor growth. *Oncogene* 2018;37:3399–3414.
- 35 Qu Y, Oloniskin T, Bain W et al. Thrombospondin-1 protects against pathogen-induced lung injury by limiting extracellular matrix proteolysis. *JCI Insight* 2018;3.
- 36 Childs BG, Li H, van Deursen JM. Senescent cells: A therapeutic target for cardiovascular disease. *J Clin Invest* 2018;128:1217–1228.
- 37 Bornstein P, Sage EH. Matricellular proteins: Extracellular modulators of cell function. *Curr Opin Cell Biol* 2002;14:608–616.
- 38 Jimenez B, Volpert OV, Crawford SE et al. Signals leading to apoptosis-dependent inhibition of neovascularization by thrombospondin-1. *Nat Med* 2000;6:41–48.
- 39 Fei P, Zaitoun I, Farnoodian M et al. Expression of thrombospondin-1 modulates the angioinflammatory phenotype of choroidal endothelial cells. *PLoS One* 2014;9:e116423.
- 40 Frangogiannis NG. The extracellular matrix in myocardial injury, repair, and remodeling. *J Clin Invest* 2017;127:1600–1612.
- 41 Lee JH, Bhang DH, Beede A et al. Lung stem cell differentiation in mice directed by endothelial cells via a BMP4-NFATc1-thrombospondin-1 axis. *Cell* 2014;156:440–455.
- 42 Hall-Glenn F, De Young RA, Huang BL et al. CCN2/connective tissue growth factor is essential for pericyte adhesion and endothelial basement membrane formation during angiogenesis. *PLoS One* 2012;7:e30562.
- 43 Rabelink TJ, van den Berg BM, Garsen M et al. Heparanase: Roles in cell survival, extracellular matrix remodelling and the development of kidney disease. *Nat Rev Nephrol* 2017;13:201–212.
- 44 Bishop JR, Schuksz M, Esko JD. Heparan sulphate proteoglycans fine-tune mammalian physiology. *Nature* 2007;446:1030–1037.
- 45 Wight TN. A role for proteoglycans in vascular disease. *Matrix Biol* 2018;71-72:396–420.
- 46 Kang I, Chang MY, Wight TN et al. Proteoglycans as immunomodulators of the innate immune response to lung infection. *J Histochem Cytochem* 2018;66:241–259.
- 47 Hering TM, Beller JA, Calout CM et al. Proteoglycans of reactive rat cortical astrocyte cultures: Abundance of N-unsubstituted glucosamine-enriched heparan sulfate. *Matrix Biol* 2015;41:8–18.
- 48 Zhang B, Xiao W, Qiu H et al. Heparan sulfate deficiency disrupts developmental angiogenesis and causes congenital diaphragmatic hernia. *J Clin Invest* 2014;124:209–221.
- 49 Higginson JR, Thompson SM, Santos-Silva A et al. Differential sulfation remodelling of heparan sulfate by extracellular 6-O-sulfatases regulates fibroblast growth factor-induced boundary formation by glial cells: implications for glial cell transplantation. *J Neurosci* 2012;32:15902–15912.
- 50 Mercer SE, Odelberg SJ, Simon HG. A dynamic spatiotemporal extracellular matrix facilitates epicardial-mediated vertebrate heart regeneration. *Dev Biol* 2013;382:457–469.
- 51 Huang ML, Smith RA, Trierer GW et al. Glycocalyx remodeling with proteoglycan mimetics promotes neural specification in embryonic stem cells. *J Am Chem Soc* 2014;136:10565–10568.
- 52 Ravi S, Haller CA, Sallach RE et al. Cell behavior on a CCN1 functionalized elastin-mimetic protein polymer. *Biomaterials* 2012;33:2431–2438.
- 53 Van Hove AH, Burke K, Antonienko E et al. Enzymatically-responsive pro-angiogenic



peptide-releasing poly(ethylene glycol) hydrogels promote vascularization in vivo. *J Control Release* 2015;217:191–201.

**54** Gjorevski N, Ranga A, Lutolf MP. Bioengineering approaches to guide stem cell-based organogenesis. *Development* 2014;141:1794–1804.

**55** Lutolf MP, Gilbert PM, Blau HM. Designing materials to direct stem-cell fate. *Nature* 2009;462:433–441.

**56** Lutolf MP, Doyonnas R, Havenstrite K et al. Perturbation of single hematopoietic stem cell fates in artificial niches. *Integr Biol* 2009;1:59–69.

**57** Ott HC, Matthiesen TS, Goh SK et al. Perfusion-decellularized matrix: Using nature's platform to engineer a bioartificial heart. *Nat Med* 2008;14:213–221.

**58** Song JJ, Guyette JP, Gilpin SE et al. Regeneration and experimental orthotopic transplantation of a bioengineered kidney. *Nat Med* 2013;19:646–651.

**59** Song JJ, Ott HC. Bioartificial lung engineering. *Am J Transplant* 2012;12:283–288.

**60** Sellaro TL, Ravindra AK, Stolz DB et al. Maintenance of hepatic sinusoidal endothelial cell phenotype in vitro using organ-specific extracellular matrix scaffolds. *Tissue Eng* 2007;13:2301–2310.

**61** Ross EA, Abrahamson DR, St John P et al. Mouse stem cells seeded into decellularized rat kidney scaffolds endothelialize and remodel basement membranes. *Organogenesis* 2012;8:49–55.

**62** Bombelli S, Meregalli C, Scalia C et al. Nephrosphere-derived cells are induced to

multilineage differentiation when cultured on human decellularized kidney scaffolds. *Am J Pathol* 2018;188:184–195.

**63** Batchelder CA, Martinez ML, Tarantal AF. Natural scaffolds for renal differentiation of human embryonic stem cells for kidney tissue engineering. *PLoS One* 2015;10:e0143849.

**64** ISSCR 2018 Annual Meeting Poster Abstract Book. 2018. [http://www.isscr.org/docs/default-source/2018-melbourne-annmtng/66670-isscr-abstracts\\_with-links.pdf?sfvrsn=4&utm\\_source=ISSCR-Infomz&utm\\_medium=email&utm\\_campaign=default](http://www.isscr.org/docs/default-source/2018-melbourne-annmtng/66670-isscr-abstracts_with-links.pdf?sfvrsn=4&utm_source=ISSCR-Infomz&utm_medium=email&utm_campaign=default). Accessed June 20, 2018.

**65** Kitano K, Schwartz DM, Zhou H et al. Bioengineering of functional human induced pluripotent stem cell-derived intestinal grafts. *Nat Commun* 2017;8:765.