



Universiteit
Leiden
The Netherlands

Who gets caught by the emotion? Attentional biases toward emotional facial expressions and their link to social anxiety and autistic traits

Folz, J.; Roth, T.S.; Nikolić, M.; Kret, M.E.

Citation

Folz, J., Roth, T. S., Nikolić, M., & Kret, M. E. (2023). Who gets caught by the emotion?: Attentional biases toward emotional facial expressions and their link to social anxiety and autistic traits. *Current Psychology*. doi:10.1007/s12144-023-04484-6

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3631035>

Note: To cite this publication please use the final published version (if applicable).



Who gets caught by the emotion? Attentional biases toward emotional facial expressions and their link to social anxiety and autistic traits

Julia Folz^{1,2} · Tom S. Roth¹ · Milica Nikolić^{1,2,3,4} · Mariska E. Kret^{1,2}

Accepted: 28 February 2023
© The Author(s) 2023

Abstract

The emotional facial expressions of other individuals are a valuable information source in adapting behaviour to situational demands, and have been found to receive prioritized attention. Yet, enhanced attentional biases, such as a bias to social threat in Social Anxiety Disorder (SAD), or blunted attention to emotional information, as assumed in Autism Spectrum Disorder (ASD), can easily become maladaptive in daily life. In order to investigate individual differences in attentional biases toward different emotional expressions (angry, happy, sad, and fearful versus neutral) and their links to social anxiety and autistic traits, we tested 104 healthy participants with an emotional dot-probe paradigm on a touch screen, and measured clinical trait levels associated with ASD and SAD. While confirming the presence of attentional biases toward all emotional expressions, we did not find robust evidence for systematic links between these biases and either clinical trait dimension. Only an exploratory Bayesian analysis pointed to a less pronounced bias towards happy facial expressions with higher autistic trait levels. Moreover, a closer examination of the attentional bias towards angry facial expressions suggested that alterations in this bias might depend on a complex interplay between both trait dimensions. Novel approaches in the assessment of attentional biases might yield the potential to describe disorder-specific biases in attention to emotions more validly.

Keywords Emotion · Facial expressions · Attention · Autistic traits · Social anxiety

Living in a world rich in (visual) stimulation, the human perceptual system requires guidance to filter the environment for crucial information. Emotional stimuli have been shown to strongly capture and hold attention in various modalities and impact subsequent behaviour (Carretié, 2014). Already

in the first years of life, infants use their caregivers' emotional expressions to evaluate the current situation, which is commonly referred to as social referencing (e.g., Möller et al., 2014). Further, also in some non-human primates, emotional expressions of conspecifics have been found to receive prioritized attention, resulting in faster responding (e.g., Kret et al., 2016; van Berlo et al., 2020). Thus, the phenomenon of attention being automatically directed to emotional expressions of others seems to be a deeply-rooted, adaptive mechanism in social animals.

Experimental paradigms have confirmed attentional biases toward various emotional facial expressions and body postures in healthy individuals (Bradley et al., 1997; Carlson & Mujica-Parodi, 2015; Valk et al., 2015; Wirth & Wentura, 2020). In individuals with a diagnosis of Autism Spectrum Disorder (ASD) or Social Anxiety Disorder (SAD), alterations in attention to emotional information have been suggested as one potential mechanism underlying social interaction difficulties (Bantin et al., 2016; Kliemann

✉ Julia Folz
j.folz@fsw.leidenuniv.nl

- ¹ Department of Cognitive Psychology, Institute of Psychology, Leiden University, Wassernaarseweg 52, 2333 AK Leiden, The Netherlands
- ² Leiden Institute for Brain and Cognition (LIBC), Leiden University, Leiden, The Netherlands
- ³ Department of Developmental and Educational Psychology, Institute of Psychology, Leiden University, Leiden, The Netherlands
- ⁴ Research Institute of Child Development and Education, University of Amsterdam, Amsterdam, The Netherlands

et al., 2010). However, research on altered attentional biases in these clinical conditions has mainly focused on social threat perception (i.e., displays of anger), whereas humans are exposed to a broad range of emotional displays in daily life. Moreover, different mechanisms have been proposed to underlie altered attention to emotion in the two clinical conditions. With the current study, we therefore aimed to take a step towards the identification of potential systematic alterations in the attention to emotional expressions in ASD and SAD by examining the link between variations in trait levels associated with the two conditions and attentional biases toward different emotional expressions.

Attention to emotional facial expressions and the dot-probe paradigm

In the non-verbal communication of affect, facial expressions are particularly salient and have been shown to effectively capture attention. Most studies so far have focussed on biases to negative stimuli and have identified an “anger superiority effect” (Hansen & Hansen, 1988). This effect describes the rapid and preferential detection of angry faces among others, highlighting the importance of threatening information. Yet, recent studies have shown that also positive stimuli, including smiling faces, attract attention compared to neutral stimuli (Pool et al., 2016). To systematically investigate biased attention to certain stimuli, namely biases to threat stimuli in individuals with a depression or anxiety disorder, the dot-probe task was developed (MacLeod et al., 1986). Since then, it has been used to investigate a variety of biases to a variety of stimuli in different populations (see van Rooijen et al., 2017). While the specific content and parameters differ between studies, they all share the general structure: a trial starts with the presentation of two stimuli for a pre-specified duration on two sides of the screen and equidistant to the centre. After they disappear, a probe appears on one of the two picture location which the participant has to react to. If the target stimulus (e.g., an emotional one) is replaced by the probe, the trial is defined as ‘congruent’. In contrast, in an ‘incongruent’ trial, the probe appears on the location of the control stimulus (e.g., a neutral one). The attentional bias is then usually calculated by looking at a difference value between reaction times during congruent and incongruent trials.

Studies looking at attentional biases to emotional facial expressions using this task have reported attentional biases to both positive (e.g., Wirth & Wentura, 2020) and negative facial emotional expressions (e.g., Bradley et al., 1997; Carlson & Mujica-Parodi, 2015). However, when contrasting different emotion categories in the dot-probe task, some studies only found a bias towards specific emotions

(e.g., Valk et al., 2015) while other studies found no bias in reaction times at all (e.g., Puls & Rothermund, 2018). Methodological differences between studies, for example, in the timing of the stimuli or the stimulus content have been suggested as potential explanations for the mixed evidence (Cooper & Langton, 2006; van Berlo et al., 2020). Importantly, inconsistent findings can also be driven by individual differences (Yiend, 2010). For example, an altered processing of emotional (facial) stimuli, including attentional processes, has been described in various mental health conditions (Kret & Ploeger, 2015).

Attention to emotion in social anxiety

Being characterized by a disproportionate and impairing fear of scrutiny in social situations, Social Anxiety Disorder (SAD) is a mental health condition that already suggests altered attentional allocation to social information by definition (DSM-V; American Psychiatric Association, 2013). The most prominent cognitive-behavioural models on the development and maintenance of SAD all describe a shift in attention to the self as a social object once a social situation is entered or even anticipated (Clark & Wells, 1995; Heimberg et al., 2010, 2014; Hofmann, 2007; Rapee & Heimberg, 1997). This altered attention entails an increased awareness of negatively biased cognitive self-representations and physiological arousal. The *cognitive-behavioral model of anxiety in social phobia* (Rapee & Heimberg, 1997; updates: Heimberg et al., 2010, 2014) additionally posits that attention is tuned to external cues in order to inform mental self-representations. Facial expressions are one example of these cues which can indicate possible negative evaluations by others, which people with SAD fear. This theoretical assumption has been confirmed in various empirical studies showing altered attention to angry facial expressions, representing social threat, in socially anxious individuals (e.g., Lazarov et al., 2021). In line with this, findings from dot-probe studies overall, yet not consistently, report an attentional bias towards threat faces (for a review, see Bantini et al., 2016). Different mechanisms have been suggested to underlie biased attention to threatening stimuli, namely (a) an initial vigilance to threat which; (b) is followed by avoidance after longer exposure (Mogg, Bradley, et al., 2004), as well as; (c) a prolonged attentional capture by threat stimuli, (i.e., difficulty in disengagement; Cisler & Koster, 2010). In the dot-probe task, the difficulty to disengage from threat with higher social anxiety was predominantly found in non-clinical samples (Salemink et al., 2007), whereas a vigilance to threat was prevalent in clinical samples (Klumpp & Amir, 2009).

Importantly, even though different types of emotional expressions appear in a social context and could provide relevant (evaluative) information about other people (Heimberg et al., 2010, 2014; Rapee & Heimberg, 1997), there is only limited research on altered attentional biases toward emotional facial expressions other than anger (Mogg, Philippot et al., 2004). In these few studies, (altered) biases, if present at all, were less pronounced compared to the anger bias. The number and breadth of these studies is, however, too limited to draw any conclusions on biased attention to emotions other than anger in social anxiety. Thus, our study aims to contribute to a better understanding of altered attention to emotional expressions associated with social anxiety by including various emotional facial expressions.

Attention to emotion in autism

Another clinical population that has been found to show alterations in attention to emotional facial expressions are autistic individuals. They tend to attend to faces and, specifically the eye and mouth region, less (Chita-Tegmark, 2016), which might contribute to deficits in identifying emotions (Kliemann et al., 2010). It has been long assumed that the active avoidance of the eye region, conveying emotional information, would be driven by an unpleasant hyperactivation of the amygdala in autistic individuals (Relevance Detection theory; Zalla & Sperduti, 2013). Recently, however, avoidance of the eye region was claimed to be the result of preventing both hypo- and hyperarousal (two-pathway model; Cuve et al., 2018). Attention to facial emotional expressions thus seems to be related to unpleasant arousal levels in autistic individuals. Yet, biased attention towards threatening faces specifically was observed in autistic children and adults (Fan et al., 2020). This led to the claim that the “anger superiority effect” (Hansen & Hansen, 1988) as basic adaptive phenomenon would be unaltered in autistic individuals (Gaigg, 2012). As an alternative explanation, given the high comorbidity between social anxiety and autism (Spain et al., 2018), the question was raised whether the threat bias observed in autistic individuals could be attributed to comorbid social anxiety. Specifically in the social domain, autistic and socially anxious individuals show similar patterns, such as choosing to be alone and avoiding or disliking social situations (White et al., 2012). Apart from one exception (Hollocks et al., 2016), experimental studies, however, have found no evidence for an influence of anxiety symptoms on the threat bias in ASD using the dot-probe task (Hollocks et al., 2013; May et al., 2015). Importantly, as most of the available dot-probe studies examine alterations in ASD, these studies were

performed with a developmental sample (children/teenagers), thus limiting the generalizability of the results.

Similarly, alterations in biases to emotional expressions other than anger have mostly been investigated in autistic children. Here, one study found no evidence of a bias to happy (nor angry) facial expressions (May et al., 2015), whereas another study found no biases toward happy or sad expressions (García-Blanco et al., 2017) in both neurotypical and autistic children. The inconclusive evidence from developmental samples is also reflected in the scarce adult literature on this topic: In one study, only adults with low but not with high autistic trait levels, showed an attentional bias to fearful expressions (Miu et al., 2012). In another study, in contrast, no differences in attentional biases between autistic and non-autistic adults were found. Both groups showed attentional biases to happy and angry faces, but not to sad faces (Monk et al., 2010). Examining all existing evidence together, it is not clear whether attentional biases to specific emotional expressions exist in autistic adults. In the current study, we attempted to diminish this knowledge gap by measuring autistic trait levels and relating them to attentional biases toward various emotional expressions.

Present study

The goal of this study was to examine whether variability in attentional biases toward different emotional facial expressions can be explained by trait levels associated with ASD and SAD in the general population. More specifically, we administered a modified version of the emotional dot-probe task on a touchscreen (see van Berlo et al., 2020) in public settings (community sample) as well as in a lab setting (student sample). To replicate as well as expand current findings on attentional biases to specific emotions, we paired angry, happy, sad, and fearful expressions with neutral expressions in the dot-probe task, and assessed autistic and social anxiety traits via self-reports. The hypotheses, including statistical models to test them, were preregistered on the Open Science Framework after data collection but before accessing the data (see <https://osf.io/8pwgy> for the preregistration, including a more detailed description of the hypotheses). In line with the proposed adaptive function of increased attention to emotions serving as communicative signals (Crivelli & Fridlund, 2018) as well as the results from most dot-probe studies to date (however, see Puls & Rothermund, 2018), we expected to observe an attentional bias toward all emotions. Given the high relevance of angry facial expressions as (social) threat signals, attentional biases might especially be pronounced for expressions of anger (e.g., Valk et al., 2015). We expected that variability in this bias could be explained by social anxiety traits, particularly that higher

social anxiety trait levels would be associated with stronger biases. The possibility of a stronger attentional bias to other emotional expressions in individuals with higher levels of social anxiety traits was additionally explored, including the goal to contrast increased initial vigilance with a difficulty to disengage from emotional expressions. Given the inconclusive literature on the relationship between autism and attentional biases in adults, we hypothesized that, due to an avoidance of the eye region, a less pronounced bias to emotional facial expressions should become apparent with higher autistic trait levels. The bias to angry faces should, however, not be linked to autistic trait levels, as suggested by past research in clinical populations (Fan et al., 2020). Further, the comorbidity between social anxiety and autism has been discussed, yet not confirmed, as being a potential source of a threat bias in individuals with ASD (e.g., May et al., 2015). Therefore, we also aimed to explore whether the link between autistic trait levels and the attentional bias towards angry faces would depend on simultaneously heightened social anxiety trait levels (i.e., a moderation effect) in our healthy sample.

Method

Participants

We tested 104 participants (75 female) with a mean age of 31.4 years ($SD = 14.5$, Range: 17–71) and the majority ($n = 95$) being right-handed. All participants reported to have no prior or current psychological or neural disorder and performed the experiment either in English ($n = 14$) or in Dutch ($n = 90$). Data was collected in three different settings in the Netherlands: the primate park ‘Apenheul’ in Apeldoorn ($n = 30$), the science festival ‘Night of Discoveries’ in Leiden ($n = 22$) and a laboratory at Leiden University ($n = 52$). One participant at the primate park and one participant at the science festival had incomplete task data and were disregarded from the analysis ($N = 102$). Sample characteristics for each location can be found in Table S1 in Online Resource 1. The total sample size was not predetermined as we could not predict the motivation of the primate park/science festival to participate in our study. Yet, the sample size for the laboratory setting was matched to the public settings to balance our sample. All participants provided informed consent prior to participation and there was no monetary reward in either setting but student participants could receive one course credit. The study was conducted in accordance with the Declaration of Helsinki and received approval by the local ethics committee of the Faculty of Social and Behavioural Sciences at Leiden University.


Stimuli and task

To create our facial emotional expression stimuli, six identities (three female) displaying acted angry, happy, sad, fearful and neutral expressions were chosen from the Nim-Stim database (Tottenham et al., 2009). The face (including the neck) of each stimulus was cut out and the remainder replaced by a grey background (RGB: 145, 145, 145), matching the colour of the task background. In order to ensure that observed effects in the dot-probe task were likely to not be caused by systematic differences in low-level features between emotion categories (e.g., de Cesarei & Codispoti, 2013), we employed the Protosc toolbox (Stuit et al., 2021) to unveil potential features significantly predicting category membership. Significant features could not be found for any of the available feature spaces (Fourier magnitudes, Fourier phases, HOGs, colour distributions and pixel intensities). The dot-probe task was programmed in E-Prime® (version 2; Schneider et al., 2002) and conducted on an Iiyama T1931SR B1 touchscreen (38 × 30 cm).

A trial was initiated by tapping on a black dot (\varnothing 6 cm) in the centre of the screen. Immediately afterwards, two pictures appeared (each 11 × 15 cm), one on the left side (7 cm from the left edge) and one on the right side (31 cm from the left edge) of the screen. Apart from the baseline trials in which only neutral stimuli were presented, one picture contained a neutral expression and the other picture contained an emotional expression of the same identity. After 300 ms (according to the recommendation by van Rooijen et al., 2017), one of the pictures was replaced by a black dot (\varnothing 6 cm) which the participant had to tap on as fast as possible. In line with the existing dot-probe terminology, a trial was labeled as congruent when an emotional (i.e., target) expression was replaced by the dot, whereas the replacement of a neutral expression was regarded as an ‘incongruent’ trial. The participant’s reaction time to this second dot was measured in each trial as the variable of interest. After participants successfully reacted to the second dot, the screen turned blank (grey) for 2 s after which the start dot of the next trial appeared (see Fig. 1). For each emotion category (angry, happy, fearful, sad) paired with a neutral stimulus, the dot could appear in the congruent or incongruent location, resulting in eight potential combinations. Each of the combinations was presented using each of the six stimulus identities and with the emotional stimulus in either the left or right location. Further, participants completed 12 trials in which two neutral stimuli were shown after which, due to a coding error, the dot always replaced the left stimulus. Thus, every participant had to complete 108 trials in total ($8 \times 6 \times 2 + 12$).

Fig. 1 Structure of a Trial in the Dot-probe Task

Note. Participants were instructed to press the dot on the touch screen, and the reaction time (RT) to the second dot (probe) was measured. With the probe appearing behind the emotional face, the displayed trial represents a congruent trial. Stimuli were taken from the NimStim set of Facial Expressions (Tottenham et al., 2009).



Procedure

After signing the informed consent form, participants received written instructions on how to perform the dot-probe task on the touchscreen. They then completed eight practice trials with flower pictures as stimuli. Due to the self-initiation of each trial by tapping on the start dot, participants could go through the experiment in a self-paced manner and take breaks whenever necessary. After performing the task (duration around 7 min), participants completed the clinical trait questionnaires and were debriefed.

Questionnaires

Social anxiety traits

We used the self-report version of the Liebowitz Social Anxiety Scale (LSAS; Liebowitz, 1987) to measure social anxiety traits in our healthy sample. The LSAS was originally developed to quantify fear and avoidance in individuals with social phobia and consists of 24 items describing different social situations that typically evoke performance or social anxiety. Fear and avoidance with regard to each item is rated separately on a 4-point Likert scale, ranging from 0 (fear rating: none; avoidance rating: never) to 3 (fear rating: severe; avoidance rating: usually). Sum scores across all items, including both fear and avoidance ratings, are calculated as an overall social anxiety trait measure, with potential scores ranging from 0 to 144. Given that one participant did not complete the questionnaire at all, we had data from 103 participants. Further, for six participants, single items were imputed (see Online Resource 2). Our

final sample's LSAS scores ($N=101$; participants without task data excluded) ranged between 4 and 83 ($M=37.1$, $SD=17.2$). The distribution of the LSAS scores had a skewness of 0.43 and a kurtosis of 2.81, thus being close to normal with a light right skew and mesokurtic shape. Internal consistency of the LSAS in our sample was excellent ($\alpha=0.91$, 95% CI [0.89, 0.94]).

Autistic traits

We used the Autism-Spectrum Quotient (AQ; Baron-Cohen et al., 2001) to measure variations in traits associated with Autism Spectrum Conditions in our sample. The AQ consists of 50 items associated with five different domains in which alterations are typically observed: social skill, attention switching, attention to detail, communication, and imagination. Ten items belong to each domain respectively and build one subscale of the AQ. In this questionnaire, respondents rate the degree to which items apply to them on a 4-point Likert scale, from 1 = *definitely agree* to 4 = *definitely disagree*. Some items are reverse coded and all scores are eventually transformed to binary values (1/2 to 0 and 3/4 to 1). A higher sum score of all items, potentially ranging between 0 and 50, reflects higher autistic trait levels. Three participants did not complete the AQ at all and we imputed missing items for four participants who had incomplete data (see Online Resource 2). The AQ sum scores encompassed values between 2 and 38 ($M=18.3$, $SD=7.6$) in our sample ($N=99$; participants without task data excluded). With a skewness of 0.69 and a kurtosis of 3.33, the AQ score distribution was also close to normal, yet slightly right-skewed

and platykurtic. The AQ in our sample showed a good internal consistency ($\alpha = 0.84$, 95% CI [0.79, 0.88]).

Data analysis

Before fitting the models, reaction times smaller than 250 ms were excluded given that they likely represent random responses (see van Berlo et al., 2020). Further, for each participant, trials exceeding their median reaction time + 2.5 median absolute deviations were excluded to filter out trials in which participants might have been distracted, thus resulting in relatively unusual high RTs. This led to an exclusion of 9.47% of all trials (angry: 8.83%, happy: 9.24%, sad: 10.25%, fearful: 10.02%, neutral: 8.59%). All analyses were performed in R 3.6.3 (R Core Team, 2020; see Online Resource 2 for further information).

Pre-registered data analysis

For all hypotheses, we aimed to fit multiple linear mixed models on reaction times. Looking at the model diagnostics, we did not spot major divergences from assumptions. The selection of the basic fixed and random effect structure for all models was informed by model comparisons based on the first model (see Online Resource 2). Eventually, we included random intercepts for *Subject* and *Trial*. The two three-way interactions *Age*Congruency*Emotion category* and *Sex*Congruency*Emotion category*, as well as all two-way interactions and main effects, were defined as predictors in all models.

To test our first hypothesis, namely that there is an attentional bias to emotional expressions and that this bias is specifically pronounced for angry expressions, we looked at the interaction between *Emotion category* and *Congruency* (controlled for by *Age* and *Sex*) as hypothesis-specific predictor in our model. Both factors, *Emotion category* and *Congruency*, were sum-coded in all analyses. To test our second and third hypotheses, namely that the attentional bias to emotions and specifically to threat expressions is enhanced with higher social anxiety traits, a three-way interaction between *Emotion category*, *Congruency* and *Social anxiety traits* was added as hypothesis-specific predictor to the model (including all two-way interactions).

Since we could not find a link between attentional biases and *Social anxiety traits* for any emotion, we did not explore a potential moderating effect of context in this interaction, as indicated in the preregistration¹. Based on

¹ To confirm that the attentional bias effect was comparable in all experimental settings, we tested for a modulation of the congruency effect by the specific location (primate park, science festival and lab) as well as by the context (public [primate park + science festival], lab). Neither interaction was significant.

the same rationale, we did not conduct the second planned exploratory analysis, which aimed to disentangle whether an alteration in attentional bias would be due to heightened vigilance for emotional expressions or a stronger difficulty to disengage from emotional expressions with elevated social anxiety traits.

In order to test the presence of a general reduction in attentional bias to emotions with higher autistic levels, with the exception of angry expressions (hypotheses 5), we added a three-way interaction between *Emotion category*, *Congruency* and *Autistic traits* as hypothesis-specific predictors to the general model (including all two-way interactions). Given that, there was indeed no link between the attentional bias towards angry expressions (as to all other expressions) and *Autistic traits*, we tested for a potential moderating effect of *Social anxiety traits* on *Autistic traits* in the prediction of the bias to angry expressions. More specifically, we fitted a model on the reaction times in trials with angry expressions with a three-way interaction between *Congruency*, *Social anxiety traits* and *Autistic traits*, including all two-way interactions and the control predictor terms.

Exploratory data analysis

As we did not observe the expected links between attentional biases and clinical trait dimension, and could not exclude that this might be due to a lack of power, we ran additional explorative data analyses, using Bayesian mixed models. Bayesian models were created in the Stan computational framework and accessed using the *brms* package (Bürkner, 2017, Bürkner, 2018), version 2.17.0. We sum coded all factorial predictors, and scaled and centered all continuous predictors. All models were run with 4 chains and 5000 iterations, of which 1000 were warmup iterations. We checked model convergence by inspecting the trace plots, histograms of the posteriors, Gelman-Rubin diagnostics, and autocorrelation plots (Depaoli & van de Schoot, 2017). We found no divergences or excessive autocorrelation.

For the exploratory analyses, we used the same dataset as for the pre-registered analyses in which extremely fast and slow reaction times were excluded by subject (see first paragraph of [Data analysis](#) section). However, for the exploratory analyses we rescaled our dependent variable in order to filter out the effect of handedness*probe location (*Probe distance*) and to ease setting a prior for the intercept. Thus, we centered the reaction times within *Subject* within *Probe distance* level (close vs. far). Thereby, we removed the distance effect and removed overall differences in reaction times between participants.

First, we explored attentional biases within each emotion category by creating a model with centered reaction time as dependent variable and *Congruency* and *Emotion Category*

and their interaction as predictors. Furthermore, we allowed the effects of all predictors to vary by *Subject*. Second, we explored whether attentional biases within each emotion category were linked to *Autistic traits* and *Social anxiety traits* by including the interactions *Congruency*Emotion Category*Autistic traits* and *Congruency*Emotion Category*Social anxiety traits*. We used regularizing Gaussian priors with $M=0$ and $SD=5$ for all fixed effects, a Gaussian prior with $M=0$ and $SD=1$ for the intercept, and default half Student t priors with 3 degrees of freedom for the random effects and residual standard deviation. We used multiple measures to summarize the posterior distribution resulting from our models: (I) the median estimate and the median absolute deviation of this estimate, (II) the 95% credible interval, and (III) the probability of direction (pd). The 95% CrI indicates the range within which the effect falls with 95% probability, while the pd indicates the proportion of the posterior distribution that is of the median's sign (Makowski et al., 2019).

Results

Pre-registered analyses

General attentional bias to emotions

As expected, we found a significant congruency effect in the general model on the reaction times in the dot-probe task, $B = -5.79$, 95% CI [-9.80, -1.78], $t(8697.27) = -2.83$, $p = .005$ (see Fig. 2A). Confirming an attentional bias to emotional

expressions, participants were on average 5.79 ms faster when the dot appeared behind the emotional expression and 5.79 ms slower when the dot appeared behind the neutral expression than their average reaction times in trial with an emotion-neutral pair. There was, however, no evidence for an enhanced attentional bias to angry compared to other facial expressions, that is, no significant interaction between *Emotion category* and *Congruency*. In addition to the general congruency effect, we also observed significant effects of the included control predictor terms. Unsurprisingly, reaction times depended on *Age*, $B = 2.94$, 95% CI [1.78, 4.10], $t(98.96) = 5.04$, $p < .001$. While older participants overall reacted more slowly to the dot probe than younger participants, this effect was less marked in trials including a happy face, $B = -0.27$, 95% CI [-0.47, -0.07], $t(8708.16) = -2.70$, $p = .007$, as revealed by the significant interaction between *Age* and *Emotion category*. Lastly, *Probe distance* significantly predicted reaction times, with slower responses to far probes (i.e., a mismatch between handedness and probe location) compared to close probes (i.e., handedness and probe location matched), $B = 30.94$, 95% CI [27.62, 34.25], $t(8703.67) = 18.30$, $p < .001$. For an overview of the model fit, see Table 1 as well as Table S3 in Online Resource 1 for the coefficients of all factor levels.

Social anxiety traits and the attentional bias to emotions

Against our hypothesis, we did not find any alteration in the overall or emotion-specific attentional bias related to social anxiety trait levels. Further, the congruency effect which we observed in the general model did not reach significance in




Fig. 2 (A) Average Reaction Times per Subject to the Probe in Congruent versus Incongruent Trials for each Emotion Category (B) Mean Predicted Attentional Bias (Difference in Reaction Times between Congruent and Incongruent Trials) per Subject across Emotion Categories and Trials

Note. (A) Boxes enclose all values between the first and third quartile (Inter-quartile range, IQR) and the whiskers extend to +/- 1.5 IQR

from the respective quartile. As outliers in reaction times were defined on an individual basis, outliers in this plot represent subjects with slow reaction times on average (see also Table S2 in the Supplementary Materials). (B) The data distribution is visualized by the violin as well as by the box (same definition as above), and single data points represent subjects. This graph shows the significant congruency effect, i.e., the general bias towards emotional facial expressions.

Table 1 Results of the Linear Mixed-Effects Models Predicting Reaction Times to the Dot Probe. Output for the Fixed and Random Effect Terms for the Model without Clinical Traits and its Interactions as Predictor, as well as for the Models with Social Anxiety Traits and Autistic Traits, Respectively, are Shown

	General model			Social anxiety traits model			Autistic traits model					
	df1	df2	F	p	df1	df2	F	p	df1	df2	F	p
Fixed effects												
Emotion category	3	8705.6	1.44	0.230	3	8620.7	1.50	0.212	3	8441.7	1.01	0.389
Congruency	1	8697.3	8.03	0.005	1	8624.9	1.92	0.166	1	8441.7	6.91	0.009
Age	1	99.0	25.41	< 0.001	1	97.0	25.47	< 0.001	1	95.0	24.69	< 0.001
Gender	1	99.0	0.16	0.687	1	97.0	0.03	0.867	1	95.0	0.21	0.645
Probe distance	1	8703.7	334.82	< 0.001	1	8613.2	331.29	< 0.001	1	8439.7	320.41	< 0.001
Clinical traits												
Emotion category*Congruency	3	8696.1	0.96	0.412	1	97.0	0.56	0.456	1	95.0	0.19	0.665
Emotion category*Age	3	8711.9	2.68	0.045	3	8607.1	0.46	0.707	3	8435.2	0.75	0.524
Congruency*Age	1	8704.0	0.91	0.339	1	8617.6	0.74	0.030	3	8443.0	3.66	0.012
Emotion category*Gender	3	8708.1	0.36	0.780	3	8617.6	0.36	0.388	1	8438.8	0.55	0.458
Congruency*Gender	1	8712.0	0.97	0.325	1	8618.9	1.37	0.781	3	8440.7	0.51	0.674
Emotion category*Clinical traits												
Congruency*Clinical traits					3	8624.8	0.86	0.462	3	8445.5	3.92	0.008
Emotion category*Congruency*Age	3	8693.2	1.17	0.321	1	8624.7	0.31	0.578	1	8454.6	1.04	0.308
Emotion category*Congruency*Gender	3	8704.0	0.40	0.754	3	8601.1	1.02	0.382	3	8427.5	0.70	0.551
Emotion category*Congruency*Clinical traits					3	8611.4	0.49	0.689	3	8437.1	0.33	0.801
Random effects			Variance				Variance				Variance	
Intercepts	Subject N = 102		7045.49		Subject N = 101		7112.70		Subject N = 99		7161.27	
	Trial N = 108		52.37		Trial N = 108		49.74		Trial N = 108		57.10	
Residual variance			6267.42				6221.84				6016.66	

Note. df1 = numerator degrees of freedom; df2 = denominator degrees of freedom.

the model including *Social anxiety traits* ($p = .166$). Yet, the effect of *Age* on reaction times to the dot probe was again significant, $B = 3.01$, 95% CI [1.83, 4.20], $t(96.96) = 5.05$, $p < .001$, as well as the interaction between *Age* and *Emotion category*. The slowing in reaction time with higher age was less pronounced if a happy facial expression (compared to the other emotions) was included in the emotion-neutral pair, independent of the probe position, $B = -0.28$, 95% CI [-0.48, -0.08], $t(8617.1) = -2.78$, $p = .005$. As observed in the general model, a far *Probe distance* was associated with slower reaction times, $B = 30.81$, 95% CI [27.50, 34.13], $t(8613.15) = 18.20$, $p < .001$. Table 2 provides an overview of the model fit and Table S4 in Online Resource 1 describes all coefficients.

Autistic traits and the attentional bias to emotions

In the attentional bias model including *Autistic traits*, there was a significant interaction between *Autistic traits* and *Emotion category*. More specifically, responses after emotion-neutral pairs with an angry facial expression were faster with higher autistic traits, $B = -0.62$, 95% CI [-1.00, -0.24], $t(8444.09) = -3.22$, $p = .001$. In contrast, emotion-neutral pairs with sad facial expressions were related to slower responses with higher autistic traits, $B = 0.39$, 95% CI [0.01, 0.78], $t(8443.17) = 2.03$, $p = .042$. Importantly, these effects were independent of the probe location (i.e., *Congruency*) and we did not observe the expected link between the attentional bias and *Autistic traits*. Further, a closer examination of the effects revealed that none of the slopes were significantly different from zero. In line with the general model

Table 2 Results of the Linear Mixed-Effects Models Predicting Reaction Times to the Dot Probe in Trials with Angry Facial Expressions

Fixed effects	df1	df2	F	p
Congruency	1	2056.91	4.20	0.041
Age	1	92.90	28.44	< 0.001
Sex	1	92.81	0.06	0.813
Probe distance	1	2056.97	113.60	< 0.001
Autistic traits	1	92.89	0.01	0.921
Social anxiety traits	1	93.07	1.44	0.233
Congruency*Age	1	2056.57	0.50	0.479
Congruency*Sex	1	2056.53	0.09	0.768
Congruency*Autistic traits	1	2056.69	3.86	0.050
Congruency*Societal anxiety traits	1	2056.71	7.51	0.006
Autistic traits*Societal anxiety traits	1	93.09	0.63	0.428
Congruency*Autistic traits*Societal anxiety traits	1	2056.63	6.37	0.012
Random effects			Variance	
Intercepts	Subject		7204.31	
	N=99			
Residual variance			6362.83	

Note. Df1 = numerator degrees of freedom; df2 = denominator degrees of freedom.

on the attentional bias to emotions, *Congruency*, $B = -7.48$, 95% CI [-13.06, -1.90], $t(8439.91) = -2.63$, $p = .009$, as well as the control terms *Probe distance*, $B = 30.11$, 95% CI [26.81, 33.41], $t(8439.65) = 17.90$, $p < .001$, and *Age*, $B = 3.12$, 95% CI [1.87, 4.37], $t(94.96) = 4.97$, $p < .001$, were significant predictors of reaction times in the model including *Autistic traits*. Participants reacted faster in congruent trials, as well as when the probe appeared on the side of their dominant hand and when they were younger. We also found a significant interaction between *Emotion category* and *Age*. Here, the slowing in reaction times with higher age was more pronounced in trials with angry facial expressions, $B = 0.24$, 95% CI [0.03, 0.44], $t(8446.75) = 2.23$, $p = .026$, and less pronounced in trials with happy facial expressions, $B = -0.32$, 95% CI [-0.52, -0.11], $t(8432.03) = -3.01$, $p = .003$. A summary of the model fit can be found in Table 2 and a closer description of all coefficients in Table S5 in Online Resource 1.

Autistic traits, social anxiety traits and the attentional bias to angry facial expressions

In line with previous research, the two questionnaire scores, indicating autistic trait and social anxiety traits, were significantly positively correlated in our sample, $r_s = 0.30$, $p < .001$. Zooming in on a potential moderating effect of *Social anxiety traits* on the link between *Autistic traits* and the attentional bias to angry facial expressions, our model revealed a significant three-way interaction between *Congruency*, *Social anxiety traits* and *Autistic traits*, $B = 0.03$, 95% CI [0.01, 0.06], $t(2056.63) = 2.52$, $p = .012$. An examination of the predicted value plots (see Fig. 3) as well as slope comparisons (see Table 3) at three different values on one of the trait dimensions (mean - 1SD, mean, mean + 1SD) suggested that this interaction is likely to be driven by a decrease in attentional bias with higher autistic traits at a relatively "high" social anxiety trait level (mean + 1SD) and/or an increase in attentional bias with higher social anxiety traits at a relatively "low" autistic trait level (mean - 1SD). In the context of this three-way interaction, the two-way interactions between each trait dimension and *Congruency* were also approaching significance (see Table 2). However, when running the same model without the three-way interaction added, this was not the case. Similar to the previous models with multiple emotion categories, *Congruency*, $B = 23.36$, 95% CI [1.01, 45.72], $t(2056.92) = 2.05$, $p = .041$, as well as the control predictors *Age*, $B = 3.48$, 95% CI [2.19, 4.78], $t(92.90) = 5.33$, $p < .001$, and *Probe distance*, $B = 36.69$, 95% CI [29.94, 43.44], $t(2056.97) = 10.66$, $p < .001$, were significant predictors in this model (see Table 2 for an overview of the model fit as well as Table S6 in Online Resource 1 for all coefficients).

Fig. 3 Effects of Levels on one Trait Dimension on Reaction Times at Distinct Levels on the Other Trait Dimension in Congruent (Red) vs. Incongruent (Blue) Trials

Note. Effects are displayed at the mean – 1SD, mean and mean + 1SD of the other trait dimension respectively.





Table 3 Difference in the Slopes Between Congruent and Incongruent Trials (Congruency Effect) with Higher Levels on one Clinical Trait Dimension at the Trait Scores Mean – 1SD, Mean and Mean + 1SD on the Respective Other Clinical Trait Dimension

	Slope difference : congruent – incongruent [SE]	95% CI	df	t	p-value
<i>Autistic traits</i>					
Mean LSAS – 1SD [19.96]	-0.82 [0.67]	-2.13, 0.48	2057	-1.24	0.216
Mean LSAS [37.26]	0.32 [0.49]	-0.65, 1.28	2056	0.64	0.520
Mean LSAS + 1SD [54.57]	1.46 [0.67]	0.15, 2.77	2056	2.18	0.030
<i>Social anxiety traits</i>					
Mean AQ – 1SD [10.64]	-0.87 [0.34]	-1.53, -0.21	2057	-2.58	0.010
Mean AQ [18.18]	-0.37 [0.23]	-0.83, 0.09	2057	-1.59	0.112
Mean AQ + 1SD [25.73]	0.13 [0.27]	-0.40, 0.66	2057	0.47	0.638

Fig. 4 Effect of the Interactions between Congruency and Social Anxiety Traits (LSAS) and Congruency and Autistic Traits (AQ) on Reaction Times
 Note. Shaded areas reflect 95% credible intervals.



Exploratory analyses

First, we explored whether participants showed attentional biases within each emotion category using a Bayesian mixed model (Table S7 in Online Resource 1). We found clear effects of *Congruency* within each level of *Emotion category* (Figure S1 in Online Resource 1), with participants responding faster on trials where the probe replaced the emotional stimulus versus neutral stimulus (**Angry**: -9.21 ms [3.39], 95% CrI [-15.86, -2.31], *pd*=1.00; **Happy**: -13.30 ms [3.27], 95% CrI [-19.59, -6.98], *pd*=1.00; **Sad**: -6.37 ms [3.13], 95% CrI [-12.57, -0.13], *pd*=0.98; **Fearful**: -6.65 ms [3.38], 95% CrI [-13.34, 0.02], *pd*=0.98). Note that the 95% CrI for Fearful spans just over zero. However, the directionality of the effect (*pd*) was clear and consistent with the other categories.

Next, we explored whether *Autistic traits* and *Social anxiety traits* moderated the effect of *Congruency* within each level of *Emotion Category* (Table S8 in Online Resource 1). We compared the effect of *Congruency* at -1SD with +1SD of the scaled *Autistic traits* and *Social anxiety traits* variables. With regard to *Autistic traits*, we did not find robust evidence that autistic trait levels moderated the effect of

Table 4 Difference in the Congruency Effect at Between -1SD and +1SD of the AQ Scale (Autistic Traits) and LSAS Scale (Social Anxiety Traits) for each Emotion Category

	Median [MAD]	95% CrI	<i>pd</i>
<i>Autistic traits (AQ)</i>			
Angry	4.57 [7.32]	-9.71, 19.41	0.74
Happy	10.12 [6.74]	-3.00, 23.61	0.94
Sad	5.28 [6.93]	-8.13, 18.71	0.78
Fearful	0.28 [7.18]	-14.03, 14.47	0.52
<i>Social anxiety traits (LSAS)</i>			
Angry	-6.49 [7.19]	-20.80, 8.01	0.81
Happy	-7.58 [6.87]	-20.97, 5.71	0.87
Sad	-6.40 [6.78]	-19.82, 7.14	0.83
Fearful	7.27 [7.28]	-7.15, 21.26	0.84

Congruency within any level of *Emotion category* (Fig. 4A; Table 4). However, we observed that participants with low autistic trait levels had a stronger attentional bias for happy faces than participants with high autistic trait levels. Although the 95% CrI spanned over 0, the directionality of the effect was clear (*pd* = 0.94). Looking at *Social anxiety traits*, we did not find robust evidence for an effect on *Congruency* (Fig. 4B; Table 4). Even though the results indicated that people who scored higher on the social anxiety

trait scale had stronger attention biases for sad, happy and angry expressions, the 95% CrIs spanned over 0, and the directionalities were relatively low (Table 4).

Discussion

In the current study, we examined attentional biases toward facial emotional expressions and their association with autistic traits and social anxiety in a general population sample. In line with most previous research, we observed attentional biases toward various emotional facial expressions, using both frequentist as well as Bayesian analysis approaches. In contrast to our hypothesis and a vast amount of literature, higher social anxiety trait levels were overall not associated with a stronger bias to angry facial expressions (i.e., social threat). There was also only an indication of a decreased attention to emotional expressions with higher autistic trait levels for happy facial expressions within our exploratory Bayesian analysis. Yet, independent of the probe location (congruent vs. incongruent), reaction times were faster with higher autistic trait levels for trials displaying an angry face, while they were slower for trials displaying a sad face. When zooming in on the attentional bias towards angry facial expressions, we found a significant interaction between autistic trait levels, social anxiety trait levels and congruency. This suggests that the link between the attentional bias to threat and the two trait dimensions might be more complex and require further exploration.

Social anxiety and the attentional bias to emotion

Surprisingly, an enhanced attentional bias to angry facial expression was not observed in people with higher levels of trait social anxiety. Possibly, the absence of this effect can be the result of the stimulus presentation duration that we employed in this study. According to a meta-analysis examining the link between social anxiety and the threat bias (Bantin et al., 2016), shorter stimulus durations (<200 ms) were associated with stronger biases. Nevertheless, attentional biases toward angry facial expressions could still be found at 500 ms and 600 ms presentation duration. Further, higher levels of social anxiety have not consistently been linked to stronger attentional biases toward angry facial expressions (Bantin et al., 2016). A recent study in a healthy student sample reported that the half of individuals with higher social anxiety trait levels were found to have the lowest attentional bias whereas the other half was partially showing vigilance toward and partially avoidance of angry faces (Neophytou & Panayiotou, 2022). Crucially, even

biases on the individual level were shown to be invariant, meaning that they change over time, and to depend on the assessment tool (MacLeod et al., 2019). The present study was the first to use a touchscreen to examine the association between social anxiety and the threat bias. Given the instability of attentional biases as well as, an interplay between various factors could explain not finding the expected effect.

When comparing the effect of social anxiety traits on the threat bias at different autistic trait levels (three-way interaction), the attentional bias only seemed to be stronger with higher social anxiety traits at low autistic trait levels. This suggests that attentional biases in social anxiety might highly depend on additional individual characteristics. As a consequence, interventions focusing on treating maladaptive attentional biases, such as Attentional Bias Modification (ABM) training (MacLeod et al., 2002), might not be beneficial for every individual. Accordingly, ABM trainings have been reported to neither consistently nor robustly result in a modification of the attentional bias toward threat (van Bockstaele & Bögels, 2014). Further, one study which found a small reduction in the bias in the visual dot-probe after ABM training, also showed that this bias is not generalizable to other tasks measuring attention to threat (van Bockstaele et al., 2017). Acknowledging the specificity of this effect, the dot-probe paradigm might not be the ideal candidate on which to directly base clinical interventions. Instead, it can be regarded as a useful additional descriptor in the complex relationship between social anxiety and the attention to emotional facial expressions.

Autistic traits and the attentional bias to emotion

We expected to observe a weaker attentional bias to all emotion facial expressions, apart from anger, with higher autistic trait levels due to a decreased processing of emotional information from the faces. In contrast to our expectations, there was only an indication of this effect for happy facial expressions in our exploratory analysis. Past research in autistic children (García-Blanco et al., 2017; May et al., 2015) as well as autistic adults (Monk et al., 2010) has found no evidence for alterations of the attentional bias to happy facial expressions. Next to the essential difference of examining a clinical population, these studies also used longer presentation times (i.e., 500 ms and/or 1500 ms) which could have allowed for a more elaborate (and less automatic) processing of the stimuli. As an alternative to arousal-related explanations (e.g., Cuve et al., 2018; Zalla & Sperduti, 2013), alterations in face processing in autistic individuals were suggested to result from “deficits” in processing social rewards, such as faces (Dawson et al.,

2005). Studies displaying happy facial expressions directed towards the participant, as used in our study, indeed suggest that those faces are associated with lower reward values in autistic individuals compared to neurotypical individuals (Dubey et al., 2015). A recent meta-analysis, however, has challenged the idea of altered social reward processing in ASD by unveiling less reward processing for both social and non-social stimuli (Bottini, 2018). In our study, we did not investigate mechanisms which could underlie a weaker attentional bias toward happy facial expressions with higher autistic trait levels. To get a better understanding of altered face perception in ASD, future studies should not only examine whether a weaker attentional bias towards happy facial expressions is present in autistic individuals, but also what the underlying mechanism of this alteration might be.

Unexpectedly, reaction times to the dot probe in trials with angry or sad facial expressions were systematically linked to autistic traits, independent of whether the probe appeared behind the emotional or the neutral expression (i.e., congruency). Higher autistic trait levels were associated with relatively faster reaction times after the presentation of an angry face and with slower reaction times after the presentation of a sad face. A potential explanation of this finding could be that the mere presence of the expressions affected the observer's arousal more strongly with higher autistic trait levels. More specifically, independent of their location, angry expressions could have elicited increases in arousal and sad expressions decreases in arousal, which would result in faster and slower reaction times, respectively. Yet, differences in reaction times between emotion categories in the general model could not be found. Further, the slopes for the links between autistic traits and reaction times in trials with angry and sad facial expressions were not significantly different from zero. Thus, the reaction times for angry and sad expressions were only different compared to the average link between reaction times to all expressions and autistic traits.

Finally, we observed a significant interaction between social anxiety traits and autistic traits in predicting the attentional bias to angry faces. Individuals with higher autistic traits showed a reduced attentional bias towards angry faces, but only when social anxiety trait levels were also high. This was in contrast to our expectations. We assumed that high social anxiety levels would go along with a stronger attentional bias to angry faces in individuals with higher autistic trait levels, as angry expression might be perceived as more threatening. Looking at the predicted value plots, the reduced attentional bias seemed to be driven by faster reaction times to the probe in incongruent trials (i.e., probe replaces neutral faces) with higher autistic traits. This potentially supports the idea of a generally heightened arousal for angry expressions. Nevertheless, given the complexity of

explaining this three-way interaction, the high overlap of confidence intervals and the lacking support of previous literature in clinical populations (Hollocks et al., 2013; May et al., 2015; Monk et al., 2010), future research should examine the interplay between autistic and social anxiety traits on attentional biases to emotion.

Limitations & future directions

While some previous research did not report an attentional bias toward (certain) emotional facial expressions (Puls & Rothermund, 2018; Valk et al., 2015), we found an attentional bias toward emotions for all emotion categories. This bias did not differ significantly between the expressions and ranged around 10 ms (6.37 ms – 13.30 ms) which is comparable with previous research (e.g., Monk et al., 2010). Thus, emotional facial expressions seem to automatically receive prioritised attention, highlighting their suggested communicative function as salient signals for conspecifics. Yet, caution has to be taken in the interpretation of results on attentional biases, as their appearance/significance might depend on various additional factors. One important factor that has been raised in the debate on the validity of attentional biases is the impact of low-level features (e.g., de Cesarei & Codispoti, 2013). Differences between stimulus categories regarding features such as spatial frequency are a general problem in interpreting results from dot-probe studies. We did not control our stimuli for these features to keep them as natural as possible and therefore cannot exclude this as a potential influence. Nevertheless, we compared our emotion categories with regard to specific low-level features (Stuit et al., 2021, see Methods section) and could not identify any significant differences. While there were no systematic differences between emotion categories, this comparison does not rule out the existence of differences in, for example, spatial frequency or colour distribution between distinct stimuli, which might have added noise to the data. Future studies investigating attentional processes in the field of emotion should try to account for those.

Another limitation of our study is that we did not collect data in clinical populations and could, therefore, only investigate the influence of trait levels associated with ASD and SAD on attentional biases. This complicates the direct comparison to previous research with clinical samples as well as the formulations of potential implications for clinical practice. Yet, the view of a “continuum of impairment” has become more popular regarding both social anxiety (Rapee & Spence, 2004) and autism (Robinson et al., 2011), with disorders lying on the extremes of clinical traits in the general population. While clinical relevance of symptoms is eventually determined by difficulties in daily life,

alterations in information processing, as well as their underlying mechanisms, may not be qualitatively different along the trait dimension. Thus, our findings in heightened trait levels may also be informative for clinical populations.

As highlighted earlier, the link between social anxiety trait levels and attentional biases to social threat seems to depend on additional individual characteristics, such as autistic traits. This indicates that the extent to which different individuals with SAD shift their attention to external evaluative cues in social situations might also vary. Socially anxious individuals who automatically shift their attention to expressions of others might indeed benefit from trainings which aim at modifying this automatic shift to prevent the perception of social threat. In socially anxious individuals who do not show disproportionate attentional biases toward external cues, other factors might contribute more strongly to the maintenance of the disorder. While this still has to be investigated further, our results generally favour a more individualized approach, which targets specific maintenance factors in the treatment of SAD (for suggestions, see Hofmann, 2007). Since we did not find evidence for a reduced attentional bias toward emotional expressions with higher autistic trait levels, difficulties in identifying others' emotions might likely not arise from altered allocation of early visual attention (300 ms in our study) toward those. Other factors might play a more important role, such as altered physiological arousal in the presence of emotional expressions, which could also explain the earlier-mentioned effects on reaction times in our study. Future studies should specifically explore these factors, as well as their link to real-life social outcomes, to inform clinical practice.

Finally, ways of capturing attentional biases more validly have recently been suggested, such as investigating trial-level attentional biases (Zvielli et al., 2015) or using response-based measures (Evans & Britton, 2018). The use of eye-tracking as an alternative technique in examining attention towards emotional versus neutral stimuli has further been encouraged to unveil individual differences in attentional processes at different stages of information processing (Clauss et al., 2022).

Conclusion

With the current study, we aimed to unveil specific links between variations in the attentional bias to emotional facial expressions and social anxiety and autistic trait levels. While an attentional bias towards all emotional facial expressions, namely angry, happy, sad and fearful, was found in our study, there was only weak evidence for systematic links between these biases and clinical traits. More specifically, our exploratory analyses suggested that only the attentional

bias to happy facial expressions was decreased with higher autistic trait levels. We did, however, additionally observe general alterations in reaction times after the presentation of certain emotional stimuli (angry, sad) with higher autistic trait levels, and the bias to angry facial expressions seemed to depend on a combination of both autistic traits and social anxiety traits in our study. Taken together, the link between clinical traits and attention, as measured by reaction times in this study, appears to be highly complex. While the dot-probe task allows to tap into general attentional tendencies toward emotional expressions, more naturalistic scenarios might be of higher informative value for revealing biases in real-life attentional processing associated with clinical conditions and build a stronger basis for clinical support.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12144-023-04484-6>

Acknowledgements We thank the primate park “Apenheul” and the “Night of Discoveries” organizers to provide us with the opportunity to collect data, all students and lab members who were involved in data collection, as well as all participants for their voluntary participation. We further thank the Dutch National Science Foundation (Nederlandse Organisatie voor Wetenschappelijk Onderzoek, NWO) for funding this research [grant number 016.VIDI.185.036].

Author Contribution **Julia Folz:** Conceptualization, Methodology, Software, Formal analysis, Data curation, Writing – Original Draft, Visualization, Project administration. **Tom S. Roth:** Formal analysis, Writing – Original Draft, Writing – review & editing. **Milica Nikolić:** Methodology, Writing – review & editing, Supervision. **Mariska E. Kret:** Conceptualization, Methodology, Resources, Writing - review & editing, Supervision, Funding acquisition.

Funding This research was funded by the Dutch National Science Foundation (Nederlandse Organisatie voor Wetenschappelijk Onderzoek, NWO) [grant number 016.VIDI.185.036].

Data Availability Following the university policy, all data from this project will be made available on the DataverseNL repository upon paper publication (<https://doi.org/10.34894/UVQHHD>).

Code Availability Code generated within this project will be made available together with the data, on the DataverseNL repository upon paper publication (<https://doi.org/10.34894/UVQHHD>).

Declarations

Competing interest The authors have no competing interests to declare that are relevant to the content of this article.

Conflict of Interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Ethical approval The study was reviewed and approved by the Psychology Ethics Committee of Leiden University (Apenheul: 2019-08-02-M.E.Kret-V1-1831; Night of Discoveries and lab setting: 2019-09-03-M.E.Kret-V1-1856).

Informed consent Written consent was obtained from all participants.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). <https://doi.org/10.1176/appi.books.9780890425596>
- Bantin, T., Stevens, S., Gerlach, A. L., & Hermann, C. (2016). What does the facial dot-probe task tell us about attentional processes in social anxiety? A systematic review. *Journal of Behavior Therapy and Experimental Psychiatry*, 50, 40–51. <https://doi.org/10.1016/j.jbtep.2015.04.009>
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning autism, males and females, scientists and Mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5–17. <https://doi.org/10.1023/A:1005653411471>
- Bottini, S. (2018). Social reward processing in individuals with autism spectrum disorder: A systematic review of the social motivation hypothesis. *Research in Autism Spectrum Disorders*, 45, 9–26. <https://doi.org/10.1016/j.rasd.2017.10.001>
- Bradley, B. P., Mogg, K., Millar, N., Bonham-Carter, C., Fergusson, E., Jenkins, J., & Parr, M. (1997). Attentional biases for emotional faces. *Cognition and Emotion*, 11(1), 25–42. <https://doi.org/10.1080/026999397380014>
- Bürkner, P. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, 80(1), 1–28. <https://doi.org/10.18637/jss.v080.i01>
- Bürkner P (2018). Advanced Bayesian Multilevel Modeling with the R Package brms. *The R Journal*, 10(1), 395–411. <https://doi.org/10.32614/RJ-2018-017>
- Carlson, J. M., & Mujica-Parodi, L. R. (2015). Facilitated attentional orienting and delayed disengagement to conscious and non-conscious fearful faces. *Journal of Nonverbal Behavior*, 39(1), 69–77. <https://doi.org/10.1007/s10919-014-0185-1>
- Carretié, L. (2014). Exogenous (automatic) attention to emotional stimuli: a review. In *Cognitive, Affective and Behavioral Neuroscience* (Vol. 14, Issue 4). <https://doi.org/10.3758/s13415-014-0270-2>
- Chita-Tegmark, M. (2016). Attention allocation in ASD: A review and Meta-analysis of Eye-Tracking Studies. *Review Journal of Autism and Developmental Disorders*, 3(3), 209–223. <https://doi.org/10.1007/s40489-016-0077-x>
- Cisler, J. M., & Koster, E. H. W. (2010). Mechanisms of attentional biases towards threat in anxiety disorders: An integrative review. *Clinical Psychology Review*, 30(2), 203–216. <https://doi.org/10.1016/j.cpr.2009.11.003>
- Clark, D. M., & Wells, A. (1995). A cognitive model of social phobia. *Social phobia: Diagnosis, assessment, and treatment* (pp. 69–93). The Guilford Press.
- Clauss, K., Gorday, J. Y., & Bardeen, J. R. (2022). Eye tracking evidence of threat-related attentional bias in anxiety- and fear-related disorders: A systematic review and meta-analysis. In *Clinical Psychology Review* (Vol. 93). Elsevier Inc. <https://doi.org/10.1016/j.cpr.2022.102142>
- Cooper, R. M., & Langton, S. R. H. (2006). Attentional bias to angry faces using the dot-probe task? It depends when you look for it. *Behaviour Research and Therapy*, 44(9), 1321–1329. <https://doi.org/10.1016/j.brat.2005.10.004>
- Crivelli, C., & Fridlund, A. J. (2018). Facial displays are tools for Social Influence. *Trends in Cognitive Sciences*, 22(5), 388–399. <https://doi.org/10.1016/j.tics.2018.02.006>
- Cuve, H. C., Gao, Y., & Fuse, A. (2018). Is it avoidance or hypoarousal? A systematic review of emotion recognition, eye-tracking, and psychophysiological studies in young adults with autism spectrum conditions. *Research in Autism Spectrum Disorders*, 55(August), 1–13. <https://doi.org/10.1016/j.rasd.2018.07.002>
- Dawson, G., Webb, S. J., & McPartland, J. (2005). Understanding the nature of face processing impairment in autism: Insights from behavioral and electrophysiological studies. In *Developmental Neuropsychology*, 27(3), 403–424. https://doi.org/10.1207/s15326942dn2703_6
- de Cesarei, A., & Codispoti, M. (2013). Spatial frequencies and emotional perception. *Reviews in the Neurosciences*, 24(1), 89–104. <https://doi.org/10.1515/revneuro-2012-0053>
- Depaoli, S., & Van de Schoot, R. (2017). Improving transparency and replication in Bayesian statistics: The WAMBS-Checklist. *Psychological methods*, 22(2), 240. <https://doi.org/10.1037/met0000065>
- de Valk, J. M., Wijnen, J. G., & Kret, M. E. (2015). Anger fosters action. Fast responses in a motor task involving approach movements toward angry faces and bodies. *Frontiers in Psychology*, 6(September), 1–7. <https://doi.org/10.3389/fpsyg.2015.01240>
- Dubey, I., Ropar, D., & de Hamilton, C., A. F (2015). Measuring the value of social engagement in adults with and without autism. *Molecular Autism*, 6(1), <https://doi.org/10.1186/s13229-015-0031-2>
- Evans, T. C., & Britton, J. C. (2018). Improving the psychometric properties of dot-probe attention measures using response-based computation. *Journal of Behavior Therapy and Experimental Psychiatry*, 60(January), 95–103. <https://doi.org/10.1016/j.jbtep.2018.01.009>
- Fan, X. Z., Duan, Y. W., Yi, L. X., & He, H. Z. (2020). Attentional bias toward threatening emotional faces in individuals with autism spectrum disorder: A meta-analysis on reaction time tasks. *Research in Autism Spectrum Disorders*, 78(September), 101646. <https://doi.org/10.1016/j.rasd.2020.101646>
- Gaigg, S. B. (2012). The interplay between emotion and cognition in Autism Spectrum Disorder: Implications for developmental theory. *Frontiers in Integrative Neuroscience*, 6(NOV), 1–35. <https://doi.org/10.3389/fnint.2012.00113>
- García-Blanco, A., López-Soler, C., Vento, M., García-Blanco, M. C., Gago, B., & Perea, M. (2017). Communication deficits and avoidance of angry faces in children with autism spectrum disorder. *Research in Developmental Disabilities*, 62, 218–226. <https://doi.org/10.1016/j.ridd.2017.02.002>
- Hansen, C. H., & Hansen, R. D. (1988). Finding the Face in the crowd: An anger superiority effect. *Journal of Personality and Social Psychology*, 54(6), 917–924. <https://doi.org/10.1037/0022-3514.54.6.917>
- Heimberg, R. G., Brozovich, F. A., & Rapee, R. M. (2010). A Cognitive Behavioral Model of Social Anxiety Disorder: Update and Extension. In S. G. Hofmann & P. M. DiBartolo (Eds.), *Social Anxiety (Second Edition)* (Second Edition, pp. 395–422). Academic Press. <https://doi.org/10.1016/B978-0-12-375096-9.00015-8>

- Heimberg, R. G., Brozovich, F. A., & Rapee, R. M. (2014). A Cognitive-Behavioral Model of Social Anxiety Disorder. In *Social Anxiety: Clinical, Developmental, and Social Perspectives* (pp. 705–728). Elsevier. <https://doi.org/10.1016/B978-0-12-394427-6.00024-8>
- Hofmann, S. G. (2007). Cognitive factors that maintain social anxiety disorder: A Comprehensive Model and its treatment implications. *Cognitive Behaviour Therapy*, 36(4), 193–209. <https://doi.org/10.1080/16506070701421313>
- Hollocks, M. J., Ozsvadjian, A., Matthews, C. E., Howlin, P., & Simonoff, E. (2013). The relationship between attentional bias and anxiety in children and adolescents with autism spectrum disorders. *Autism Research*, 6(4), 237–247. <https://doi.org/10.1002/aur.1285>
- Hollocks, M. J., Pickles, A., Howlin, P., & Simonoff, E. (2016). Dual cognitive and biological correlates of anxiety in Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 46(10), 3295–3307. <https://doi.org/10.1007/s10803-016-2878-2>
- Kliemann, D., Dziobek, I., Hatri, A., Steimke, R., & Heekeren, H. R. (2010). Atypical reflexive gaze patterns on emotional faces in autism spectrum disorders. *Journal of Neuroscience*, 30(37), 12281–12287. <https://doi.org/10.1523/JNEUROSCI.0688-10.2010>
- Klumpp, H., & Amir, N. (2009). Examination of vigilance and disengagement of threat in social anxiety with a probe detection task. *Anxiety Stress and Coping*, 22(3), 283–296. <https://doi.org/10.1080/10615800802449602>
- Kret, M. E., Jaasma, L., Bionda, T., & Wijnen, J. G. (2016). Bonobos (*Pan paniscus*) show an attentional bias toward conspecifics' emotions. *Proceedings of the National Academy of Sciences of the United States of America*, 113(14), 3761–3766. <https://doi.org/10.1073/pnas.1522060113>
- Kret, M. E., & Ploeger, A. (2015). Emotion processing deficits: A liability spectrum providing insight into comorbidity of mental disorders. *Neuroscience and Biobehavioral Reviews*, 52, 153–171. <https://doi.org/10.1016/j.neubiorev.2015.02.011>
- Lazarov, A., Basel, D., Dolan, S., Dillon, D. G., Pizzagalli, D. A., & Schneier, F. R. (2021). Increased attention allocation to socially threatening faces in social anxiety disorder: A replication study. *Journal of Affective Disorders*, 290, 169–177. <https://doi.org/10.1016/j.jad.2021.04.063>
- Liebowitz, M. R. (1987). Social phobia. *Modern Problems in Pharmacopsychiatry*, 22, 141–173. <https://doi.org/10.1159/000414022>
- MacLeod, C., Grafton, B., & Notebaert, L. (2019). Anxiety-linked attentional Bias: Is it Reliable? *Annu Rev Clin Psychol*, 15, 529–554. <https://doi.org/10.1146/annurev-clinpsy-050718>
- MacLeod, C., Mathews, A., & Tata, P. (1986). Attentional Bias in Emotional Disorders. *Journal of Abnormal Psychology*, 95(1), 15–20. <https://doi.org/10.1037/0021-843X.95.1.15>
- MacLeod, C., Rutherford, E., Campbell, L., Ebsworthy, G., & Holker, L. (2002). Selective attention and emotional vulnerability: Assessing the causal basis of their association through the experimental manipulation of attentional bias. *Journal of Abnormal Psychology*, 111(1), 107–123. <https://doi.org/10.1037/0021-843X.111.1.107>
- Makowski, D., Ben-Shachar, M. S., Chen, S. A., & Lüdtke, D. (2019). Indices of effect existence and significance in the Bayesian framework. *Frontiers in psychology*, 10, 2767. <https://doi.org/10.3389/fpsyg.2019.02767>
- May, T., Cornish, K., & Rinehart, N. J. (2015). Mechanisms of anxiety related attentional biases in children with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 45(10), 3339–3350. <https://doi.org/10.1007/s10803-015-2500-z>
- Miu, A. C., Paná, S. E., & Avram, J. (2012). Emotional face processing in neurotypicals with autistic traits: Implications for the broad autism phenotype. *Psychiatry Research*, 198(3), 489–494. <https://doi.org/10.1016/j.psychres.2012.01.024>
- Möller, E. L., Majdandžić, M., & Bögels, S. M. (2014). Fathers' versus mothers' social referencing signals in relation to infant anxiety and avoidance: A visual cliff experiment. *Developmental Science*, 17(6), 1012–1028. <https://doi.org/10.1111/desc.12194>
- Mogg, K., Bradley, B. P., Miles, F., & Dixon, R. (2004a). Time course of attentional bias for threat scenes: Testing the vigilance-avoidance hypothesis. *Cognition and Emotion*, 18(5), 689–700. <https://doi.org/10.1080/02699930341000158>
- Mogg, K., Philippot, P., & Bradley, B. P. (2004b). Selective attention to angry faces in clinical Social Phobia. *Journal of Abnormal Psychology*, 113(1), 160–165. <https://doi.org/10.1037/0021-843X.113.1.160>
- Monk, C. S., Weng, S. J., Wiggins, J. L., Kurapati, N., Louro, H. M. C., Carrasco, M., Maslowsky, J., Risi, S., & Lord, C. (2010). Neural circuitry of emotional face processing in autism spectrum disorders. *Journal of Psychiatry and Neuroscience*, 35(2), 105–114. <https://doi.org/10.1503/jpn.090085>
- Neophytou, K., & Panayiotou, G. (2022). Does attention bias modification reduce anxiety in socially anxious college students? An experimental study of potential moderators and considerations for implementation. *Plos One*, 17(2 February). <https://doi.org/10.1371/journal.pone.0264256>
- Pool, E., Brosch, T., Delplanque, S., & Sander, D. (2016). Attentional bias for positive emotional stimuli: A meta-analytic investigation. *Psychological Bulletin*, 142(1), 79–106. <https://doi.org/10.1037/bul0000026>
- Puls, S., & Rothermund, K. (2018). Attending to emotional expressions: No evidence for automatic capture in the dot-probe task. *Cognition and Emotion*, 32(3), 450–463. <https://doi.org/10.1080/02699931.2017.1314932>
- Rapee, R. M., & Heimberg, R. G. (1997). A cognitive-behavioral model of anxiety in social phobia. *Behaviour Research and Therapy*, 35(8), 741–756. [https://doi.org/10.1016/S0005-7967\(97\)00022-3](https://doi.org/10.1016/S0005-7967(97)00022-3)
- Rapee, R. M., & Spence, S. H. (2004). The etiology of social phobia: Empirical evidence and an initial model. *Clinical Psychology Review*, 24(7), 737–767. <https://doi.org/10.1016/j.cpr.2004.06.004>
- Robinson, E. B., Munir, K., Munafò, M. R., Hughes, M., McCormick, M. C., & Koenen, K. C. (2011). Stability of autistic traits in the General Population: Further evidence for a Continuum of Impairment. *Journal of the American Academy of Child and Adolescent Psychiatry*, 50(4), 376–384. www.jaacap.org
- Salemink, E., van den Hout, M. A., & Kindt, M. (2007). Selective attention and threat: Quick orienting versus slow disengagement and two versions of the dot probe task. *Behaviour Research and Therapy*, 45(3), 607–615. <https://doi.org/10.1016/j.brat.2006.04.004>
- Schneider, W., Eschman, A., & Zuccolotto, A. (2002). E-Prime (Version 2.0). [Computer software and manual]. Pittsburgh, PA: Psychology Software Tools Inc.
- Spain, D., Sin, J., Linder, K. B., McMahon, J., & Happé, F. (2018). Social anxiety in autism spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, 52, 51–68. <https://doi.org/10.1016/j.rasd.2018.04.007>
- Stuit, S. M., Paffen, C. L. E., & van der Stigchel, S. (2021). Introducing the prototypical stimulus characteristics Toolbox: Protocol. *Behavior Research Methods*. <https://doi.org/10.3758/s13428-021-01737-9>
- Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., Marcus, D. J., Westerlund, A., Casey, B. J., & Nelson, C. (2009). The NimStim set of facial expressions: Judgments from untrained research participants. *Psychiatry Research*, 168(3), 242–249. <https://doi.org/10.1016/j.psychres.2008.05.006>
- van Berlo, E., Bionda, T., & Kret, M. (2020). Attention Towards Emotions is Modulated by Familiarity with the Expressor: A Comparison Between Bonobos and Humans. 0–2. <https://doi.org/10.1101/2020.05.11.089813>

- van Bockstaele, B., & Bögels, S. M. (2014). Mindfulness-Based Therapy for Social Anxiety Disorder. In *Social Anxiety: Clinical, Developmental, and Social Perspectives* (pp. 729–751). Elsevier. <https://doi.org/10.1016/B978-0-12-394427-6.00025-X>
- van Bockstaele, B., Salemink, E., Bögels, S. M., & Wiers, R. W. (2017). Limited generalisation of changes in attentional bias following attentional bias modification with the visual probe task. *Cognition and Emotion*, *31*(2), 369–376. <https://doi.org/10.1080/02699931.2015.1092418>
- van Rooijen, R., Ploeger, A., & Kret, M. E. (2017). The dot-probe task to measure emotional attention: A suitable measure in comparative studies? *Psychonomic Bulletin and Review*, *24*(6), 1686–1717. <https://doi.org/10.3758/s13423-016-1224-1>
- White, S. W., Bray, B. C., & Ollendick, T. H. (2012). Examining shared and unique aspects of social anxiety disorder and autism spectrum disorder using factor analysis. *Journal of Autism and Developmental Disorders*, *42*(5), 874–884. <https://doi.org/10.1007/s10803-011-1325-7>
- Wirth, B. E., & Wentura, D. (2020). It occurs after all: Attentional bias towards happy faces in the dot-probe task. *Attention Perception and Psychophysics*, *82*(5), 2463–2481. <https://doi.org/10.3758/s13414-020-02017-y>
- Yiend, J. (2010). The effects of emotion on attention: A review of attentional processing of emotional information. *Cognition and Emotion*, *24*(1), 3–47. <https://doi.org/10.1080/02699930903205698>
- Zalla, T., & Sperduti, M. (2013). The amygdala and the relevance detection theory of autism: An evolutionary perspective. *Frontiers in Human Neuroscience*, *7*(DEC), 1–15. <https://doi.org/10.3389/fnhum.2013.00894>
- Zvielli, A., Bernstein, A., & Koster, E. H. W. (2015). Temporal Dynamics of Attentional Bias. *Clinical Psychological Science*, *3*(5), 772–788. <https://doi.org/10.1177/2167702614551572>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.