

# **Tropical moments of tropical Jacobians**

Jong, R.S. de; Shokrieh, F.

# Citation

Jong, R. S. de, & Shokrieh, F. (2022). Tropical moments of tropical Jacobians. *Canadian Journal Of Mathematics*, *75*(4), 1045-1075. doi:10.4153/S0008414X22000220

Version:	Publisher's Version
License:	Creative Commons CC BY 4.0 license
Downloaded from:	https://hdl.handle.net/1887/3563405

**Note:** To cite this publication please use the final published version (if applicable).

Canad. J. Math. Vol. 75 (4), 2023 pp. 1045–1075 http://dx.doi.org/10.4153/S0008414X22000220

© The Author(s), 2022. Published by Cambridge University Press on behalf of The Canadian Mathematical Society. This is an Open Access article, distributed



under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/ by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Tropical moments of tropical Jacobians

# Robin de Jong<sup>®</sup> and Farbod Shokrieh<sup>®</sup>

*Abstract.* Each metric graph has canonically associated to it a polarized real torus called its tropical Jacobian. A fundamental real-valued invariant associated to each polarized real torus is its tropical moment. We give an explicit and efficiently computable formula for the tropical moment of a tropical Jacobian in terms of potential theory on the underlying metric graph. We show that there exists a universal linear relation between the tropical moment, a certain capacity called the tau invariant, and the total length of a metric graph. To put our formula in a broader context, we relate our work to the computation of heights attached to principally polarized abelian varieties.

# 1 Introduction

# 1.1 Background and aim

A *lattice* is the datum of a finitely generated free abelian group H together with a positive definite bilinear form  $[\cdot, \cdot]$  on the real vector space  $H_{\mathbb{R}} = H \otimes_{\mathbb{Z}} \mathbb{R}$ . An alternative way of packaging the data  $(H, [\cdot, \cdot])$  is to consider a real torus  $\mathbb{T} = H_{\mathbb{R}}/H$  equipped with a *polarization*, that is, a positive definite bilinear form on its tangent space. A lattice  $(H, [\cdot, \cdot])$  (or equivalently, a polarized real torus) canonically determines a compact convex region in  $H_{\mathbb{R}}$  given by

$$Vor(0) = \{z \in H_{\mathbb{R}} : \text{ for all } \lambda \in H : [z, z] \le [z - \lambda, z - \lambda]\},\$$

known as the *Voronoi polytope* of  $(H, [\cdot, \cdot])$  centered at the origin. Let  $\mu_L$  denote a Lebesgue measure on  $H_{\mathbb{R}}$ . In this work, we are interested in the second moment of  $(H, [\cdot, \cdot])$ , defined as the value in  $\mathbb{R}$  of the integral

(1.1) 
$$\int_{\operatorname{Vor}(0)} [z, z] \, \mathrm{d} \mu_L(z),$$

or rather its normalized version, which we call the *tropical moment*:

(1.2) 
$$I(H, [\cdot, \cdot]) = I(\mathbb{T}) = \frac{\int_{\operatorname{Vor}(0)} [z, z] \, d\mu_L(z)}{\int_{\operatorname{Vor}(0)} d\mu_L(z)}.$$

The tropical moment is closely related to other lattice invariants such as the *covering radius* and the *packing radius*. The celebrated book [13] by Conway and Sloane contains

Received by the editors September 28, 2021; revised May 3, 2022; accepted May 11, 2022.

Published online on Cambridge Core May 24, 2022.

The second-named author was partially supported by the NSF CAREER grant DMS-2044564. AMS subject classification: 14T15, 14G40, 11G50, 94C05, 52C45.

Keywords: Tropical Jacobian, moment, random spanning trees, Voronoi polytope, zonotope, potential theory, tau invariant, heights of abelian varieties, Arakelov invariants.

an entire chapter (Chapter 21) devoted to the tropical moment, full of examples and of ways of computing it. In information theory, the tropical moment determines the quality of a lattice as a vector quantizer.

In this paper, we will study lattices canonically associated to *metric graphs*. Our purpose is to show that the tropical moment of these lattices allows an explicit expression in terms of the capacity attached to the effective resistance function. All ingredients of this expression are moreover efficiently computable. Up to dimension 3, every lattice arises from a metric graph, so that, in particular, one can recover the moment computations in [9, 13] from our expression. We mention that explicit formulas for the tropical moments of lattices in dimension 4 were obtained by Zimmermann in [32].

A metric graph is a compact, connected length metric space  $\Gamma$  homeomorphic to a topological graph. The lattice that we associate to a metric graph  $\Gamma$  has as underlying abelian group H its *first homology group*  $H = H_1(\Gamma, \mathbb{Z})$ . The real vector space  $H_{\mathbb{R}} = H_1(\Gamma, \mathbb{R})$  is equipped with a standard inner product, determined by the lengths of the edges in a model of  $\Gamma$ . See Section 3.3. The resulting "homology lattice" is independent of the choice of a model. The polarized real torus  $H_{\mathbb{R}}/H$  associated to  $\Gamma$  is commonly known as the *tropical Jacobian* of  $\Gamma$ , and denoted by Jac( $\Gamma$ ).

#### 1.2 Main result

It is well known that one can think of a metric graph  $\Gamma$  as an electrical network. For  $x, y, z \in \Gamma$ , we define  $j_z(x, y)$  to be the electric potential at x if one unit of current enters the network at y and exits at z, with z "grounded" (i.e., has zero potential). The *j*-function provides a fundamental solution of the Laplacian operator on  $\Gamma$ , so naturally it is an important function in studying harmonic analysis on  $\Gamma$ . The *effective resistance* between x and y is defined as  $r(x, y) = j_y(x, x)$ , which has the expected physical meaning in terms of electrical networks. See Section 4.2.

It is convenient to fix a vertex set (which is a finite nonempty set containing all the branch points of  $\Gamma$ ) and to think of the metric graph  $\Gamma$  as being obtained from a finite (combinatorial) graph *G*. The metric data will be encoded by positive real numbers  $\ell(e)$  for each edge *e* of *G*. We call the finite weighted graph arising in this way a *model* of  $\Gamma$ . If *e* is an edge of *G*, the *Foster coefficient* of *e* is defined by  $\Gamma(e) = 1 - r(u, v)/\ell(e)$ , where *u* and *v* are the two endpoints of the edge  $e = \{u, v\}$  (see Definition 4.4).

Our main result is the following formula for the tropical moment of the homology lattice of  $\Gamma$ .

**Theorem** A (=Theorem 8.1) Let  $\Gamma$  be a metric graph. Fix a model G of  $\Gamma$ , and fix a vertex q of G. Then, for the tropical moment  $I(Jac(\Gamma))$  of the tropical Jacobian of  $\Gamma$ , one has

$$I(\operatorname{Jac}(\Gamma)) = \frac{1}{12} \sum_{e} \mathsf{F}(e)^{2} \ell(e) + \frac{1}{4} \sum_{e=\{u,v\}} \left( r(u,v) - \frac{j_{u}(v,q)^{2} + j_{v}(u,q)^{2}}{\ell(e)} \right),$$

where the sums are over all edges  $e \in E(G)$ .

### Tropical moments of tropical Jacobians

As we explain in Remark 8.2(ii), the summands in this formula can all be computed in matrix multiplication time  $O(n^{\omega})$ , where *n* is the number of vertices in the given model *G* of  $\Gamma$ , and  $\omega$  is the exponent for matrix multiplication (currently  $\omega < 2.38$ ). It is generally believed that to compute the second moment of a lattice is a hard problem. For example, if one wants to compute the tropical moment via the general "simplex method" (see, e.g., [13, Chapter 21, Section 2]), then one needs to know the set of vertices of Vor(0). However, as is shown in [17], even computing the *number* of vertices of Vor(0) is already #P-hard. Interestingly, the example that is used to show this is given by the homology lattice of a graph.

#### **1.3** Sketch of the proof of the main result

The proof of Theorem A is very subtle. Our strategy is as follows. To handle the integral in (1.2), we provide an explicit polytopal decomposition of Vor(0). Given a base point  $q \in V(G)$ , there is a full-dimensional polytope  $\sigma_T + \mathcal{C}_T$  in our decomposition attached to each *spanning tree* T of G. Here,  $\mathcal{C}_T$  is a centrally symmetric polytope, which makes  $\sigma_T$  into the center of  $\sigma_T + \mathcal{C}_T$  (see Theorem 7.5). The desired integral over Vor(0) is then a sum of integrals over each  $\sigma_T + \mathcal{C}_T$ .

The contribution from the centrally symmetric polytopes  $C_T$  is rather easy to handle. The real difficulty comes in handling the contributions from the centers  $\sigma_T$ . We introduce the notion of *energy level* of rooted spanning trees (see Definition 6.1). A crucial ingredient is the notion of *cross ratio* introduced in [16] for electrical networks. We prove that the weighted average of energy levels over all spanning trees has a remarkably simple expression in terms of values of the *j*-function (see Theorem 6.3). Proving this, in turn, uses some subtle computations related to functions arising from *random spanning trees* (see Section 5). We also repeatedly use our generalized (and quantitative) version of Rayleigh's law in electrical networks, as developed in the companion paper [16].

Recall that the homology group of a graph naturally gives rise to a regular matroid. In principle, most of our approach leading to the computation of tropical moments should generalize to the setting of regular matroids. One can replace the subdivision in Theorem 7.5 with an arbitrary *tight coherent subdivision* in the sense of [10, Section 4]. We do not expect a clean expression as in Theorem 6.3 for general regular matroids. However, we note that already Lemma 6.5 yields an efficient algorithm for the weighted average of energy levels of bases of regular matroids.

#### 1.4 Applications and context for our formula

Theorem A can be used to give a simple connection between  $I(\text{Jac}(\Gamma))$  and a wellknown potential theoretic capacity associated to  $\Gamma$  called the *tau invariant*, denoted by  $\tau(\Gamma)$  (see Section 11 for its definition). The invariant  $\tau(\Gamma)$  can be traced back to the fundamental work of Chinburg and Rumely [11] in their study of the Arakelov geometry of arithmetic surfaces at non-archimedean places. We have the following result. **Theorem B** (=Theorem 11.4) Let  $\Gamma$  be a metric graph. Let  $\tau(\Gamma)$  denote the tau invariant of  $\Gamma$ , and let  $\ell(\Gamma)$  denote its total length. Then the identity

$$\frac{1}{2}\tau(\Gamma) + I(\operatorname{Jac}(\Gamma)) = \frac{1}{8}\ell(\Gamma)$$

holds in  $\mathbb{R}$ .

Next, we will explain in Section 9 how our work can be used for the computation of the *stable Faltings height* of principally polarized abelian varieties defined over number fields. For a discussion of an explicit example in this context, we refer to Section 10. As we will see there, our work gives a complete conceptual explanation of all entries in a table, found by Autissier [3], related to the calculation of local non-archimedean terms in a formula for the stable Faltings height of curves of genus 2.

Finally, we note that Theorem B may be thought of as an analogue, in the nonarchimedean setting, of a remarkable identity established by Wilms [28, Theorem 1.1] between analytic invariants of Riemann surfaces. In fact, in [15, 29], Theorem B is used together with [28, Theorem 1.1] to derive a formula for the asymptotic behavior of the so-called Zhang–Kawazumi invariant [20, 21, 31] in arbitrary one-parameter semistable degenerations of Riemann surfaces.

#### **1.5** Structure of the paper

In Section 2, we review the notion of polarized real tori and define the notion of tropical moments. In Section 3, we review the notions of weighted graphs and of metric graphs and their models. Moreover, we introduce the tropical Jacobian of a metric graph. In Section 4, we review potential theory and harmonic analysis on metric graphs, mainly from the perspective of our companion paper [16]. In Section 5, we study two functions that arise from the theory of random spanning trees. In Section 6, we introduce the notion of energy levels of rooted spanning trees, and prove that the average of energy levels has a simple expression in terms of the *j*-function. In Section 7, we study the combinatorics of the Voronoi polytopes arising from graphs, and present our suitable polytopal decomposition. In Section 8, we prove Theorem A. In Section 9 we discuss our application to the computation of stable Faltings heights. In Section 10, we elaborate upon an example related to Jacobian varieties in dimension 2. In Section 11, we introduce the tau invariant and prove Theorem B.

# 2 Polarized real tori and tropical moments

The purpose of this section is to set notations and terminology related to polarized real tori and their tropical moments.

#### 2.1 Polarized real tori

A (*Euclidean*) *lattice* is a pair  $(H, [\cdot, \cdot])$  consisting of a finitely generated free  $\mathbb{Z}$ -module H and a positive definite symmetric bilinear form  $[\cdot, \cdot]$  on the real vector space  $H_{\mathbb{R}} = H \otimes_{\mathbb{Z}} \mathbb{R}$ . Attached to each lattice  $(H, [\cdot, \cdot])$ , one has a real torus  $\mathbb{T} = H_{\mathbb{R}}/H$ , equipped with a natural structure of compact Riemannian manifold. We refer to the Riemannian manifold  $\mathbb{T}$  as a *polarized real torus*. The tropical Jacobian of a metric

graph (see Section 3.3) is an example of a polarized real torus. Clearly, "lattice" and "polarized real torus" are equivalent notions. We will mainly prefer the terminology of polarized real tori.

#### 2.2 Voronoi decompositions and tropical moment

Let  $\mathbb{T}$  be a polarized real torus coming from a lattice  $(H, [\cdot, \cdot])$  as above. For each  $\lambda \in H$ , we denote by  $Vor(\lambda)$  the Voronoi polytope of the lattice  $(H, [\cdot, \cdot])$  around  $\lambda$ :

 $\operatorname{Vor}(\lambda) \coloneqq \{z \in H_{\mathbb{R}} \colon \text{for all } \lambda' \in H \colon [z - \lambda, z - \lambda] \leq [z - \lambda', z - \lambda'] \}.$ 

Note that, for each  $\lambda \in H$ , we have  $Vor(\lambda) = Vor(0) + \lambda$ . Moreover, Vor(0), up to some identifications on its boundary, is a fundamental domain for the translation action of H on  $H_{\mathbb{R}}$ .

**Definition 2.1** (cf. [13, Chapter 21]) The *tropical moment* of the polarized real torus  $\mathbb{T}$  is set to be the value of the integral

(2.1) 
$$I(\mathbb{T}) \coloneqq \int_{\operatorname{Vor}(0)} [z, z] \, \mathrm{d}\mu_L(z)$$

Here,  $\mu_L$  is the Lebesgue measure on  $H_{\mathbb{R}}$ , normalized to have  $\mu_L(Vor(0)) = 1$ .

# 3 Metric graphs, models, and tropical Jacobians

The purpose of this section is to set notations and terminology related to weighted graphs, metric graphs, and their models. We also define the tropical Jacobian of a metric graph (see Section 3.3). Most of the material in this section is straightforward, and we leave details to the interested reader.

#### 3.1 Weighted graphs

By a *weighted graph*, we mean a finite weighted connected multigraph *G* with no loop edges. The set of vertices of *G* is denoted by V(G), and the set of edges of *G* is denoted by E(G). We let n = |V(G)| and m = |E(G)|. An edge *e* is called a *bridge* if  $G \setminus e$  is disconnected. The weights of edges are determined by a *length* function  $\ell: E(G) \to \mathbb{R}_{>0}$ . We let  $\mathbb{E}(G) = \{e, \bar{e}: e \in E(G)\}$  denote the set of *oriented edges*. We have  $\bar{\bar{e}} = e$ . For each subset  $\mathcal{A} \subseteq \mathbb{E}(G)$ , we define  $\overline{\mathcal{A}} = \{\bar{e}: e \in \mathcal{A}\}$ . An *orientation*  $\mathcal{O}$  on *G* is a partition  $\mathbb{E}(G) = \mathcal{O} \cup \overline{\mathcal{O}}$ . We have an obvious extension of the length function  $\ell: \mathbb{E}(G) \to \mathbb{R}_{>0}$  by requiring  $\ell(e) = \ell(\bar{e})$ . There is a natural map  $\mathbb{E}(G) \to V(G) \times V(G)$  sending an oriented edge *e* to  $(e^+, e^-)$ , where  $e^-$  is the start point of *e* and  $e^+$  is the end point of *e*.

### Notation

For  $e \in E(G)$ , we sometimes refer to its endpoints by  $e^+$ ,  $e^-$  even when an orientation is not fixed, so  $e = \{e^+, e^-\}$ . We only allow ourselves to do this if the underlying expression is symmetric with respect to  $e^+$  and  $e^-$ , so there is no danger of confusion. The reader is welcome to fix an orientation  $\mathbb{O}$  and think of  $e^+$  and  $e^-$  in the sense explained above.

R. de Jong and F. Shokrieh



Figure 1:

(a) A metric graph Γ.

(b) A weighted graph model G of  $\Gamma$ .

(c) A rooted spanning tree (T, q) of *G* and the orientation  $\mathcal{T}_q$ .

A spanning tree T of G is a maximal subset of E(G) that contains no circuit (closed simple path). Equivalently, T is a minimal subset of E(G) that connects all vertices of G.

For a fixed  $q \in V(G)$  and spanning tree *T* of *G*, we will refer to the pair (T, q) as a *spanning tree with a root at q* (or just a *rooted spanning tree*). The choice of *q* imposes a preferred orientation on all edges of *T*. Namely, one can require that all edges are oriented *away from q* on the spanning tree *T* (see Figure 1c). We denote this orientation on *T* by  $\mathcal{T}_q \subseteq \mathbb{E}(G)$ .

Given a commutative ring *R*, it is convenient to define the 1-chains with coefficients in *R* as the free module

$$C_1(G,R) := \frac{\bigoplus_{e \in \mathbb{E}(G)} Re}{\langle e + \bar{e} : e \in \mathcal{O} \rangle}$$

Note that  $\bar{e} = -e$  in  $C_1(G, R)$ . For each orientation  $\mathcal{O}$  on G, we have an isomorphism  $C_1(G, R) \simeq \bigoplus_{e \in \mathcal{O}} Re$ . For each subset  $\mathcal{A} \subseteq \mathbb{E}(G)$ , we define its *associated* 1-chain as  $\gamma_{\mathcal{A}} = \sum_{e \in \mathcal{A}} e$ .

#### 3.2 Metric graphs and models

A *metric graph* (or *metrized graph*) is a pair  $(\Gamma, d)$  consisting of a compact connected topological graph  $\Gamma$ , together with an *inner metric d*. Equivalently, if  $\Gamma$  is not a one-point space, then a metric graph is a compact connected metric space  $\Gamma$  which has the property that every point has an open neighborhood isometric to a star-shaped set, endowed with the path metric.

The points of  $\Gamma$  that have valency different from 2 are called *branch points* of  $\Gamma$ . A *vertex set* for  $\Gamma$  is a finite set *V* of points of  $\Gamma$  containing all the branch points of  $\Gamma$  with the property that for each connected component *c* of  $\Gamma \setminus V$ , the closure of *c* in  $\Gamma$  is isometric with a closed interval.

A vertex set *V* for  $\Gamma$  naturally determines a weighted graph *G* by setting V(G) = V, and by setting E(G) to be the set of connected components of  $\Gamma \setminus V$ . We call *G* a *model* of  $\Gamma$ . An *edge segment* (based on the choice of a vertex set *V*) is the closure in  $\Gamma$  of a connected component of  $\Gamma \setminus V$ . Note that there is a natural bijective correspondence between E(G) and the edge segments of  $\Gamma$  determined by *V*. By a small abuse of



Figure 2: The electrical network N corresponding to the graphs in Figure 1a,b.

terminology, we will refer to the elements of E(G) also as edge segments of  $\Gamma$ . Given an edge segment  $e \subset \Gamma$  (based on the choice of a vertex set *V*), we denote its boundary  $\partial e \subset V$  by  $\partial e = \{e^-, e^+\}$ . In particular, we will also use the notation  $\{e^-, e^+\}$  for the boundary set of an edge segment *e* if there is no (preferred) orientation present. We hope that this does not lead to confusion.

Conversely, every weighted graph *G* naturally determines a metric graph  $\Gamma_G$  containing V(G) by gluing closed intervals  $[0, \ell(e)]$  for  $e \in E(G)$  according to the incidence relations. Note that V(G) is naturally a vertex set of  $\Gamma_G$ , and the associated model is precisely *G*. See Figure 1a,b.

#### 3.3 Tropical Jacobians

Let  $\Gamma$  be a metric graph. Fix a model G of  $\Gamma$ . Let  $\mathcal{O} = \{e_1, \ldots, e_m\}$  be a labeling of an orientation  $\mathcal{O}$  on G. The real vector space  $C_1(G, \mathbb{R}) \simeq \bigoplus_{i=1}^m \mathbb{R}e_i$  has a canonical inner product defined by  $[e_i, e_j] = \delta_i(j)\ell(e_i)$ . Here,  $\delta_i$  denotes the delta (Dirac) measure on  $\{1, 2, \ldots, m\}$  centered at *i*. The resulting inner product space  $(C_1(G, \mathbb{R}), [\cdot, \cdot])$  is independent of the choice of  $\mathcal{O}$  and its labeling.

The inner product  $[\cdot, \cdot]$  restricts to an inner product, also denoted by  $[\cdot, \cdot]$ , on the homology lattice  $H = H_1(G, \mathbb{Z}) \subset C_1(G, \mathbb{Z})$ . The pair  $(H, [\cdot, \cdot])$  is a canonical lattice associated to  $\Gamma$  (independent of the choice of the model *G*), and we have a canonical identification  $H \simeq H_1(\Gamma, \mathbb{Z})$ . Note that  $H_{\mathbb{R}} \simeq H_1(\Gamma, \mathbb{R})$ . The associated polarized real torus  $H_1(\Gamma, \mathbb{R})/H_1(\Gamma, \mathbb{Z})$  is called the *tropical Jacobian* of  $\Gamma$  [23, 25], and denoted by Jac( $\Gamma$ ).

# 4 Potential theory on metric graphs

In this section, we closely follow [16] and review those results that are needed in this paper.

#### 4.1 Graphs as electrical networks

Let  $\Gamma$  be a metric graph, and let G be a model of  $\Gamma$ . We may think of  $\Gamma$  (or G) as an electrical network in which each edge  $e \in E(G)$  is a resistor having resistance  $\ell(e)$ . See Figure 2.

When studying the "potential theory" on a metric graph  $\Gamma$ , it is convenient to always fix an (arbitrary) model *G*, and think of it as an electrical network.

#### 4.2 Laplacians and *j*-functions

Let  $\Gamma$  be a metric graph, and let *G* be a model of  $\Gamma$ . We have the distributional Laplacian operator (see [16, Section 3.1])

$$\Delta: PL(\Gamma) \rightarrow DMeas_0(\Gamma),$$

where PL( $\Gamma$ ) is the real vector space consisting of all continuous piecewise affine real valued functions on  $\Gamma$  that can change slope finitely many times on each closed edge segment, and DMeas<sub>0</sub>( $\Gamma$ ) is the real vector space of discrete measures  $\nu$  on  $\Gamma$  with  $\nu(\Gamma) = 0$ . We also have the combinatorial Laplacian operator (see [16, Section 3.2])

$$\Delta: \mathcal{M}(G) \to \mathrm{DMeas}_0(G),$$

where  $\mathcal{M}(G)$  is the real vector space of real-valued functions on V(G), and  $DMeas_0(G)$  is the real vector space of discrete measures v on V(G) with v(V(G)) = 0. The distributional Laplacian  $\Delta$  and the combinatorial Laplacian  $\Delta$  are compatible in the sense described in [16, Section 3.3]. Moreover, the combinatorial Laplacian on *G* can be conveniently presented by its *Laplacian matrix*; let  $\{v_1, \ldots, v_n\}$  be a labeling of V(G). The Laplacian matrix  $\mathbf{Q}$  associated to *G* is the  $n \times n$  matrix  $\mathbf{Q} = (q_{ij})$  where, for  $i \neq j$ , we have  $q_{ij} = -\sum_{e = \{v_i, v_j\} \in E(G)} 1/\ell(e)$ . The diagonal entries are determined by forcing the matrix to have zero-sum rows.

The Laplacian matrix of *G* can also be expressed in terms of the *incidence matrix* of *G*. Let  $\{v_1, \ldots, v_n\}$  be a labeling of V(G) as before. Fix an orientation  $\mathcal{O} = \{e_1, \ldots, e_m\}$  on *G*. The incidence matrix **B** associated to *G* is the  $n \times m$  matrix **B** =  $(b_{ij})$ , where  $b_{ij} = +1$  if  $e_j^+ = v_i$  and  $b_{ij} = -1$  if  $e_j^- = v_i$  and  $b_{ij} = 0$  otherwise. Let **D** denote the  $m \times m$  diagonal matrix with diagonal entries  $\ell(e_i)$  for  $e_i \in \mathcal{O}$ . We have  $\mathbf{Q} = \mathbf{B}\mathbf{D}^{-1}\mathbf{B}^{\mathrm{T}}$ , where  $(\cdot)^{\mathrm{T}}$  denotes the matrix transpose operation.

A fundamental solution of the Laplacian is given by *j*-functions. We follow the notation of [11]. See [16, Section 4] and the references therein for more details. Let  $\Gamma$  be a metric graph and fix two points  $y, z \in \Gamma$ . We denote by  $j_z(\cdot, y; \Gamma)$  the unique function in PL( $\Gamma$ ) satisfying: (i)  $\Delta(j_z(\cdot, y; \Gamma)) = \delta_y - \delta_z$ , and (ii)  $j_z(z, y; \Gamma) = 0$ . If the metric graph  $\Gamma$  is clear from the context, we write  $j_z(x, y)$  instead of  $j_z(x, y; \Gamma)$ . The *j*-function exists and is unique, and satisfies the following basic properties:

- $i_z(x, y)$  is jointly continuous in all three variables *x*, *y*, *z* ∈ Γ.
- $\diamond \ j_z(x,y) = j_z(y,x).$
- $\diamond \ 0 \leq j_z(x,y) \leq j_z(x,x).$

$$\diamond \ j_z(x,x) = j_x(z,z).$$

The *effective resistance* between two points  $x, y \in \Gamma$  is  $r(x, y) := j_y(x, x)$ . If we want to clarify the underlying metric graph  $\Gamma$ , we use the notation  $r(x, y; \Gamma)$ .

Let *G* be an arbitrary model of  $\Gamma$ . One can explicitly compute the quantities  $j_q(p, v) \in \mathbb{R}$  for  $q, p, v \in V(G)$  using linear algebra (see [8, Section 3]) as follows. Fix a labeling of V(G) as before, and let **Q** be the corresponding Laplacian matrix. Let  $\mathbf{Q}_q$  be the  $(n-1) \times (n-1)$  matrix obtained from **Q** by deleting the row and column corresponding to  $q \in V(G)$  from **Q**. It is well known that  $\mathbf{Q}_q$  is invertible. Let  $\mathbf{L}_q$  be the  $n \times n$  matrix obtained from  $\mathbf{Q}_q^{-1}$  by inserting zeros in the row and column corresponding to q. One can easily check that  $\mathbf{QL}_q = \mathbf{I} + \mathbf{R}_q$ , where **I** is the

 $n \times n$  identity matrix and  $\mathbf{R}_q$  has all -1 entries in the row corresponding to q and has zeros elsewhere. It follows from the compatibility of the two Laplacians that  $\mathbf{L}_q = (j_q(p, v))_{p, v \in V(G)}$ . The matrix  $\mathbf{L}_q$  is a *generalized inverse* of  $\mathbf{Q}$ , in the sense that  $\mathbf{QL}_q \mathbf{Q} = \mathbf{Q}$ .

*Remark 4.1* Computing  $L_q$  takes time at most  $O(n^{\omega})$ , where  $\omega$  is the exponent for matrix multiplication (currently  $\omega < 2.38$ ).

#### 4.3 Cross ratios

Let  $\Gamma$  be a metric graph and fix  $q \in \Gamma$ . As in [16], we define the *cross ratio* function (with respect to the base point q)  $\xi_q \colon \Gamma^4 \to \mathbb{R}$  by

$$\xi_q(x, y, z, w) := j_q(x, z) + j_q(y, w) - j_q(x, w) - j_q(y, z).$$

If we want to clarify the graph  $\Gamma$ , we use the notation  $\xi_q(x, y, z, w; \Gamma)$  instead. As is observed in [16, Remark 6.1(i)], we have the identity

$$-2\xi_q(x, y, z, w) = r(x, z) + r(y, w) - r(x, w) - r(y, z).$$

*Remark 4.2* We borrowed the *cross ratio* terminology in [16] from the book of Baker and Rumely [7, Appendix B]. These cross ratios on the Berkovich hyperbolic space and on its natural extension to the Berkovich projective line also play an important role in the work of Favre and Rivera-Letelier [18, Section 6.3]. The terminology is also justified in the context of Gromov hyperbolic spaces, where this is sometimes called "cross difference" (see, for example, [27, Section 4.5]).

Cross ratios satisfy the following basic properties:

- ◊  $ξ(x, y, z, w) := ξ_q(x, y, z, w)$  is independent of the choice of *q*.
- $\diamond \ \xi(x, y, z, w) = \xi(z, w, x, y).$

$$\diamond \ \xi(y,x,z,w) = -\xi(x,y,z,w).$$

(4.1) 
$$\langle v_1, v_2 \rangle_{\mathrm{en}} \coloneqq \int_{\Gamma \times \Gamma} j_q(x, y) \mathrm{d} v_1(x) \mathrm{d} v_2(y) .$$

*Example 4.3* The following identities will be useful for our computations. It follows from  $\xi_x(q, x, q, y) = \xi_y(q, x, q, y)$  that

(4.2) 
$$r(x,q) - r(y,q) = j_x(y,q) - j_y(x,q).$$

It follows from  $\xi_x(x, y, x, q) = \xi_q(x, y, x, q)$  that

(4.3) 
$$j_x(y,q) = j_q(x,x) - j_q(x,y).$$

It follows from  $\xi_x(x, y, x, q) = \xi_y(x, y, x, q)$  that

(4.4) 
$$r(x, y) = j_x(y, q) + j_y(x, q).$$

It follows from  $\xi_x(x, y, x, y) = \xi_q(x, y, x, y)$  that

(4.5) 
$$r(x, y) = j_q(x, x) + j_q(y, y) - 2j_q(x, y).$$

#### 4.4 **Projections**

Let *G* be a weighted graph. Fix an orientation O. Let *T* be a spanning tree of *G*. The weight of *T* is the product  $w(T) \coloneqq \prod_{e \notin T} \ell(e)$ . The coweight of *T* is the product  $w'(T) \coloneqq \prod_{e \in T} \ell^{-1}(e)$ . The weight and coweight of *G* are  $w(G) \coloneqq \sum_T w(T)$  and  $w'(G) \coloneqq \sum_T w'(T)$ , where the sums are over all spanning trees of *G*. The quantity w(G) depends only on the underlying metric graph  $\Gamma$ .

Let  $\mathbf{M}_T$  be the  $m \times m$  matrix whose columns are obtained from 1-chains circ(T, e) associated to fundamental circuits of T, and let  $\mathbf{N}_T$  be the  $m \times m$  matrix whose columns are obtained from 1-chains cocirc(T, e) associated to fundamental cocircuits of T (see [16, Section 7.2]). Consider the following matrix averages:

$$\mathbf{P} = \sum_{T} \frac{w(T)}{w(G)} \mathbf{M}_{T}, \quad \mathbf{P}' = \sum_{T} \frac{w'(T)}{w'(G)} \mathbf{N}_{T},$$

the sums being over all spanning trees *T* of *G*. It is a classical theorem of Kirchhoff [22] that the matrix of  $\pi$ :  $C_1(G, \mathbb{R}) \twoheadrightarrow H_1(G, \mathbb{R})$ , with respect to  $\mathcal{O}$ , is **P**. Similarly, the matrix of  $\pi'$ :  $C_1(G, \mathbb{R}) \twoheadrightarrow H_1(G, \mathbb{R})^{\perp}$ , with respect to  $\mathcal{O}$ , is  $(\mathbf{P}')^{\mathrm{T}}$ .

Let  $\Xi$  be the  $m \times m$  matrix of cross ratios:

$$\Xi := (\xi(e^{-}, e^{+}, f^{-}, f^{+}))_{e, f \in \mathcal{O}}.$$

Let **L** be any generalized inverse of **Q** (i.e., **QLQ** = **Q**). Then we have  $\Xi = \mathbf{B}^{\mathrm{T}}\mathbf{LB}$ . It is shown in [16, Theorem 7.5] that the matrix of  $\pi: C_1(G, \mathbb{R}) \twoheadrightarrow H_1(G, \mathbb{R})$ , with respect to  $\mathcal{O}$ , is  $\mathbf{I} - \mathbf{D}^{-1}\Xi$ , and the matrix of  $\pi': C_1(G, \mathbb{R}) \twoheadrightarrow H_1(G, \mathbb{R})^{\perp}$ , with respect to  $\mathcal{O}$ , is  $\mathbf{D}^{-1}\Xi$ . In particular, for each  $f \in \mathcal{O}$ , we have

(4.6) 
$$\mathsf{F}(e,f) \coloneqq \begin{cases} 1 - r(e^{-}, e^{+})/\ell(e), & \text{if } e = f, \\ -\xi(e^{-}, e^{+}, f^{-}, f^{+})/\ell(e), & \text{if } e \neq f. \end{cases}$$

$$\mathsf{F}'(e,f) = \xi(e^-, e^+, f^-, f^+)/\ell(e).$$

Moreover, we have equalities:

$$\mathbf{P} = \mathbf{I} - \mathbf{D}^{-1} \Xi, \quad \mathbf{P}' = \Xi \mathbf{D}^{-1}.$$

**Definition 4.4** The Foster coefficient of  $e \in \mathbb{E}(G)$  is, by definition,

$$F(e) := F(e, e) = 1 - \frac{r(e^{-}, e^{+})}{\ell(e)}.$$

Clearly,  $F(e) = F(\bar{e})$ , so F(e) is also well defined for  $e \in E(G)$ .

#### Remark 4.5

(i) It follows from (4.7) that

$$\mathsf{F}(e) = \sum_{T \neq e} \frac{w(T)}{w(G)},$$

the sum being over all spanning trees T of G not containing e.

(ii) It is a consequence of "Rayleigh's monotonicity law" that  $0 \le F(e) < 1$ , and the equality F(e) = 0 holds if and only if *e* is a bridge. (See [16] and the references therein for more details.)

Fix an arbitrary path  $\gamma$  from  $\gamma$  to x. Let  $\gamma_{\gamma x}$  denote the associated 1-chain. Then, by [16, Corollary 7.10], we have

(4.8) 
$$r(x, y) = [\boldsymbol{\gamma}_{yx}, \pi'(\boldsymbol{\gamma}_{yx})].$$

*Example 4.6* The following observation will be useful for computations. Let  $e = \{u, v\}$  denote an edge segment in a metric graph  $\Gamma$ , and let  $p \in e$  be a point with distance *x* from *u* and distance  $\ell(e) - x$  from *v*. Then, for each point  $q \in \Gamma$ , we have

$$r(p,q) = \frac{\ell(e) - x}{\ell(e)} r(u,q) + \frac{x}{\ell(e)} r(v,q) + \mathsf{F}(e) \frac{(\ell(e) - x)x}{\ell(e)}$$

This follows, for example, by a direct computation using (4.8). We leave the details to the interested reader.

#### 4.5 Generalized Rayleigh's laws

We will need the following two corollaries of [16, Theorem B]. Let  $\Gamma$  be a metric graph. Let *e* be an edge segment of  $\Gamma$  with boundary points  $\partial e = \{e^-, e^+\}$ . Let  $\Gamma/e$  denote the metric graph obtained by contracting *e* (equivalently, by setting  $\ell(e) = 0$ ). Then

(4.9) 
$$j_z(x, y; \Gamma/e) = j_z(x, y; \Gamma) - \frac{\xi(x, z, e^-, e^+; \Gamma) \,\xi(y, z, e^-, e^+; \Gamma)}{r(e^-, e^+; \Gamma)},$$

and

(4.10) 
$$r(x, y; \Gamma/e) = r(x, y; \Gamma) - \frac{\xi(x, y, e^{-}, e^{+}; \Gamma)^{2}}{r(e^{-}, e^{+}; \Gamma)}.$$

Note that (4.10) is, in fact, a special case of (4.9).

#### 4.6 Contractions and models

Let *G* be a weighted graph, and let  $e \in E(G)$  be an edge of *G*. We denote by G/e the weighted graph obtained from *G* by contracting the edge *e* and removing all loops that might be created in the process. Assume that *G* is a model of the metric graph  $\Gamma$ . In particular, we may view *e* as an edge segment of  $\Gamma$ . We then observe that, for  $x, y, z, w \in V(G)$ , the cross ratio  $\xi(x, y, z, w; \Gamma/e)$  measured on  $\Gamma/e$  is equal to the cross ratio  $\xi(x, y, z, w; G/e)$  measured on (the metric graph canonically associated to) G/e. A similar remark pertains to the *j*-function  $j_z(x, y; \Gamma/e)$  and the effective resistance function  $r(x, y; \Gamma/e)$ . We leave the details to the reader.

# 5 Calculus of random spanning trees

In this section, we study two functions arising from random spanning trees.

**Definition 5.1** Let *G* be a model of a metric graph  $\Gamma$ . For edges  $e = \{e^-, e^+\}$  and  $f = \{f^-, f^+\}$  of *G*, we define

$$P(e, f) := \begin{cases} r(e^-, e^+; G)/\ell(e), & \text{if } e = f, \\ r(e^-, e^+; G)/\ell(e) \times r(f^-, f^+; G/e)/\ell(f), & \text{if } e \neq f. \end{cases}$$

We use the notation P(e) := P(e, e). If we want to clarify the underlying model *G*, we use the notations P(e, f; G) and P(e; G).

By Definition 4.4 and Remark 4.5(i), we know

(5.1) 
$$P(e) = 1 - F(e) = \sum_{T \ni e} \frac{w(T)}{w(G)},$$

the sum being over all spanning trees *T* of *G* containing *e*. So P(e) is the probability of *e* being present in a *random spanning tree*, where a spanning tree *T* is chosen with probability w(T)/w(G).

A similar probabilistic interpretation holds for P(e, f) when  $e \neq f$ . Namely, since P(e, f) = P(e; G)P(f; G/e), it represents the probability of both *e* and *f* being present in a random spanning tree. In other words,

(5.2) 
$$P(e,f) = \sum_{T \ni e,f} \frac{w(T)}{w(G)}.$$

It follows that P(e, f) = P(f, e). One can use (4.10) (alternatively, the "transfercurrent theorem"—see [24, Section 4.2]) to compute P(e, f) directly in terms of invariants of *G*.

#### Definition 5.2

(i) Let  $\mathfrak{s}: V(G) \to \mathbb{R}$  be the function defined by sending  $p \in V(G)$  to

$$\mathfrak{s}(p) \coloneqq \sum_{e = \{p, x\}} \mathrm{P}(e),$$

the sum being over all edges *e* incident to *p* in *G* (i.e., the *star* of *p*). (ii) Let  $\mathfrak{t}: V(G) \times E(G) \to \mathbb{R}$  be the function defined by sending (p, e) to

$$\mathfrak{t}(p,e) \coloneqq \sum_{f=\{p,x\}} \mathbb{P}(e,f),$$

the sum being over all edges *f* incident to *p* in *G*.

**Proposition 5.3** Fix a vertex  $q \in V(G)$ . We have

$$\mathfrak{s}(p) = \sum_{e=\{p,x\}} \frac{r(x,q) - r(p,q)}{\ell(e)} + 2 - 2\,\delta_q(p),$$

the sum being over all edges  $e \in E(G)$  incident to p.



*Figure 3*: Contracting the edge  $e = \{u, v\}$ .

**Proof** By (4.5), we may write  $r(p, x) = j_q(x, x) - j_q(p, p) + 2(j_q(p, p) - j_q(x, p))$ . Therefore,

$$\begin{split} \mathfrak{s}(p) &= \sum_{e = \{p, x\}} \frac{r(p, x)}{\ell(e)} \\ &= \sum_{e = \{p, x\}} \frac{j_q(x, x) - j_q(p, p)}{\ell(e)} + 2 \sum_{e = \{p, x\}} \frac{j_q(p, p) - j_q(x, p)}{\ell(e)} \\ &= \sum_{e = \{p, x\}} \frac{j_q(x, x) - j_q(p, p)}{\ell(e)} + 2\Delta \left( j_q(\cdot, p) \right) (p) \\ &= \sum_{e = \{p, x\}} \frac{j_q(x, x) - j_q(p, p)}{\ell(e)} + 2(\delta_p(p) - \delta_q(p)). \end{split}$$

Recall from Section 4.6 the weighted graph G/e obtained by contracting the edge  $e = \{u, v\} \in E(G)$  and removing all loops that might be created in the process. Let  $Par(e) \subseteq E(G)$  denote the set of all edges *parallel* to *e* (i.e., connecting *u* and *v*). We make the identification  $E(G/e) = E(G) \setminus Par(e)$ , and the two vertices  $u, v \in V(G)$  will be identified with a single vertex  $v_e \in V(G/e)$  (see Figure 3), and  $V(G) \setminus \{u, v\} = V(G/e) \setminus \{v_e\}$ .

**Theorem 5.4** Fix a vertex  $q \in V(G)$ . Let  $p \in V(G)$  and  $e = \{u, v\} \in E(G)$ . Then

$$t(p,e) = \frac{r(u,v;G)}{\ell(e)} \sum_{\substack{f = \{p,x\}\\ f \in E(G) \setminus Par(e)}} \frac{r(x,q;G/e) - r(p,q;G/e)}{\ell(f)} + 2\frac{r(u,v;G)}{\ell(e)} \left(1 - \delta_q(p) - \frac{\xi(u,v,u,q;G)}{r(u,v;G)} \delta_u(p) - \frac{\xi(v,u,v,q;G)}{r(u,v;G)} \delta_v(p)\right).$$

The sum is over all edges  $f \in E(G) \setminus Par(e)$  incident to p in G.

Proof By definition,

$$\mathbf{P}(e,f) = \frac{r(u,v;G)}{\ell(e)}\mathbf{P}(f;G/e).$$

If  $p \notin \{u, v\}$ , the result follows immediately from Proposition 5.3:

$$\mathfrak{t}(p,e) = \frac{r(u,v;G)}{\ell(e)} \left( \sum_{\substack{f = \{p,x\}\\f \in E(G)}} \frac{r(x,q;G/e) - r(p,q;G/e)}{\ell(e)} + 2 - 2\,\delta_q(p) \right),$$

the sum being over all edges  $f \in E(G)$  incident to p in G (equivalently, all edges  $f \in E(G/e)$  incident to p in G/e).

So, by symmetry, it remains to show the equality for p = u. Recall that we denote the vertex obtained by identifying u and v in G/e by  $v_e$  (Figure 3). As in the proof of Proposition 5.3, we first compute

$$t(u, e) = \frac{r(u, v; G)}{\ell(e)} \left( \sum_{\substack{f = \{u, x\}\\f \in E(G) \setminus Par(e)}} \frac{j_q(x, x; G/e) - j_q(v_e, v_e; G/e)}{\ell(f)} \right)$$
(5.3)
$$+ 2 \frac{r(u, v; G)}{\ell(e)} \left( \sum_{\substack{f = \{u, x\}\\f \in E(G) \setminus Par(e)}} \frac{j_q(v_e, v_e; G/e) - j_q(x, v_e; G/e)}{\ell(f)} \right),$$

where the sums are over all edges  $f \in E(G) \setminus Par(e)$  incident to u in G. Unlike in the proof of Proposition 5.3, we *cannot* interpret the second sum as the Laplacian of the *j*-function on G/e because the summation is not over all edges incident to  $v_e$  in G/e (e.g., in Figure 3, only edges on the left of  $v_e$  appear in the summation). We proceed by "lifting" the problem to G using generalized Rayleigh's laws (4.9) and (4.10). We find

$$\frac{j_q(v_e, v_e; G/e) - j_q(x, v_e; G/e)}{\ell(f)} = \frac{j_q(u, u; G) - j_q(x, u; G)}{\ell(f)} + \frac{\xi(u, v, u, q; G)}{r(u, v; G)} \left(\frac{\xi(u, v, x, q; G) - \xi(u, v, u, q; G)}{\ell(f)}\right).$$

It is easily checked that the right-hand side is zero for x = v. Moreover, by the definition of cross ratios, we compute

$$\xi(u,v,x,q;G) - \xi(u,v,u,q;G) = \xi(u,v,x,u;G).$$

So we have

(5.4)

$$\sum_{\substack{f = \{u, x\}\\f \in E(G) \setminus Par(e)}} \frac{j_q(v_e, v_e; G/e) - j_q(x, v_e; G/e)}{\ell(f)} = \sum_{\substack{f = \{u, x\}\\f \in E(G)}} \frac{j_q(u, u; G) - j_q(x, u; G)}{\ell(f)}$$
$$+ \frac{\xi(u, v, u, q; G)}{r(u, v; G)} \sum_{\substack{f = \{u, x\}\\f \in E(G)}} \frac{\xi(u, v, x, u; G)}{\ell(f)}.$$

As in the proof of Proposition 5.3, we have

(5.5) 
$$\sum_{\substack{f=\{u,x\}\\f\in E(G)}} \frac{j_q(u,u;G) - j_q(x,u;G)}{\ell(f)} = \Delta(j_q(\cdot,u))(u)$$
$$= \delta_u(u) - \delta_q(u) = 1 - \delta_q(u).$$

By expanding with respect to the base point *q*, we also compute

(5.6)  

$$\sum_{\substack{f = \{u, x\} \\ f \in E(G)}} \frac{\xi(u, v, x, u; G)}{\ell(f)} = \sum_{\substack{f = \{u, x\} \\ f \in E(G)}} \frac{j_q(v, u; G) - j_q(v, x; G)}{\ell(f)}$$

$$- \sum_{\substack{f = \{u, x\} \\ f \in E(G)}} \frac{j_q(u, u; G) - j_q(u, x; G)}{\ell(f)}$$

$$= \Delta(j_q(v, \cdot))(u) - \Delta(j_q(u, \cdot))(u)$$

$$= (\delta_v(u) - \delta_q(u)) - (\delta_u(u) - \delta_q(u))$$

$$= -1.$$

The result for p = u follows by putting together (5.3)–(5.6).

# 6 Energy levels of rooted spanning trees

In this section, we prove a very subtle identity for cross ratios (Theorem 6.3).

#### 6.1 Energy levels

Recall that a rooted spanning tree (T, q) of *G* comes with a preferred orientation  $\mathcal{T}_q \subseteq \mathbb{E}(G)$ , where all edges are oriented *away from q* on the spanning tree *T* (see Section 3.1).

**Definition 6.1** We define the *energy level* of a rooted spanning tree (T, q) to be

$$\boldsymbol{\rho}(T,q) \coloneqq \sum_{e,f\in\mathbb{T}_q} \xi(e^-,e^+,f^-,f^+).$$

The following result is a justification for our terminology, and is useful in our later computation. Let  $\deg_T(v)$  denote the number of edges incident with  $v \in V(G)$  in the spanning tree *T*. Consider the canonical element  $v_{(T,q)} \in DMeas_0(G)$  associated to the rooted spanning tree (T, q) of *G* defined by

$$\nu_{(T,q)} = \sum_{\nu \in V(G)} \left( \deg_T(\nu) - 2 \right) \delta_{\nu} + 2\delta_q.$$

*Lemma 6.2 We have* 

(6.1)

$$\rho(T,q) = \langle v_{(T,q)}, v_{(T,q)} \rangle_{\text{en}} = \sum_{\nu, w \in V(G)} \left( \deg_T(\nu) - 2 \right) \left( \deg_T(w) - 2 \right) j_q(\nu, w).$$

**Proof** By the definition of the preferred orientation  $\mathcal{T}_q \subseteq \mathbb{E}(G)$ , we have

$$\sum_{e\in \mathfrak{T}_q} \left( \delta_{e^-} - \delta_{e^+} \right) = v_{(T,q)}.$$

This is because every vertex  $v \neq q$  has exactly one incoming edge and  $\deg_T(v) - 1$  outgoing edges in  $\mathcal{T}_q$ . At q, however, there is no incoming edge in  $\mathcal{T}_q$ .

The result now follows directly from bilinearity of the energy pairing:

$$\rho(T,q) = \sum_{e,f\in\mathcal{T}_q} \xi(e^-, e^+, f^-, f^+) = \sum_{e,f\in\mathcal{T}_q} \langle \delta_{e^-} - \delta_{e^+}, \delta_{f^-} - \delta_{f^+} \rangle_{\mathrm{en}}$$
$$= \langle \sum_{e\in\mathcal{T}_q} (\delta_{e^-} - \delta_{e^+}), \sum_{f\in\mathcal{T}_q} (\delta_{f^-} - \delta_{f^+}) \rangle_{\mathrm{en}} = \langle v_{(T,q)}, v_{(T,q)} \rangle_{\mathrm{en}}.$$

The second equality in (6.1) follows from (4.1), because  $j_q(\cdot, q) = j_q(q, \cdot) = 0$ .

### 6.2 The average of energy levels

We can now state the most technical ingredient in computing tropical moments of tropical Jacobians.

**Theorem 6.3** Fix a vertex  $q \in V(G)$ . We have the equality

$$\frac{1}{w(G)}\sum_{T}w(T)\rho(T,q) = \sum_{e=\{u,v\}}\frac{j_{u}(v,q)^{2}+j_{v}(u,q)^{2}}{\ell(e)},$$

where the sum on the left is over all spanning trees *T* of *G*, and the sum on the right is over all edges  $e \in E(G)$ .

**Proof** The result follows by putting together Lemmas 6.5–6.7.

*Lemma* 6.4 *Let* (T, q) *be a rooted spanning tree of G. We have the equality* 

$$\begin{split} \rho(T,q) &= 4 \sum_{x,y \in V(G)} j_q(x,y) - 4 \sum_{e \in T} \sum_{x \in V(G)} \left( j_q(e^-,x) + j_q(e^+,x) \right) \\ &+ \sum_{e,f \in T} \left( j_q(e^-,f^-) + j_q(e^+,f^+) + j_q(e^-,f^+) + j_q(e^+,f^-) \right). \end{split}$$

**Proof** The result follows from Lemma 6.2, and an application of a (generalized) *handshaking lemma*: for each  $\psi \in \mathcal{M}(G)$ , we have

$$\sum_{x \in V(G)} \deg_T(x) \psi(x) = \sum_{e \in T} \left( \psi(e^-) + \psi(e^+) \right).$$

*Lemma 6.5* Fix a vertex  $q \in V(G)$ . We have the equality

$$\frac{1}{w(G)} \sum_{T} w(T) \rho(T,q) = 4 \sum_{x,y \in V(G)} j_q(x,y) - 4 \sum_{e \in E(G)} \sum_{x \in V(G)} \left( j_q(e^-,x) + j_q(e^+,x) \right) P(e)$$
  
+ 
$$\sum_{e,f \in E(G)} \left( j_q(e^-,f^-) + j_q(e^+,f^+) + j_q(e^-,f^+) + j_q(e^+,f^-) \right) P(e,f).$$

**Proof** This follows from Lemma 6.4: one only needs to change the order of summations and use (5.1) and (5.2).

To prove the next two results, it is convenient to work with matrices. We fix labelings  $V(G) = \{v_1, ..., v_n\}$  and  $E(G) = \{e_1, ..., e_m\}$ , and we introduce the following matrices:

- ♦  $\mathbf{1}_n \in \mathbb{R}^n$  and  $\mathbf{1}_m \in \mathbb{R}^m$  denote the all-ones vectors.
- $\delta_{v} \in \mathbb{R}^{n}$  (resp.  $\delta_{e} \in \mathbb{R}^{m}$ ) denotes the characteristic vector of  $v \in V(G)$  (resp.  $e \in E(G)$ ).
- ♦ **A** =  $(h_{ij})$  denotes the *n* × *m* unsigned incidence matrix of *G*, where  $h_{ij} = 1$  if  $e_j^+ = v_i$  or  $e_j^- = v_i$ , and  $h_{ij} = 0$  otherwise.
- ♦ **X** is the *n* × *n* diagonal matrix with diagonal (*i*, *i*)-entries  $j_q(v_i, v_i) = r(v_i, q)$ .
- ♦ **Y** is the *m* × *m* diagonal matrix with diagonal (i, i)-entries P $(e_i)$ .
- ◇ **Z** is the *m* × *m* matrix whose diagonal entries are zero, and the (i, j)-entries  $(i \neq j)$  are P $(e_i, e_j)$ .

We will also use the matrices  $\mathbf{Q}$ ,  $\mathbf{L}_q$ ,  $\mathbf{R}_q$ , and  $\mathbf{I}$  introduced in Section 4.2.

*Lemma 6.6* We have the equality

(6.2)  
$$\sum_{e \in E(G)} \sum_{x \in V(G)} \left( j_q(e^-, x) + j_q(e^+, x) \right) \mathbb{P}(e) = 2 \sum_{x, y \in V(G)} j_q(x, y) - \sum_{x \in V(G)} j_q(x, x).$$

**Proof** We start by writing the left-hand side of (6.2) in terms of our matrices:

$$\sum_{e \in E(G)} \sum_{x \in V(G)} \left( j_q(e^-, x) + j_q(e^+, x) \right) \mathbf{P}(e) = \mathbf{1}_n^{\mathrm{T}} \mathbf{L}_q \mathbf{A} \mathbf{Y} \mathbf{1}_m$$

By definitions, we observe  $\mathbf{AYI}_m = (\mathfrak{s}(v_1), \dots, \mathfrak{s}(v_n))^{\mathrm{T}}$ . So, by Proposition 5.3, we have

$$\mathbf{AY1}_m = -\mathbf{QX1}_n + 2\left(\mathbf{1}_n - \boldsymbol{\delta}_q\right).$$

Therefore,

$$\mathbf{1}_{n}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{A}\mathbf{Y}\mathbf{1}_{m} = -\mathbf{1}_{n}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{Q}\mathbf{X}\mathbf{1}_{n} + 2\mathbf{1}_{n}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{1}_{n} - 2\mathbf{1}_{n}^{\mathrm{T}}\mathbf{L}_{q}\boldsymbol{\delta}_{q}.$$

Note that  $\mathbf{L}_q \boldsymbol{\delta}_q = \mathbf{0}$  and  $\mathbf{L}_q \mathbf{Q} = \mathbf{I} + \mathbf{R}_q^{\mathrm{T}}$  (see Section 4.2). We also have  $\mathbf{R}_q^{\mathrm{T}} \mathbf{X} = \mathbf{0}$ . Therefore,

$$\mathbf{1}_{n}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{A}\mathbf{Y}\mathbf{1}_{m} = 2\mathbf{1}_{n}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{1}_{n} - \mathbf{1}_{n}^{\mathrm{T}}\mathbf{X}\mathbf{1}_{n},$$

which is the right-hand side of (6.2).

For our next computation, it is convenient to use the notion of *Hadamard–Schur* products of matrices: for two  $k \times m$  matrices A, B, the Hadamard–Schur product, denoted by  $A \circ B$ , is a matrix of the same dimension as A and B with entries given by  $(A \circ B)_{ii} = (A)_{ii}(B)_{ii}$ .

One useful (and easy to prove) fact about Hadamard-Schur products is the following:

(6.3) 
$$\mathbf{1}_{k}^{\mathrm{T}}\left(A\circ B\right)\mathbf{1}_{m}=\mathrm{Trace}\left(A^{\mathrm{T}}B\right).$$

*Lemma 6.7* We have the equality

$$\sum_{e,f \in E(G)} \left( j_q(e^-, f^-) + j_q(e^+, f^+) + j_q(e^-, f^+) + j_q(e^+, f^-) \right) \mathbb{P}(e, f)$$
$$= \sum_{e = \{u,v\}} \frac{j_u(v, q)^2 + j_v(u, q)^2}{\ell(e)} + 4 \sum_{x,y \in V(G)} j_q(x, y) - 4 \sum_{x \in V(G)} j_q(x, x).$$

**Proof** We first write

(6.4)

$$\sum_{e,f\in E(G)} \left( j_q(e^-,f^-) + j_q(e^+,f^+) + j_q(e^-,f^+) + j_q(e^+,f^-) \right) \mathbb{P}(e,f) \\ = \sum_{e\in E(G)} \left( j_q(e^-,e^-) + j_q(e^+,e^+) + 2j_q(e^-,e^+) \right) \mathbb{P}(e) \\ + \sum_{\substack{e,f\in E(G)\\e\neq f}} \left( j_q(e^-,f^-) + j_q(e^+,f^+) + j_q(e^-,f^+) + j_q(e^+,f^-) \right) \mathbb{P}(e,f).$$

We write the second sum in (6.4) in terms of our matrices:

(6.5) 
$$\sum_{\substack{e,f \in E(G) \\ e \neq f}} \left( j_q(e^-, f^-) + j_q(e^+, f^+) + j_q(e^-, f^+) + j_q(e^+, f^-) \right) \mathbb{P}(e, f)$$
$$= \mathbf{1}_m^{\mathrm{T}} \left( \left( \mathbf{A}^{\mathrm{T}} \mathbf{L}_q \mathbf{A} \right) \circ \mathbf{Z} \right) \mathbf{1}_m.$$

By (6.3), we know

$$\mathbf{l}_{m}^{\mathrm{T}}\left(\left(\mathbf{A}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{A}\right)\circ\mathbf{Z}\right)\mathbf{l}_{m}=\operatorname{Trace}\left(\mathbf{A}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{A}\mathbf{Z}\right)=\sum_{e\in E(G)}\boldsymbol{\delta}_{e}^{\mathrm{T}}\left(\mathbf{A}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{A}\mathbf{Z}\right)\boldsymbol{\delta}_{e}.$$

By definitions, one observes  $\mathbf{AZ}\boldsymbol{\delta}_e = (\mathfrak{t}(v_1, e), \dots, \mathfrak{t}(v_n, e))^{\mathrm{T}}$ . Let

$$e = \{u, v\}$$
,  $\beta_u = \frac{\xi(u, v, u, q)}{r(u, v)}$ ,  $\beta_v = \frac{\xi(v, u, v, q)}{r(u, v)}$ ,  $P(e) = \frac{r(u, v)}{\ell(e)}$ .

Theorem 5.4 states

$$\mathbf{AZ}\boldsymbol{\delta}_{e}=\mathbf{P}(e)\left(-\mathbf{Q}\widetilde{\mathbf{X}}\mathbf{1}_{n}+2\left(\mathbf{1}_{n}-\boldsymbol{\delta}_{q}-\beta_{u}\boldsymbol{\delta}_{u}-\beta_{v}\boldsymbol{\delta}_{v}\right)\right).$$

Here,  $\widetilde{\mathbf{X}}$  is the  $n \times n$  diagonal matrix with diagonal (i, i)-entries  $j_q(v_i, v_i; G/e)$  for  $v_i \notin \{u, v\}$ . The diagonal entries corresponding to both u and v are  $j_q(v_e, v_e; G/e)$ . Thus, we find

$$\boldsymbol{\delta}_{e}^{\mathrm{T}}\left(\mathbf{A}^{\mathrm{T}}\mathbf{L}_{q}\mathbf{A}\mathbf{Z}\right)\boldsymbol{\delta}_{e}=\mathrm{P}(e)\left(\boldsymbol{\delta}_{u}+\boldsymbol{\delta}_{v}\right)^{\mathrm{T}}\mathbf{L}_{q}\left(-\mathbf{Q}\widetilde{\mathbf{X}}\mathbf{1}_{n}+2\left(\mathbf{1}_{n}-\boldsymbol{\delta}_{q}-\boldsymbol{\beta}_{u}\boldsymbol{\delta}_{u}-\boldsymbol{\beta}_{v}\boldsymbol{\delta}_{v}\right)\right).$$

Note that  $\mathbf{L}_q \boldsymbol{\delta}_q = \mathbf{0}, \mathbf{L}_q \mathbf{Q} = \mathbf{I} + \mathbf{R}_q^{\mathrm{T}}$ , and  $\mathbf{R}_q^{\mathrm{T}} \widetilde{\mathbf{X}} = \mathbf{0}$ . We obtain

(6.6)  

$$\boldsymbol{\delta}_{e}^{\mathrm{T}} \left( \mathbf{A}^{\mathrm{T}} \mathbf{L}_{q} \mathbf{A} \mathbf{Z} \right) \boldsymbol{\delta}_{e} = \mathrm{P}(e) \left( \boldsymbol{\delta}_{u} + \boldsymbol{\delta}_{v} \right)^{\mathrm{T}} \left( -\widetilde{\mathbf{X}} \mathbf{1}_{n} + 2 \mathbf{L}_{q} \left( \mathbf{1}_{n} - \beta_{u} \boldsymbol{\delta}_{u} - \beta_{v} \boldsymbol{\delta}_{v} \right) \right)$$

$$= -\mathrm{P}(e) \left( j_{q} (v_{e}, v_{e}; G/e) + j_{q} (v_{e}, v_{e}; G/e) \right) + 2 \mathrm{P}(e) \sum_{x \in V(G)} \left( j_{q} (u, x) + j_{q} (v, x) \right)$$

$$- 2 \left( \frac{\xi(u, v, u, q)}{\ell(e)} \left( j_{q} (u, u) + j_{q} (v, u) \right) + \frac{\xi(v, u, v, q)}{\ell(e)} \left( j_{q} (u, v) + j_{q} (v, v) \right) \right).$$

We now use our generalized Rayleigh's law (4.10) twice and write everything in terms of invariants of G:

(6.7)  
$$j_q(v_e, v_e; G/e) = j_q(u, u) - \frac{\xi(u, q, u, v)^2}{r(u, v)} = j_q(u, u) - \frac{j_u(v, q)^2}{r(u, v)},$$
$$j_q(v_e, v_e; G/e) = j_q(v, v) - \frac{\xi(v, q, u, v)^2}{r(u, v)} = j_q(v, v) - \frac{j_v(u, q)^2}{r(u, v)}.$$

We also use the definition of cross ratios to compute

(6.8) 
$$\xi(u, v, u, q) = j_q(u, u) - j_q(u, v)$$
,  $\xi(v, u, v, q) = j_q(v, v) - j_q(u, v)$ .

Putting together (6.4)–(6.8), with a simple computation, we obtain

(6.9)  

$$\sum_{e,f\in E(G)} \left( j_q(e^-, f^-) + j_q(e^+, f^+) + j_q(e^-, f^+) + j_q(e^+, f^-) \right) P(e, f)$$

$$= \sum_{e=\{u,v\}} \frac{j_u(v,q)^2 + j_v(u,q)^2}{\ell(e)} + 2 \sum_{e=\{u,v\}} \sum_{x\in V(G)} P(e) \left( j_q(u,x) + j_q(v,x) \right)$$

$$+ 2 \sum_{e=\{u,v\}} P(e) j_q(u,v) - 2 \sum_{e=\{u,v\}} \frac{j_q(u,u)^2 + j_q(v,v)^2 - 2j_q(u,v)^2}{\ell(e)}.$$

By Lemma 6.6, the second term in (6.9) is simplified as

(6.10)  

$$\sum_{e=\{u,v\}} \sum_{x \in V(G)} \mathbb{P}(e) \left( j_q(u,x) + j_q(v,x) \right) = 2 \sum_{x,y \in V(G)} j_q(x,y) - \sum_{x \in V(G)} j_q(x,x).$$

The third and fourth terms in (6.9) are simplified as follows:

$$\sum_{e=\{u,v\}} P(e)j_q(u,v) - \sum_{e=\{u,v\}} \frac{j_q(u,u)^2 + j_q(v,v)^2 - 2j_q(u,v)^2}{\ell(e)}$$
$$= \sum_{e=\{u,v\}} \left( \frac{j_q(u,u) + j_q(v,v) - 2j_q(u,v)}{\ell(e)} j_q(u,v) - \frac{j_q(u,u)^2 + j_q(v,v)^2 - 2j_q(u,v)^2}{\ell(e)} \right)$$

R. de Jong and F. Shokrieh

$$\begin{split} &= -\sum_{e=\{u,v\}} \left( j_q(u,u) \frac{j_q(u,u) - j_q(u,v)}{\ell(e)} + j_q(v,v) \frac{j_q(v,v) - j_q(u,v)}{\ell(e)} \right) \\ &= -\sum_{x \in V(G)} j_q(x,x) \sum_{f=\{x,y\}} \frac{j_q(x,x) - j_q(x,y)}{\ell(f)} = -\sum_{x \in V(G)} j_q(x,x) \Delta(j_q(x,\cdot))(x) \\ &= -\sum_{x \in V(G)} j_q(x,x) \left( \delta_x(x) - \delta_q(x) \right) = -\sum_{x \in V(G)} j_q(x,x). \end{split}$$

For the first equality, we used (4.5). The third equality is by a (generalized) handshaking lemma. The result now follows by putting together (6.9)-(6.11).

#### 6.3 Average of energy levels, a variation

For our main application, we will need the following slight variation of Theorem 6.3. Let  $\pi: C_1(G, \mathbb{R}) \twoheadrightarrow H_1(G, \mathbb{R})$  denote the orthogonal projection (as defined in Section 4.4).

**Definition 6.8** We define the *center* of a rooted spanning tree (T, q) to be

$$\sigma_T \coloneqq \frac{1}{2} \sum_{e \in \mathcal{T}_q} \pi(e).$$

*Theorem 6.9* We have the equality

$$\frac{1}{w(G)}\sum_{T}w(T)[\sigma_{T},\sigma_{T}] = \frac{1}{4}\sum_{e=\{u,v\}}\left(r(u,v) - \frac{j_{u}(v,q)^{2} + j_{v}(u,q)^{2}}{\ell(e)}\right),$$

where the sum on the left is over all spanning trees *T* of *G*, and the sum on the right is over all edges  $e \in E(G)$ .

**Proof** Using (4.6) and Definition 6.1, we compute (6.12)

$$\begin{split} & \left[\sigma_{T}, \sigma_{T}\right] = \frac{1}{4} \left[\sum_{e \in \mathcal{T}_{q}} \pi(e), \sum_{e \in \mathcal{T}_{q}} \pi(e)\right] = \frac{1}{4} \left(\sum_{e \in \mathcal{T}_{q}} \left[\pi(e), \pi(e)\right] + \sum_{\substack{e, f \in \mathcal{T}_{q} \\ e \neq f}} \left[\pi(e), \pi(f)\right]\right) \\ & = \frac{1}{4} \left(\sum_{e \in T} \mathsf{F}(e)\ell(e) + \sum_{\substack{e, f \in \mathcal{T}_{q} \\ e \neq f}} \mathsf{F}(e, f)\ell(e)\right) = \frac{1}{4} \left(\sum_{e \in T} \mathsf{F}(e)\ell(e) - \sum_{\substack{e, f \in \mathcal{T}_{q} \\ e \neq f}} \xi(e^{-}, e^{+}, f^{-}, f^{+})\right) \\ & = \frac{1}{4} \left(\sum_{e \in T} \mathsf{F}(e)\ell(e) + \sum_{e \in \mathcal{T}_{q}} r(e^{-}, e^{+}) - \sum_{e, f \in \mathcal{T}_{q}} \xi(e^{-}, e^{+}, f^{-}, f^{+})\right) \\ & = \frac{1}{4} \sum_{\substack{e \in T \\ e = \{u, v\}}} \left(\mathsf{F}(e)\ell(e) + r(u, v)\right) - \frac{1}{4}\rho(T, q). \end{split}$$

By Definition 4.4 and (5.1) and by changing the order of summations, we compute

(6.13)  

$$\sum_{T} \frac{w(T)}{w(G)} \sum_{\substack{e \in T \\ e = \{u,v\}}} (\mathsf{F}(e)\ell(e) + r(u,v)) = \sum_{e = \{u,v\}} (\mathsf{F}(e)\ell(e) + r(u,v)) \sum_{T \ni e} \frac{w(T)}{w(G)}$$

$$= \sum_{e = \{u,v\}} (\mathsf{F}(e)\ell(e) + r(u,v)) \operatorname{P}(e) = \sum_{e = \{u,v\}} r(u,v).$$

The result now follows from (6.12), (6.13), and Theorem 6.3.

# 7 Combinatorics of Voronoi polytopes

Throughout this section, we fix a metric graph  $\Gamma$  and a model *G*. We are interested in the combinatorics of the lattice  $(H_1(G,\mathbb{Z}), [\cdot, \cdot])$ . More specifically, we study the combinatorics of the Voronoi polytopes  $Vor(\lambda)$  (as defined in Section 2.2) for the lattice  $(H_1(G,\mathbb{Z}), [\cdot, \cdot])$ . Since  $Vor(\lambda) = Vor(0) + \lambda$  for all  $\lambda \in H_1(G,\mathbb{Z})$ , it suffices to understand the Voronoi polytope Vor(0) around the origin.

Let Vol (·) denote the volume measure induced by the bilinear form  $[\cdot, \cdot]$  on  $H_1(G, \mathbb{R})$ . Let w(G) be the weight of *G* (as in Section 4.4). Put  $g = \dim_{\mathbb{R}} H_1(G, \mathbb{R})$ .

Lemma 7.1  $\operatorname{Vol}(\operatorname{Vor}(0)) = \sqrt{w(G)}$ .

**Proof** The Voronoi polytopes  $\{Vor(0) + \lambda: \lambda \in H_1(G, \mathbb{Z})\}$  induce a periodic polytopal decomposition of  $H_1(G, \mathbb{R})$ . Therefore, Vor(0), up to some identifications on its boundary, gives a fundamental domain for the translation action of  $H_1(G, \mathbb{Z})$  on  $H_1(G, \mathbb{R})$ . Therefore,  $Vol(Vor(0)) = \sqrt{\det(\mathbf{G})}$  where **G** is any Gram matrix for the lattice  $(H_1(G, \mathbb{Z}), [\cdot, \cdot])$ .

Fix an orientation  $\mathcal{O}$  on G, and fix a spanning tree T of G. It is well known that  $\{\operatorname{circ}(T, e): e \in \mathcal{O} \setminus T\}$  is a basis for  $H_1(G, \mathbb{R})$ . Let  $\mathbf{C}_T$  denote the totally unimodular  $g \times m$  matrix whose rows correspond to these basis elements. Then a Gram matrix for the lattice  $(H_1(G, \mathbb{Z}), [\cdot, \cdot])$  is  $\mathbf{G}_T = \mathbf{C}_T \mathbf{D} \mathbf{C}_T^T$ . The result now follows from a standard application of the Cauchy–Binet formula for determinants.

*Remark 7.2* A geometric proof of Lemma 7.1 can be found in [2, Section 5].

Let  $\pi: C_1(G, \mathbb{R}) \twoheadrightarrow H_1(G, \mathbb{R})$  denote the orthogonal projection (as defined in Section 4.4). Each finite collection of 1-chains  $V = \{\mathbf{v}_1, \dots, \mathbf{v}_k\} \subset C_1(G, \mathbb{R})$  generates a *zonotope*  $\mathcal{Z}(V)$  defined as

$$\mathcal{Z}(V) = \left\{ \sum_{i=1}^{k} \alpha_i \mathbf{v}_i : -1 \le \alpha_i \le 1 \right\} \subset C_1(G, \mathbb{R}).$$

**Proposition 7.3** Fix an orientation O on G. Then

$$\operatorname{Vor}(0) = \frac{1}{2} \mathcal{Z}(\{\pi(e) : e \in \mathcal{O}\}).$$

**Proof** This is well known. To the best of our knowledge, this was first proved in [26, Proposition 5.2]. See also [17, Theorem 2] for a different proof. ■

Fix an orientation  $\bigcirc$  on *G*. Let *T* be a spanning tree of *G*. We define

$$\mathcal{C}_T \coloneqq \frac{1}{2} \mathcal{Z} \left( \left\{ \pi(e) \colon e \in \mathcal{O} \setminus T \right\} \right).$$

Note that  $\mathcal{C}_T$  is independent of the choice of  $\mathcal{O}$ .

#### Lemma 7.4

- (a)  $C_T$  is a g-dimensional parallelotope. Equivalently,  $\{\pi(e): e \in \mathbb{O} \setminus T\}$  is a basis for  $H_1(G, \mathbb{R})$ .
- (b)  $\operatorname{Vol}(\mathcal{C}_T) = w(T)/\sqrt{w(G)}$ .

**Proof** (a) Note that, for  $e, e' \in \mathbb{O} \setminus T$ , we have  $[\pi(e), \operatorname{circ}(T, e')] = \ell(e)\delta_e(e')$ . It is well known that  $\{\operatorname{circ}(T, e): e \in \mathbb{O} \setminus T\}$  is a basis for  $H_1(G, \mathbb{R})$ .

(b) This is proved in [2, Proposition 5.4].

Now, fix  $q \in V(G)$ . Recall (see Section 3.1) that the rooted spanning tree (T, q) comes with a preferred orientation  $\mathcal{T}_q \subseteq \mathbb{E}(G)$  (where edges are oriented away from q on the spanning tree T). Let  $\sigma_T := \frac{1}{2} \sum_{e \in \mathcal{T}_q} \pi(e)$  be the center of (T, q) (Definition 6.8). We now state and prove the main result of this section.

Theorem 7.5 The collection of parallelotopes

 $\{\sigma_T + \mathcal{C}_T: T \text{ is a spanning tree of } G\}$ 

induces a polytopal decomposition of Vor(0):

- (i)  $\operatorname{Vor}(0) = \bigcup_T (\sigma_T + \mathcal{C}_T)$ , the union being over all spanning trees T of G.
- (ii) If  $T \neq T'$  are two spanning trees such that

$$\mathcal{F} \coloneqq (\sigma_T + \mathcal{C}_T) \cap (\sigma_{T'} + \mathcal{C}_{T'})$$

*is nonempty, then*  $\mathcal{F}$  *is a face of both* ( $\sigma_T + \mathcal{C}_T$ ) *and* ( $\sigma_{T'} + \mathcal{C}_{T'}$ ).

**Proof** It follows from Proposition 7.3 that, for all spanning trees *T* of *G*, we have

(7.1) 
$$\sigma_T + \mathcal{C}_T \subseteq \operatorname{Vor}(0).$$

Moreover, by Lemmas 7.1 and 7.4(b), we know

(7.2) 
$$\sum_{T} \operatorname{Vol} \left( \sigma_{T} + \mathcal{C}_{T} \right) = \operatorname{Vol} \left( \operatorname{Vor}(0) \right),$$

the sum being over all spanning trees T of G.

Next, we describe how these parallelotopes can intersect each other. Let  $T \neq T'$  be two spanning trees. We choose an orientation  $\mathcal{O}$  on G that agrees with  $\mathcal{T}_q$  for  $e \in T$ , disagrees with  $\mathcal{T}'_q$  for  $e \in T' \setminus T$ , and is arbitrary outside  $T \cup T'$ . We may partition  $\mathcal{O}$ into the following (possibly empty) subsets:

$$\begin{split} \mathbb{S}_1 &\coloneqq \mathbb{T}_q \setminus \left( \mathbb{T}_q' \cup \overline{\mathbb{T}_q'} \right), \ \mathbb{S}_2 &\coloneqq \overline{\mathbb{T}_q'} \setminus \left( \mathbb{T}_q \cup \overline{\mathbb{T}_q} \right), \ \mathbb{S}_3 &\coloneqq \mathbb{T}_q \cap \overline{\mathbb{T}_q'} \ , \ \mathbb{S}_4 &\coloneqq \mathbb{T}_q \cap \mathbb{T}_q', \\ \mathbb{S}_5 &\coloneqq \mathbb{O} \setminus \left( \mathbb{S}_1 \cup \mathbb{S}_2 \cup \mathbb{S}_3 \cup \mathbb{S}_4 \right). \end{split}$$

We can now describe the corresponding parallelotopes as follows:

$$\sigma_{T} + \mathcal{C}_{T} = \frac{1}{2} \mathcal{Z} \left( \left\{ \pi(e) : e \in \mathcal{S}_{2} \cup \mathcal{S}_{5} \right\} \right) + \frac{1}{2} \sum_{e \in \mathcal{S}_{1} \cup \mathcal{S}_{3} \cup \mathcal{S}_{4}} \pi(e),$$
  
$$\sigma_{T'} + \mathcal{C}_{T'} = \frac{1}{2} \mathcal{Z} \left( \left\{ \pi(e) : e \in \mathcal{S}_{1} \cup \mathcal{S}_{5} \right\} \right) - \frac{1}{2} \sum_{e \in \mathcal{S}_{2} \cup \mathcal{S}_{3}} \pi(e) + \frac{1}{2} \sum_{e \in \mathcal{S}_{4}} \pi(e).$$

Consider the oriented subgraph  $\mathcal{D} := S_1 \cup S_2 \cup S_3$ . Note that the oriented subgraph  $\mathcal{D}$  is obtained from the edge set  $T \cup T'$  by deleting those edges in  $T \cap T'$  that have the same orientation in  $\mathcal{T}_q$  and in  $\mathcal{T}'_q$ , and by orienting the remaining edges according to  $\mathcal{O}$ . One can extend  $\mathcal{D}$  to a *totally cyclic* (also known as *strongly connected*) subgraph of *G*, after possibly adding some oriented edges to  $\mathcal{D}$  chosen from the (unoriented) edges of  $S_4$ . This is because  $\mathcal{D}$ , by construction, has no directed cocircuit. This implies that there exists a vector  $\mathbf{v} \in H_1(G, \mathbb{R})$  such that

$$[\mathbf{v}, \pi(e)] > 0, \qquad \text{if } e \in \mathbb{S}_1 \cup \mathbb{S}_2 \cup \mathbb{S}_3, \\ [\mathbf{v}, \pi(e)] = 0, \qquad \text{if } e \in \mathbb{S}_5.$$

To see this, consider the oriented (cographic) hyperplane arrangement with normal vectors  $\{\pi(e): e \in \mathcal{O}\}$  in the vector space  $H_1(G, \mathbb{R})$ . It is well known (see, e.g., [19, Lemma 8.2]) that there is a one-to-one correspondence between cells of this hyperplane arrangement and totally cyclic subgraphs of *G* (see also [1]). Our desired vector **v** is any vector in the cell corresponding to any totally cyclic extension of  $\mathcal{D}$ .

Consider the function  $\mathfrak{h}: H_1(G, \mathbb{R}) \to \mathbb{R}$  defined by

$$\mathfrak{h}(z) = [\mathbf{v}, z + \sum_{e \in \mathbb{S}_2} \frac{1}{2}\pi(e) - \sum_{e \in \mathbb{S}_1 \cup \mathbb{S}_4} \frac{1}{2}\pi(e)].$$

Let  $\mathcal{F} = (\sigma_T + \mathcal{C}_T) \cap (\sigma_{T'} + \mathcal{C}_{T'})$ . We compute

$$\mathfrak{h}(\sigma_{T} + \mathfrak{C}_{T}) = \{ 0 \le x \le \sum_{e \in \mathfrak{S}_{2}} [\mathbf{v}, \pi(e)] \} + \frac{1}{2} \sum_{e \in \mathfrak{S}_{3}} [\mathbf{v}, \pi(e)] \subseteq \mathbb{R}_{\ge 0},$$
$$\mathfrak{h}(\sigma_{T'} + \mathfrak{C}_{T'}) = \{ -\sum_{e \in \mathfrak{S}_{1}} [\mathbf{v}, \pi(e)] \le x \le 0 \} - \frac{1}{2} \sum_{e \in \mathfrak{S}_{3}} [\mathbf{v}, \pi(e)] \subseteq \mathbb{R}_{\le 0}.$$

- If  $S_3 \neq \emptyset$ , then  $\mathfrak{h}(\sigma_T + \mathfrak{C}_T) \cap \mathfrak{h}(\sigma_{T'} + \mathfrak{C}_{T'}) = \emptyset$  and, therefore,  $\mathfrak{F} = \emptyset$ .
- If  $S_3 = \emptyset$ , then  $\mathfrak{h}(\sigma_T + \mathfrak{C}_T) \cap \mathfrak{h}(\sigma_{T'} + \mathfrak{C}_{T'}) = \{0\}$ . Consider the hyperplane  $\mathcal{H} = \mathfrak{h}^{-1}(0)$  in  $H_1(G, \mathbb{R})$ . One computes

$$\begin{aligned} \mathcal{H} \cap (\sigma_T + \mathcal{C}_T) &= \mathcal{H} \cap (\sigma_{T'} + \mathcal{C}_{T'}) \\ &= \frac{1}{2} \mathcal{Z} \left( \left\{ \pi(e) : e \in \mathcal{S}_5 \right\} \right) + \frac{1}{2} \sum_{e \in \mathcal{S}_1 \cup \mathcal{S}_4} \pi(e) - \frac{1}{2} \sum_{e \in \mathcal{S}_2} \pi(e) \\ &= \mathcal{F}. \end{aligned}$$

The result now follows from these intersection patterns, together with (7.1) and (7.2).

# 8 The tropical moment of a tropical Jacobian

In this section, we prove our promised potential theoretic expression for the tropical moment of a tropical Jacobian.

**Theorem 8.1** Let  $\Gamma$  be a metric graph. Fix a model G of  $\Gamma$ , and fix a point  $q \in V(G)$ . Then

$$I(\operatorname{Jac}(\Gamma)) = \frac{1}{12} \sum_{e} \mathsf{F}(e)^{2} \ell(e) + \frac{1}{4} \sum_{e=\{u,v\}} \left( r(u,v) - \frac{j_{u}(v,q)^{2} + j_{v}(u,q)^{2}}{\ell(e)} \right),$$

where the sums are over all edges  $e \in E(G)$ .

Proof Recall

$$I(\operatorname{Jac}(\Gamma)) = \int_{\operatorname{Vor}(0)} [z, z] d\mu_L(z),$$

where  $\mu_L$  is the Lebesgue measure on  $H_1(G, \mathbb{R})$ , normalized to have  $\mu_L(Vor(0)) = 1$ . By Theorem 7.5, we obtain

(8.1) 
$$I(\operatorname{Jac}(\Gamma)) = \sum_{T} \int_{\sigma_{T}+\mathcal{C}_{T}} [z,z] \, \mathrm{d}\mu_{L}(z),$$

the sums being over all spanning trees T of G. We also have

(8.2)  

$$\int_{\sigma_T + \mathcal{C}_T} [z, z] d\mu_L(z) = \int_{\mathcal{C}_T} [y + \sigma_T, y + \sigma_T] d\mu_L(y)$$

$$= \int_{\mathcal{C}_T} ([y, y] + 2[\sigma_T, y] + [\sigma_T, \sigma_T]) d\mu_L(y)$$

$$= \int_{\mathcal{C}_T} [y, y] d\mu_L(y) + \int_{\mathcal{C}_T} [\sigma_T, \sigma_T] d\mu_L(y).$$

The last equality is because  $\mathcal{C}_T$  is centrally symmetric and  $[\sigma_T, y]$  is odd.

By Lemmas 7.1 and 7.4(b), we have

(8.3) 
$$\int_{\mathcal{C}_T} [\sigma_T, \sigma_T] d\mu_L(y) = \frac{\operatorname{Vol}(\mathcal{C}_T)}{\operatorname{Vol}(\operatorname{Vor}(0))} [\sigma_T, \sigma_T] = \frac{w(T)}{w(G)} [\sigma_T, \sigma_T].$$

On the other hand,

(8.4)  

$$\int_{\mathcal{C}_{T}} [y, y] d\mu_{L}(y) = \frac{\operatorname{Vol}(\mathcal{C}_{T})}{\operatorname{Vol}(\operatorname{Vor}(0))} \int_{[-\frac{1}{2}, \frac{1}{2}]^{g}} [\sum_{e \notin T} \alpha_{e} \pi(e), \sum_{e \notin T} \alpha_{e} \pi(e)] d\alpha$$

$$= \frac{w(T)}{w(G)} \int_{[-\frac{1}{2}, \frac{1}{2}]^{g}} \left( \sum_{\substack{e, f \notin T \\ e \neq f}} \alpha_{e} \alpha_{f} [\pi(e), \pi(f)] + \sum_{e \notin T} \alpha_{e}^{2} [\pi(e), \pi(e)] \right) d\alpha$$

$$= \frac{w(T)}{w(G)} \sum_{e \notin T} [\pi(e), \pi(e)] \int_{[-\frac{1}{2}, \frac{1}{2}]^{g}} \alpha_{e}^{2} d\alpha.$$



*Figure 4*: A banana graph  $\Gamma$ , consisting of two branch points and *m* edges  $\{e_1, \ldots, e_m\}$  with  $\ell(e_i) = x_i$ .

Here,  $d\alpha = \prod_{e \notin T} d\alpha_e$  denotes the usual Lebesgue measure on  $\mathbb{R}^g$ . The first equality is by the *change of variables* theorem, and the last equality holds because  $\alpha_e \alpha_f$  is an odd function on the symmetric domain  $\left[-\frac{1}{2}, \frac{1}{2}\right]^g$ . Clearly (see Section 4.4),

(8.5) 
$$[\pi(e), \pi(e)] = \mathsf{F}(e)\ell(e), \quad \int_{[-\frac{1}{2}, \frac{1}{2}]^g} \alpha_e^2 \mathrm{d}\alpha = \int_{-\frac{1}{2}}^{\frac{1}{2}} \alpha_e^2 \mathrm{d}\alpha_e = \frac{1}{12}.$$

Putting (8.1)–(8.5) together, we obtain

(8.6) 
$$I(\operatorname{Jac}(\Gamma)) = \frac{1}{12} \sum_{T} \frac{w(T)}{w(G)} \sum_{e \notin T} \mathsf{F}(e)\ell(e) + \sum_{T} \frac{w(T)}{w(G)} [\sigma_T, \sigma_T].$$

The first sum is simplified immediately, after changing the order of summations (see Remark 4.5(i)):

(8.7) 
$$\sum_{T} \frac{w(T)}{w(G)} \sum_{e \notin T} \mathsf{F}(e)\ell(e) = \sum_{e \in E(G)} \mathsf{F}(e)\ell(e) \sum_{T \neq e} \frac{w(T)}{w(G)} = \sum_{e \in E(G)} \mathsf{F}(e)^{2}\ell(e).$$

The result now follows from (8.6), (8.7), and Theorem 6.9.

#### Remark 8.2

- (i) If e = {u, v} is a bridge, then one observes that r(u, v) = ℓ(e) and F(e) = 0 and {j<sub>u</sub>(v, q), j<sub>v</sub>(u, q)} = {0, ℓ(e)}. So e contributes 0 to I(Jac(Γ)). This is expected, as bridges do not contribute to the first homology.
- (ii) By (4.3)–(4.5), it is clear that everything in the formula for I(Jac(Γ)) in Theorem 8.1 can be expressed in terms of the entries of L<sub>q</sub> and edge lengths. As mentioned in Remark 4.1, all entries of L<sub>q</sub> can be computed in matrix multiplication time O(n<sup>ω</sup>), where n is the number of vertices in a model of Γ, and ω is the exponent for matrix multiplication.

*Example 8.3* Consider the metric graph  $\Gamma$  in Figure 4.

Fix the minimal model *G* whose vertex set consists of the two branch points *u* and *v*, and let q = v. Let R denote the effective resistance between *u* and *v*:

$$\mathbf{R}^{-1} = \sum_{i=1}^{m} x_i^{-1}.$$

Then, for each edge  $e_i = \{u, v\}$ , we have

$$F(e_i) = 1 - R/x_i$$
,  $r(u, v) = R$ ,  $j_u(v, q) = R$ ,  $j_v(u, q) = 0$ .

R. de Jong and F. Shokrieh

By Theorem 8.1, one computes

$$I(\operatorname{Jac}(\Gamma)) = \frac{1}{12} \left( \sum_{i=1}^{m} x_i + \frac{m-2}{\sum_{i=1}^{m} x_i^{-1}} \right).$$

In the special case  $x_1 = \cdots = x_m = 1$ , the lattice  $(H_1(\Gamma, \mathbb{Z}), [\cdot, \cdot])$  corresponds to the root lattice  $A_{m-1}$ . So one recovers the formula in [13, p. 460]:

$$I(A_{m-1}) = \frac{m^2 + m - 2}{12m}.$$

We record two other special cases for our application in Section 10:

• If m = 2, then the metric graph  $\Gamma$  is just a circle, and

$$I(\operatorname{Jac}(\Gamma)) = \frac{1}{12}\ell,$$

where  $\ell$  is the total length of the circle.

• If m = 3, then we have

$$I(\operatorname{Jac}(\Gamma)) = \frac{1}{12} \left( x_1 + x_2 + x_3 + \frac{x_1 x_2 x_3}{x_1 x_2 + x_2 x_3 + x_3 x_1} \right)$$

The case m = 3 corresponds to a lattice of rank 2, with Gram matrix

$$\left(\begin{array}{cc} x_1 + x_2 & x_1 \\ x_1 & x_1 + x_3 \end{array}\right)$$

on a suitable basis. As each two-dimensional lattice has a Gram matrix on a suitable basis of this shape, our formula gives a "quick" expression for the tropical moment of any two-dimensional lattice in terms of a Gram matrix.

# 9 Connection with arithmetic geometry

A metric graph may be canonically interpreted as a *skeleton* of a *Berkovich curve*, and the tropical Jacobian of a metric graph can be interpreted as the *canonical skeleton* of a *Berkovich Jacobian variety* [5]. More generally, one may view the canonical skeleton of any Berkovich polarized abelian variety as a polarized real torus in a canonical way, via *non-archimedean uniformization*. Based on this connection, our results can be applied in the study of polarized abelian varieties. For example, we have the following application concerning the computation of *Arakelov heights* attached to principally polarized abelian variet to generally. For more background and for terminology used in this section, we refer to [14].

Let *k* be a number field, and let  $M(k)_0$  and  $M(k)_\infty$  denote the set of nonarchimedean places and the set of complex embeddings of *k*. Let  $(A, \lambda)$  be a principally polarized abelian variety defined over *k*. Assume that *A* has semistable reduction over *k*. For  $v \in M(k)_\infty$ , we let  $I(A_v, \lambda_v)$  denote the *I*-invariant, as defined in [3], of the principally polarized complex abelian variety  $(A_v, \lambda_v)$  obtained by extending scalars to  $\bar{k}_v \simeq \mathbb{C}$ . For  $v \in M(k)_0$ , we let  $I(A_v, \lambda_v)$  denote the tropical moment of the canonical skeleton of the Berkovich analytification of  $(A, \lambda)$  at *v*, viewed as a polarized real torus. Let Nv be the cardinality of the residue field at  $v \in M(k)_0$ .

**Theorem** [14, Theorem A] Let  $\Theta$  be a symmetric effective divisor on A that defines the polarization  $\lambda$ , and put  $L = \mathcal{O}_A(\Theta)$ . Let  $h'_L(\Theta)$  denote the Néron–Tate height of the cycle  $\Theta$ , and let  $h_F(A)$  denote the stable Faltings height of A. Set  $g = \dim(A)$ . Then the equality

(9.1)

$$h_F(A) = 2g h'_L(\Theta) - \kappa_0 g + \frac{1}{[k:\mathbb{Q}]} \left( \sum_{\nu \in M(k)_0} I(A_\nu, \lambda_\nu) \log N\nu + 2 \sum_{\nu \in M(k)_\infty} I(A_\nu, \lambda_\nu) \right)$$

holds in  $\mathbb{R}$ .

Assume that  $v \in M(k)_0$  is a finite place such that the canonical skeleton of the Berkovich analytification of  $(A, \lambda)$  at v can be realized as the tropical Jacobian of some (explicitly given) metric graph. For example,  $(A, \lambda)$  could be the Jacobian variety of a smooth projective geometrically connected curve with semistable reduction over k. Then Theorem 8.1 can be applied to compute the local term  $I(A_v, \lambda_v)$  efficiently. We shall illustrate this in Section 10 by discussing the case of Jacobian varieties of dimension 2 in some detail.

# 10 Example: heights of Jacobians in dimension 2

In this section, we specialize (9.1) to the case where the principally polarized abelian variety  $(A, \lambda)$  is a Jacobian variety of dimension 2. We use (9.1) together with Example 8.3 to give a conceptual explanation of a result due to Autissier [3].

Let *k* be a number field. Let *X* be a smooth projective geometrically connected curve of genus 2 with semistable reduction over *k*. Let Jac(X) be the Jacobian variety of *X*. As in Section 9, we denote by  $M(k)_0$  and  $M(k)_\infty$  the set of non-archimedean places and the set of complex embeddings of *k*. For each  $v \in M(k)_0$ , we have a metric graph  $\Gamma_v$ canonically associated to *X* at *v* by taking the dual graph of the geometric special fiber of the stable model of *X* at *v* (see, for example, [30]). The tropical Jacobian Jac $(\Gamma_v)$  is canonically isometric with the canonical skeleton of the Berkovich analytification of Jac(X) at *v*.

Equation (9.1) specializes into the identity

(10.1)  
$$h_{F}(\operatorname{Jac}(X)) = 4 h_{L}'(\Theta) - 2 \kappa_{0} + \frac{1}{[k:\mathbb{Q}]} \sum_{\nu \in M(k)_{0}} I(\operatorname{Jac}(\Gamma_{\nu})) \log N\nu$$
$$+ \frac{2}{[k:\mathbb{Q}]} \sum_{\nu \in M(k)_{\infty}} I(\operatorname{Jac}(X_{\nu})).$$

In [3, Théorème 5.1], Autissier proves an identity

(10.2)  
$$h_{F}(Jac(X)) = 4 h'_{L}(\Theta) - 2 \kappa_{0} + \frac{1}{[k:\mathbb{Q}]} \sum_{\nu \in M(k)_{0}} \alpha_{\nu} \log N\nu$$
$$+ \frac{2}{[k:\mathbb{Q}]} \sum_{\nu \in M(k)_{\infty}} I(Jac(X_{\nu})),$$

where, for each  $v \in M(k)_0$ , the local invariant  $\alpha_v$  is given explicitly in terms of the metric graph  $\Gamma_v$  by means of a table. See the Remarque following the proof of [3, Théorème 5.1] where, for each of the seven possible topological types of stable dual graphs in genus 2, the invariant  $\alpha_v$  is given in terms of the edge lengths of a minimal model of  $\Gamma_v$ .

A simple explicit calculation using Example 8.3 shows that for all seven topological types as listed in Autissier's table, the equality  $\alpha_v = I(Jac(\Gamma_v))$  is verified. For example, consider case VII in the last row of Autissier's table, corresponding to a banana graph with three edges. The table in this case gives

$$\alpha_{\nu} = \frac{1}{12} \left( x_1 + x_2 + x_3 + \frac{x_1 x_2 x_3}{x_1 x_2 + x_2 x_3 + x_3 x_1} \right)$$

where  $x_1, x_2, x_3$  are the three edge lengths. By Example 8.3, with m = 3, we see immediately that  $\alpha_v = I(Jac(\Gamma_v))$  for case VII. The cases I–VI are very similar, and in fact simpler, as the irreducible components of the corresponding graphs are either bridges, which by Remark 8.2(ii) contribute zero to  $I(Jac(\Gamma_v))$ , or circles, which by Example 8.3 with m = 2 contribute 1/12 of their total length to  $I(Jac(\Gamma_v))$ . We thus have a complete conceptual explanation of all entries in Autissier's table.

# 11 Connection with the tau invariant of a metric graph

Let  $\Gamma$  be a metric graph. The notion of Arakelov–Green's function  $g_{\mu}(x, y)$  associated to a measure  $\mu$  on  $\Gamma$  is introduced in [11, 30]. It can be shown [11, Theorem 2.11] that there exists a unique measure  $\mu_{can}$  on  $\Gamma$  having total mass 1, such that  $g_{\mu_{can}}(x, x)$  is a constant. This constant is by definition  $\tau(\Gamma)$ . Alternatively,  $\tau(\Gamma)$  can be interpreted as a certain "capacity," with equilibrium measure  $\mu_{can}$  and with potential kernel (1/2 times) the effective resistance function r(x, y) [6, Corollary 14.2]. See also [4, 6, 12] for more background, examples, and formulas.

We will work with the following definition (see, e.g., [6, Lemma 14.4]) in terms of the effective resistance function.

**Definition 11.1** Fix a point  $q \in \Gamma$ . Let  $f(x) = \frac{1}{2}r(x,q)$ . We put  $\tau(\Gamma) \coloneqq \int_{\Gamma} (f'(x))^2 dx$ ,

where dx denotes the (piecewise) Lebesgue measure on  $\Gamma$ .

It is elementary and well known that the real number  $\tau(\Gamma)$  is independent of the choice of  $q \in \Gamma$ .

We have the following explicit (and efficiently computable) formula for  $\tau(\Gamma)$ . See also [12, Proposition 2.9] for an equivalent form of this formula. Our proof is somewhat different and avoids "circuit reduction theory."

**Theorem 11.2** Let  $\Gamma$  be a metric graph. Fix a model G of  $\Gamma$ , and fix a point  $q \in V(G)$ . Then

$$\tau(\Gamma) = \frac{1}{12} \sum_{e} \mathsf{F}(e)^{2} \ell(e) + \frac{1}{4} \sum_{e=\{u,v\}} \frac{(r(u,q) - r(v,q))^{2}}{\ell(e)},$$

where the sums are over all edges  $e \in E(G)$ .

**Proof** Let f(x) be as in Definition 11.1. Let *G* be a model of  $\Gamma$ . Then

$$\tau(\Gamma) = \sum_{e \in E(G)} \int_{e} \left(f'(x)\right)^2 \mathrm{d}x.$$

We identify each edge segment *e* in *E*(*G*) with an interval  $[0, \ell(e)]$  and represent a point  $x \in e$  by  $x \in [0, \ell(e)]$ . By Example 4.6, we have

$$2f'(x) = \mathsf{F}(e) - \frac{r(u,q) - r(v,q)}{\ell(e)} - 2\frac{\mathsf{F}(e)}{\ell(e)}x.$$

With a direct computation, we obtain

$$\int_{e} \left(f'(x)\right)^{2} \mathrm{d}x = \int_{0}^{\ell(e)} \left(f'(x)\right)^{2} \mathrm{d}x = \frac{1}{12} \mathsf{F}(e)^{2} \ell(e) + \frac{1}{4} \frac{\left(r(u,q) - r(v,q)\right)^{2}}{\ell(e)},$$

and the result follows.

The reader will notice the similarity between the right-hand side in Theorem 11.2 and the right-hand side in Theorem 8.1. In fact, we can now prove a simple linear relation between  $\tau(\Gamma)$ , the tropical moment  $I(\text{Jac}(\Gamma))$ , and the *total length* of  $\Gamma$ . The result was announced as Theorem B in Section 1.

**Definition 11.3** Let  $\Gamma$  be a metric graph, and fix a model *G* of  $\Gamma$ . The *total length* of  $\Gamma$  is defined by

$$\ell(\Gamma) \coloneqq \sum_{e \in E(G)} \ell(e).$$

It is easily seen that  $\ell(\Gamma)$  is independent of the choice of the model *G*.

**Theorem 11.4** Let  $\Gamma$  be a metric graph. The identity

$$\frac{1}{2}\tau(\Gamma) + I(\operatorname{Jac}(\Gamma)) = \frac{1}{8}\ell(\Gamma)$$

holds in  $\mathbb{R}$ .

**Proof** Fix a model *G* of  $\Gamma$ . By Theorem 11.2 and (4.2), we have

(11.1) 
$$\tau(\Gamma) = \frac{1}{12} \sum_{e} \mathsf{F}(e)^2 \ell(e) + \frac{1}{4} \sum_{e=\{u,v\}} \frac{(j_u(v,q) - j_v(u,q))^2}{\ell(e)},$$

where the sums are over all edges  $e \in E(G)$ . The result follows from (4.4), (11.1), Theorem 8.1, and the following direct computation:

$$\frac{1}{2}\tau(\Gamma) + I(\operatorname{Jac}(\Gamma))$$
  
=  $\frac{1}{8}\sum_{e=\{u,v\}} \left( \mathsf{F}(e)^2 \ell(e) + \frac{(j_u(v,q) - j_v(u,q))^2}{\ell(e)} + 2r(u,v) - 2\frac{j_u(v,q)^2 + j_v(u,q)^2}{\ell(e)} \right)$ 

$$= \frac{1}{8} \sum_{e=\{u,v\}} \left( \mathsf{F}(e)^2 \ell(e) + 2r(u,v) - \frac{(j_u(v,q) + j_v(u,q))^2}{\ell(e)} \right)$$
$$= \frac{1}{8} \sum_{e=\{u,v\}} \left( \left( 1 - \frac{r(u,v)}{\ell(e)} \right)^2 \ell(e) + 2r(u,v) - \frac{r(u,v)^2}{\ell(e)} \right)$$
$$= \frac{1}{8} \sum_{e=\{u,v\}} \ell(e) = \frac{1}{8} \ell(\Gamma).$$

Acknowledgment We thank the anonymous referee for helpful suggestions.

# References

- O. Amini, Lattice of integer flows and poset of strongly connected orientations. Preprint, 2010. arXiv:1007.2456
- [2] Y. An, M. Baker, G. Kuperberg, and F. Shokrieh, Canonical representatives for divisor classes on tropical curves and the matrix-tree theorem. Forum Math. Sigma. 2(2014), e24, 25 pp.
- [3] P. Autissier, Hauteur de Faltings et hauteur de Néron-Tate du diviseur thêta. Compos. Math. 142(2006), no. 6, 1451–1458.
- [4] M. Baker and X. Faber, *Metrized graphs, Laplacian operators, and electrical networks*. In: Quantum graphs and their applications, Contemporary Mathematics, 415, American Mathematical Society, Providence, RI, 2006, pp. 15–33.
- [5] M. Baker and J. Rabinoff, The skeleton of the Jacobian, the Jacobian of the skeleton, and lifting meromorphic functions from tropical to algebraic curves. Int. Math. Res. Not. IMRN. 16(2015), 7436–7472.
- [6] M. Baker and R. Rumely, Harmonic analysis on metrized graphs. Can. J. Math. 59(2007), no. 2, 225–275.
- [7] M. Baker and R. Rumely, Potential theory and dynamics on the Berkovich projective line, Mathematical Surveys and Monographs, 159, American Mathematical Society, Providence, RI, 2010.
- [8] M. Baker and F. Shokrieh, *Chip-firing games, potential theory on graphs, and spanning trees.* J. Combin. Theory Ser. A. 120(2013), no. 1, 164–182.
- [9] E. S. Barnes and N. J. A. Sloane, *The optimal lattice quantizer in three dimensions*. SIAM J. Algebraic Discrete Methods. 4(1983), no. 1, 30–41.
- [10] L. J. Billera and B. Sturmfels, Fiber polytopes. Ann. Math. 135(1992), no. 3, 527-549.
- [11] T. Chinburg and R. Rumely, The capacity pairing. J. Reine Angew. Math. 434(1993), 1-44.
- [12] Z. Cinkir, The tau constant of a metrized graph and its behavior under graph operations. Electron. J. Combin. 18(2011), no. 1, Article no. 81, 42 pp.
- [13] J. H. Conway and N. J. A. Sloane (1999). Sphere packings, lattices and groups. 3rd ed., Grundlehren der Mathematischen Wissenschaften, 290, Springer, New York.
- [14] R. de Jong and F. Shokrieh, Faltings height and Néron-Tate height of a theta divisor. Compos. Math. 158(2022a), no. 1, 1–32.
- [15] R. de Jong and F. Shokrieh, Jumps in the height of the Ceresa cycle. To appear in J. Differ. Geom. Preprint, 2022b. arXiv:2104.10060
- [16] R. de Jong and F. Shokrieh, Metric graphs, cross ratios, and Rayleigh's laws. To appear in Rocky Mt. J. Math. Preprint, 2022c. arXiv:1810.02638
- [17] M. Dutour Sikirić, A. Schürmann, and F. Vallentin, Complexity and algorithms for computing Voronoi cells of lattices. Math. Comput. 78(2009), no. 267, 1713–1731.
- [18] C. Favre and J. Rivera-Letelier, Équidistribution quantitative des points de petite hauteur Sur la droite projective. Math. Ann. 335(2006), no. 2, 311–361.
- [19] C. Greene and T. Zaslavsky, On the interpretation of Whitney numbers through arrangements of hyperplanes, zonotopes, non-radon partitions, and orientations of graphs. Trans. Amer. Math. Soc. 280(1983), no. 1, 97–126.
- [20] N. Kawazumi, Johnson's homomorphisms and the Arakelov–Green function. Preprint, 2008. arXiv:0801.4218

- [21] N. Kawazumi, Canonical 2-forms on the moduli space of Riemann surfaces. In: Handbook of Teichmüller theory. Vol. II, IRMA Lectures in Mathematics and Theoretical Physics, 13, European Mathematical Society, Zürich, 2009, pp. 217–237.
- [22] G. Kirchhoff, Ueber die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Vertheilung galvanischer Ströme geführt wird. Ann. Phys. 148(1847), no. 12, 497–508.
- [23] M. Kotani and T. Sunada, Jacobian tori associated with a finite graph and its abelian covering graphs. Adv. Appl. Math. 24(2000), no. 2, 89–110.
- [24] R. Lyons and Y. Peres, Probability on trees and networks, Cambridge Series in Statistical and Probabilistic Mathematics, 42, Cambridge University Press, New York, 2016.
- [25] G. Mikhalkin and I. Zharkov, *Tropical curves, their Jacobians and theta functions*. In: *Curves and abelian varieties*, Contemporary Mathematics, 465, American Mathematical Society, Providence, RI, 2008, pp. 203–230.
- [26] T. Oda and C. S. Seshadri, Compactifications of the generalized Jacobian variety. Trans. Amer. Math. Soc. 253(1979), 1–90.
- [27] J. Väisälä, Gromov hyperbolic spaces. Expo. Math. 23(2005), no. 3, 187-231.
- [28] R. Wilms, New explicit formulas for Faltings' delta-invariant. Invent. Math. 209(2017), no. 2, 481–539.
- [29] R. Wilms, Degeneration of Riemann theta functions and of the Zhang-Kawazumi invariant with applications to a uniform Bogomolov conjecture. Preprint, 2021. arXiv:2101.04024
- [30] S. Zhang, Admissible pairing on a curve. Invent. Math. 112(1993), no. 1, 171–193.
- [31] S.-W. Zhang, Gross-Schoen cycles and dualising sheaves. Invent. Math. 179(2010), no. 1, 1–73.
- [32] M. C. Zimmermann, Configurations of sublattices and Dirichlet-Voronoi cells of periodic point sets. Ph.D. thesis, Technischen Universität Dortmund, 2017. https://doi.org/10.17877/DE290R-18235

Mathematics Institute, Leiden University, Leiden, The Netherlands e-mail: rdejong@math.leidenuniv.nl

Department of Mathematics, University of Washington, Seattle, WA, USA e-mail: farbod@uw.edu