



Universiteit
Leiden
The Netherlands

Circadian control of white and brown adipose tissues

Yue, K.M.; Rensen, P.C.N.; Kooijman, S.

Citation

Yue, K. M., Rensen, P. C. N., & Kooijman, S. (2023). Circadian control of white and brown adipose tissues. *Current Opinion In Genetics And Development*, 80.
doi:10.1016/j.gde.2023.102056

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3634252>

Note: To cite this publication please use the final published version (if applicable).



Circadian control of white and brown adipose tissues

Kaiming Yue, Patrick CN Rensen* and Sander Kooijman*

White and brown adipose tissues are highly dynamic organs anticipating and responding to changes in the environment. The circadian timing system facilitates anticipation, and it is therefore not surprising that circadian disturbances, a prominent feature of modern 24/7 society, increase the risk for (cardio)metabolic diseases. In this mini-review, we will address mechanisms and strategies to mitigate disease risk associated with circadian disturbances. In addition, we discuss the opportunities arising from the knowledge we gained about circadian rhythms in these adipose tissues, including the application of chronotherapy, optimizing endogenous circadian rhythms to allow for more effective intervention, and the identification of novel therapeutic targets.

Address

Department of Medicine, Division of Endocrinology, and Einthoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, the Netherlands

Corresponding author: Kooijman, Sander (S.Kooijman@lumc.nl)

* Twitter account: [@Rensen_Lab](#), [@KooijmanSander](#)

Current Opinion in Genetics & Development 2023, **80**:102056

This review comes from a themed issue on **Molecular and Genetic Basis of Disease**

Edited by **Zach Gerhart-Hines** and **Ruth Loos**

For complete overview of the section, please refer to the article collection, "[Molecular and Genetic Basis of Disease \(2023\)](#)"

Available online 25 May 2023

<https://doi.org/10.1016/j.gde.2023.102056>

0959-437X/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

The concept of adipose tissue has radically changed in the past decades. While the classical view of white adipose tissue (WAT) is a passive lipid-storing tissue, it actually is a very dynamic organ anticipating and responding to changes in nutrient availability. Adding to this, the presence of the thermogenic and lipid-combusting brown adipose tissue (BAT) in adult humans was only established 15 years ago, and more recent insight into the physiology also here suggests anticipation and response to, for example, changes in environmental temperature.

Anticipation and adaptation of adipose tissue

Dealing with daily and seasonal fluctuations in nutrient availability and environmental temperature is, for a large part, facilitated through the circadian timing system. For this, adipocytes and all other cell types in adipose tissue possess a cell-autonomous transcription–translation feedback loop. Core clock genes serve as the molecular basis for endogenous oscillations under constant environmental conditions. Rhythms are fine-tuned to run at exactly 24 h through, among others, post-translational modifications as a result of variation in cellular energy status, and by synchronization to the outside world via hormones, the autonomic nervous system, temperature, and nutrient availability. In turn, circadian variation in WAT function leads to, for example, a circadian rhythm in leptin levels independent of meal timing, and the circadian activity in BAT likely contributes to the rhythm in body core temperature to which other bodily organs and even the central clock in the brain respond. Interestingly, circadian rhythms in BAT also seem to adapt to (seasonal) variation in daily light exposure [1], thereby possibly anticipating upcoming changes in environmental temperature. And while our knowledge on circadian rhythms in specific BAT and WAT functions is increasing, a more or less understudied phenomenon is the circadian rhythm in more general housekeeping processes such as cell proliferation. Notably, both the circadian clock machinery and diurnal feeding patterns have been implicated in adipocyte progenitor proliferation, and may therefore be critical determinants of adipose tissue expansion and health over time [2].

Implications of circadian disturbance

Regulation of circadian rhythms at multiple levels, including the entrainment to the environment via so-called Zeitgebers (e.g. light and food cues), on the one hand, facilitates anticipation and adaptation, but on the other hand, also makes us vulnerable for circadian disturbances, often with consequences for metabolic and cardiovascular disease risk. In addition, modern-day society characterized by all-day electrical lighting, continuous access to food, and need for shift work causes misalignment of behavior and endogenous anticipatory circadian rhythms. As an example, WAT shows mild insulin resistance during the night to redirect glucose produced by the liver to the brain as an essential energy source. The implication, however, is that eating at night or even late-evening meals have been shown to cause spikes in blood glucose levels and increase the risk of type-2 diabetes [3]. To

make things worse, circadian disruption additionally attenuates glucose-stimulated insulin secretion by the pancreas [4]. Thus, eating during a night shift, especially readily digestible carbohydrates in sweets and soft drinks, constitutes a strong risk for cardiometabolic disorders due to a combination of insufficient insulin secretion and adipose tissue responding to a lesser extent to this insulin. Interestingly, time-restricted eating appears to be a feasible strategy to alleviate cardiometabolic risks associated with shift work [5], and also daytime eating prevents circadian misalignment and glucose intolerance in simulated shift work [6]. Nevertheless, the multifactorial disease risk in shift workers likely requires a personalized, multidisciplinary approach [7].

Identification of the mechanisms driving circadian rhythms in adipose tissue may help us to select promising prevention or intervention strategies. In subcutaneous adipose tissue biopsies of healthy subjects, approximately 800 rhythmic transcripts (2% of target genes in the assay) can be detected under 'constant routine' conditions [8]. In such a protocol, exogenous time cues are minimized through exposure to constant light, evenly distributed food intake, and no sleep for at least 24 h. The number of oscillating transcripts dramatically increases under less-stringent diurnal conditions. With 'normal' meal sleep timing, at least 25% of the genes show different expression levels during the course of the day, and time-restricted eating additionally changes the rhythmicity of 450 genes [9]. Specifically, circadian expression of core clock genes, glucoregulatory genes, and genes involved in cell proliferation and differentiation are altered in WAT upon time-restricted eating in humans [9], patterns that have been linked to healthy adipose function and expansion in mice [2]. In murine BAT, approximately 40% of all expressed genes was found to be oscillating in unrestricted conditions [10]. These numbers indicate a role for core clock genes, and possibly an even bigger role for external factors. Indeed, a landmark study by Turek et al. [11] demonstrated that whole-body circadian locomotor output cycle kaput (CLOCK) mutant mice are obese and develop features of the metabolic syndrome when fed a high-fat diet. Adipocyte-specific loss of brain and muscle ARNT-like 1 (BMAL1) has also been reported to increase adiposity due to defective lipolytic activity in WAT [12]. Nuclear receptor subfamily-1 group-D member 1 (NR1D1), also known as REV-ERB α , was identified as one of the main regulators of diurnal variation in body temperature and BAT activity through its suppressive effects on uncoupling protein-1 (UCP1) expression [13].

With regard to the environmental factors, we previously demonstrated in mice that prolonged daily light exposure increases adiposity through attenuation of BAT activity [14]. In line with these data, it has been reported

that artificial light at night while sleeping is associated with risk of weight gain and obesity in humans [15]. Mechanistically, we identified in mice a physiological rhythm in glucocorticoids as an important regulator of diurnal BAT function, although this effect was likely mediated via altered sympathetic outflow toward BAT [16]. Interestingly, others have also reported weight gain upon flattening of daily glucocorticoid oscillations in mice, and attributed this to a direct effect of continuous versus pulsatile glucocorticoids on peroxisome proliferator-activated receptor gamma (PPAR γ)-mediated white adipocyte differentiation [17]. Thus, beyond the well-known effects of hypercorticism (e.g. in chronic stress) on adipose tissue function, a circadian rhythm in glucocorticoids also seems critical for metabolic health. Nocturnin, a circadian-regulated deadenylase, has also been reported to regulate adipocyte differentiation in a PPAR γ -dependent manner [18], and nocturnin-deficient mice are resistant to diet-induced obesity. Likewise, period 2 was found to directly interact with PPAR γ to regulate adipogenesis [19]. Most strikingly, time-restricted feeding was shown to protect from obesity and metabolic syndrome in the context of circadian disruption due to genetic loss of clock genes in mice, highlighting the relative importance of external timing cues [20].

Beyond the effects of genetic or environmental circadian disruption, it is tempting to speculate that alterations in the circadian system also contribute to the increased risk of cardiometabolic diseases with increasing age. Aging is typically associated with reduced amplitude or even loss of rhythmicity. In WAT and BAT of aged mice, the fraction of oscillating lipids decreases from approximately 22–14% and 18%, respectively [21]. In line with this, we recently reported that peak abundance of lipoprotein lipase (LPL), a critical regulator of triglyceride-derived fatty acid uptake, was strongly attenuated in BAT of middle-aged female mice [22]. Combined with the lower expression of genes involved in fat oxidation, the net result was increased adiposity and attenuated rhythms in core body temperature. Provided these findings can be translated to humans, these data suggest BAT-targeted interventions may particularly benefit the aged population.

Opportunities arising from circadian rhythms in adipose tissue

Before addressing the question whether circadian cycles in adipose tissue should be considered when therapeutically targeting the tissue, it is imperative to note that the existence of circadian variation in WAT and BAT function also implies that potential rhythmic activity should be taken into account when performing measurements. As an example, we previously demonstrated that angiotensin-like 4 (ANGPTL4), a protein

that inhibits LPL activity, redirects triglycerides from WAT to BAT uptake during cold [23]. Those experiments were, mostly for practical reasons, performed during the light phase. Only later, we realized that when experiments would have been performed during the dark phase, when ANGPTL4 levels are low/absent [1,10], this effect would probably have been missed. Thus, knowledge of circadian rhythms in the body is useful, if not critical, in the design and interpretation of experiments, but also when interpreting patient data.

The presence of metabolically active BAT is associated with cardiometabolic health in humans [24], indicating that stimulating BAT — despite its relative low abundance in humans — might have therapeutic potential. Cold exposure is the most effective way to activate thermogenesis in BAT and simultaneously promotes lipolysis in WAT, but comes with practical limitations. We therefore recently asked ourselves the question whether time of day is an important determinant in the metabolic effects of cold [25]. In males, cold-induced energy expenditure was found to be higher in the morning than in the evening. This effect was accompanied by an elevation in free fatty acids in the morning, but not evening. In females, there was no (statistical) difference in cold-induced energy expenditure between morning and evening, but free fatty acids as well as triglycerides and cholesterol levels were increased after cold to a higher extent in the morning than in the evening. Whether these differences have implications for the therapeutic potential of cold strategies remains to be investigated.

Chronotherapy can be considered to optimize therapeutic effectiveness as shown for cold exposure, and/or to minimize side effects. In the context of adipose-targeted therapy, the latter may be particularly relevant when using sympathomimetics to stimulate lipolysis in WAT and thermogenesis in BAT. In humans, both processes are primarily driven by signaling through the β 2-adrenergic receptor [26], but this receptor also exerts prominent functions in the vasculature. It would be of high interest to investigate if, for example, β 2-adrenergic receptor-mediated vasodilation also shows a circadian rhythm.

As an alternative strategy to the timing of therapeutic interventions, one might also consider optimizing endogenous circadian rhythms to promote resilience against circadian disruption or to allow for more effective pharmacological intervention. A fascinating development in treatment of cardiometabolic disorders is the generation of dual- or even triple agonists, with recent successes in clinical trials reported for combined glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-1 (GLP-1) receptor agonism. The impressive weight-lowering effect of dual-GIP and GLP-1 receptor agonism is most likely mediated via

promotion of satiety and a reduction in food intake [27], but preclinical evidence also suggests improvement of the lipid-buffering capacity of WAT, with increased lipid uptake in postprandial state and potentiated lipolysis in the fasted state contributing to better overall metabolic health (reviewed in [28]). Given the differential regulation of WAT in the fast-fed state, it would be of interest to investigate if time-restricted eating can potentiate the health benefits of dual-GIP/GLP-1 receptor agonism. Of note, time-restricted eating does not seem to be more beneficial than calorie restriction when it comes to promoting weight loss in humans [29]. However, it does attenuate daily glucose levels in blood accompanied by increased GIP levels following dinner and > 300 genes being differentially regulated in subcutaneous adipose tissue [30]. Late eating on the other hand increases hunger and decreases energy expenditure as (partly) explained by lower adipose-derived leptin levels and changes in the adipose transcriptome consistent with decreased lipolysis and increased adipogenesis [31].

Last but not least, one could ask the question if defining the molecular on- and off-switches in diurnal WAT or BAT functioning would yield novel therapeutic targets. To explore this further, we recently performed RNA sequencing in BAT samples collected from C57BL/6J mice at three-hour interval throughout a 24-hour period [10]. In the gene cluster that shows expression patterns overlapping with the previously reported rhythm in metabolic BAT activity [11], we found marked enrichment for lipid catabolic processes, with *Lpl* as the gene with the largest amplitude. Of the known LPL modulators, *Angptl4* showed the largest diurnal amplitude in antiphase with *Lpl* expression, and both *Angptl4* knockout and overexpression attenuated oscillations in LPL and triglyceride-derived fatty acid uptake by BAT. These data suggest that time of day is essential when targeting LPL activity through ANGPTL4 modulation. In addition, transcription factor enrichment analysis identified PPAR γ as one of the potential drivers of the circadian transcriptional activity in BAT, and chromatin immunoprecipitation (ChIP) sequencing revealed six rhythmic binding sites for PPAR γ in the promoter region of *Lpl* [10]. Next, questions to address are how rhythms in PPAR γ -mediated transcription are generated, and if a PPAR γ agonist with a short half-life can be used to stimulate thermogenic activity.

Concluding remarks

Taken together, adipose tissues show strong variation in activity during the course of the day with implications for nutrient handling and cardiometabolic disease risk. Circadian disruption following shift work, or attenuated circadian amplitudes as occurring during aging are associated with adipose dysfunction and accelerated disease

development. Incremental evidence suggests that circadian rhythms in adipose tissue should be considered when measuring WAT or BAT function, and when targeting these tissues in the treatment of cardiometabolic disorders.

CRedit authorship contribution statement

Kaiming Yue, Patrick Rensen, Sander Kooijman:

Writing – original draft, Writing – review & editing.

Data Availability

No data were used for the research described in the article.

Declaration of Competing Interest

Nothing declared.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. van den Berg R, Kooijman S, Noordam R, et al.: **A diurnal rhythm in brown adipose tissue causes rapid clearance and combustion of plasma lipids at wakening.** *Cell Rep* 2018, **22**:3521-3533.
2. Ribas-Latre A, Santos RB, Fekry B, et al.: **Cellular and physiological circadian mechanisms drive diurnal cell proliferation and expansion of white adipose tissue.** *Nat Commun* 2021, **12**:3482.
- Both the circadian clock machinery and diurnal feeding patterns were identified as important contributors of adipocyte progenitor proliferation, and may therefore be critical determinants of adipose tissue expansion and health over time.
3. Dashti HS, Gomez-Abellan P, Qian J, et al.: **Late eating is associated with cardiometabolic risk traits, obesogenic behaviors, and impaired weight loss.** *Am J Clin Nutr* 2021, **113**:154-161.
4. Qian J, Dalla Man C, Morris CJ, et al.: **Differential effects of the circadian system and circadian misalignment on insulin sensitivity and insulin secretion in humans.** *Diabetes Obes Metab* 2018, **20**:2481-2485.
5. Manoogian ENC, Zadourian A, Lo HC, et al.: **Feasibility of time-restricted eating and impacts on cardiometabolic health in 24-h shift workers: the Healthy Heroes randomized control trial.** *Cell Metab* 2022, **34**:e1447.
- Time-restricted eating was shown as a feasible strategy in shift workers with potential benefits for cardiometabolic health.
6. Chellappa SL, Qian J, Vujovic N, et al.: **Daytime eating prevents internal circadian misalignment and glucose intolerance in night work.** *Sci Adv* 2021, **7**:eabg9910.
7. Schilperoort M, Rensen PCN, Kooijman S: **Time for novel strategies to mitigate cardiometabolic risk in shift workers.** *Trends Endocrinol Metab* 2020, **31**:952-964.
8. Christou S, Wehrens SMT, Isherwood C, et al.: **Circadian regulation in human white adipose tissue revealed by transcriptome and metabolic network analysis.** *Sci Rep* 2019, **9**:2641.
9. Zhao L, Hutchison AT, Liu B, et al.: **Time-restricted eating alters the 24-hour profile of adipose tissue transcriptome in men with obesity.** *Obesity* 2022, **31** Suppl 1:63-74.
10. van Eenige R, In Het Panhuis W, Schonke M, et al.: **Angiopoietin-like 4 governs diurnal lipoprotein lipase activity in brown adipose tissue.** *Mol Metab* 2022, **60**:101497.
- Here we addressed the question if defining the molecular on- and off-switches in diurnal BAT functioning would yield novel therapeutic approaches. We identified potential involvement of PPAR γ and concluded that time of day is essential when targeting LPL activity in BAT to improve cardiometabolic health.
11. Turek FW, Joshu C, Kohsaka A, et al.: **Obesity and metabolic syndrome in circadian Clock mutant mice.** *Science* 2005, **308**:1043-1045.
12. Paschos GK, Ibrahim S, Song WL, et al.: **Obesity in mice with adipocyte-specific deletion of clock component Arntl.** *Nat Med* 2012, **18**:1768-1777.
13. Gerhart-Hines Z, Feng D, Emmett MJ, et al.: **The nuclear receptor Rev-erb α controls circadian thermogenic plasticity.** *Nature* 2013, **503**:410-413.
14. Kooijman S, van den Berg R, Ramkisoensing A, et al.: **Prolonged daily light exposure increases body fat mass through attenuation of brown adipose tissue activity.** *Proc Natl Acad Sci USA* 2015, **112**:6748-6753.
15. Park YM, White AJ, Jackson CL, et al.: **Association of exposure to artificial light at night while sleeping with risk of obesity in women.** *JAMA Intern Med* 2019, **179**:1061-1071.
16. Kroon J, Schilperoort M, In Het Panhuis W, et al.: **A physiological glucocorticoid rhythm is an important regulator of brown adipose tissue function.** *Mol Metab* 2021, **47**:101179.
17. Bahrami-Nejad Z, Zhao ML, Tholen S, et al.: **A transcriptional circuit filters oscillating circadian hormonal inputs to regulate fat cell differentiation.** *Cell Metab* 2018, **27**:e858.
18. Kawai M, Green CB, Lecka-Czernik B, et al.: **A circadian-regulated gene, Nocturnin, promotes adipogenesis by stimulating PPAR γ nuclear translocation.** *Proc Natl Acad Sci USA* 2010, **107**:10508-10513.
19. Grimaldi B, Bellet MM, Katada S, et al.: **PER2 controls lipid metabolism by direct regulation of PPAR γ .** *Cell Metab* 2010, **12**:509-520.
20. Chaix A, Lin T, Le HD, et al.: **Time-restricted feeding prevents obesity and metabolic syndrome in mice lacking a circadian clock.** *Cell Metab* 2019, **29**:e304.
21. Held NM, Buijink MR, Elfrink HL, et al.: **Aging selectively dampens oscillation of lipid abundance in white and brown adipose tissue.** *Sci Rep* 2021, **11**:5932.
22. (In), Het Panhuis W, Schonke M, Siebeler R, et al.: **Aging attenuates diurnal lipid uptake by brown adipose tissue.** *Aging (Albany NY)* 2022, **14**:7734-7751.
23. Dijk W, Heine M, Vergnes L, et al.: **ANGPTL4 mediates shuttling of lipid fuel to brown adipose tissue during sustained cold exposure.** *Elife* 2015, **4**:e08428.
24. Becher T, Palanisamy S, Kramer DJ, et al.: **Brown adipose tissue is associated with cardiometabolic health.** *Nat Med* 2021, **27**:58-65.
- Based on retrospectively analysis of 134 529 18F-FDG PET-CT scans from 52 487 patients the authors report that individuals with BAT had lower prevalences of cardiometabolic diseases. Findings were supported by improved blood glucose and triglyceride values, and the protective effects of BAT were more pronounced in individuals with overweight or obesity, indicating that BAT might play a role in mitigating the deleterious effects of obesity.
25. Straat ME, Martinez-Tellez B, Sardjoe Mishre A, et al.: **Cold-induced thermogenesis shows a diurnal variation that unfolds differently in males and females.** *J Clin Endocrinol Metab* 2022, **107**:1626-1635.
- Here we report time-dependent effects of cold exposure on energy expenditure and BAT activity as measured by passive-infrared thermography.
26. Blondin DP, Nielsen S, Kuipers EN, et al.: **Human brown adipocyte thermogenesis is driven by beta2-AR stimulation.** *Cell Metab* 2020, **32**:e287.
27. Jastreboff AM, Aronne LJ, Ahmad NN, et al.: **Tirzepatide once weekly for the treatment of obesity.** *N Engl J Med* 2022, **387**:205-216.

Dual GIPR/GLP1R agonism was shown promote impressive weight-lowering effects in humans.

28. Samms RJ, Coghlan MP, Sloop KW: **How may GIP enhance the therapeutic efficacy of GLP-1?** *Trends Endocrinol Metab* 2020, **31**:410-421.
29. Liu D, Huang Y, Huang C, *et al.*: **Calorie restriction with or without time-restricted eating in weight loss.** *N Engl J Med* 2022, **386**:1495-1504.
30. Zhao L, Hutchison AT, Liu B, *et al.*: **Time-restricted eating improves glycemic control and dampens energy-consuming pathways in human adipose tissue.** *Nutrition* 2022, **96**:111583.
31. Vujovic N, Piron MJ, Qian J, *et al.*: **Late isocaloric eating increases hunger, decreases energy expenditure, and modifies metabolic pathways in adults with overweight and obesity.** *Cell Metab* 2022, **34**:e1487.