



Universiteit
Leiden
The Netherlands

Association between organizational characteristics and adequate pain management at the intensive care unit

Roos-Blom, M.J.; Dongelmans, D.; Stilma, W.; Spijkstra, J.J.; Jonge, E. de; Keizer, N. de

Citation

Roos-Blom, M. J., Dongelmans, D., Stilma, W., Spijkstra, J. J., Jonge, E. de, & Keizer, N. de. (2020). Association between organizational characteristics and adequate pain management at the intensive care unit. *Journal Of Critical Care*, 56, 1-5. doi:10.1016/j.jcrc.2019.11.010

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3278244>

Note: To cite this publication please use the final published version (if applicable).



Association between organizational characteristics and adequate pain management at the intensive care unit

Marie-José Roos-Blom, MSc^{a,b,*}, Dave Dongelmans, MD, PhD^{b,c}, Willemke Stilma, RN, MSc^{c,d}, Jan Jaap Spijkstra, MD^{b,e}, Evert de Jonge, MD, PhD^f, Nicolette de Keizer, PhD^{a,b}

^a Amsterdam UMC, University of Amsterdam, Department of Medical Informatics, Amsterdam Public Health research institute, Amsterdam, The Netherlands

^b National Intensive Care Evaluation (NICE) foundation, Amsterdam, the Netherlands

^c Amsterdam UMC, location AMC, University of Amsterdam, Department of Intensive Care Medicine, Amsterdam, The Netherlands

^d ACHIEVE, Center of Applied Research, Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands

^e Amsterdam UMC, location VUmc, Vrije Universiteit Amsterdam, Department of Intensive Care Medicine, Amsterdam, The Netherlands

^f Leiden University Medical Center, Department of Intensive Care Medicine, Leiden, The Netherlands

ARTICLE INFO

Keywords:

Pain
Intensive care units
Organizational characteristics
Nurse to patient ratio
Bed occupancy rate
Intensivist presence

ABSTRACT

Purpose: Half of the patients experience pain during their ICU stay which is known to influence their outcomes. Nurses and physicians encounter organizational barriers towards pain assessment and treatment. We aimed to evaluate the association between adequate pain management and nurse to patient ratio, bed occupancy rate, and fulltime presence of an intensivist.

Materials and methods: We performed unadjusted and case-mix adjusted mixed-effect logistic regression modeling on data from thirteen Dutch ICUs to investigate the association between ICU organizational characteristics and adequate pain management, i.e. patient-shift observations in which patients' pain was measured and acceptable, or unacceptable and normalized within 1 h.

All ICU patients admitted between December 2017 and June 2018 were included, excluding patients who were delirious, comatose or had a Glasgow coma score < 8 at the first day of ICU admission.

Results: Case-mix adjusted nurse to patient ratios of 0.70 to 0.80 and over 0.80 were significantly associated with adequate pain management (OR [95% confidence interval] of respectively 1.14 [1.07–1.21] and 1.16 [1.08–1.24]). Bed occupancy rate and intensivist presence showed no association.

Conclusion: Higher nurse to patient ratios increase the percentage of patients with adequate pain management especially in medical and mechanically ventilated patients.

© 2019 Elsevier Inc. All rights reserved.

1. Introduction

Pain experienced during ICU stay is a major issue and known to influence patient outcomes. Pain decreases comfort and sleep and increases morbidity, mortality and length of stay [1–3]. During their stay half of the ICU patients experience moderate to severe pain [4,5]. Pain increases during procedures such as turning, chest tube removal, wound drain removal, and arterial line insertion. Furthermore pain experienced before a procedure is associated with greater pain during a procedure [6,7]. The frequent assessment and appropriate treatment

of pain is associated with decreased incidence of pain, need for sedatives, duration of mechanical ventilation, ICU and hospital length of stay, and mortality [3,8,9]. Therefore, it should be encouraged that pain during ICU admission is optimally managed. However nurses and physicians encounter barriers towards pain assessment and treatment, for example pain is not always easy to measure, especially in mechanically ventilated or sedated patients [10]. Furthermore, qualitative studies investigating health professionals' practices and barriers regarding pain management in ICUs indicated that inadequate staffing levels to meet workload and the need of doctor's approval for prescribing proper pain medication hampered nurses to provide adequate pain management [11–13]. In addition, studies that described pain experience of ICU patients showed that most patients waited for the nurse to ask them about their pain before communicating its presence [14]. Therefore, it can be hypothesized that a lower nurse to patient ratio, higher bed occupancy rate, and a reduced presence of intensivists are associated with less patients receiving adequate pain management. To our knowledge no studies have quantitatively investigated the association

Abbreviations: Fte, full-time equivalent; NICE, national intensive care evaluation; VAS, Visual Analog Scale; NRS, Numeric Rating Scale; BPS, Behavioral Pain Scale; CPOT, Critical-Care Pain Observation Tool; EHR, Electronic Health Record; CPOE, computerized provider order entry.

* Corresponding author at: Amsterdam UMC, University of Amsterdam, Department of Medical Informatics, Room J1B-113.2, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.

E-mail address: m.blom@amsterdamumc.nl (M.-J. Roos-Blom).

Table 1
Description of patient characteristics of the 13 included intensive care units.

Patient characteristic	Admission count (%)	Patient-shifts with at least one pain measurement, %
ICU admissions	8136	81.2 (65,565/80743)
Gender male	5140 (63.2)	81.1 (42,940/52971)
Admission type		
Medical	3714 (45.6)	79.8 (40,745/51068)
Surgical	4422 (54.4)	83.6 (24,820/29675)
Mechanical ventilation first 24 h	4497 (55.3)	80.7 (44,527/55170)
Age in years ^a	Median (IQR) 67.0 (57.0–74.0)	
APACHE IV mortality probability ^a	Mean (SD) 0.17 (0.23)	

^a Presented as median with interquartile range (IQR) or mean and standard deviation (SD), but included as splines in the regression analysis

between these ICU organizational characteristics and pain management before. Within this multicenter study we aim to gain more insight into the organizational characteristics associated with adequate pain management in Dutch ICU patients during their admission.

2. Materials and methods

2.1. Data collection

In the Netherlands, the National Intensive Care Evaluation (NICE) registry, enables ICUs to monitor and improve their quality of care. The NICE registry provides all 83 Dutch ICUs with audit and feedback (A&F) information on patient outcomes such as mortality and length of stay, and optional on topics such as organizational characteristics, complications, sepsis and sequential organ failure [15]. In 2017, the NICE registry developed and implemented a web based module with four actionable quality indicators in order to improve quality of pain management [16]. The data needed to calculate the pain indicators consist of date and time of pain assessment, pain score, and type of assessment tool [17]. Pain measurements were performed in patients at rest, usually by nurses, but also by physicians, and measured with validated measurement instruments. The Visual Analog Scale (VAS) or the Numeric Rating Scale (NRS) was used in patients able to self-report pain and the Behavioral Pain Scale (BPS) or Critical-Care Pain Observation Tool (CPOT) in patients not able to self-report, e.g. sedated or mechanically ventilated patients [8,18,19].

We used data from thirteen (15.7%) mixed medical-surgical ICUs that voluntarily engaged in the pain management module of the NICE registry and for which data on organizational characteristics and pain management was available for all patients admitted between December 2017 and June 2018. The ICUs extracted these data from their electronic health record (EHR) in addition to their regular uploaded NICE data on patient demographics, physiological and diagnostic data such as comorbidities and reason of admission [15]. Patients who were not delirious or comatose and had a Glasgow coma score ≥ 8 at the first day of ICU admission were eligible for inclusion, because the pain instruments are only validated for these patients [20].

2.2. Outcome measure

Our outcome measure was the proportion of ICU patients per shift with adequate pain management. Our unit of observation is a patient shift as the availability of nurses and intensivists, and bed occupancy rate change over shifts. Adequate pain management for a patient during a shift (day, evening or night shift) was defined as ‘yes (1)’ when at least one pain measurement was performed during that shift and all measurements performed had acceptable pain scores OR in case of an

unacceptable pain score the pain was re-measured and normalized within 1 h [16,21]. A pain measurement had an unacceptable score when VAS/NRS > 3 , CPOT > 2 and BPS > 5 [8,18,19]. We excluded the first shift of patients’ ICU admission because presence of pain is not in control of the ICU when patients from the operating room or from the emergency room arrive at the ICU with high pain scores.

2.3. Organizational determinants of adequate pain management

Based on literature, expert opinion and the availability of data in the NICE registry we explored the impact of four potential organizational determinants of adequate pain management: nurse to patient ratio, bed occupancy rate, intensivist presence (hours) during workdays and intensivist presence (hours) during weekends. Nurse to patient ratio was determined by dividing the number of full-time equivalent (Fte) ICU certified nurses during a shift by the maximum number of patients present at the ICU during that shift. Bed occupancy rate was determined by dividing the maximum number of patients present at the ICU during a shift by the number of available ICU beds during that shift, multiplied by 100.

2.4. Data analysis

We performed unadjusted – i.e. univariate and case-mix adjusted – i.e. multivariate mixed-effect logistic regression modeling to investigate the association between each of the four organizational characteristics and adequate pain management. Nurse to patient ratio and bed occupancy rate were included in the models as quintiles and intensivist presence during workdays and weekends as dichotomous variables (< 24 h vs. full time coverage i.e. 24 h).

In 2007 Chanques et al. [4] showed that the intensity of NRS scores at rest were higher in ICU medical patients as compared to surgical-trauma patients. Other factors that have been shown to be associated with higher pain scores in ICU patients are a young age, number of comorbidities, colon cancer, and abdominal surgery [22–25]. Therefore, in each of the multivariate models we adjusted the effect of the organizational characteristic for age, gender, admission type (medical vs. surgical) and Acute Physiology and Chronic Health Evaluation (APACHE) IV mortality probability. We included APACHE IV mortality probability as an overall measure for severity of illness as it includes the combined information on comorbidities, physiological disturbance, and admission diagnosis and because the limited number of included ICUs ($n = 13$) requires a strict policy on the number of covariates in the model. Age and APACHE IV mortality probability were included in the model as restricted cubic splines to allow a non-linear relationship with adequate pain.

We included a random intercept for ‘ICU’ and for ‘patient admission’ to account for clustering effects of observations within ICUs and for repeated measurements within patients. We tested with ANOVA whether the model including a specific organizational characteristic improved compared to a model with only the case-mix variables and random

Table 2
Characteristics of the 13 included intensive care units.

Organizational characteristic	n = 13
Number of admissions mechanically ventilated in first 24 h ^a	182 (59–651)
Nurse to patient ratio, average per shift ^a	0.75 (0.68–0.84)
Bed occupancy rate, average % per shift ^a	73.9 (67.8–84.1)
Intensivist presence (hours)	
Working days, n (%)	
< 24.0	6 (46.2)
24.0	7 (53.8)
Weekends, n (%)	
< 24.0	6 (46.2)
24.0	7 (53.8)

^a Presented as median (interquartile range; IQR)




Fig. 1. Distribution of percentage of patients with adequate pain management per shift across the included intensive care units over a 6-month period.

intercepts for 'ICU' and 'patient' included. We defined improvement as a *p*-value smaller than 0.05.

As post-hoc analysis we repeated the multivariate analyses that resulted in a significant association for subgroups medical vs. surgical admissions and for patients mechanically ventilated in the first 24 h vs. not mechanically ventilated in the first 24 h. Statistical analysis were performed using R version 3.4.3 (R Foundation for Statistical Computing; Vienna, Austria).

3. Results

Of the thirteen included ICUs six were located in a general hospital, five in a teaching hospital and two in a university affiliated hospital. The median number of ICU beds was 14.0 (interquartile range (IQR); 8.0–30.2). Supplementary material 1 shows the flow of patient inclusion. Table 1 shows characteristics of the 8136 included patients together with the percentage of patients per shift with at least one pain measurement. Table 2 describes per organizational characteristic the median and interquartile range (IQR) of the thirteen included ICUs. For all ICUs the presence of intensivists appeared to be equal for weekdays and weekends, therefore this variable was combined in the regression analysis. The percentage of patients with adequate pain management during a specific shift ranged from 57.6% to 84.8% in the thirteen ICUs (Fig. 1).

Table 3 presents the unadjusted and adjusted odds ratios for the associations between the organizational determinants and adequate pain

management. The unadjusted nurse to patient ratio of 0.55 or higher was significantly associated with a higher percentage of patients with adequate pain management as compared to a nurse to patient ratio of 0.55 or less. This association remained significant for the nurse to patient ratios of 0.70 to 0.80 and over 0.80 after case-mix adjustment. Adding nurse to patient ratio to the model that only included the case-mix variables and random intercepts improved the model significantly (ANOVA $p < .001$). Bed occupancy rate and intensivist presence were unadjusted and adjusted not associated with adequate pain management and did not improve the model that only included the case-mix variables and random intercepts.

The post-hoc analysis (Table 4) showed that a case-mix adjusted nurse to patient ratio of 0.63 or higher was significantly associated with a higher percentage of patients with adequate pain management in medical and mechanically ventilated patients. For surgical and non-mechanically ventilated patients we found no association.

4. Discussion

We examined the association between four ICU organizational characteristics and adequate pain management. This study shows that nurse to patient ratio is significantly associated with adequate pain management in Dutch ICUs especially in medical and mechanically ventilated patients. For bed occupancy rate and presence of intensivists we did not find an association with adequate pain management.

Our finding that a higher nurse to patient ratio increases the percentage of patients with adequate pain management per shift confirms the results from earlier mostly qualitative studies in which ICU nurses indicated that a lack of manpower and time withheld them from adequate pain management [11,12,26]. However, this does not necessarily mean that patients admitted to an ICU with lower nurse to patients ratios experience more pain. An alternative explanation for our finding is that pain is treated appropriate at all ICUs, but ICUs with a higher nurse to patient ratio may have more time to record the normalized pain score into the EHR. However, complete registration is part of adequate pain management to avoid excessive use or side effects of pain medication due to missing normalized pain scores [27]. Our hypothesis was that a high bed occupancy rate would be associated with a decreased percentage of patients with adequate pain management during a shift. However, bed occupancy rate did not show to be associated with adequate pain management. A possible explanation is the fact that the bed occupancy rate does not account for patient turnover. Patient turnover is a factor within the ICU work environment that disrupts workflows associated with the nursing process resulting in increased nursing workload. Consequently, when the time cost of patient turnover exceeds the time available within a nurse's schedule this might result in less adequate

Table 3
Odds ratios (ORs) for adequate pain management of unadjusted and case-mix adjusted analyses.

Organizational characteristic	OR (95% CI)	p-value	Adjusted ^a OR (95% CI)	p-value
Nurse to patient ratio				
Up to 0.55	Reference		Reference	
0.55 to 0.63	1.06 (1.00–1.12)	0.047	1.06 (1.00–1.12)	0.05
0.63 to 0.70	1.07 (1.00–1.14)	0.04	1.06 (1.00–1.13)	0.06
0.70 to 0.80	1.14 (1.07–1.22)	<0.001	1.14 (1.07–1.21)	<0.001
Over 0.80	1.17 (1.09–1.25)	<0.001	1.16 (1.08–1.24)	<0.001
Bed occupancy rate				
Up to 69.0	Reference		Reference	
69.0 to 80.0	1.01 (0.95–1.08)	0.77	1.01 (0.95–1.08)	0.68
80.0 to 88.2	1.03 (0.96–1.11)	0.34	1.04 (0.97–1.11)	0.29
88.2 to 96.3	1.01 (0.94–1.09)	0.72	1.02 (0.94–1.09)	0.67
Over 96.3	1.02 (0.95–1.10)	0.53	1.02 (0.95–1.10)	0.53
Intensivist presence (hours)^b				
<24.0	Reference		Reference	
24	0.88 (0.41–1.88)	0.74	0.94 (0.43–2.08)	0.88

^a Adjusted for the case-mix variables gender, age, admission type (medical or surgical), and Acute Physiology and Chronic Health Evaluation (APACHE) IV mortality probability.

^b The results for working days and weekends are presented together because ICUs with an intensivist present 24 hours per day during working days had this also during weekends.

Table 4
Odds ratios (ORs) for adequate pain management of case-mix adjusted analyses within medical vs. surgical and mechanically ventilated vs. non-mechanically ventilated patients.

Organizational characteristic	Medical admission		Surgical admission		Mechanically ventilated		Non-mechanically ventilated	
	Adjusted ^a OR (95% CI)	p-value	Adjusted ^a OR (95% CI)	p-value	Adjusted ^b OR (95% CI)	p-value	Adjusted ^b OR (95% CI)	p-value
Nurse to patient ratio								
Up to 0.55	Reference		Reference		Reference		Reference	
0.55 to 0.63	1.05 (0.98–1.13)	0.19	1.08 (0.98–1.18)	0.12	1.06 (0.99–1.14)	0.08	1.06 (0.95–1.18)	0.28
0.63 to 0.70	1.10 (1.01–1.19)	0.02	1.01 (0.91–1.12)	0.83	1.08 (1.00–1.16)	0.05	1.04 (0.92–1.17)	0.52
0.70 to 0.80	1.20 (1.11–1.30)	<0.001	1.04 (0.94–1.15)	0.47	1.16 (1.07–1.25)	<0.001	1.10 (0.99–1.24)	0.09
Over 0.80	1.25 (1.14–1.35)	<0.001	1.01 (0.91–1.12)	0.88	1.22 (1.13–1.33)	<0.001	1.04 (0.93–1.17)	0.49

^a Adjusted for the case-mix variables gender, age and Acute Physiology and Chronic Health Evaluation (APACHE) IV mortality probability.

^b Adjusted for the case-mix variables gender, age, admission type (medical or surgical), and Acute Physiology and Chronic Health Evaluation (APACHE) IV mortality probability.

pain management [28]. However, when the increased need for nursing care can be accommodated by an increase in nursing staff, patient turnover is not a problem. We hypothesized that presence of intensivists was associated with adequate pain management as physicians prescribe pain medication which thereafter can be administered by the ICU nurses. The absence of an association between intensivist presence and adequate pain management might be explained by the availability of nurse-driven protocols. All included ICUs mentioned they had a pain protocol, but unfortunately we do not have any data in our dataset on whether this were nurse-driven protocols. A nurse-driven protocol enables ICU nurses to make decisions on their own e.g. about administering pain medication, without or with less consultation of the attending intensivist [29,30]. Another explanation might be the use of computerized provider order entry (CPOE) as these systems can improve pain control by reducing medication prescription errors or initiation of prompts should a intensivist fail to order pain medication for a patient who reports pain [31].

A strength of this study is that we performed multicenter analysis and we believe that our results are likely to be generalizable to other ICUs with similar organizational factors. Next, by analyzing the association of nurse to patient ratio and bed occupancy rate with adequate pain management per shift we took into account that these factors can differ per shift. A limitation of this study might be that selection bias has occurred because ICUs participated on a voluntarily basis. The ICUs that participated are more likely to have their pain management practice and organization well-arranged than those that did not participate. Another possible limitation is that we did not have information on measures that directly influence the process of pain management such as the availability of decision support in the electronic health record of the ICUs that prompt to measure or re-measure pain, or to evaluate the pain medication. Decision support may improve pain management [32,33], but it can also result in alert fatigue – i.e. health professionals ignoring the prompts [34]. Furthermore, the effectiveness of the prompts also depends on the quality of it. Future research on this type and other organizational characteristics might further unravel how ICUs might improve pain management.

5. Conclusion

We found a positive association between nurse to patient ratio and adequate pain management especially in medical and mechanically ventilated patients. This finding confirms the believes of nurses that a lack of manpower withheld them from adequate pain management and underpins the importance of sufficient nurse staffing levels.

Funding

The board of the NICE foundation financed this study and approved the study design, data collection and analysis, interpretation of the data.

Declaration of Competing Interest

None.

Acknowledgements

The authors would like to thank all the ICUs who participated in and delivered data to the NICE registry. Furthermore we would like to acknowledge F. Termorshuizen for his statistical advice.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jcrrc.2019.11.010>.

References

- [1] Jacobi J, Fraser GL, Coursin DB, Riker RR, Fontaine D, Wittbrodt ET, et al. Clinical practice guidelines for the sustained use of sedatives and analgesics in the critically ill adult. *Crit Care Med* 2002;30(1):119–41.
- [2] Barr J, Fraser GL, Puntillo K, Ely EW, Gelinas C, Dasta JF, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med* 2013;41(1):263–306.
- [3] Payen JF, Bosson JL, Chanques G, Mantz J, Labarere J, DOLOREA Investigators. Pain assessment is associated with decreased duration of mechanical ventilation in the intensive care unit: a post hoc analysis of the DOLOREA study. *Anesthesiology* 2009;111(6):1308–16.
- [4] Chanques G, Sebbane M, Barbotte E, Viel E, Eledjam JJ, Jaber S. A prospective study of pain at rest: incidence and characteristics of an unrecognized symptom in surgical and trauma versus medical intensive care unit patients. *Anesthesiology* 2007;107(5):858–60.
- [5] Payen JF, Chanques G, Mantz J, Hercule C, Auriant I, Leguillou JL, et al. Current practices in sedation and analgesia for mechanically ventilated critically ill patients: a prospective multicenter patient-based study. *Anesthesiology* 2007;106(4):687–95 (quiz 891–2).
- [6] Puntillo KA, Max A, Timsit JF, Vignoud L, Chanques G, Robleda G, et al. Determinants of procedural pain intensity in the intensive care unit. The Europain(R) study. *Am J Respir Crit Care Med* 2014;189(1):39–47.
- [7] Puntillo KA, White C, Morris AB, Perdue ST, Stanik-Hutt J, Thompson CL, et al. Patients' perceptions and responses to procedural pain: results from thunder project II. *Am J Crit Care* 2001;10(4):238–51.
- [8] Chanques G, Jaber S, Barbotte E, Violet S, Sebbane M, Perrigault PF, et al. Impact of systematic evaluation of pain and agitation in an intensive care unit. *Crit Care Med* 2006;34(6):1691–9.
- [9] Gelinas C, Arbour C, Michaud C, Vaillant F, Desjardins S. Implementation of the critical-care pain observation tool on pain assessment/management nursing practices in an intensive care unit with nonverbal critically ill adults: a before and after study. *Int J Nurs Stud* 2011;48(12):1495–504.
- [10] Joffe AM, Hallman M, Gelinas C, Herr DL, Puntillo K. Evaluation and treatment of pain in critically ill adults. *Semin Respir Crit Care Med* 2013;34(2):189–200.
- [11] Stevens B, Riahi S, Cardoso R, Ballantyne M, Yamada J, Beyene J, et al. The influence of context on pain practices in the NICU: perceptions of health care professionals. *Qual Health Res* 2011;21(6):757–70.
- [12] Wang HL, Tsai YF. Nurses' knowledge and barriers regarding pain management in intensive care units. *J Clin Nurs* 2010;19(21–22):3188–96.
- [13] Kizza JB, Mulira JK. Nurses' pain assessment practices with critically ill adult patients. *Int Nurs Rev* 2015;62(4):573–82.
- [14] Gelinas C. Management of pain in cardiac surgery ICU patients: have we improved over time? *Intensive Crit Care Nurs* 2007;23(5):298–303.
- [15] van de Klundert N, Holman R, Dongelmans DA, de Keizer NF. Data resource profile: the Dutch National Intensive Care Evaluation (NICE) registry of admissions to adult intensive care units. *Int J Epidemiol* 2015;44(6):1850–h.
- [16] Roos-Blom MJ, Gude WT, Spijksra JJ, de Jonge E, Dongelmans D, de Keizer NF. Measuring quality indicators to improve pain management in critically ill patients. *J Crit Care* 2019;49:136–42.
- [17] Dutch National Intensive Care Evaluation foundation (NICE). Data dictionary: pain indicators. [https://stichting-nice.nl/dd/#793](https://stichting-nice.nl/dd/#793;); 2017. [accessed October 5, 2017].

- [18] Gelinas C, Harel F, Fillion L, Puntillo KA, Johnston CC. Sensitivity and specificity of the critical-care pain observation tool for the detection of pain in intubated adults after cardiac surgery. *J Pain Symptom Manage* 2009;37(1):58–67.
- [19] Ahlers SJ, van Gulik L, van der Veen AM, van Dongen HP, Bruins P, Belitser SV, et al. Comparison of different pain scoring systems in critically ill patients in a general ICU. *Crit Care* 2008;12(1):R15.
- [20] Gelinas C, Puntillo KA, Joffe AM, Barr J. A validated approach to evaluating psychometric properties of pain assessment tools for use in nonverbal critically ill adults. *Semin Respir Crit Care Med* 2013;34(2):153–68.
- [21] Gude WT, Roos-Blom MJ, van der Veer SN, de Jonge E, Peek N, Dongelmans DA, et al. Electronic audit and feedback intervention with action implementation toolbox to improve pain management in intensive care: protocol for a laboratory experiment and cluster randomised trial. *Implement Sci* 2017;12(1):68.
- [22] Al Sutari MM, Abdalrahim MS, Hamdan-Mansour AM, Ayasrah SM. Pain among mechanically ventilated patients in critical care units. *J Res Med Sci* 2014;19(8):726–32.
- [23] Puntillo K, Weiss SJ. Pain: its mediators and associated morbidity in critically ill cardiovascular surgical patients. *Nurs Res* 1994;43(1):31–6.
- [24] Desbiens NA, Wu AW, Broste SK, Wenger NS, Connors Jr AF, Lynn J, et al. Pain and satisfaction with pain control in seriously ill hospitalized adults: findings from the SUPPORT research investigations. For the SUPPORT investigators Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. *Crit Care Med* 1996;24(12):1953–61.
- [25] Carroll KC, Atkins PJ, Herold GR, Mlcek CA, Shively M, Clopton P, et al. Pain assessment and management in critically ill postoperative and trauma patients: a multisite study. *Am J Crit Care* 1999;8(2):105–17.
- [26] Shannon K, Bucknall T. Pain assessment in critical care: what have we learnt from research. *Intensive Crit Care Nurs* 2003;19(3):154–62.
- [27] Skrobik Y. Pain may be inevitable; inadequate management is not. *Crit Care* 2008;12(2):142.
- [28] VanFosson CA, Yoder LH, Jones TL. Patient turnover: a concept analysis. *ANS Adv Nurs Sci* 2017;40(3):298–310.
- [29] Durant DJ. Nurse-driven protocols and the prevention of catheter-associated urinary tract infections: a systematic review. *Am J Infect Control* 2017;45(12):1331–41.
- [30] van Valen R, van Vuuren H, van Domburg RT, van der Woerd D, Hofland J, Bogers AJ. Pain management after cardiac surgery: experience with a nurse-driven pain protocol. *Eur J Cardiovasc Nurs* 2012;11(1):62–9.
- [31] Blankenship JF, Rogers L, White J, Carey A, Fosnocht D, Hopkins C, et al. Prospective evaluation of the treatment of pain in the ED using computerized physician order entry. *Am J Emerg Med* 2012;30(8):1613–6.
- [32] Carrothers KM, Barr J, Spurlock B, Ridgely MS, Damberg CL, Ely EW. Contextual issues influencing implementation and outcomes associated with an integrated approach to managing pain, agitation, and delirium in adult ICUs. *Crit Care Med* 2013;41(9 Suppl 1):S128–35.
- [33] Grol R, Grimshaw J. From best evidence to best practice: effective implementation of change in patients' care. *Lancet* 2003;362(9391):1225–30.
- [34] Backman R, Bayliss S, Moore D, Litchfield I. Clinical reminder alert fatigue in healthcare: a systematic literature review protocol using qualitative evidence. *Syst Rev* 2017;6(1):255.