



Universiteit
Leiden
The Netherlands

Mechanical stress and inflammation have opposite effects on Wnt signaling in human chondrocytes

Timmermans, R.G.M.; Blom, A.B.; Nelissen, R.G.H.H.; Broekhuis, D.; Kraan, P.M. van der; Meulenbelt, I.; ... ; Ramos, Y.F.M.

Citation

Timmermans, R. G. M., Blom, A. B., Nelissen, R. G. H. H., Broekhuis, D., Kraan, P. M. van der, Meulenbelt, I., ... Ramos, Y. F. M. (2023). Mechanical stress and inflammation have opposite effects on Wnt signaling in human chondrocytes. *Journal Of Orthopaedic Research*. doi:10.1002/jor.25673

Version: Publisher's Version

License: [Creative Commons CC BY-NC-ND 4.0 license](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Downloaded from:

Note: To cite this publication please use the final published version (if applicable).

RESEARCH ARTICLE

Mechanical stress and inflammation have opposite effects on Wnt signaling in human chondrocytes

Ritchie G. M. Timmermans^{1,2}  | Arjen B. Blom²  | Rob G. H. H. Nelissen³  |
Demiën Broekhuis³  | Peter M. van der Kraan²  | Ingrid Meulenbelt¹  |
Martijn H. J. van den Bosch²  | Yolande F. M. Ramos¹ 

¹Department of Biomedical Data Sciences,
Section Molecular Epidemiology, Leiden
University Medical Center, Leiden,
The Netherlands

²Experimental Rheumatology, Radboud
University Medical Center, Nijmegen,
The Netherlands

³Department of Orthopedics, Leiden
University Medical Center, Leiden,
The Netherlands

Correspondence

Yolande F. M. Ramos, Department of
Biomedical Data Sciences, Section Molecular
Epidemiology, Leiden University Medical
Center, Einthovenweg 20, 2333 ZC Leiden,
The Netherlands.
Email: Y.F.M.Ramos@LUMC.nl

Funding information

Dutch Arthritis Society,
Grant/Award Number: DAF-17-1-401

Abstract

Dysregulation of Wingless and Int-1 (Wnt) signaling has been strongly associated with development and progression of osteoarthritis (OA). Here, we set out to investigate the independent effects of either mechanical stress (MS) or inflammation on Wnt signaling in human neocartilage pellets, and to relate this Wnt signaling to OA pathophysiology. OA synovium-conditioned media (OAS-CM) was collected after incubating synovium from human end-stage OA joints for 24 h in medium. Cytokine levels in the OAS-CM were determined with a multiplex immunoassay (Luminex). Human neocartilage pellets were exposed to 20% MS, 2% OAS-CM or 1 ng/mL Interleukin-1 β (IL-1 β). Effects on expression levels of Wnt signaling members were determined by reverse transcription-quantitative polymerase chain reaction. Additionally, the expression of these members in articular cartilage from human OA joints was analyzed in association with joint space narrowing (JSN) and osteophyte scores. Protein levels of IL-1 β , IL-6, IL-8, IL-10, tumor necrosis factor α , and granulocyte-macrophage colony-stimulating factor positively correlated with each other. MS increased noncanonical *WNT5A* and *FOS* expression. In contrast, these genes were downregulated upon stimulation with OAS-CM or IL-1 β . Furthermore, Wnt inhibitors *DKK1* and *FRZB* decreased in response to OAS-CM or IL-1 β exposure. Finally, expression of *WNT5A* in OA articular cartilage was associated with increased JSN scores, but not osteophyte scores. Our results demonstrate that MS and inflammatory stimuli have opposite effects on canonical and noncanonical Wnt signaling in human neocartilage. Considering the extent to which MS and inflammation contribute to OA in individual patients, we hypothesize that targeting specific Wnt pathways offers a more effective, individualized approach.

KEYWORDS

human neocartilage, inflammation, mechanical stress, osteoarthritis, Wnt signaling

Martijn H. J. van den Bosch and Yolande F. M. Ramos shared last authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Journal of Orthopaedic Research*® published by Wiley Periodicals LLC on behalf of Orthopaedic Research Society.

1 | INTRODUCTION

Although considered a whole joint disease, the most prominent hallmark of osteoarthritis (OA) is the progressive degradation of the articular cartilage.¹ Articular cartilage is a highly specialized tissue, responsible for smooth movement of the joints and distribution of mechanical pressure.² The extracellular matrix (ECM) of the cartilage contains primarily collagens and proteoglycans. The negative charge of the proteoglycans attracts water into the cartilage that acts as a buffer zone for the underlying subchondral bone. With cartilage loading, water is being pushed out of the cartilage, causing a strain on the tissue. While physiological mechanical strains promote beneficial effects on the cartilage, hyperphysiological mechanical strains or mechanical stress (MS) can cause damage and dysregulate critical underlying pathways leading to a catabolic and proinflammatory response.³ This is one of the causes of inflammation of the synovium, which occurs in >70% of patients suffering from all grades of knee OA.⁴ Together, MS and inflammation create microtraumas in the articular cartilage.⁵ However, the degree to which MS and synovial inflammation contribute to cartilage degeneration in OA can be expected to highly differ between patients, making these important parameters to study separately to open new avenues for more personalized care and treatment.

The Wntless and Int-1 (Wnt) signaling pathway is an evolutionary highly conserved pathway that plays a critical role in both embryonic development and tissue homeostasis throughout life. Wnt proteins activate different Wnt pathways, depending on the Wnt ligand, frizzled receptor, and coreceptors. Either the β -catenin-dependent canonical Wnt pathway or the noncanonical Wnt pathway is activated, where the latter can be further divided into the planar cell polarity (PCP) and the Wnt/ Ca^{2+} pathway.^{6–8} Wnt signaling is usually tightly regulated and dysregulation of Wnt signaling has been linked to severe skeletal disorders.⁹ Furthermore, it is evident that aberrant Wnt signaling in the joint can play a key role in development and progression of OA.^{8,10–15} Overactive Wnt signaling, due to decreased levels of Wnt inhibitors, such as *FRZB* and *DKK1* and increased levels of Wnt ligands, such as *WNT5A* and *WNT5B*, can lead to chondrocyte hypertrophy, increased catabolic activity and decreases anabolic activity, resulting in cartilage degeneration.^{6,13} Whereas it is clear that Wnt signaling is dysregulated in OA, determining the extent to which MS and inflammation individually cause aberrant Wnt signaling may contribute to a better understanding of the effectiveness of Wnt-based treatments in patients prone to develop OA, or decrease progression of patients with early to moderate OA.

Here, we set out to investigate how MS and inflammation individually affect Wnt signaling. Thereto, we employed human primary articular OA chondrocytes into a three-dimensional (3D) neocartilage pellet model with high-quality physiological cartilage. We exposed neocartilage to MS, which induced cartilage damage as was previously demonstrated,¹⁶ or exposed neocartilage to OA synovium-conditioned media (OAS-CM) or Interleukin-1 β (IL-1 β). For the OAS-CM, we characterized the inflammatory status based on the

concentration of different OA-relevant cytokines. Subsequently, we determined the effects of these OA-related stimuli on the expression of ligands, inhibitors, transcription factors, and receptors of canonical and noncanonical Wnt signaling that are dysregulated in OA.¹⁷ Finally, we investigated the relation between messenger RNA (mRNA) levels of these Wnt members in OA articular cartilage and radiographic scores.

2 | METHODS

2.1 | Conditioned media from OA synovium

OAS-CM was obtained as described previously.¹⁸ In short, synovium from anonymous patients undergoing total knee joint replacement surgery as a result of end-stage OA ($n = 10$) was collected in the Radboud University Medical Center and cultured for 24 h in DMEM with 0.1% BSA (0.3 g tissue/mL). Debris was removed by centrifugation and aliquots of OAS-CM were stored at -20°C until further use.

2.2 | Immunoassays

Protein levels of IL-1 β , IL-4, IL-6, IL-8, IL-10, IL-17, tumor necrosis factor α (TNF- α), interferon- γ (IFN- γ), granulocyte-macrophage colony-stimulating factor (GM-CSF), and MCP-1 in the OAS-CM were measured with a multiplex immunoassay on a Bio-Plex 200 system using a multiplex cytokine kit (Luminex, Bio-Rad Laboratories), according to the manufacturer's protocol.

2.3 | Cell culture

The effects of MS and a proinflammatory environment on Wnt signaling were investigated in our previously established human 3D in vitro neocartilage pellet model.^{16,19} Human primary articular chondrocytes were isolated from macroscopically preserved areas of anonymized patients ($n = 12$) undergoing total knee joint replacement due to end-stage OA included in the Research in Articular Osteoarthritis Cartilage (RAAK) study²⁰ at Leiden University Medical Center (Leiden, The Netherlands), or at Radboud University Medical Center (Nijmegen, the Netherlands). There was no overlap between donors of synovium tissue and primary chondrocytes. Human primary articular chondrocytes were expanded in culture for two passages in DMEM (high glucose, Gibco), supplemented with 10% fetal bovine serum (FBS, Biowest), antibiotics (100 U/mL penicillin and 100 $\mu\text{g/mL}$ streptomycin, Gibco), and 5 ng/mL bFGF (Peprotech). Subsequently, pellets (2.5×10^5 cells/pellet) were created by centrifugation. The following day, the medium was replaced with a serum-free chondrogenic medium (CDM), consisting of DMEM (high glucose, Gibco), supplemented with 50 $\mu\text{g/mL}$ ascorbic acid (Sigma-Aldrich), 40 $\mu\text{g/mL}$ L-proline (Sigma-Aldrich), 100 $\mu\text{g/mL}$ sodium pyruvate (Sigma-Aldrich), 0.1 μM dexamethasone (Sigma-Aldrich), 1% insulin-transferrin-selenium +, antibiotics (100 U/mL

penicillin and 100 µg/mL streptomycin, Gibco), and 10 ng/mL human transforming growth factor-β1 (hTGF-β1) (Peprotech) and refreshed every 3–4 days, as previously described.^{16,19}

2.4 | Exposure to MS, IL-1β, and OAS-CM

MS was applied using a Mach-1 mechanical testing system (Biomomentum) as previously described.¹⁶ At two consecutive days, neocartilage pellets (*n*=12 donors) were transferred to a sterile 0.9% phosphate buffered saline solution (B. Braun) and subjected twice to MS (10 min, 20% strain, frequency of 5 Hz, with a 10-min recovery interval between MS), and pellets were collected 12 h after MS. To largely reach a uniformity of strain for each donor, the diameters of the neocartilage pellets within each donor were measured and averaged, where the standard deviation of the average diameter was less than 5% of the average diameter. Before 24-h incubation with OAS-CM (2%) or IL-1β (1 ng/mL, Bio-Techne), pellets were cultured in the absence of dexamethasone (4 days prior) and TGF-β (1 day prior) to avoid the inhibition of an inflammatory response. To reflect biological variation across patients, three neocartilage pellets from each donor were exposed separately to 2% OAS-CM of three independent donors.

2.5 | Reverse transcription-quantitative polymerase chain reaction (RT-qPCR)

Per donor, two replicate neocartilage pellets were collected in TRIzol (Invitrogen). RNA was isolated using the RNeasy Mini Kit (Qiagen)

according to the manufacturer's protocol, and was treated with RNase-Free DNase (Qiagen) to remove DNA contamination. RNA concentrations and quality were measured using the NanoDrop. RNA was reverse-transcribed into complementary DNA (cDNA) using the Transcriptor First Strand cDNA Synthesis Kit (Roche). Subsequently, cDNA was amplified using FastStart SYBR Green Master (Roche) and mRNA expression was measured in triplicates in a MicroAmp Optical 384-Well Reaction Plate (ThermoFisher Scientific), using the QuantStudio Flex Real-Time PCR system (Applied Biosystems), with the following cycling conditions: 10 min 95°C; 10 s 95°C, 30 s 60°C, 20 s 72°C (45 cycles); 1 min 65°C and 15 s 95°C. Primer efficiency was tested using a cDNA dilution series, and primers with an efficiency between 90% and 110% were considered reliable. *GADPH* and *SDHA* were used as housekeeping genes. Fold changes (FCs) were calculated using the $2^{-\Delta\Delta C_t}$ method with $\Delta C_t = C_{t_{\text{gene of interest}}} - C_{t_{\text{average housekeeping genes}}}$ and $\Delta\Delta C_t = \Delta C_{t_{\text{condition}}} - \Delta C_{t_{\text{Control}}}$. Genes were selected based on their differential expression in OA articular cartilage and/or were representative of either canonical or non-canonical Wnt signaling.¹⁷ Primer sequences are listed in Table 1.

2.6 | Association of mRNA levels in preserved OA cartilage with radiographic scores

Gene expression levels of Wnt signaling members from macroscopically unaffected or preserved OA cartilage (RAAK study; *N* = 37) were extracted from our previously established RNA sequencing data set (Table 2).^{17,19,21} Radiographic scores (Kellgren–Lawrence, joint space narrowing [JSN], and osteophytes) were previously determined

TABLE 1 Primer sequences.

Gene	Forward sequence	Reverse sequence
<i>GAPDH</i>	5'-TGCCATGTAGACCCCTTGAAG-3'	5'-ATGGTACATGACAAGGTGCGG-3'
<i>SDHA</i>	5'-TGGAGCTGCAGAACCTGATG-3'	5'-TGTAAGTCTTCCTTGGCATGC-3'
<i>WNT5A</i>	5'-TCGCCAAGGAGTTTGTGGAT-3'	5'-CCATCTTATACACAGCCCTGC-3'
<i>WNT5B</i>	5'-TCGCCAAGGAGTTTGTGGAT-3'	5'-CCATCTTATACACAGCCCTGC-3'
<i>WNT9A</i>	5'-GTTCCAGTTCGCTTTGAGC-3'	5'-AAGGCAGTCTCCTTGAAGCC-3'
<i>WNT16</i>	5'-AACACCACGGGCAAAGAAAAC-3'	5'-ATCAACTTGCGACAGCCT-3'
<i>TCF7L1</i>	5'-CGATCCAAAGACAGGAATCCCC-3'	5'-GATTGTGTCGACAGCTCCG-3'
<i>TCF7L2</i>	5'-CCGACGTAGACCCCAAAACA-3'	5'-ATTGTCTACGGTGCCAGG-3'
<i>LEF1</i>	5'-GGTAGCAATTGGCAGCCCTA-3'	5'-CAAACCAGTCTGCTGAACGC-3'
<i>CCN4</i>	5'-CCTGCTTGGGGTTCATAGGA-3'	5'-TGGGTCTCTCAAGGCTCTGT-3'
<i>JUN</i>	5'-GAGCTGGAGCGCCTGATAAT-3'	5'-CCCTCCTGCTCATCTGTAC-3'
<i>FOS</i>	5'-GGAGAATCCGAAGGGAAGGA-3'	5'-AGTTGGTCTGTCTCCGCTTG-3'
<i>DKK1</i>	5'-TGACAACCTACCAGCCGTACC-3'	5'-CAGGCGAGACAGATTTGCAC-3'
<i>FRZB</i>	5'-ATTGACTTCCAGCACGAGCC-3'	5'-CGAGTGGCGGTACTTGATGAG-3'
<i>FZD1</i>	5'-AGTTCGTGCCAATCCTGACA-3'	5'-TGTTCAAAGGAGCCTGCGAA-3'
<i>FZD5</i>	5'-AGCACCATCGTTGTGTCTTGA-3'	5'-TCCACCAATCACCATCGCTT-3'

TABLE 2 Baseline characteristics of RAAK samples with OA phenotypes.

Characteristic	n = 37
Age-year	
Mean (SD)	69.8 (7.7)
Sex-no (%)	
Female	29 (78%)
Male	8 (22%)
Joint type-no (%)	
Hip	12 (32%)
Knee	25 (68%)
KL score (SD)	2.7 (0.6)
JSN score (SD) ^a	4.7 (1.2)
OP score (SD) ^a	2.8 (2.1)

Abbreviations: JSN, joint space narrowing; KL score, Kellgren–Lawrence score; OA, osteophytosis; OP, osteophytes; RAAK, Research in Articular Osteoarthritis Cartilage.

^aThe JSN score (0–6) or OP score (0–6) is defined as the sum of JSN grades (0–3) or OP grades (0–3), determined at the superior and inferior areas of the acetabular–femoral hip joint or at the medial and lateral areas of the tibiofemoral knee joint, respectively.

by two experienced readers prior and independent of the current study.²¹ In short, osteophytes and JSN were graded using the OARSI atlas, with 0–3 scores for lateral and medial compartments of the tibiofemoral knee joints and for inferior and superior compartments of the acetabular–femoral hip joint by consensus opinion. Association of gene expression levels with radiographic scores was determined by applying a generalized estimation equation (GEE).

2.7 | Statistical analysis

Graph creation and statistics were carried out in GraphPad Prism v.9, unless stated otherwise. Correlations between the cytokines were carried out using a Spearman correlation test. Statistical significance of mean differences in $-\Delta Ct$ levels between MS, OAS-CM, and IL-1 β with their respective controls was estimated using a GEE. Since perfect pairs were absent, by applying a GEE, we could adjust for dependencies among donors by adding a random effect for sample donors. We followed a linear GEE model, with $-\Delta Ct$ levels as dependent variable and condition as a factor. Associations of mRNA levels in preserved OA cartilage with JSN score and osteophytosis (OP) score were carried out with a linear GEE model, using age, sex, KL score, and joint type as covariates, and are depicted as odds ratios (OR).^{16,22} GEE analyses were carried out using IBM SPSS Statistics 25. The cytokine concentrations and FCs are described using the mean $\pm$ difference between 95% CI and the mean. ORs are described using the mean and the 95% CI. Differences in expression levels of Wnt genes in preserved OA articular cartilage from knees and hips

were calculated with a linear GEE model, using age and sex as covariates. *p* Values of the correlation analysis, RT-qPCR analysis, associations of Wnt signaling with radiographic scores, and differences in gene expression between knees and hips were corrected for multiple testing using the Benjamini–Hochberg method, with a significant cutoff value of 0.05, as is indicated by the false discovery rate (FDR).

3 | RESULTS

3.1 | Cytokine levels in undiluted conditioned media derived from OA synovium

To gain insight into the level of inflammatory factors secreted from OA synovium, we performed a Luminex assay of the OAS-CM to determine concentrations of 10 different OA-relevant cytokines. As shown in Figure 1, high levels of IL-6 ($1.31 \times 10^5 \pm 9.74 \times 10^4$ pg/mL) and IL-8 ($1.56 \times 10^5 \pm 1.39 \times 10^5$ pg/mL) were observed. Relative to these two cytokines, we detected low levels of IL-1 β (190.7 ± 158.1 pg/mL), IL-10 (100.6 ± 62.5 pg/mL), TNF- α ($1.16 \times 10^3 \pm 1.01 \times 10^3$ pg/mL), IFN- γ (0.9 ± 0.6 pg/mL), GM-CSF (29.3 ± 41.3 pg/mL), and monocyte chemoattractant protein-1 (MCP-1) (623.1 ± 74.6 pg/mL). Furthermore, IL-4 and IL-17 were too low to detect (*data not shown*).

To investigate the coherency of secreted cytokine levels between patients, we explored correlations between the different cytokines. Overall, this showed significant positive correlations between levels of IL-1 β , IL-6, IL-8, IL-10, TNF- α , and GM-CSF (Figure 2). Notably, MCP-1 did not show any significant correlation with the other cytokines. Some of the strongest correlations were observed for IL-8 with IL-6 ($\rho = 0.92$, $FDR = 1.22 \times 10^{-2}$) and GM-CSF ($\rho = 0.89$, $FDR = 1.22 \times 10^{-2}$).

3.2 | Effects of MS and inflammation on canonical and noncanonical Wnt signaling

To determine the individual effects of MS and inflammation on Wnt signaling, we measured mRNA levels of genes involved in Wnt signaling by RT-qPCR. As shown in Figure 3, upon applying 20% MS, specifically mRNA expression of noncanonical Wnt ligand *WNT5A* ($FC = 3.62 \pm 3.10$, $FDR = 4.35 \times 10^{-2}$) and *WNT5B* ($FC = 1.85 \pm 0.66$, $FDR = 8.12 \times 10^{-2}$) was consistently upregulated, albeit not FDR significant for *WNT5B* (Figure 3A). Moreover, we observed a high and significant upregulation of transcription factor *FOS* ($FC = 18.12 \pm 19.47$, $FDR = 1.68 \times 10^{-6}$, Figure 3C). Notable in Figure 3D is the absence of response of Wnt ligand scavenger *FRZB* and canonical Wnt inhibitor *DKK1*, which are usually among the most consistent downregulated genes associated with OA pathophysiology.^{11,15}

On the other hand, exposure to 2% OAS-CM downregulated expression of noncanonical Wnt ligands *WNT5A* ($FC = 0.65 \pm 0.19$, $FDR = 9.27 \times 10^{-4}$), *WNT5B* ($FC = 0.49 \pm 0.11$, $FDR = 1.89 \times 10^{-3}$), and *WNT16* ($FC = 0.47 \pm 0.42$, $FDR = 1.87 \times 10^{-3}$, Figure 3A), and




FIGURE 1 Cytokine levels in undiluted OAS-CM. Levels of IL-1 β , IL-4, IL-6, IL-8, IL-10, IL-17, TNF- α , IFN- γ , GM-CSF and MCP-1, measured by Luminex in OAS-CM ($n = 10$). Graphs depict the mean with 95% confidence intervals, and each colored dot represents a single OAS-CM. GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN- γ , interferon- γ ; IL-1 β , Interleukin-1 β ; MCP-1, monocyte chemoattractant protein-1; OA, osteophytosis; OAS-CM, OA synovium-conditioned media; TNF- α , tumor necrosis factor α .




FIGURE 2 Correlation matrix between cytokine levels in OAS-CM ($n = 10$). Correlations were calculated using a Spearman correlation test. p Values were corrected for multiple testing using the Benjamini-Hochberg method. Only significant correlations ($FDR \leq 0.05$) display the Spearman (ρ) correlation coefficient. FDR, false discovery rate; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN- γ , interferon- γ ; IL, Interleukin; MCP-1, monocyte chemoattractant protein-1; OAS-CM, OA synovium-conditioned media; TNF- α , tumor necrosis factor α .

noncanonical transcription factor *JUN* ($FC = 0.64 \pm 0.16$, $FDR = 9.27 \times 10^{-4}$, Figure 3C). Concurrently, exposure to 2% OAS-CM decreased expression of the canonical transcription factor *TCF7L1* ($FC = 0.63 \pm 0.18$, $FDR = 2.68 \times 10^{-3}$, Figure 3B), *DKK1* ($FC = 0.66 \pm 0.14$, $FDR = 3.44 \times 10^{-2}$), and *FRZB* ($FC = 0.51 \pm 0.13$, $FDR = 2.48 \times 10^{-4}$) (Figure 3D). Moreover, frizzled receptors *FZD1* ($FC = 0.66 \pm 0.16$, $FDR = 2.15 \times 10^{-4}$) and *FZD5* ($FC = 0.71 \pm 0.29$, $FDR = 9.98 \times 10^{-5}$, Figure 3E) were down-regulated upon exposure to 2% OAS-CM while the response to MS was less consistent. Furthermore, we compared the effects of OAS-CM on Wnt signaling to IL-1 β , a cytokine commonly used to mimic OA-associated processes in vitro although our data indicate its levels are relatively low in OAS-CM. We observed overall similar effects of OAS-CM and IL-1 β on the expression of Wnt ligands and inhibitors (Supporting Information Figure S1).

3.3 | Associations of Wnt mRNA levels in OA articular cartilage of joints with JSN and osteophytes

Given the fact that the responses of members of Wnt signaling to the OA-related triggers (20% MS and 2% OAS-CM) are highly specific and different, we next wanted to explore whether these different responses could mark endotypes of that pathophysiological process in preserved cartilage. Hereto, we related expression patterns of these Wnt members in preserved regions of OA articular cartilage to




FIGURE 3 Mechanical stress and OAS-CM have opposite effects on Wnt signaling in human neocartilage. (A) Wnt ligands. (B) Canonical TFs and canonical target *CCN4/WISP1*. (C) Noncanonical TF *JUN* and its AP-1 associated TF *FOS*. (D) Canonical Wnt inhibitor *DKK1* and Wnt ligand scavenger *FRZB*. (E) Frizzled receptors 1 and 5. Lines represent differences in mRNA expression levels ($n = 12$) between control and 20% MS or control and OAS-CM within a single donor. *FDR ≤ 0.05 , **FDR ≤ 0.01 , and ***FDR ≤ 0.001 . AP-1, activator protein 1; FDR, false discovery rate; mRNA, messenger RNA; MS, mechanical stress; OA, osteophytosis; OAS-CM, OA synovium-conditioned media; OAS-CM, OA synovial conditioned media; TF, transcription factor; Wnt, Wingless and Int-1.

TABLE 3 Associations of mRNA levels of Wnt signaling members with hallmarks of radiographic OA.

Gene	JSN score ^a					OP score ^a				
	OR ^b	95% CI		p Value ^b	FDR ^c	OR ^b	95% CI		p Value ^b	FDR ^c
		Lower	Upper				Lower	Upper		
WNT5A	1.65	1.27	2.15	0.00018	0.0050	1.05	0.88	1.25	0.6100	0.6832
WNT5B	1.18	0.97	1.44	0.1010	0.2724	1.07	0.96	1.19	0.2420	0.3640
WNT9A	0.87	0.71	1.05	0.1490	0.2980	1.15	1.02	1.29	0.0230	0.1073
WNT16	0.81	0.50	1.31	0.3820	0.5213	1.08	0.83	1.41	0.5570	0.6498
TCF7L1	1.04	0.91	1.20	0.5550	0.6498	1.00	0.92	1.08	0.9220	0.9220
TCF7L2	1.09	1.00	1.18	0.0450	0.1575	0.98	0.93	1.03	0.4680	0.5956
LEF1	1.63	1.07	2.47	0.0220	0.1073	1.14	0.91	1.42	0.2470	0.3640
CCN4	1.20	0.96	1.49	0.1070	0.2724	1.05	0.94	1.18	0.3910	0.5213
JUN	1.16	0.91	1.49	0.2240	0.3640	1.01	0.86	1.19	0.8670	0.8991
FOS	1.10	0.94	1.28	0.2430	0.3640	1.02	0.92	1.14	0.6930	0.7463
DKK1	0.74	0.55	0.98	0.0360	0.1440	0.89	0.75	1.04	0.1380	0.2972
FRZB	0.65	0.46	0.93	0.0170	0.1073	0.88	0.72	1.09	0.2310	0.3640
FZD1	1.16	0.98	1.38	0.0940	0.2724	1.12	1.02	1.23	0.0140	0.1073
FZD5	0.89	0.76	1.04	0.1380	0.2972	0.90	0.82	0.99	0.0220	0.1073

Abbreviations: CI, confidence interval; FDR, false discovery rate; JSN, joint space narrowing; KL score, Kellgren–Lawrence score; mRNA, messenger RNA; OA, osteophytosis; OP, osteophytes; OR, odds ratio; Wnt, Wingless and Int-1.

^aThe JSN score or OP score is defined as the sum of JSN grades (0–3) or OP grades (0–3), determined at the superior and inferior areas of the acetabular–femoral hip joint or at the medial and lateral areas of the tibiofemoral knee joint, respectively.

^bOdds ratios describe the association between mRNA levels and JSN grades or OP grades. Odds ratios and *p* values were generated by generalized estimating equation, with mRNA expression level in preserved OA cartilage as dependent variable, and age, sex, KL score, and joint type as covariates.

^c*p* Values were corrected for multiple testing using the Benjamini–Hochberg method, as indicated by the false discovery rate (FDR). FDR ≤ 0.05 are shown in bold.

hallmarks of radiographic OA (i.e., JSN and osteophytes scores; Table 3). We observed that *WNT5A* mRNA levels were significantly positively associated with JSN scores (OR = 1.65, 95% CI = 1.27, 2.15), independent of age, sex, KL score, and joint type. Albeit not FDR significant, decreased mRNA levels of *FRZB* (OR = 0.65, 95% CI = 0.46, 0.93) and *DKK1* (OR = 0.74, 95% CI = 0.55, 0.98) were also associated with higher JSN scores.

4 | DISCUSSION

In this study, we investigated the effects of two processes relevant for OA etiology, namely, MS and inflammation, on Wnt signaling in human primary chondrocyte neocartilage pellets. We observed that MS activated transcription of noncanonical Wnt ligands in neocartilage pellets, while exposure to proinflammatory stimuli resulted in inhibition. Furthermore, the inflammatory stimuli decreased the expression of (canonical) Wnt inhibitors, which could result in the activation of Wnt signaling. Finally, we associated *WNT5A* mRNA levels in preserved articular cartilage from OA joints with higher JSN scores.

As a multifactorial and complex disease, OA shows high heterogeneity in disease pathophysiology between patients. Many

but not all patients develop synovial inflammation in the joint, and physically active patients can be very different from inactive patients, considering the response of cartilage to mechanical events.²³ Moreover, while it is clear that overactive Wnt signaling extensively contributes to OA (progression), we now observed that MS and inflammation have different effects on canonical and noncanonical Wnt signaling. This is interesting in light of developing Wnt inhibitors specific for canonical and noncanonical Wnt signaling, which may be more effective in some patients as compared with others, due to the heterogeneous nature of OA. The latter is probably also related to the different effects that MS and synovial inflammation and the combination have on articular cartilage.

Here, we applied 2% OAS-CM which previously has been shown to reduce sulfated glycosaminoglycan (sGAG) production by 50% in pellet cultures of bone marrow-derived mesenchymal stem cells that were differentiated into chondrocytes.²⁴ Furthermore, the addition of tofacitinib and oxozaenol, in the presence of OAS-CM, has previously been demonstrated to rescue the sGAG content, confirming that factors secreted by the OA synovium are responsible for the observed effects.¹⁸ The high levels of IL-6 and IL-8 in the OAS-CM and the presence of other cytokines indicate a strong proinflammatory potential of the OAS-CM. Although the importance and relevance of IL-1β in OA is under discussion,²⁵ we observed

similar effects of exposure to OAS-CM and IL-1 β alone on most members of Wnt signaling. Therefore, IL-1 β seems a good model cytokine to study the effects of synovial inflammation on Wnt signaling in case OAS-CM is not available.

MS resulted in upregulation of *WNT5A* and *WNT5B*, with the latter not being FDR significant. These two ligands signal through the noncanonical pathway. While *WNT5A* and *WNT5B* are highly similar in structure, they have different functions in chondrocytes.²⁶ *WNT5A* induces cytokine production and senescence, whereas *WNT5B* stimulates collagen type I secretion and is responsible for inhibiting the production of ECM components and chondrocyte differentiation.^{27,28} In all donors, we observed an increase in expression of the transcription factor *FOS* and in 10 out of 12 donors for *JUN*. Although mostly *JUN* is associated with the noncanonical PCP/JNK pathway, *FOS* and *JUN* dimerize, forming the activator protein 1 (AP-1) transcription factor. AP-1, and can via inhibition of microRNA-30a (miR-30a) stimulate expression of the aggrecanase *ADAMTS5*, a matrix-degrading enzyme that plays an important role in cartilage degradation and OA.²⁹ We previously showed the MS-induced *ADAMTS5* expression in the neocartilage.¹⁶ The applied MS in this model may upregulate *ADAMTS5* via AP-1 and miR-30a, however, this remains to be established. In contrast, we observed opposite effects on noncanonical Wnt signaling after exposure to proinflammatory stimuli compared with MS. Both OAS-CM and IL-1 β exposure led to strong downregulation of the canonical Wnt inhibitor *DKK1* and Wnt ligand scavenger *FRZB*, which has previously been demonstrated for IL-1 β .¹¹ Multiple genome-wide association studies have identified a nonsynonymous single-nucleotide polymorphism residing in the *FRZB* gene that reduces the ability of *FRZB* to antagonize Wnt signaling, and linked this to increased risks of OA.^{15,30} Moreover, high levels of both *FRZB* and *DKK1* have been linked to reduced progression of OA.³¹ In line with this, we observed that both *DKK1* and *FRZB* mRNA levels, although not FDR significant, were negatively associated with JSN scores. Reduced *DKK1* and *FRZB* levels can lead to excessive canonical Wnt signaling with chondrocyte hypertrophy and increased protease activity as a consequence.⁶ We did not observe any effects on *DKK1* and *FRZB* in the mechanically stressed neocartilage. This is different compared with mechanically injured cartilage explants, which previously were shown to have decreased *DKK1* and *FRZB* expression.¹⁴ Chondrocytes may respond differently to certain types of MS and should be taken into consideration when selecting preclinical models for drug testing.³

WNT16 is a ligand that activates mostly noncanonical Wnt signaling and can only weakly activate canonical Wnt signaling in high doses. In certain situations, *WNT16* can counteract excessive canonical Wnt signaling. Additionally, *Wnt16*^{-/-} mice produce lower levels of lubricin and exhibit increased chondrocyte apoptosis, resulting in more severe OA, which suggests a protective effect of *Wnt16* on cartilage homeostasis.³² Moreover, *WNT16* is strongly upregulated in end-stage OA articular cartilage.¹⁷ However, we observed reduced expression of *WNT16* after exposure to OAS-CM and IL- β , while no effect of MS was observed. Late *WNT16* upregulation may be an attempt of the chondrocyte to protect

cartilage from excessive damage, while in early phases it may remain unchanged.^{33,34}

The members of the TCF/LEF family are transcription factors regulating the transcription of target genes from the β -catenin-dependent/canonical Wnt pathway. After exposure to IL-1 β , we observed that *TCF7L1* expression was decreased while *LEF1* was increased. *LEF1* is thought to act as a strong transcriptional activator and *TCF7L1* is considered a transcriptional repressor.³⁵ We speculate that exposure to IL-1 β may cause more active transcription of canonical Wnt target genes.

Previously, we have demonstrated the involvement of *CCN4/WISP1* in OA. *CCN4/WISP1* is upregulated in both cartilage and synovium in mice with experimental OA, and aggravates cartilage degeneration in multiple OA mice models.^{10,12} Furthermore, *CCN4/WISP1* expression is increased in lesioned areas from human OA cartilage compared with macroscopically preserved areas, and mostly present near cartilage lesions. Surprisingly, we did not find any effects of MS nor the proinflammatory stimuli on *CCN4/WISP1* in the neocartilage. Possibly, to detect *CCN4/WISP1* changes a different timing is required. Alternatively, in in vivo OA models, the *CCN4/WISP1* response is very likely a combination of MS and inflammation, which we here aimed to separate.

Commonly, CDM contains dexamethasone to promote chondrogenesis. Dexamethasone, however, has strong anti-inflammatory properties, and has been shown to inhibit Wnt signaling.³⁶ Therefore, to prevent that the dexamethasone in our study would counteract the inflammatory effects of the OAS-CM and IL-1 β , neocartilage pellets were cultured without dexamethasone 4 days before the exposure. This difference in composition of the CDM with OAS-CM as compared with the mechanically stressed neocartilage may have resulted in differences in basal expression levels of several Wnt genes in the control groups, such as *WNT9A* and *LEF1* (Figure 3) and should be taken into account in future studies.

Associating Wnt signaling in preserved OA cartilage to JSN scores and OP scores revealed, only higher *WNT5A* mRNA levels to be associated with higher JSN scores. We further observed that in knees, expression of 10 out of 14 genes was significantly lower (e.g., *WNT5A* and *WNT5B*) or higher (e.g., *FRZB*) compared with hips (Supporting Information Figure S2). In the neocartilage, MS increased *WNT5A* expression but not exposure to OAS-CM. Given these observations, it could be promising to further investigate the potential of *WNT5A* as a marker for MS. This would require more in-depth research measuring the long-term effects of OA-related pathologies on *WNT5A* in cartilage (e.g., ex vivo explants). Increased *WNT5A* levels have been shown to repress chondrogenic differentiation and synthesis of cartilage matrix components.³⁷ It is tempting to speculate that the increased levels of *WNT5A* we observed as the result of MS will affect the propensity of preserved cartilage to enter an OA disease state.

Taken together, MS activated the transcription of noncanonical Wnt signaling, while the inflammatory stimuli can promote canonical Wnt signaling by downregulating *DKK1* and *FRZB*. In addition, we found that *WNT5A* mRNA levels in preserved OA articular cartilage

are associated with a specific radiographic phenotype, namely, JSN. On the basis of these results, we think that during OA, certain patients may experience more MS-based events while others experience more inflammatory-based events. Therefore, the efficacy of inhibitors that specifically target canonical or noncanonical Wnt signaling may depend on the underlying etiology of the OA subtype and we advocate that targeting specific Wnt pathways could offer a more effective, individualized approach.

AUTHOR CONTRIBUTIONS

Study concept and design: Ritchie G. M. Timmermans, Arjen B. Blom, Ingrid Meulenbelt, Martijn H. J. van den Bosch, and Yolande F. M. Ramos. **Acquisition of material and data:** Ritchie G. M. Timmermans, Rob G. H. H. Nelissen, and Demiën Broekhuis. **Data analysis:** Ritchie G. M. Timmermans, Arjen B. Blom, Ingrid Meulenbelt, Martijn H. J. van den Bosch, and Yolande F. M. Ramos. **Preparation of the manuscript:** Ritchie G. M. Timmermans, Arjen B. Blom, Peter M. van der Kraan, Ingrid Meulenbelt, Martijn H. J. van den Bosch, and Yolande F. M. Ramos. **Critical reviewing and approval of the manuscript:** All authors.

ACKNOWLEDGMENTS

We thank all study participants of the RAAK study. The Leiden University Medical Center has and is supporting the RAAK. The RAAK study has been approved by the medical ethical committee of the Leiden University Medical Center (P08.239/P19.013). Written informed consent was obtained from all patients, and patients had the right to withdraw at any time. In the Radboud University Medical Center, patients were informed about the anonymized use of the material for research and had the right to decline the use of their material for research (2018-4319). We especially thank Henk van Beuningen (providing the OAS-CM), Monique Helsen (Luminex), and Niek Bloks (applying MS to the neocartilage). We thank all the members of our groups. We also thank Robert van der Wal, Enrike van der Linden, Peter van Schie, Shaho Hasan, Maartje Meijer, Daisy Latijnhouwers, Anika Rabelink-Hoogenstraaten, and Geert Spierenburg for their contribution to the collection of the joint tissue. Data are generated within the scope of the Medical Delta programs Regenerative Medicine 4D: Generating complex tissues with stem cells and printing technology and Improving Mobility with Technology. Furthermore, the research leading to these results has received funding from the Dutch Arthritis Society (DAF-17-1-401). All authors declare that they have nothing to disclose.

ORCID

Ritchie G. M. Timmermans  <http://orcid.org/0000-0002-0787-0856>
Arjen B. Blom  <https://orcid.org/0000-0001-9613-5070>
Rob G. H. H. Nelissen  <https://orcid.org/0000-0003-1228-4162>
Demiën Broekhuis  <https://orcid.org/0000-0002-3125-5028>
Peter M. van der Kraan  <https://orcid.org/0000-0002-2262-3802>
Ingrid Meulenbelt  <https://orcid.org/0000-0001-7786-7081>

Martijn H. J. van den Bosch  <https://orcid.org/0000-0001-8074-826X>

Yolande F. M. Ramos  <http://orcid.org/0000-0003-1459-413X>

REFERENCES

- Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the joint as an organ. *Arthritis Rheum.* 2012;64(6):1697-1707.
- Carballo CB, Nakagawa Y, Sekiya I, Rodeo SA. Basic science of articular cartilage. *Clin Sports Med.* 2017;36(3):413-425.
- Sanchez-Adams J, Leddy HA, McNulty AL, O'Connor CJ, Guilak F. The mechanobiology of articular cartilage: bearing the burden of osteoarthritis. *Curr Rheumatol Rep.* 2014;16(10):451.
- Guerhazi A, Hayashi D, Roemer FW, et al. Synovitis in knee osteoarthritis assessed by contrast-enhanced magnetic resonance imaging (MRI) is associated with radiographic tibiofemoral osteoarthritis and MRI-detected widespread cartilage damage: the MOST study. *J Rheumatol.* 2014;41(3):501-508.
- Li G, Yin J, Gao J, et al. Subchondral bone in osteoarthritis: insight into risk factors and microstructural changes. *Arthritis Res Ther.* 2013;15(6):223.
- Wang Y, Fan X, Xing L, Tian F. Wnt signaling: a promising target for osteoarthritis therapy. *Cell Commun Signaling.* 2019;17(1):97.
- van den Bosch MH, Gleissl TA, Blom AB, van den Berg WB, van Lent PL, van der Kraan PM. Wnts talking with the TGF- β superfamily: WISPers about modulation of osteoarthritis. *Rheumatology.* 2015;55(9):1536-1547.
- Lories RJ, Monteagudo S. Review article: is Wnt signaling an attractive target for the treatment of osteoarthritis? *Rheumatol Ther.* 2020;7(2):259-270.
- Huybrechts Y, Mortier G, Boudin E, Van Hul W. WNT signaling and bone: lessons from skeletal dysplasias and disorders. *Front Endocrinol.* 2020;11:165.
- Blom AB, Brockbank SM, van Lent PL, et al. Involvement of the Wnt signaling pathway in experimental and human osteoarthritis: prominent role of Wnt-induced signaling protein 1. *Arthritis Rheum.* 2009;60(2):501-512.
- Leijten JC, Bos SD, Landman EB, et al. GREM1, FRZB and DKK1 mRNA levels correlate with osteoarthritis and are regulated by osteoarthritis-associated factors. *Arthritis Res Ther.* 2013;15(5):R126.
- van den Bosch MH, Blom AB, Kram V, et al. WISP1/CCN4 aggravates cartilage degeneration in experimental osteoarthritis. *Osteoarthritis Cartilage.* 2017;25(11):1900-1911.
- van den Bosch MH, Ramos YFM, den Hollander W, et al. Increased WISP1 expression in human osteoarthritic articular cartilage is epigenetically regulated and decreases cartilage matrix production. *Rheumatology.* 2019;58(6):1065-1074.
- Dell'Accio F, De Bari C, El Tawil NM, et al. Activation of WNT and BMP signaling in adult human articular cartilage following mechanical injury. *Arthritis Res Ther.* 2006;8(5):R139.
- Loughlin J, Dowling B, Chapman K, et al. Functional variants within the secreted frizzled-related protein 3 gene are associated with hip osteoarthritis in females. *Proc Natl Acad Sci.* 2004;101(26):9757-9762.
- Timmermans RGM, Bloks NGC, Tuerlings M, et al. A human in vitro 3D neo-cartilage model to explore the response of OA risk genes to hyper-physiological mechanical stress. *Osteoarthritis Cartilage Open.* 2022;4(1):100231.
- Coutinho de Almeida R, Ramos YFM, Mahfouz A, et al. RNA sequencing data integration reveals a miRNA interactome of osteoarthritis cartilage. *Ann Rheum Dis.* 2019;78(2):270-277.
- van Beuningen HM, de Vries-van Melle ML, Vitters EL, et al. Inhibition of TAK1 and/or JAK can rescue impaired chondrogenic

- differentiation of human mesenchymal stem cells in osteoarthritis-like conditions. *Tissue Eng Part A*. 2014;20(15-16):2243-2252.
19. Bomer N, den Hollander W, Ramos YFM, et al. Underlying molecular mechanisms of DIO2 susceptibility in symptomatic osteoarthritis. *Ann Rheum Dis*. 2015;74(8):1571-1579.
 20. Ramos YFM, den Hollander W, Bovée JVMG, et al. Genes involved in the osteoarthritis process identified through genome wide expression analysis in articular cartilage; the RAAK study. *PLoS ONE*. 2014;9(7):e103056.
 21. Coutinho de Almeida R, Mahfouz A, Mei H, et al. Identification and characterization of two consistent osteoarthritis subtypes by transcriptome and clinical data integration. *Rheumatology*. 2020; 60(3):1166-1175.
 22. Houtman E, Coutinho de Almeida R, Tuerlings M, et al. Characterization of dynamic changes in Matrix Gla Protein (MGP) gene expression as function of genetic risk alleles, osteoarthritis relevant stimuli, and the vitamin K inhibitor warfarin. *Osteoarthritis Cartilage*. 2021;29(8):1193-1202.
 23. Griffin TM, Guilak F. The role of mechanical loading in the onset and progression of osteoarthritis. *Exerc Sport Sci Rev*. 2005;33(4):195-200.
 24. van Geffen EW, van Caam APM, Vitters EL, et al. Interleukin-37 protects stem cell-based cartilage formation in an inflammatory osteoarthritis-like microenvironment. *Tissue Eng Part A*. 2019;25(15-16):1155-1166.
 25. Vincent TL. IL-1 in osteoarthritis: time for a critical review of the literature. *F1000Research*. 2019;8:934.
 26. Suthon S, Perkins RS, Bryja V, et al. Corrigendum: WNT5B in physiology and disease. *Front Cell Dev Biol*. 2021;9:667581.
 27. Huang J, Chen C, Liang C, et al. Dysregulation of the Wnt signaling pathway and synovial stem cell dysfunction in osteoarthritis development. *Stem Cells Dev*. 2020;29(7):401-413.
 28. Tao SC, Yuan T, Zhang YL, Yin WJ, Guo SC, Zhang CQ. Exosomes derived from miR-140-5p-overexpressing human synovial mesenchymal stem cells enhance cartilage tissue regeneration and prevent osteoarthritis of the knee in a rat model. *Theranostics*. 2017;7(1):180-195.
 29. Ji Q, Xu X, Zhang Q, et al. The IL-1 β /AP-1/miR-30a/ADAMTS-5 axis regulates cartilage matrix degradation in human osteoarthritis. *J Mol Med*. 2016;94(7):771-785.
 30. Evangelou E, Chapman K, Meulenbelt I, et al. Large-scale analysis of association between GDF5 and FRZB variants and osteoarthritis of the hip, knee, and hand. *Arthritis Rheum*. 2009;60(6):1710-1721.
 31. Lane NE, Nevitt MC, Lui LY, de Leon P, Corr M. Wnt signaling antagonists are potential prognostic biomarkers for the progression of radiographic hip osteoarthritis in elderly Caucasian women. *Arthritis Rheum*. 2007;56(10):3319-3325.
 32. Nalesso G, Thomas BL, Sherwood JC, et al. WNT16 antagonises excessive canonical WNT activation and protects cartilage in osteoarthritis. *Ann Rheum Dis*. 2017;76(1):218-226.
 33. Karlsson C, Dehne T, Lindahl A, et al. Genome-wide expression profiling reveals new candidate genes associated with osteoarthritis. *Osteoarthritis Cartilage*. 2010;18(4):581-592.
 34. Xu Y, Barter MJ, Swan DC, et al. Identification of the pathogenic pathways in osteoarthritic hip cartilage: commonality and discord between hip and knee OA. *Osteoarthritis Cartilage*. 2012;20(9): 1029-1038.
 35. Sierra RA, Hoverter NP, Ramirez RN, et al. TCF7L1 suppresses primitive streak gene expression to support human embryonic stem cell pluripotency. *Development*. 2018;145(4):dev161075.
 36. Naito M, Ohashi A, Takahashi T. Dexamethasone inhibits chondrocyte differentiation by suppression of Wnt/ β -catenin signaling in the chondrogenic cell line ATDC5. *Histochem Cell Biol*. 2015;144(3): 261-272.
 37. Mao G, Zhang Z, Hu S, et al. Exosomes derived from miR-92a-3p-overexpressing human mesenchymal stem cells enhance chondrogenesis and suppress cartilage degradation via targeting WNT5A. *Stem Cell Res Ther*. 2018;9(1):247.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Timmermans RGM, Blom AB, Nelissen RGHH, et al. Mechanical stress and inflammation have opposite effects on Wnt signaling in human chondrocytes. *J Orthop Res*. 2023;1-10.
[doi:10.1002/jor.25673](https://doi.org/10.1002/jor.25673)