

Binge-eating disorder in the Arabic world and the Netherlands, assessment, etiology, efficacy, effectiveness and economic evaluation of psychological interventions

Melisse, B.

Citation

Melisse, B. (2023, September 13). Binge-eating disorder in the Arabic world and the Netherlands, assessment, etiology, efficacy, effectiveness and economic evaluation of psychological interventions.

Version: Publisher's Version

License: License agreement concerning inclusion of doctoral thesis in the

Institutional Repository of the University of Leiden

Downloaded from:

Note: To cite this publication please use the final published version (if applicable).

1. Chapter 1 General introduction

Background

Eating disorders have a significant impact on the psychological, social and physical wellbeing and quality of life of affected patients (Agh et al., 2015; Hay et al., 2015; Rojo-Moreno et al., 2015) including anxiety, depression, and even high rates of mortality (Smink et al., 2012; Watson et al., 2012). The conceptualization of eating disorders has expanded over the last 10 years (Hay, 2020). The fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM 5) specifies four types of eating disorders: anorexia nervosa (AN), bulimia nervosa (BN), binge eating disorder (BED) and otherwise specified feeding or eating disorder (OSFED). The OSFED category comprises patients who do not meet the behavioral frequency or all other criteria of AN, BN or BED (APA, 2013). However, until 2013 BED was a provisional diagnosis and therefore not included in the DSM. Therefore, BED is the most recently acknowledged eating disorder. Main reason for the revision of the DSM was the high frequency of eating disorder not otherwise specified (EDNOS) diagnoses. All other eating disorders of clinical significance that did not meet the criteria of AN or BN received the diagnosis EDNOS, even patients that barely missed the diagnostic threshold for AN or BN, or patients that met the criteria for BED (Smink et al., 2013; Walsh et al., 2015). Consequently, the diagnostic criteria for AN and BN were elaborated upon, and new disorders, such as BED were added in the DSM-5 (APA, 2013). The broader criteria of eating disorders are more likely to increase the clinical utility of the DSM-5 (Hay, 2020).

Eating disorders can either be approached from a diagnostic perspective, stipulating the differences among the various eating disorders, or from a transdiagnostic perspective. The transdiagnostic model for eating disorders assumes that all disorders share the same maintaining mechanisms (Fairburn et al., 2003), rather than each having a unique underlying

factor. The core overarching mechanism of all eating disorders is evaluating one's self-worth predominantly in terms of control over weight, shape and eating (Cooper & Fairburn, 2011; Fairburn et al., 2003). The diagnostic approach states that the differences among the various eating disorders are that AN is characterized by a severely restrictive food pattern, malnourishment, being underweight and sometimes the occurrence of binges followed by compensatory behavior such as laxative misuse and purging behavior. In contrast, BN is characterized by a normal weight, binges followed by compensatory behavior and a restrictive, irregular food pattern. Lastly, BED is characterized by an irregular food pattern, recurrent episodes of binge eating accompanied by a sense of lack of control, marked distress and absence of compensatory behavior. The binges are followed by feelings of shame, guilt and disgust (APA, 2013; Hay, 2020). Finally, OSFED refers to eating disorders which do not meet the criteria for AN, BN or BED, such as atypical AN, atypical BN, atypical BED, purging disorder or night eating syndrome. Life-time prevalence of AN is estimated at 0.16% and 0.63% for BN (Qian et al., 2021). BED and OSFED are the most common eating disorders with estimated life-time prevalence of 2% (Kessler et al., 2013) and 5% (Hay, 2020), respectively. However, these numbers might be outdated as recently, and related to the Covid-19 crisis, the estimated prevalence of BED has increased and patients seeking help display more severe symptoms (Termorshuizen et al., 2020).

Though BED is the most common eating disorder, BED remains underrecognized in comparison to AN and BN (Mitchison & Hay, 2014) leading to undertreatment of this disorder in clinical practice in the Netherlands. At the Novarum center for eating disorders we regularly admit patients who suffer from BED complaints for decades. Usually, they have been referred for weight loss treatments in the past. Consequently, the number of binges increase which leads to weight gain (Fairburn, 2008).

In conclusion, there is still a lack of sufficient data about clinical characteristics and the clinical course of BED. Moreover, due to its recent acknowledgement, data regarding BED examining the etiology, prevalence, correlates, and response to treatment are scarce. Therefore, this dissertation will focus on BED. It will focus on several knowledge gaps regarding assessment, prevalence, and correlates of BED. Another focus area will be treatment for BED: its efficacy, effectiveness, and cost-effectiveness. In short, three main subjects will be addressed in this dissertation: assessment, correlates and treatment.

Eating disorders occur globally

Though data regarding efficacy and cost-effectiveness of treatments for BED are scarce in the western world, etiology, prevalence, and correlates are unknown of several Arab countries such as Saudi Arabia. This may be because traditional Arab notions of beauty differ from the western ideal, with a curvy body being associated with fertility and wealth (Ford et al., 1990). However, since the rise of the thin ideal (Gordon, 2001) there are indications that also evaluation of self-worth predominantly in terms of control over weight and shape is increasingly prevalent in Saudi Arabia. After observation of the large diet (Imarcgroup, 2023) and cosmetic surgery (Bell, 2019) industries and from personal communications, it could cautiously be concluded that overevaluation of shape and weight is present in modernized Saudi Arabia. In addition, recent studies show that eating disorders occur globally (Keel & Klump, 2003; Thomas et al., 2016). However, research regarding eating disorders including BED is still in its infancy in Saudi Arabia. In addition, the lack of popular knowledge about eating disorders leads to stigmatization, and delayed help seeking (Griffiths et al., 2015). Since it is culturally unacceptable to discuss personal matters with non-family members, only 0.3% of Saudis with an eating disorder seek help, compared to 20% of eating disorder patients in western countries (Latzer et al., 2009). Saudis ae more prone to seek help when they

experience physical complaints, such as diabetes mellitus or kidney failure. Furthermore, eating disorders are not perceived as diseases of primary care and therefore barely recognized in Saudi Arabia (Alkhadari et al., 2016). Saudi Arabia faces an absence of assessment tools and specialized treatment centers (Darcy et al., 2012; Qadan, 2009). Moreover, Saudi Arabia is a widespread country, covering 2,150,000 km², including large desert plains. Therefore, even when specialist facilities would be available, Saudis would potentially deal with geographical barriers for treatment (Abrahamsson et al., 2018). Research regarding eating disorders in Saudi Arabia is further complicated by Saudi being a relatively culturally reclusive society, hard to access, and dealing with several taboos regarding mental health problems. Moreover, Saudi citizens are very sensitive to how they are viewed by others, and therefore, less likely to participate in surveys, questioning them on their personal beliefs and values (Al-Darmaki, 2003). It is challenging to build stepping stones for the investigation of eating disorder pathology in Saudi Arabia. It is deemed necessary to increase awareness on eating disorders in the Saudi community (Thomas et al., 2014).

Correlates of eating disorder pathology

The Saudi culture is currently undergoing rapid changes. Since the oil boom of the 1970's Saudi Arabia is dealing with an increased affluence resulting in rapid sociocultural and socioeconomic changes (Pike et al., 2014; Thomas et al., 2018). Furthermore, research shows that countries dealing with rapid sociocultural changes are impacted by an increase in eating disorder prevalence (Gordon, 2001; Pavlova et al., 2010). Various explanations have been put forth. Sociocultural changes come along with a sedentary lifestyle (WHO, 2006), and a dietary shift towards western types of foods, higher in salt, sugar and fat, all instrumental in the rise of excess weight (Hawks et al., 2004; Madanat et al., 2007). Currently, Saudi Arabia has one of the highest prevalence rates of excess weight (WHO, 2006), which is known to be

associated with eating disorder pathology, especially with recurrent binge eating and BED (Unikel Santoncini et al., 2018; Van der Horst et al., 2019; Van Son et al., 2013). Moreover, excess weight appears to be associated with body-shape dissatisfaction, and Saudis who are dissatisfied with their bodies have an increased risk for unhealthy dietary habits (Stice & Shaw, 2002), which may increase the risk of developing eating disorder pathology (Cooper et al., 1987). Other consequences of the rapid sociocultural changes in Saudi Arabia are associated with eating disorder pathology as well. The arrival of western companies, western expatriates (Thomas et al., 2018; Zeeni et al., 2013), increased exposure to western cultures (Pike & Dunne, 2015), and western social media (Madanat et al., 2007; Musaiger & Al-Mannai, 2013, 2014) led to a rise in the popularity of the "thin ideal" (Ford et al., 1990; Thomas et al., 2014). Consequently, levels of dieting, body-shape dissatisfaction and eating disorder pathology increased (Gow et al., 2020; Haines & Neumark-Sztainer, 2006; Pike & Dunne, 2015). Furthermore, rapid sociocultural changes might be accompanied by low selfesteem and increased stress. Both appear associated with eating disorder pathology (Gow et al., 2020; van der Valk et al., 2018; Yoneda et al., 2021). In addition, excess weight strengthens the association between eating disorder pathology and its correlates (Friedman et al., 2002; Madanat et al., 2007; Musaiger & Al-Mannai, 2013). However, since Saudi Arabia deals with a high prevalence of excess weight it is unclear whether media use, body-shape dissatisfaction, low self-esteem, increased socio-economic status, westernization and perceived stress are still correlates of eating disorder pathology after controlling for the effect of body mass index (BMI, kg/m²) (Gow et al., 2020; Haines & Neumark-Sztainer, 2006; van der Valk et al., 2018; Yoneda et al., 2021). The prevalence of eating disorder pathology could potentially be predominantly associated with the rates of excess weight (Unikel Santoncini et al., 2018). Identification of correlates of eating disorder pathology, including BED and the covariating role of high BMI may facilitate the identification of individuals at risk for

developing an eating disorder and who may benefit from participation in prevention programs (Stice et al., 2010; Stice et al., 2007). Consequently, reliable and valid assessment tools adapted to the Saudi language and culture are needed, in order to detect Saudis at high risk of eating disorders and body-shape dissatisfaction (Latzer et al., 2009; Musaiger et al., 2013). Therefore, this dissertation will first focus on assessment of eating disorder pathology.

Assessment

Various interview and self-report measures are developed to screen for eating disorders and body-shape dissatisfaction. The Eating Disorder Examination (EDE), a semistructured interview is the most widely used assessment tool to assess eating disorders and eating disorder symptoms such as binge eating and shape concern (Cooper & Fairburn, 1993). The EDE is generally considered reliable and valid, has good internal consistency (Cooper et al., 1989), test-retest reliability (Rizvi et al., 2000), and discriminative and concurrent validity are supported (Rosen et al., 1990). However, as administration of the EDE is time consuming and the instrument needs to be administered by trained professionals, self-report measures can be a less time consuming and more practical alternative. The Body Shape Questionnaire (BSQ) (Cooper et al., 1987) is most often used to measure body-shape dissatisfaction (Rosen et al., 1996). The original British BSQ (Cooper et al., 1987) is translated and adapted to various western cultures, such as Sweden, Norway, Portugal, Spain, Italy and various Latin-American countries such as Brazil, Colombia and Mexico (Ayensa et al., 2002; Castrillón Moreno et al., 2007; da Silva et al., 2014; Kapstad, 2015; Pook, 2008; Rosen et al., 1996; Stefanile et al., 2009; Welch et al., 2012). In addition, the Eating Disorder Examination-Questionnaire (EDE-Q) (Fairburn & Beglin, 1994) is a self-report measure based on the EDE (Cooper & Fairburn, 1993). The EDE-Q and EDE show acceptable levels of agreement, and therefore, when necessary, the EDE-Q can be used instead of the EDE (Reas et al., 2006).

Therefore, the EDE-Q is the most widely used self-report assessment tool to screen for eating disorder pathology and to assess its severity (Fairburn & Beglin, 1994). The EDE-Q is currently available in the Netherlands, Finland, Spain, Australia, Malaysia, Japan, Mexico and Israel, and various other countries (Aardoom et al., 2012; Calugi et al., 2017; Isomaa et al., 2016; Peláez-Fernández et al., 2012; Unikel Santoncini et al., 2018; Zohar et al., 2017). Although the BSQ and EDE-Q are valid and reliable assessment tools with good psychometric properties, such as high internal consistency and test-retest reliability, factorial and convergent validity (da Silva et al., 2014; Isomaa et al., 2016; Kapstad, 2015; Luce et al., 2008; Pook, 2008; Rosen et al., 1990; Welch et al., 2012), the BSQ and EDE-Q are currently not available in an Arabic version adapted for use in Saudi Arabia. Absence of assessment tools to measure body-shape dissatisfaction and eating disorder pathology may lead to overlooking or underdiagnosing BED and other eating disorders, which hampers timely proper treatment (Alkhadari et al., 2016). In addition, the ability to screen for body-shape dissatisfaction is an important first step to help prevent eating disorders through intervention programs, since targeted programs are more effective than universal preventative programs (Stice et al., 2019). Due to cultural differences, norms for Western and Arabic populations may differ (Welch et al., 2011). Therefore, the present thesis aims to adapt the BSQ and EDE-Q to the Saudi language and culture and evaluate its psychometric properties. In addition, valid assessment tools could also be used to measure reduction of eating disorder symptoms after eating disorder treatment, and therefore to assess efficacy and effectiveness of the various eating disorder treatments across the globe. This dissertation will also focus on efficacy and effectiveness of eating disorder treatment.

Treatment

Data regarding efficacy, effectiveness and cost-effectiveness of treatments for BED are scarce in the western world. Currently there is a range of treatments available for BED and other eating disorders, such as Dialectical Behavioral Therapy, Interpersonal Therapy and Cognitive Behavioral Therapy (ANZAED, 2014; Fairburn, 2008; NICE, 2017). Treatments for eating disorders based on cognitive behavioral principles display superior treatment outcomes (Agras et al., 2000). However, in 2003 Fairburn suggested that eating disorder pathology is maintained by a set of common mechanisms and that treatments addressing these mechanisms should be effective for various eating disorder presentations (Fairburn et al., 2003). Therefore, Fairburn modified the most leading evidence-based treatment for BN (CBT-BN) (Cooper & Fairburn, 1993) to make it suitable for treatment of all eating disorders. This resulted in Enhanced Cognitive Behavioral Therapy (CBT-E) (Fairburn, 2008), of which efficacy is proven in several randomized controlled trials. CBT-E appeared effective among transdiagnostic samples and for adult patients suffering from BN and atypical AN (Fairburn et al., 2003). In addition, CBT-E appeared also affective among adolescents with BED. Therefore, the Dutch national guideline recommends "Cognitive Behavioral Therapy-Enhanced" (CBT-E) (Zorgstandaard, 2021). However, data are scarce on how adult patients diagnosed with BED respond to CBT-E when offered individually (Berg et al., 2020; Fairburn et al., 2015; Fairburn et al., 2009; Wade et al., 2017). In addition, efficacy of CBT-E is mostly examined in randomized controlled trials, conducted under ideal circumstances, while effectiveness studies involve a naturalistic design and measure the degree of beneficial effect in real world, clinical settings (Andrews, 1999; Nordon et al., 2016). Therefore, as efficacy of CBT-E is well documented, its effectiveness among patients suffering from BED should be investigated. Furthermore, comparing the effectiveness of CBT-E between various eating disorders could potentially confirm its transdiagnostic nature. Moreover, determination of factors predicting CBT-E treatment outcome would provide prognostic information about for

whom CBT-E is likely to be beneficial and for whom less so. Treatment outcomes can potentially be maximized by understanding its predictors as more targeted treatments can be offered and potentially add-ons can be developed, thus improving clinical decision making (Kraemer, 2013).

Towards more efficiency: Guided self-help treatment

CBT-E is effective, but it is a time consuming treatment, requesting intense therapist involvement. In order to offer more efficient treatments, several new technologies are implemented, such as online guided self-help interventions or eMental Health (Donker & Kleiboer, 2018; Eysenbach, 2001). Therefore, the availability of eMental Health interventions has grown extensively over the past years. During eMental Health interventions patients will complete home assignments and upload them in an interactive treatment platform. eMental Health interventions can be unguided or therapist guided. During guided interventions patients and therapist interact through telephone calls, videoconferencing or chat. In addition, guided eMental Health interventions are more effective than unguided eMental Health interventions (Andersson & Cuijpers, 2009; Palmqvist et al., 2007; Spek et al., 2007). Furthermore, eMental Health interventions are effective for alcohol abuse (Cunningham et al., 2009; Eysenbach et al., 2010), depression (Andersson & Cuijpers, 2009; Andrews et al., 2010; Spek et al., 2007) and anxiety disorders (Andrews et al., 2010; Palmqvist et al., 2007; Spek et al., 2007). However, it is yet unknown whether CBT-E is effective and indeed efficient when offered by an online treatment modality. In addition, many authors mention the present gap between treatment needs and treatment provision (Abrahamsson et al., 2018; Becker et al., 2010; Fairburn & Patel, 2017; Shafran et al., 2009). There is a lack of specialized therapists and consequently, eating disorder patients have to deal with long waiting time to commence treatment (Fairburn & Peveler, 1990; Shafran et al., 2009). For instance, over all treatment

facilities in Amsterdam, the average wait time for an intial intake session is 11 weeks, and a further 20 weeks for treatment (NZA Nederlandse zorg autoriteit). Thus, there is a need to increase access for patients to specialized eating disorder treatment (Abrahamsson et al., 2018). The shortage in treatment availability can be covered by offering treatment as eMental Health. eMental Health offered as a guided self-help intervention may reduce the amount of intensive therapy, the amount of specialists time invested in a single treatment and, therefore, potentially reduce waiting time duration (Abrahamsson et al., 2018; Fairburn & Peveler, 1990; Shafran et al., 2009). Furthermore, guided self-help increases access to specialized treatment (Abrahamsson et al., 2018), which may improve quality of life (Fairburn & Peveler, 1990; Shafran et al., 2009). Guided self-help has also several additional advantages for the patient, such as the removal of geographical barriers, the possibility to hold sessions within the patient's own environment, reduced travel costs and travel time, and to communicate with their therapist wherever they are (Abrahamsson et al., 2018; Becker et al., 2010; Evans et al., 2011). Furthermore, according to a stepped care principle (Bower & Gilbody, 2005), guided self-help should be the first treatment of choice for patients diagnosed with BED, as preferably, the least resource-intensive treatment is delivered first (Mitchell et al., 2011). As international guidelines recommend guided self-help based on cognitive behavioral principles for BED (ANZAED, 2014; NICE, 2017), offering a guided self-help version of CBT-E including 12 weekly brief telephone sessions first might be an efficient treatment modality for BED.

Responding to the absence of a CBT-E based guided self-help treatment in Dutch,
Novarum, Center for Eating Disorders in the Netherlands, developed a guided self-help
version of CBT-E, based on the self-help section (Part two) of *Overcoming Binge Eating, The Proven Program to Learn Why You Binge and How You Can Stop* (Fairburn, 2013; Fairburn,
2016). As a first step of evaluating this treatment modality for BED, its efficacy should be

examined in a randomized controlled trial by comparing outcomes of an experimental treatment group to a waiting list control group. Thus, efficacy will be examined in comparison to a delayed treatment control condition. It is hypothesized that guided self-help CBT-E is superior to the control condition with regard to a decrease in binges, full recovery and diminished clinical impairment. Guided self-help CBT-E is also expected to be superior to the waiting list with regard to an increase in self-reported quality of life.

When the efficacy of guided self-help CBT-E is demonstrated, it could potentially mitigate the scarcity of specialized therapists, since guided self-help CBT-E requires less therapist involvement. In addition, if proven efficacious guided self-help CBT-E could be a cost-effective alternative compared to treatment as usual (ANZAED, 2014; NICE, 2017). Guided self-help CBT-E is briefer than in-person CBT-E and therefore potentially associated with lower costs of offering treatment. For example, costs of offering guided self-help based on regular CBT vary between €630- €900, whereas the costs for in person CBT-E are around €3,500 (Berg et al., 2020; Jenkins, 2021; König et al., 2018). In addition, guided self-help CBT-E potentially decreases societal burdens (Weissman, 2017), since BED is associated with non communicable diseases such as hypertension and diabetes mellitus (Eapen et al., 2006) and decreased work place productivity (Jenkins, 2021).

Conclusion

While studies regarding eating disorders are in their infancy in Saudi Arabia, knowledge in the Netherlands is more advanced. The Netherlands has several validated assessment tools available, such as the EDE (Cooper & Fairburn, 1993) and the EDE-Q (Fairburn & Beglin, 1994) as well as several kind of evidence based psychological treatments for eating disorders. Example of such an effective treatment is CBT-E (Berg et al., 2020; Fairburn et al., 2015).

However, effectiveness of CBT-E for BED is currently unknown, as well as efficacy and cost-effectiveness of the recently developed guided self-help version of CBT-E.

Saudi Arabia currently deals with an absence of culturally validated assessment tools and specialized treatment centers. It is of importance to increase awareness regarding eating disorders since Saudi Arabia currently deals with rapid sociocultural changes, which is associated with eating disorder pathology (Pike et al., 2014; Thomas et al., 2018). Cautiously, it can be concluded that the current state of knowledge concerning eating disorders in Saudi Arabia appears comparable to those of eating disorders in the Netherlands around the 1980's (Alkhadari et al., 2016; Qadan, 2009).

Aims

In this dissertation several knowledge gaps of aspects of eating disorders, mainly BED in the Netherlands and Saudi Arabia will be investigated. However, due to the major differences regarding the current state of knowledge, studies in the Netherlands will focus on different aspects of BED than in Saudi Arabia. Each chapter in this thesis focuses on different aspects of eating disorders which include screening, identification of correlates and therefore individuals at potential risk, treatment effectiveness, treatment efficacy, and cost-effectiveness.

Aim of the studies in Saudi Arabia is two-fold: (1) to adapt and examine psychometric properties of the two most used assessment tools to examine body dissatisfaction and eating disorder pathology: the BSQ and EDE-Q; (2) to investigate potential correlates of eating disorder pathology.

Main purpose of the studies in the Netherlands is to examine the effectiveness and efficacy of two treatment modalities of CBT-E for BED. The first treatment modality will be

in-person CBT-E for BED. As several trials showed that the efficacy of CBT-E is well documented among BN and transdiagnostic samples, its effectiveness in clinical practice using ROM data will be evaluated for patients with BED. Through a naturalistic design, outcome predictors of CBT-E will also be examined. The second treatment modality for BED will be eMental Health, offered as an online guided self-help version of CBT-E. Therefore, next efficacy of guided self-help CBT-E for BED will be examined through a randomized controlled trial (RCT) and an economic evaluation alongside the RCT will be undertaken.

Outline of this thesis

Chapter 2 reports on the evaluation of an adapted version of the EDE-Q in order to make it suitable for use among Saudis.

Chapter 3 describes the development of a Saudi version of the BSQ.

Chapter 4 presents a comprehensive synthesis regarding rapid sociocultural changes in Saudi Arabia and its consequences, including correlates and risk factors for BED. It will offer guidance for further studies investigating correlates of eating disorders.

Chapter 5 investigates the associations between expressions of sociocultural changes and eating disorder pathology in Saudi Arabia.

Chapter 6 will examine the effectiveness of CBT-E in everyday clinical practice. It investigates and compares effectiveness of CBT-E among patients with BED to patients diagnosed with BN and OSFED. Furthermore, predictors of treatment outcome are evaluated.

Chapter 7 will present the study protocol of the study described in Chapter 8.

Chapter 8 reports on the efficacy of online guided self-help CBT-E in a randomized controlled trial.

Chapter 9 will focus on cost-effectiveness and cost-utility of guided self-help CBT-E utilizing the outcome data of the randomized controlled trial.

Finally, **chapter 10** contains a summary and general discussion of the main findings.

Furthermore, the strengths and limitations will be discussed, and directions for future studies will be considered. Last, implications of the findings of the studies for clinical practice will be presented.

References

- Aardoom, J. J., Dingemans, A. E., Slof Op't Landt, M. C., & Van Furth, E. F. (2012). Norms and discriminative validity of the Eating Disorder Examination Questionnaire (EDE-Q). *Eating behaviors*, *13*(4), 305-309. https://doi.org/10.1016/j.eatbeh.2012.09.002
- Abrahamsson, N., Ahlund, L., Ahrin, E., & Alfonsson, S. (2018). Video-based CBT-E improves eating patterns in obese patients with eating disorder: A single case multiple baseline study. *J Behav Ther Exp Psychiatry*, *61*, 104-112. https://doi.org/10.1016/j.jbtep.2018.06.010
- Agh, T., Kovacs, G., Pawaskar, M., Supina, D., Inotai, A., & Voko, Z. (2015). Epidemiology, health-related quality of life and economic burden of binge eating disorder: a systematic literature review. *Eat Weight Disord*, 20(1), 1-12. https://doi.org/10.1007/s40519-014-0173-9
- Agras, W. S., Crow, S. J., Halmi, K. A., Mitchell, J. E., Wilson, G. T., & Kraemer, H. C. (2000). Outcome predictors for the cognitive behavior treatment of bulimia nervosa: data from a multisite study. *The American journal of psychiatry*, 157(8), 1302-1308.
- Al-Darmaki, F. R. (2003). Attitudes towards seeking professional psychological help: what really counts for United Arab Emirates university students *Social Behavior and Personality: an international journal*, *31*(5), 497-508. https://doi.org/10.2224/sbp.2003.31.5.497
- Alkhadari, S., Alsabbrri, A. O., Mohammad, I. H. A., Atwan, A. A., Alqudaihi, F., & Zahid, M. A. (2016). Prevalence of psychiatric morbidity in the primary health clinic attendees in Kuwait. *Journal of Affective Disorders*, 195, 15-20. https://doi.org/https://doi.org/10.1016/j.jad.2016.01.037
- Andersson, G., & Cuijpers, P. (2009). Internet-Based and Other Computerized Psychological Treatments for Adult Depression: A Meta-Analysis. *Cognitive Behaviour Therapy*, 38(4), 196-205. https://doi.org/10.1080/16506070903318960
- Andrews. (1999). Efficacy, effectiveness and efficiency in mental health service delivery. *Australian and New Zealand Journal of Psychiatry*, *33*(3), 316-322. https://doi.org/10.1046/j.1440-1614.1999.00581.x
- Andrews, G., Cuijpers, P., Craske, M. G., McEvoy, P., & Titov, N. (2010). Computer therapy for the anxiety and depressive disorders is effective, acceptable and practical health care: a meta-analysis. *PloS one*, *5*(10), e13196. https://doi.org/10.1371/journal.pone.0013196
- ANZAED. (2014). The Australian Psychiatric Workforce: RANZCP Workforce Survey Report (Australian Membership). In: RANZCP Melbourne, VIC
- APA. (2013). American Psychiatric Association. Diagnostic and Statistical manual of Mental Disorders fifth edition: DSM- 5. American Psychiatric Publishing.
- Ayensa, J. I. B., Grima, F. G., & Landívar, E. G. (2002). Insatisfacción corporal en adolescentes medida con el Body Shape Questionnaire (BSQ): efecto del anonimato, el sexo y la edad. *International Journal of Clinical and Health Psychology*, 2(3), 439-450. https://doi.org/https://www.redalyc.org/articulo.oa?id=33720304
- Becker, A. E., Hadley Arrindell, A., Perloe, A., Fay, K., & Striegel-Moore, R. H. (2010). A qualitative study of perceived social barriers to care for eating disorders: perspectives from ethnically diverse health care consumers. *Int J Eat Disord*, *43*(7), 633-647. https://doi.org/10.1002/eat.20755
- Bell, J. (2019, 26-01-2019). Nip and tuck: Why more Saudis are opting for cosmetic surgery than ever before. *Arab News*. https://www.arabnews.com/node/1442116/saudi-arabia
- Berg, E., Schlochtermeier, D., Koenders, J., Mooij, L., Goudriaan, A., Blankers, M., . . . Dekker, J. (2020). Implementing cognitive behavioral therapy-enhanced in a routine inpatient and outpatient setting: Comparing effectiveness and treatment costs in two

- consecutive cohorts. *International Journal of eating disorders*, *53*(3), 461-471. https://doi.org/10.1002/eat.23229
- Bower, P., & Gilbody, S. (2005). Stepped care in psychological therapies: access, effectiveness and efficiency: narrative literature review. *The British Journal of Psychiatry*, 186(1), 11-17.
- Calugi, S., Milanese, C., Sartirana, M., El Ghoch, M., Sartori, F., Geccherle, E., . . . Dalle Grave, R. (2017). The Eating Disorder Examination Questionnaire: reliability and validity of the Italian version. *Eating and weight disorders : EWD*, 22(3), 509-514. https://doi.org/10.1007/s40519-016-0276-6
- Castrillón Moreno, D., Luna Montaño, I., Avendaño Prieto, G., & Pérez-Acosta, A. M. (2007). Validación del Body Shape Questionnaire (Cuestionario de la Figura Corporal) BSQ para la población colombiana. *Acta colombiana de psicología*, 10(1), 15-23.
- Cooper, P. J., & Fairburn, C. G. (1993). Confusion over the core psychopathology of bulimia nervosa. *Int J Eat Disord*, *13*(4), 385-389. https://doi.org/https://doi.org/10.1002/1098-108X(199305)13:4%3C385::AID-EAT2260130406%3E3.0.CO;2-W
- Cooper, P. J., Taylor, M. J., Cooper, Z., & Fairbum, C. G. J. I. J. o. e. d. (1987). The development and validation of the Body Shape Questionnaire. In (Vol. 6, pp. 485-494).https://doi.org/10.1002/1098-108X(198707)6:4<485::AID-EAT2260060405>3.0.CO;2-O.
- Cooper, Z., Cooper, P. J., & Fairburn, C. G. (1989). The validity of the eating disorder examination and its subscales. *Br J Psychiatry*, *154*(6), 807-812. https://doi.org/10.1192/bjp.154.6.807
- Cooper, Z., & Fairburn, C. G. (2011). The Evolution of "Enhanced" Cognitive Behavior Therapy for Eating Disorders: Learning From Treatment Nonresponse. *Cognitive and Behavioral Practice*, *18*(3), 394-402. https://doi.org/10.1016/j.cbpra.2010.07.007
- Cunningham, J. A., Wild, T. C., Cordingley, J., Van Mierlo, T., & Humphreys, K. (2009). A randomized controlled trial of an internet-based intervention for alcohol abusers. *Addiction*, 104(12), 2023-2032. https://doi.org/10.1111/j.1360-0443.2009.02726.x
- da Silva, W. R., Dias, J. C. R., Maroco, J. o., & Campos, J. A. D. B. (2014). Confirmatory factor analysis of different versions of the Body Shape Questionnaire applied to Brazilian university students. *Body Image*, 11(4), 384-390. https://doi.org/10.1016/j.bodyim.2014.06.001
- Darcy, A. M., Doyle, A. C., Lock, J., Peebles, R., Doyle, P., & Le Grange, D. (2012). The eating disorders examination in adolescent males with anorexia nervosa: How does it compare to adolescent females? *International Journal of eating disorders*, 45(1), 110-114. https://doi.org/10.1002/eat.20896
- Donker, T., & Kleiboer, A. (2018). e-health innovations for global mental health. *Global mental health*, *5*, e5.
- Eapen, V. M., Mabrouk, A. A., & bin Othman, S. (2006). Disordered Eating Attitudes and symptomatology among adolescent girls in the United Arab Emirates. *Eating behaviors*, 7(1), 53-60. https://doi.org/https://doi.org/10.1016/j.eatbeh.2005.07.001
- Evans, E. J., Hay, P. J., Mond, J., Paxton, S. J., Quirk, F., Rodgers, B., . . . Sawoniewska, M. A. (2011). Barriers to help-seeking in young women with eating disorders: a qualitative exploration in a longitudinal community survey. *Eat Disord*, *19*(3), 270-285. https://doi.org/10.1080/10640266.2011.566152
- Eysenbach, G. (2001). What is e-health? *Journal of medical Internet research*, 3(2), e833.
- Eysenbach, G., van Mierlo, T., Spijkerman, R., Postel, M. G., de Haan, H. A., ter Huurne, E. D., . . . de Jong, C. A. J. (2010). Effectiveness of a Web-based Intervention for

- Problem Drinkers and Reasons for Dropout: Randomized Controlled Trial. *Journal of medical Internet research*, 12(4). https://doi.org/10.2196/jmir.1642
- Fairburn, C. (2008). Cognitive behavior therapy and eating disorders. Guilford Press.
- Fairburn, C., & Beglin, S. J. (1994). Assessment of eating disorders: interview or self-report questionnaire? *Int J Eat Disord*, *16*(4), 363-370. https://doi.org/https://doi.org/10.1002/1098-108X(199412)16:4<363::AID-EAT2260160405>3.0.CO;2-%23
- Fairburn, C. G. (2013). Overcoming binge eating. Guilford Press.
- Fairburn, C. G. (2016). *Overwin je eetbuien, waarom je te veel eet en hoe je daarmee kunt stoppen* (L. Berkhuizen, Trans.). Uitgeverij Nieuwezijds.
- Fairburn, C. G., Bailey-Straebler, S., Basden, S., Doll, H. A., Jones, R., Murphy, R., . . . Cooper, Z. (2015). A transdiagnostic comparison of enhanced cognitive behaviour therapy (CBT-E) and interpersonal psychotherapy in the treatment of eating disorders. *Behaviour research and therapy*, 70, 64-71. https://doi.org/10.1016/j.brat.2015.04.010
- Fairburn, C. G., Cooper, Z., Doll, H. A., O'Connor, M. E., Bohn, K., Hawker, D. M., . . . Palmer, R. L. (2009). Transdiagnostic cognitive-behavioral therapy for patients with eating disorders: a two-site trial with 60-week follow-up. *American Journal of Psychiatry*, *166*(3), 311-319. https://doi.org/10.1176/appi.ajp.2008.08040608
- Fairburn, C. G., Cooper, Z., & Shafran, R. (2003). Cognitive behaviour therapy for eating disorders: A "transdiagnostic" theory and treatment. *Behaviour research and therapy*, 41(5), 509-528. https://doi.org/10.1016/s0005-7967(02)00088-8
- Fairburn, C. G., & Patel, V. (2017). The impact of digital technology on psychological treatments and their dissemination. *Behav Res Ther*, 88, 19-25. https://doi.org/10.1016/j.brat.2016.08.012
- Fairburn, C. G., & Peveler, R. C. (1990). Bulimia nervosa and a stepped care approach to management. *Gut*, *31*(11), 1220-1222.
- Ford, K. A., Dolan, B. M., & Evans, C. (1990). Cultural factors in the eating disorders: a study of body shape preferences of Arab students. *J Psychosom Res*, *34*(5), 501-507. https://doi.org/10.1016/0022-3999(90)90024-x
- Friedman, K. E., Reichmann, S. K., Costanzo, P. R., & Musante, G. J. (2002). Body Image Partially Mediates the Relationship between Obesity and Psychological Distress. *Obesity Research*, 10(1), 33-41. https://doi.org/10.1038/oby.2002.5
- Gordon, R. A. (2001). Eating disorders East and West: A culture-bound syndrome unbound. *International Journal of Eating disorders*, *1*, 16.
- Gow, M. L., Tee, M. S. Y., Garnett, S. P., Baur, L. A., Aldwell, K., Thomas, S., . . . Jebeile, H. (2020). Pediatric obesity treatment, self-esteem, and body image: A systematic review with meta-analysis. *Pediatric obesity*, *15*(3), e12600. https://doi.org/10.1111/ijpo.12600
- Haines, J., & Neumark-Sztainer, D. (2006). Prevention of obesity and eating disorders: a consideration of shared risk factors. *Health education research*, 21(6), 770-782.
- Hay, P. (2020). Current approach to eating disorders: a clinical update. *Internal medicine journal*, *50*(1), 24-29. https://doi.org/10.1111/imj.14691
- Hay, P., Girosi, F., & Mond, J. (2015). Prevalence and sociodemographic correlates of DSM-5 eating disorders in the Australian population. *Journal of Eating disorders*, *3*(1), 19. https://doi.org/10.1186/s40337-015-0056-0
- Imarcgroup. (2023). *UAE Weight Loss Market: Industry Trends, Share, Size, Growth, Opportunity and Forecast 2022-2027*. UAE Weight Loss Market Report. Retrieved 18-1 from https://www.imarcgroup.com/uae-weight-loss-market
- Isomaa, R., Lukkarila, I. L., Ollila, T., Nenonen, H., Charpentier, P., Sinikallio, S., & Karhunen, L. (2016). Development and preliminary validation of a Finnish version of

- the Eating Disorder Examination Questionnaire (EDE-Q). *Nordic journal of psychiatry*,, 70(7), 542-546.
- https://doi.org/https://doi.org/10.1080/08039488.2016.1179340
- Jenkins, P. E. (2021). Clinical and costeffectiveness of two ways of delivering guided selfhelp for people with an eating disorder: A multiarm randomized controlled trial. *International Journal of eating disorders*, 54(7), 1224.
- Kapstad, H. (2015). Validation of the Norwegian short version of the Body Shape Questionnaire (BSQ-14). *Nordic journal of psychiatry*, *69*(7), 509. https://doi.org/https://doi.org/10.3109/08039488.2015.1009486
- Keel, P. K., & Klump, K. L. (2003). Are eating disorders culture- bound syndromes? Implications for conceptualizing their etiology. *Psychological Bulletin*, *129*(5), 747-769. https://doi.org/10.1037/0033-2909.129.5.747
- Kessler, R. C., Berglund, P. A., Chiu, W. T., Deitz, A. C., Hudson, J. I., Shahly, V., . . . Xavier, M. (2013). The Prevalence and Correlates of Binge Eating Disorder in the World Health Organization World Mental Health Surveys. *Biological Psychiatry*, 73(9), 904-914. https://doi.org/10.1016/j.biopsych.2012.11.020
- König, H.-H., Bleibler, F., Friederich, H.-C., Herpertz, S., Lam, T., Mayr, A., . . . Egger, N. (2018). Economic evaluation of cognitive behavioral therapy and Internet-based guided self-help for binge-eating disorder. *International Journal of eating disorders*, 51(2), 155-164. https://doi.org/10.1002/eat.22822
- Kraemer, H. C. (2013). Discovering, comparing, and combining moderators of treatment on outcome after randomized clinical trials: a parametric approach. *Statistics in medicine*, 32(11), 1964-1973. https://doi.org/10.1002/sim.5734
- Latzer, Y., Azaiza, F., & Tzischinsky, O. (2009). Eating attitudes and dieting behavior among religious subgroups of Israeli- Arab adolescent females. *Journal of Religion and Health*, 48(2), 189-199. https://doi.org/https://doi.org/10.1007/s10943-008-9189-7
- Luce, K. H., Crowther, J. H., & Pole, M. (2008). Eating Disorder Examination Questionnaire (EDE-Q): norms for undergraduate women. *Int J Eat Disord*, *41*(3), 273-276. https://doi.org/10.1002/eat.20504
- Madanat, H. N., Brown, R. B., & Hawks, S. R. (2007). The impact of body mass index and Western advertising and media on eating style, body image and nutrition transition among Jordanian women. *Public Health Nutrition*, *10*(10), 1039- 1046. https://doi.org/https://doi.org/10.1017/S1368980007666713
- Mitchell, J. E., Agras, S., Crow, S., Halmi, K., Fairburn, C. G., Bryson, S., & Kraemer, H. (2011). Stepped care and cognitive-behavioural therapy for bulimia nervosa: randomised trial. *British Journal of Psychiatry*, *198*(5), 391-397. https://doi.org/10.1192/bjp.bp.110.082172
- Mitchison, D., & Hay, P. J. (2014). The epidemiology of eating disorders: genetic, environmental, and societal factors. *Clinical epidemiology*, *6*, 89-97. https://doi.org/10.2147/CLEP.S40841
- Musaiger, A. O., & Al- Mannai, M. (2013). Role of obesity and media in body weight concern among female university students in Kuwait. *Eating behaviors*, *14*(1), 229-232. https://doi.org/https://doi.org/10.1016/j.appet.2012.10.012
- Musaiger, A. O., & Al- Mannai, M. (2014). Association between exposure to media and body weight concern among female university students in five Arab countries: A preliminary cross- cultural study. *Journal of Biosocial Sciences*, 46(2), 240-247. https://doi.org/https://doi.org/10.1017/S0021932013000278
- Musaiger, A. O., Al-Mannai, M., Tayyem, R., Al-Lalla, O., Ali, E. Y., Kalam, F., . . . Chirane, M. (2013). Risk of disordered eating attitudes among adolescents in seven Arab

- countries by gender and obesity: a cross-cultural study. *Appetite*, 60(1), 162-167. https://doi.org/https://doi.org/10.1016/j.appet.2012.10.012
- NICE. (2017). *Eating disorders: Recognition and treatment*. National Institute for Health Care Excellence
- Nordon, C., Karcher, H., Groenwold, R. H. H., Ankarfeldt, M. Z. I., Pichler, F., Chevrou-Severac, H., . . . D, C. M. F. (2016). The "Efficacy-Effectiveness Gap": Historical Background and Current Conceptualization. *Value in Health*, *19*(1), 75-81. https://doi.org/10.1016/j.jval.2015.09.2938
- Palmqvist, B. r., Carlbring, P., & Andersson, G. (2007). Internet-delivered treatments with or without therapist input: does the therapist factor have implications for efficacy and cost? *Expert Review of Pharmacoeconomics & Outcomes Research*, 7(3), 291-297. https://doi.org/10.1586/14737167.7.3.291
- Pavlova, B., Uher, R., Dragomirecka, E., & Papezova, H. (2010). Trends in hospital admissions for eating disorders in a country undergoing a socio-cultural transition, the Czech Republic 1981-2005. *Soc Psychiatry Psychiatr Epidemiol*, *45*(5), 541-550. https://doi.org/10.1007/s00127-009-0092-7
- Peláez-Fernández, M. A., Javier Labrador, F., & Raich, R. M. (2012). Validation of eating disorder examination questionnaire (EDE-Q)--Spanish version--for screening eating disorders. *The Spanish journal of psychology*, *15*(2), 817-824.
- Pike, K., & Dunne, P. (2015). The rise of eating disorders in Asia: a review. *Journal of Eating disorders*, 3(1), 1-14.
- Pike, K. M., Hoek, H. W., & Dunne, P. E. (2014). Cultural trends and eating disorders. *Curr Opin Psychiatry*, 27(6), 436-442. https://doi.org/10.1097/YCO.00000000000100
- Pook, M. (2008). Evaluation and comparison of different versions of the Body Shape Questionnaire. *Psychiatry Research*, *158*(1), 67. https://doi.org/10.1016/j.psychres.2006.08.002
- Qadan, L. (2009). Anorexia nervosa: beyond boundaries. *Int J Eat Disord*, 42(5), 479-481. https://doi.org/10.1002/eat.20641
- Qian, J., Wu, Y., Liu, F., Zhu, Y., Jin, H., Zhang, H., . . . Yu, D. (2021). An update on the prevalence of eating disorders in the general population: a systematic review and meta-analysis. *Eating and Weight Disorders-Studies on Anorexia*, *Bulimia and Obesity*, 1-14.
- Reas, D. L., Grilo, C. M., & Masheb, R. M. (2006). Reliability of the Eating Disorder Examination-Questionnaire in patients with binge eating disorder. *Behaviour research and therapy*, 44(1), 43-51. https://doi.org/10.1016/j.brat.2005.01.004
- Rizvi, S. L., Peterson, C. B., Crow, S. J., & Agras, W. S. (2000). Test-retest reliability of the eating disorder examination. *Int J Eat Disord*, 28(3), 311-316. https://doi.org/10.1002/1098-108x(200011)28:3<311::aid-eat8>3.0.co;2-k
- Rojo-Moreno, L., Arribas, P., Plumed, J., Gimeno, N., García-Blanco, A., Vaz-Leal, F., . . . Livianos, L. (2015). Prevalence and comorbidity of eating disorders among a community sample of adolescents: 2-year follow-up. *Psychiatry Research*, 227(1), 52-57.
 - https://www.sciencedirect.com/science/article/abs/pii/S0165178115001079?via%3Dih ub
- Rosen, J. C., Jones, A., Ramirez, E., & Waxman, S. (1996). Body Shape Questionnaire: studies of validity and reliability. *Int J Eat Disord*, 20(3), 315-319. https://doi.org/10.1002/(SICI)1098-108X(199611)20:3<315::AID-EAT11>3.0.CO;2-Z
- Rosen, J. C., Vara, L., Wendt, S., & Leitenberg, H. J. I. J. o. E. D. (1990). Validity studies of the eating disorder examination. *9*(5), 519-528.

- https://doi.org/https://doi.org/10.1002/1098-108X(199009)9:5%3C519::AID-EAT2260090507%3E3.0.CO;2-K
- Shafran, R., Clark, D. M., Fairburn, C. G., Arntz, A., Barlow, D. H., Ehlers, A., . . . Wilson, G. T. (2009). Mind the gap: Improving the dissemination of CBT. *Behav Res Ther*, 47(11), 902-909. https://doi.org/10.1016/j.brat.2009.07.003
- Smink, F. R., van Hoeken, D., & Hoek, H. W. (2012). Epidemiology of eating disorders: incidence, prevalence and mortality rates. *Curr Psychiatry Rep*, *14*(4), 406-414. https://doi.org/10.1007/s11920-012-0282-y
- Smink, F. R., van Hoeken, D., & Hoek, H. W. (2013). Epidemiology, course, and outcome of eating disorders. *Curr Opin Psychiatry*, 26(6), 543-548. https://doi.org/10.1097/YCO.0b013e328365a24f
- Spek, V., Cuijpers, P., Nyklícek, I., Riper, H., Keyzer, J., & Pop, V. (2007). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: a meta-analysis. *Psychological medicine*, *37*(3), 319-328.
- Stefanile, C., Matera, C., & Pisani, E. (2009). Body shape questionnaire (BSQ-14): An italian version. *Journal of Eating disorders*, 6, 485-494.
- Stice, E., Johnson, S., & Turgon, R. (2019). Eating Disorder Prevention. *The Psychiatric clinics of North America*, 42(2), 309-318. https://doi.org/10.1016/j.psc.2019.01.012
- Stice, E., Ng, J., & Shaw, H. (2010). Risk factors and prodromal eating pathology. *Journal of Child Psychology and Psychiatry*, *51*(4), 518-525. https://doi.org/https://doi.org/10.1111/j.1469-7610.2010.02212.x
- Stice, E., Shaw, H., & Marti, C. N. (2007). A Meta-Analytic Review of Eating Disorder Prevention Programs: Encouraging Findings. *Annual Review of Clinical Psychology*, 3, 207-231. https://doi.org/10.1146/annurev.clinpsy.3.022806.091447
- Stice, E., & Shaw, H. E. (2002). Role of body dissatisfaction in the onset and maintenance of eating pathology: a synthesis of research findings. *Journal of psychosomatic research*, 53(5), 985-993. https://doi.org/10.1016/s0022-3999(02)00488-9
- Termorshuizen, J. D., Watson, H. J., Thornton, L. M., Borg, S., Flatt, R. E., MacDermod, C. M., . . . Bulik, C. M. (2020). Early impact of COVID-19 on individuals with self-reported eating disorders: A survey of ~1,000 individuals in the United States and the Netherlands. *The International journal of eating disorders*, *53*(11), 1780-1790. https://doi.org/10.1002/eat.23353
- Thomas, J., Al Marzooqi, F. H., Tahboub- Schulte, S., & Furber, S. W. (2014). Changing physical appearance preferences in the United Arab Emirates. *Mental Health, Religion & Culture*, *17*(6), 594-600. https://doi.org/https://doi.org/10.1080/13674676.2013.871242
- Thomas, J., O'Hara, L., Quadflieg, S., & Weissgerber, S. C. (2018). Acculturation, out-group positivity and eating disorders symptoms among Emirati women. *Eat Weight Disord*, 23(2), 241-246. https://doi.org/10.1007/s40519-016-0358-5
- Thomas, J., Quadflieg, S., & O'Hara, L. (2016). Implicit out-group preference is associated with eating disorders symptoms amongst Emirati females. *Eating behaviors*, 21, 48-53. https://doi.org/https://doi.org/10.1016/j.eatbeh.2015.12.005
- Unikel Santoncini, C., Bojorquez Chapela, I., Díaz de León Vázquez, C. n., Vázquez Velázquez, V. n., Rivera Márquez, J. A., Galván Sánchez, G., & Rocha Velis, I. (2018). Validation of eating disorders examination questionnaire in Mexican women. *International Journal of eating disorders*, 51(2), 146-154. https://doi.org/10.1002/eat.22819
- Van der Horst, H., Van Os, J., van der Horst, H. E., & van Elburg, A. A. (2019). De patiënt met een eetprobleem. In: Bohn Stafleu van Loghumhttp://hdl.handle.net/1874/388648

- van der Valk, E. S., Savas, M., & van Rossum, E. F. C. (2018). Stress and Obesity: Are There More Susceptible Individuals? *Current Obesity Reports*, 7(2), 193-203. https://doi.org/10.1007/s13679-018-0306-y
- Van Son, G. E., van der Meer, P. A. M., & Van Furth, E. F. (2013). Correlates and associations between weight suppression and binge eating symptomatology in a population-based sample. *Eating behaviors*, *14*(2), 102-106. https://doi.org/10.1016/j.eatbeh.2012.11.003
- Wade, S., Byrne, S., & Allen, K. (2017). Enhanced cognitive behavioral therapy for eating disorders adapted for a group setting. *International Journal of eating disorders*, 50(8), 863-872. https://doi.org/https://doi.org/10.1002/eat.22723
- Walsh, B. T., Attia, E., Glasofer, D. R., & Sysko, R. (2015). *Handbook of assessment and treatment of eating disorders*. American Psychiatric Pub. https://books.google.nl/books?hl=nl&lr=&id=EYYJDAAAQBAJ&oi=fnd&pg=PP1&dq=walsh+2016+eating+disorders&ots=D2sZ6qWqo4&sig=ASTcfJsFf89DbuW9GnGEuH-bVQw#v=onepage&q=walsh%202016%20eating%20disorders&f=false
- Watson, H. J., Allen, K., Fursland, A., Byrne, S. M., & Nathan, P. R. (2012). Does Enhanced Cognitive Behaviour Therapy for Eating Disorders Improve Quality of Life? *European Eating disorders Review*, 20(5), 393-399. https://doi.org/10.1002/erv.2186
- Weissman, R. S. (2017). Reducing the burden of suffering from eating disorders: Unmet treatment needs, cost of illness, and the quest for cost-effectiveness. *Behaviour research and therapy*, 88, 49.
- Welch, E., Birgegard, A., Parling, T., & Ghaderi, A. (2011). Eating disorder examination questionnaire and clinical impairment assessment questionnaire: general population and clinical norms for young adult women in Sweden. *Behav Res Ther*, 49(2), 85-91. https://doi.org/10.1016/j.brat.2010.10.010
- Welch, E., Lagerström, M., & Ghaderi, A. (2012). Body Shape Questionnaire: Psychometric properties of the short version (BSQ-8C) and norms from the general Swedish population. *Body Image*, *9*(4), 547-550. https://doi.org/10.1016/j.bodyim.2012.04.009
- WHO. (2006). Country cooperation strategy for WHO and the United Arab Emirates 2005-2009
- Yoneda, M., Kubota, M., Watanabe, H., & Egusa, G. (2021). Westernization of Lifestyle and Atherosclerosis in the Japanese: Lessons from the Hawaii Los Angeles Hiroshima Study. *Journal of atherosclerosis and thrombosis*, 28(3), 214-222. https://doi.org/10.5551/jat.RV17046
- Zeeni, N., Gharibeh, N., & Katsounari, I. (2013). The influence of sociocultural factors on the eating attitudes of Lebanese and Cypriot students: a cross-cultural study. *J Hum Nutr Diet*, 26 Suppl 1, 45-52. https://doi.org/10.1111/jhn.12059
- Zohar, A. H., Lev-Ari, L., & Bachner-Melman, R. (2017). The EDE-Q in Hebrew: Structural and Convergent/Divergent Validity in a Population Sample. *Isr J Psychiatry Relat Sci*, 54(3), 15-20. https://www.ncbi.nlm.nih.gov/pubmed/29735808