

## Insights from scanning tunneling microscopy experiments into correlated electron systems

Benschop, T.

## Citation

Benschop, T. (2023, September 26). *Insights from scanning tunneling microscopy experiments into correlated electron systems. Casimir PhD Series*. Retrieved from https://hdl.handle.net/1887/3642190

Version:	Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/3642190

Note: To cite this publication please use the final published version (if applicable).

## Propositions

## Accompanying the thesis

"Insights from Scanning Tunneling Microscopy Experiments into Correlated Electron Systems"

- 1. The breakdown of the superconducting state in overdoped cuprates is not a result of a diminishing pairing interaction. [chapter 2]
- 2. In terms of twist angle homogeneity, open twisted bilayer graphene devices can rival the quality of fully encapsulated devices. [chapter 3]
- 3. The lack of evidence for superconductivity in open twisted bilayer graphene devices suggests a critical role for the encapsulating hBN layers in establishing superconductivity. [chapter 3]
- 4. The noise enhancements measured on the surface of  $Sr_2IrO_4$  can be explained if substantial random telegraph noise is present in addition to the shot noise of tunneling electrons. [chapter 5]
- 5. The work by Kopp *et al.* claims that "ferromagnetic fluctuations in the superconducting state will act as pair breakers and [thereby] explain the demise of superconductivity in overdoped cuprates", however, this is in contradiction with our observations. [Kopp *et al.*, PNAS **104**, 6123 (2007)]
- 6. The point contact spectroscopy measurements in the work by Oh *et al.* are no longer local, and therefore are not definitive proof that the spectroscopic gap measured with conventional STS is a result of superconductivity. [Oh *et al.*, Nature **600**, 240 (2021)]
- 7. The electronic effects attributed to tip-induced strain in the work of Mesple *et al.* can also (partially) be explained by tip-induced band bending effects. [Mesple *et al.*, PRL **127**, 126405 (2021)]
- 8. Regarding the work by Yan *et al.*, because the lack of intermittent atomic resolution images between deposition cycles, it is hard to attribute the observed V-shaped spectra to intrinsic physics of  $Sr_2IrO_4$ . [Yan *et al.*, PRX **5**, 041018 (2015)]
- 9. The scientific community should be an example for the global society, where everybody, regardless of age, gender, ethnicity or political background, can participate.

Tjerk Benschop Leiden, 26 September 2023