



**Universiteit
Leiden**
The Netherlands

Letter to the editor: how large a study is needed to detect TKA revision rate reductions attributable to robotic or navigated technologies? A simulation-based power analysis

Koster, L.A.; Kaptein, B.L.; Pijls, B.G.; Nelissen, R.G.H.H.

Citation

Koster, L. A., Kaptein, B. L., Pijls, B. G., & Nelissen, R. G. H. H. (2022). Letter to the editor: how large a study is needed to detect TKA revision rate reductions attributable to robotic or navigated technologies? A simulation-based power analysis, *480*(4), 836-837.
doi:10.1097/CORR.0000000000002123

Version: Publisher's Version

License: [Creative Commons CC BY-NC-ND 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3281391>

Note: To cite this publication please use the final published version (if applicable).

Letter to the Editor: How Large a Study is Needed to Detect TKA Revision Rate Reductions Attributable to Robotic or Navigated Technologies? A Simulation-based Power Analysis

Lennard A. Koster MSc¹ , Bart L. Kaptein MSc, PhD¹, Bart G. Pijls MD, PhD¹, Rob G.H.H. Nelissen MD, PhD¹

To the Editor,

We read the study, “How Large a Study is Needed to Detect TKA Revision Rate Reductions Attributable to Robotic or Navigated Technologies? A Simulation-based Power Analysis” by Hickey et al. [1] with great interest. Their main conclusion was that it would take an impractically large number of patients (and a very long time) to answer the question of whether robotic or

navigated TKA approaches are advantageous, which suggests that any advantage of those technologies in terms of reducing revision risk is likely to be quite small, if it is present at all. The situation described by the authors is a common problem for most prosthetic joint replacement studies using revision as an endpoint. We see this in studies on new arthroplasty designs, fixation methods, and surgical techniques [3].

One of the major causes of revision arthroplasty is aseptic loosening, as pointed out by Hickey et al. [1]. It may take 10 years or more before loosening causes symptoms, becomes visible on conventional radiographs, or potentially leads to revision surgery. However, radiostereometric analysis (RSA) makes it possible to detect implant loosening early postoperatively (6 months to 12 months), which is why it's been used for more than 40 years [5, 7]. Indeed, RSA allows in vivo three-dimensional measurement of implant migration relative to the bone with an accuracy of 0.2 mm for translations and 0.5° for rotations [2]. Therefore, the sample size and follow-up time can be greatly reduced. Only a relatively small number of patients followed for a relatively short period of time (2 years) is required to

produce similar results as long-term studies (10 years or more) with thousands of patients [5, 8]. For instance, knowing what we know today, it would have been possible to detect Boneloc (Biomet Inc) bone cement as a potential disaster at 6-month follow-up using RSA [5].

To our knowledge, two randomized controlled trials (RCTs) [4, 9] on computer-assisted surgery (CAS) in TKA used RSA to measure tibial baseplate migration. van Strien et al. [9] and Petursson et al. [4] studied the effect on tibial baseplate migration 2 years after surgery randomizing 57 and 54 patients, respectively. They found no clinically relevant differences in tibial migration. The authors of both studies did not expect to find a difference between groups in the long term. A meta-analysis by Rhee et al. [6] using nine RCTs with long-term follow-up confirmed this assumption by finding no differences in functional outcomes and survivorship between CAS and conventional TKA [6]. The findings of these RSA-based studies [4, 9] are supportive of the main findings of Hickey et al. [1] in that they suggest there is likely to be little, if any, clinical benefit from robotic surgery in terms of the clinically important endpoint of tibial revision.

Work in an RSA setting can provide answers like the above without the substantial delay associated with large, long-term randomized trials, which

(RE: Hickey MD, Anglin C, Masri B, Hodgson AJ. How Large a Study is Needed to Detect TKA Revision Rate Reductions Attributable to Robotic or Navigated Technologies? A Simulation-based Power Analysis. *Clin Orthop Relat Res*. 2021;479: 2350–2361.)

The authors certify that there are no funding or commercial associations (consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted article related to the author or any immediate family members.

All ICMJE Conflict of Interest Forms for authors and *Clinical Orthopaedics and Related Research*® editors and board members are on file with the publication and can be viewed on request.

The opinions expressed are those of the writer, and do not reflect the opinion or policy of *CORR*® or The Association of Bone and Joint Surgeons®.

L. A. Koster , Department of Orthopaedic Surgery, Leiden University Medical Center, Albinusdreef 2, PO Box 9600, 2300 RC, Leiden, the Netherlands, Email: l.a.koster@lumc.nl

¹marked for authors Department of Orthopaedics, Leiden University Medical Center, Leiden, the Netherlands

Letter to the Editor

would be expensive and impractical to conduct. Based on this, we believe RSA can and should play an important role in the evaluation and phased introduction of new implants, fixation approaches, and surgical techniques.

References

1. Hickey MD, Anglin C, Masri B, Hodgson AJ. How large a study is needed to detect TKA revision rate reductions attributable to robotic or navigated technologies? A simulation-based power analysis. *Clin Orthop Relat Res.* 2021; 479:2350-2361.
2. Kaptein BL, Valstar ER, Stoel BC, Reiber HC, Nelissen RG. Clinical validation of model-based RSA for a total knee prosthesis. *Clin Orthop Relat Res.* 2007; 464:205-209.
3. Michelson JD, Riley LH Jr. Considerations in the comparison of cemented and cementless total hip prostheses. *J Arthroplasty.* 1989;4:327-334.
4. Petursson G, Fenstad AM, Gothesen O, et al. Similar migration in computer-assisted and conventional total knee arthroplasty. *Acta Orthop.* 2017;88: 166-172.
5. Pijls BG, Plevier JWM, Nelissen R. RSA migration of total knee replacements. *Acta Orthop.* 2018;89:1-9.
6. Rhee SJ, Kim HJ, Lee CR, Kim CW, Gwak HC, Kim JH. A comparison of long-term outcomes of computer-navigated and conventional total knee arthroplasty: a meta-analysis of randomized controlled trials. *J Bone Joint Surg Am.* 2019;101: 1875-1885.
7. Ryd L, Albrektsson BEJ, Carlsson L, et al. Rentgen stereophotogrammetric analysis as a predictor of mechanical loosening of knee prostheses. *J Bone Joint Surg Br.* 1995;77B:377-383.
8. Valstar ER, Gill R, Ryd L, Flivik G, Borlin N, Karrholm J. Guidelines for standardization of radiostereometry (RSA) of implants. *Acta Orthop.* 2005; 76:563-572.
9. van Strien T, van der Linden-van der Zwaag E, Kaptein B, van Erkel A, Valstar E, Nelissen R. Computer assisted versus conventional cemented total knee prostheses alignment accuracy and micro-motion of the tibial component. *Int Orthop.* 2009;33:1255-1261.