



Universiteit
Leiden

The Netherlands

Urbanization in Indonesia and its impact on non-communicable diseases: a clinical, epidemiological, and immunological study

Kurniawan, F.

Citation

Kurniawan, F. (2023, October 19). *Urbanization in Indonesia and its impact on non-communicable diseases: a clinical, epidemiological, and immunological study*. Retrieved from <https://hdl.handle.net/1887/3644030>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3644030>

Note: To cite this publication please use the final published version (if applicable).



Chapter 3

IMPACT OF RURAL-URBAN ENVIRONMENT ON METABOLIC PROFILE AND RESPONSE TO A 5-DAY HIGH-FAT DIET

Dicky L Tahapary^{1,2,3,4#*}, Karin de Ruiter^{2#}, Farid Kurniawan^{1,4#}, Yenny Djuardi^{3,5}, Yanan Wang⁶, Siti M.E. Nurdin⁷, Elisa Iskandar^{3,5}, Dominikus Minggu⁸, Em Yunir^{1,4}, Bruno Guigas², Taniawati Supali^{3,5}, Patrick C.N. Rensen⁶, Erliyani Sartono², Pradana Soewondo^{1,4\$}, Dante S Harbuwono^{1,4\$}, Johannes WA Smit^{4,9\$}, Maria Yazdanbakhsh^{2\$*}

#\$These authors contributed equally; *Corresponding author

(Scientific Reports 2018;8:8149. doi:10.1038/s41598-018-25092-6)

ABSTRACT

Epidemiological studies have indicated that rural living might be protective against type 2 diabetes development. We compared the metabolic profile and response to a short-term high-fat high-calorie diet (HFD) of men with the same genetic background living in an urban and rural area of Indonesia. First, we recruited 154 Floresian male subjects (18-65 years old), of whom 105 lived in a rural area (Flores) and 49 had migrated and lived in urban area (Jakarta) for more than 1 year. The urban group had significantly higher whole-body insulin resistance (IR), as assessed by homeostatic-model-assessment of IR (HOMA-IR), [mean difference (95%CI), p-value: 0.10 (0.02 - 0.17), p=0.010]. Next, we recruited 17 urban and 17 rural age-and-BMI-matched healthy-young-male volunteers for a 5-day HFD challenge. The HOMA-IR increased in both groups similarly [-0.77 (-2.03 - 0.49), p=0.223]. Neither rural living nor factors associated with rural living such as current helminth infection and total IgE were associated with protection against acute induction of IR by HFD.

INTRODUCTION

The prevalence of obesity and type 2 diabetes (T2D) is increasing worldwide, especially in low and middle-income countries (LMIC) that are currently facing rapid rate of urbanization.[1, 2] Rural-to-urban migration has indeed been shown to be associated with increased obesity and other cardiovascular (CV) risk factors, such as dyslipidemia and hypertension,[3-11] suggesting that living in rural environment might be protective against T2D development.

In addition to changes towards a sedentary lifestyle and increased dietary fat intake, migration to an urban environment is also associated with a reduction exposure to microorganism and parasites, such as helminth infections, which are still endemic in many rural areas of LMIC.[12] Recent data suggests that helminth infections might confer a protection against the development of obesity and T2D,[13-16] presumably by promoting type-2 and regulatory immune responses and subsequent reduction in systemic inflammation.[17-19] However, it is worth mentioning that the relative contribution of helminth infections in comparison to the more established factors such as a sedentary lifestyle and diet remains to be clarified.

Urban subjects have been reported to have longer sedentary periods and shorter active periods compared to those living in rural areas.[20] Furthermore, an increase in dietary fat intake, commonly observed upon rural-to-urban migration,[7,20] has been reported to be associated with impaired insulin resistance (IR) and glucose homeostasis.[21] Mice on high-fat diet (HFD) have provided models to study obesity and the development of IR.[22,23] Similarly, in humans, short-term HFD has been utilized to study the susceptibility to the development of IR.[24-28] Using this model, it has been possible to show how risk of IR is dependent on whether the participant is Caucasian or South Asian.[25,28] Short-term HFD has also been shown to induce organ-specific and systemic inflammation as evidenced by the increase in plasma cholesteryl ester transfer protein (CETP) levels,[24,29] predominantly produced by Kupffer cells (KC),[30] and plasma C-reactive protein (CRP) levels.[24]

Taken together, the chronic increase of energy rich diet, in addition to a more sedentary lifestyle, among people who migrate from a rural to urban areas,[20] might lead to the development of IR and T2D. However, there is still incomplete insight into the pathophysiology of the development of IR and T2D in rural-to-urban migration. In addition, there has been no study comparing the metabolic response towards a short-term HFD in terms of changes in glucose homeostasis and inflammation, between people living in urban and rural areas.

As some metabolic differences between individuals living in rural and urban area can be due to genetic differences, this study compared the metabolic profile between individuals with the same genetic background living in urban and rural areas. We also compared the metabolic and inflammatory response of individuals living in a rural and an urban area to a 5-day high-fat high-calorie (HFD) diet. Furthermore, since rural areas often go hand in hand with helminth infections and its associated IgE responses, we aimed to assess their contribution to metabolic profile. We hypothesized that individuals living in rural area would have a better metabolic profile and would be relatively more protected from the induction of IR and inflammation by the HFD compared to those living in an urban area.

RESULTS

The metabolic profile of rural and urban study participants.

The mean length of stay of urban subjects in Jakarta was 20.7 (range: 1-40) years. The differences in metabolic profile between subjects living in rural and urban are summarized in **Table 1**. Urban subjects had a significantly higher homeostatic model assessment (HOMA)-IR compared to rural subjects [1.45 (1.06 - 1.90) vs. 0.96 (0.80 - 1.13), respectively, $p = 0.01$]. Similarly, other metabolic parameters, such as 2-hour blood glucose, hemoglobin A1c (HbA1c), body mass index (BMI), waist circumference, and leptin level were significantly higher in urban subjects (**Table 1**). Interestingly, independent of age, increasing length of stay in urban area (in years) was positively associated with increasing BMI (in kg/m²) [estimate (95% CI), 0.15 (0.04 - 0.27), $p = 0.01$, **Figure 1A**], waist circumference (in cm) [0.45 (0.14 - 0.76) cm, $p=0.006$, **Figure 1B**], but not HOMA-IR [0.005 (-0.003 - 0.013), $p = 0.18$]. Increasing length of stay in urban area was also associated with a trend of increase in leptin level (in ng/mL) [0.013 (-0.001 - 0.027), $p = 0.07$].

The prevalence of soil-transmitted helminth (STH) was significantly lower in the urban compared to rural subjects [5% (2/42) vs. 57% (52/92), respectively, $p < 0.0001$]. Similarly, the levels of total IgE, often driven by STH infections,[31] were lower in the urban compared to rural subjects (168 (105–271) IU/mL vs. 931 (702–1,235) IU/mL, respectively, $p < 0.0001$) (**Table 1**). As the number of subjects with current STH infections in urban area was very low ($n = 2$), it was not possible to assess the contribution of current STH infections to the HOMA-IR difference between urban and rural subjects. Therefore, we used the total IgE level as a proxy for past and current exposure to STH. The age-adjusted difference in HOMA-IR between urban and rural subjects was slightly attenuated [from estimated mean differences (95% CI), 0.09 (0.02–0.17), $p = 0.001$ to 0.08 (–0.00–0.17), $p = 0.06$] after further adjustment for total IgE level (**Table 2**). Moreover, adjustment for total IgE level also attenuated the age-adjusted difference in waist circumference [from 7.2 (2.0–11.3) cm, $p = 0.001$ to 4.2 (–0.5–8.8) cm, $p = 0.08$] and leptin level [from 0.36 (0.18–0.55) ng/mL, $p < 0.0001$ to 0.10 (–0.03–0.24) ng/mL, $p = 0.14$] (**Table 2**). When assessing the contribution of adiposity and leptin levels to the difference in HOMA-IR between urban and rural subjects, we observed that adjustment for waist circumference [to 0.02 (–0.04–0.08), $p = 0.55$] or both waist circumference and leptin level [to 0.01 (–0.06–0.07), $p = 0.77$] strongly attenuated the difference in HOMA-IR (**Table 2**).

Table 1. Comparison of metabolic profiles between subjects living in urban and rural area.

Variables	Urban (n=49)	Rural (n=105)
Duration in urban (in years)	20.7 (1.0-40.0)	-
Age (in years)	39.3 (13.5)	44.5 (12.2)*
HOMA-IR	1.45 (1.06 - 1.90)	0.96 (0.80 - 1.13)*
Fasting Insulin (mU/L)	4.9 (3.8 - 6.4)	3.1 (2.5 - 3.8)**
Fasting Blood Glucose (mmol/L)	5.7 (1.4)	5.4 (0.9)
2h-Blood Glucose (mmol/L)	7.7 (3.2)	5.9 (1.9)**
HbA1c ^a (mmol/L)	37.9 (14.3)	32.3 (6.6)*
HbA1c ^a (%)	5.6 (1.3)	5.1 (0.6)*
Body Mass Index (kg/m ²)	24.3 (4.9)	22.7 (4.0)*
Waist Circumference (cm)	84.9 (13.8)	79.3 (11.9)*
Adiponectin (µg/mL)	4.38 (3.31 - 5.78)	3.54 (3.09 - 4.07)
Leptin (ng/mL)	5.62 (3.98 - 7.92)	2.64 (2.06 - 3.38)*
CRP (mg/L)	1.57 (1.17 - 2.05)	1.67 (1.29 - 2.11)
Total IgE (IU/mL)	168 (105 - 271)	931 (702 - 1,235)**
Prevalence of STH (% , n/N)	5 (2/42)	57 (52/92)**

All variables are presented as mean and its standard deviation, however, HOMA-IR, fasting insulin, adiponectin, leptin, CRP, and total IgE level are presented as geomean (95%CI) and were log transformed for analysis, while duration in urban is presented as mean (range). Analysis for the difference between urban and rural group was performed using independent t-test (*p<0.05, **p<0.0001) ^aHba1c measurements were available in 42 and 95 of urban and rural subjects respectively. Abbreviation: HOMA-IR=the homeostatic model assessment of insulin resistance, CRP=C-reactive protein, STH=soil-transmitted helminth.

In addition, we stratified rural and urban subjects based on STH infection status into three groups, resulting in an urban group without STH infections, a rural group without STH infections, and a rural group with STH infections. The highest mean level of HOMA-IR, waist circumference, and leptin was observed in the urban group without STH infections, followed by the rural group without STH infections and the lowest among the rural group with STH infections (**Figure S1**). The opposite relationship was observed for total IgE level (**Figure S1**).




Figure 1. The association between length of stay in urban area with adiposity. The association between length of time in urban area with (A) body mass index (BMI) and (B) waist circumference are presented in scatter plot graphs (n=49), and analysed using age-adjusted linear regression. Each year increase of a time spent in urban area was associated with a significant increase in both (a) BMI [0.152 (0.036 - 0.269) kg/m², p=0.012] and (b) Waist Circumference [0.449 (0.135 - 0.762) cm, p=0.006].

Table 2. Associations between living in urban and rural area with HOMA-IR, leptin, and waist circumference.

Variables	Differences for each variable between urban and rural (rural group as the reference group)*					
	Crude	Model 1 (Age)	Model 2 (Age+Total IgE)	Model 3 (Age+Waist)	Model 4 (Age+Total IgE+Waist)	Model 5 (Age+Waist+Leptin)
HOMA-IR [§]	0.10 (0.02 - 0.17) p=0.010	0.09 (0.02 - 0.17) p=0.016	0.08 (-0.00 - 0.17) p=0.061	0.02 (-0.04 - 0.08) p=0.545	0.04 (-0.03 - 0.11) p=0.294	0.01 (-0.06 - 0.07) p=0.774
Leptin (ng/mL) [§]	0.33 (0.14 - 0.51) p=0.001	0.36 (0.18 - 0.55) p<0.0001	0.10 (-0.03 - 0.24) p=0.137	0.11 (-0.01 - 0.22) p=0.076	0.08 (-0.05 - 0.21) p=0.216	-
Waist Circumference (cm)	5.6 (1.3 - 9.9) p=0.010	7.2 (3.0 - 11.3) p=0.001	4.2 (-0.5 - 8.8) p=0.077	-	-	-

*Beta coefficient (95% CI) from linear regression. [§]HOMA-IR and leptin level were log transformed for analysis. Model 1: adjusted for age. Model 2: adjusted for model 1 plus total IgE level. Model 3: adjusted for model 1 plus waist circumference. Model 4: adjusted for model 2 plus waist circumference. Model 5: adjusted for model 3 plus leptin level. Abbreviation: HOMA-IR = the homeostatic model assessment of insulin resistance.

Comparison of metabolic responses after a short-term HFD intervention between subjects living in an urban and rural area

Among subjects who were included in the interventional part of the study ($n = 34$), we observed no significant differences between the age-and-BMI-matched urban ($n = 17$) and rural group ($n = 17$) in terms of HOMA-IR, adipose-IR index, CRP, and lipid levels at D-0 (Pre HFD). At this time point, serum CETP levels were significantly lower in the urban group [1.96 (0.58) $\mu\text{g/mL}$ vs 2.59 (0.64) $\mu\text{g/mL}$, in urban and rural group respectively, $p = 0.006$]. Both groups showed a good compliance in terms of dietary intervention, all participants consumed all the cream provided and maintained their regular diet, resulting in a mean daily calorie intake that was $\sim 60\%$ higher compared to their regular diet, and $\sim 56\%$ of energy was derived from fat. The details of the dietary composition are shown in the **Supplementary Table S1**.

Intervention with a 5-day HFD resulted in a significant increase of HOMA-IR in both the urban [from 0.78 (0.51 - 1.09) to 1.13 (0.78 - 1.57), $p = 0.03$] and rural group [from 0.87 (0.59 - 1.21) to 1.69 (1.01 - 2.45), $p = 0.001$] (**Figure 2A, Table S1**), which was mainly driven by the increase in fasting insulin level in both urban [from 4.05 (2.98 - 5.52) mU/L to 5.59 (4.18 - 7.47) mU/L , $p = 0.02$] and rural group [from 4.63 (3.42 - 6.26) mU/L to 7.68 (5.70 - 10.34) mU/L , $p = 0.001$] (**Table S1**). Comparing the changes in IR before and after intervention between urban and rural group, we observed no significant differences for either HOMA-IR [estimated mean differences (95% CI), -0.77 (-1.95 - 0.41), $p = 0.21$] (**Figure 2A, Table S2**) or adipose-IR index [-41.2 (-115.1 - 32.7), $p = 0.28$] (**Figure 2B, Table S2**).

Interestingly, we observed a significant increase in CETP levels after HFD intervention in the urban group only [from 1.96 (0.58) $\mu\text{g/mL}$ to 2.28 (0.63) $\mu\text{g/mL}$, $p = 0.004$ in urban group vs from 2.59 (0.64) $\mu\text{g/mL}$ to 2.58 (0.72) $\mu\text{g/mL}$, $p = 0.93$ in rural group) (**Figure 2C**). Therefore, in comparison to the rural group, the increase in CETP level was significantly higher in urban group [0.33 (0.06 - 0.60) $\mu\text{g/mL}$, $p = 0.02$] (**Figure 2C, Table S2**). However, as indicated above, the CETP levels were already much higher in the rural group at D-0 (Pre HFD), even higher than the D-6 (post-HFD) CETP level in the urban group. Intervention with the HFD did not significantly increase CRP levels

in the two groups (Table S2). When assessing the effects of HFD on lipid levels, we observed no significant difference in changes in total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and triglyceride (TG) levels between the urban and rural group, while the increase in high-density lipoprotein cholesterol (HDL-C) after intervention was significantly higher in the urban group compared to the rural group [0.09 (0.01-0.17) mmol/L, $p = 0.04$] (Table S2).




Figure 2. Comparison of metabolic responses to high-fat diet. HOMA-IR and adipose-IR index are presented as geometric mean and its corresponding 95% confidence interval, while CETP levels are presented as mean with its standard deviation. There were no significant differences in the increase of HOMA-IR (A), adipose-IR index (B) between urban and rural group, however, the increase in CETP level (C) was higher in the urban group. Furthermore, in rural group, there were no significant differences in the increase of HOMA-IR (D), adipose-IR index (E), and CETP level (F) between STH-infected and uninfected group. The difference between before and after intervention for each group was analysed using paired t-test, while the difference in the magnitude of changes for each parameter was analysed using linear mixed model (* $p < 0.05$, NS: $p > 0.05$).

The effect of current STH infections on the metabolic responses upon short-term HFD intervention.

Next, due to the very low prevalence of STH infections in the urban group [6% (1/17)], the effect of current STH infections on the metabolic response towards a short-term HFD intervention was only assessed in the rural group of which 50% was positive for STH infection (8/16). Thus, our study was underpowered (power of 56%) to detect any differences in metabolic responses between STH-infected and uninfected subjects.

Despite a significantly lower baseline body weight in STH-infected subjects in comparison to STH-uninfected subjects [51.1 (11.0) kg vs 63.3 (10.2) kg, $p = 0.04$], there was no significant difference in the magnitude of increase in HOMA-IR [-1.08 (-3.38-1.22), $p = 0.36$] (**Figure 2D**), adipose-IR [-87.8 (-222.1-46.4), $p = 0.21$] (**Figure 2E**), or CETP level [-0.21 (-0.62-0.20) $\mu\text{g/mL}$, $p = 0.32$] (**Figure 2F**) after HFD intervention. Interestingly, we observed a significantly higher increase in LDL-C level [0.27 (0.05-0.49) mmol/L, $p = 0.03$] after HFD intervention among STH-infected subjects in comparison to STH-uninfected subjects (**Table S3**). However, the level of LDL-C was much lower in the STH-infected group at D-0 in comparison to the STH-uninfected group [2.22 (0.31) mmol/L vs 2.97 (0.63) mmol/L, $p = 0.01$], and the level of LDL-C at D-6 in STH-infected group [2.44 (0.28) mmol/L] did not reach the level of LDL-C in the STH-uninfected group at D-0 (**Table S3**).

DISCUSSION

Our study showed that, in comparison to individuals living in a rural area, those living in an urban area had higher whole-body IR, as assessed by HOMA-IR, adiposity, and leptin levels. Whereas the higher whole-body IR was mainly mediated by the higher adiposity and leptin levels observed in urban individuals, differences in exposures to STH infection between urban and rural individuals, might contribute to a small extent to the differences observed in whole-body IR, adiposity or leptin levels. Intervention with a short-term HFD increased whole-body IR in both the urban and rural group. In comparison to the rural group, the CETP level was lower in the urban group, and the HFD intervention induced a stronger increase in CETP levels in this group. The presence of STH infections did not seem to have a

protective effect on the acute induction of IR from short-term HFD. However, it has to be noted that our study was underpowered to detect an effect of STH.

Our study found that the higher whole-body IR in individuals living in urban area was mediated by the higher adiposity, as well as a higher leptin level, a pro-inflammatory adipokine, which has been previously reported to be associated with glucose metabolism.[8,32] The increase of adiposity and, to a lesser extent, leptin level, was positively associated with the duration of time spent in the urban environment. This suggests that a higher degree of acculturation in terms of urban lifestyle, drifting away from their traditional lifestyle,[11] could lead to a positive energy balance,[20] hence increasing adiposity over time. In addition, reduced exposures to environmental factors, such as to STH infections, which have been shown to have beneficial metabolic effects[13] partly through the induction of type-2 and regulatory immune response,[18,19] might contribute to the difference in whole-body-IR, adiposity, and leptin level between urban and rural individuals. This was supported by our finding that the difference in whole-body IR, adiposity, and leptin level between urban and rural individuals was attenuated, but only slightly, after adjustment for total IgE level, a general marker for type-2 immune responses, and a proxy for past and current STH exposures.[31]

As expected, the overall metabolic profile of individuals living in a rural area, in term of adiposity and whole-body IR, was better in comparison to those living in an urban area. However, in contrast to our hypothesis, a short-term 5-day HFD intervention induced a similar increase of IR in both urban and rural individuals. As both groups were BMI-matched, these findings suggest that the direct protective metabolic effect of a combined past and current environmental exposures to helminths,[13] independent of their effect on adiposity, might be relatively weak in comparison to the strong induction of IR by the HFD intervention. Indeed, our group has recently reported that the increased IR in STH-infected subjects after deworming was mainly mediated by the increased adiposity.[16] Thus, adjusting for adiposity, in a way, remove the possible main pathway for STH-associated protection against the development for IR.

Although our study was underpowered to assess the effect of current STH

infection, it is possible that the presence of current STH infections might not be sufficient to protect against a strong induction of IR by short-term HFD, as in rural subjects, the increase in IR after HFD in STH-infected and STH-uninfected subjects was similar. However, it is also possible that the HFD intervention in STH-infected subjects with lower body weight would have a stronger impact than in STH-uninfected subjects, thereby masking any protective effects of STH infections.

Interestingly, we observed that the baseline serum CETP level was significantly lower in urban subjects. As CETP is mainly produced by KCs, higher CETP level may represent an increase in hepatic macrophage (KC) content, hence liver inflammation. [30] Also, environmental factors in the rural area, mainly exposure to various infectious agents, such as microorganisms and parasites, may explain the increased CETP level. For instance, it has been shown that subjects with chronic hepatitis C virus infection have elevated serum CETP levels.[33] Supporting this, the prevalence of hepatitis in our rural study area was higher than our urban study area (4.3% vs 0.8%).[34] However, currently, there are no available data connecting macrophage polarization status to CETP level and therefore further studies are needed.[35] In contrast to what is seen in urban subjects,[24,29] we found no increase in CETP levels in rural subjects after the HFD intervention. It is possible that the lack of an increase in CETP levels in rural subjects was caused by the already high baseline CETP levels, thus precluding its further increase after HFD intervention. Our results suggest an inflammation-independent mechanism of short-term HFD-associated induction of IR[23] as there was no significant increase in CRP following HFD. Studies on the role of inflammation in HFD-associated induction of IR have shown conflicting results. In one study, an increase in CRP and expression of M1 macrophage markers in skeletal muscle was reported,[24] while in another, no increase was seen in circulating pro-inflammatory cytokines.[36]

In terms of lipid levels, while no significant changes in lipid levels were observed in rural group, HFD intervention significantly increased HDL-C level in urban group. Our study and other observed that urban subjects had a relatively higher fat intake than rural subjects[20] at baseline. Thus, the relative difference in the changes of dietary composition before and after intervention[37-39] between

urban and rural individuals might potentially contribute to the difference in HDL-C level changes after intervention. In the rural group, we observed no significant changes in LDL-C in STH-uninfected subjects, whereas the HFD intervention resulted in a significant increase in LDL-C in STH-infected subjects. This might be related to the lower baseline LDL-C level and body weight in STH-infected subjects.

Our study is the first to compare the metabolic profile between people with the same genetic background, living in different environments (urban and rural) and to assess the metabolic responses to an intervention with a standardized short-term HFD. However, our study has several limitations. First, our study was only performed in male subjects, and potential differences in the outcomes might be observed in females. Next, due to the low prevalence of STH in urban area, our study could only assess the effect of current STH infections on HFD-induced IR in rural subjects. We also used a calculated HOMA-IR instead of the gold standard glycemic clamp to assess IR. In addition, there was no data available on physical activity, there were no biopsies of specific metabolic tissues (liver, muscle, adipose tissue), and we did not analyze the gut microbiota, all known to play an important role in metabolic profile and response.

In conclusion, in comparison to their rural ethnic counterparts, individuals living in an urban area had a higher whole-body IR, which was mainly mediated by their higher adiposity. The differences between urban and rural individuals in terms of past and current exposures to STH seem to have a relatively small contribution to the difference in whole-body IR. Contrary to our hypothesis, intervention with a short-term HFD induced similar increase in IR, in urban and rural individuals, and in helminth infected and uninfected subjects. However, well-powered larger studies are needed to determine which factors in terms of urbanization contribute to IR.

METHODS

Study Design and Population

The present study consisted of a cross-sectional and an interventional study. The cross-sectional study was performed in an urban (Jakarta) and a rural area (Nangapanda, Ende, Flores island) in Indonesia. We recruited 49 males (18-65

years old) with Floresian ethnic background who had migrated from Flores island and lived in Jakarta for more than 1 year (urban group). As their rural counterparts, we recruited 105 Floresian males with a similar age range, randomly selected from three villages in Nangapanda with age stratification, as described previously.[40]

For the HFD intervention study, 17 from urban and 17 from rural area, age-and-BMI-matched healthy young male volunteers (18-40 years old) were recruited via local healthcare workers who informed their community, in both Nangapanda and Jakarta, of the study. BMI-matching was performed to assess whether the difference between urban and rural in term of past or current exposure to STH infections affect the HFD-associated increase in IR, independent of adiposity. Exclusion criteria were T2D, recent body weight changes, intake of medication that could affect inflammation or IR.

The study was approved by the Medical Ethical Committee of the Faculty of Medicine, Universitas Indonesia (556/H2.F1/ETIK/2014) and performed in accordance with the principles of the revised Declaration of Helsinki. All volunteers gave written informed consent before participation.

Cross sectional Study

In the cross-sectional study, we invited all subjects to come to the Field Study Centre (FSC) in both rural and urban area to undergo clinical measurements and blood sample collections. Stool samples were also collected. All clinical measurements and blood sample collections were performed after an overnight fast. Anthropometric measurements of body weight, height, and waist circumference were performed. BMI was calculated as weight in kg divided by square of height in meter.

After collection of fasting blood samples, we performed an oral glucose tolerance test (OGTT), in which blood glucose levels were re-measured 2 hours after subjects were given 75g glucose dissolved in 200 mL of water (2h-BG). In this cross sectional study, we calculated HOMA-IR (homeostatic model assessment of insulin resistance), a well-validated measure of whole-body IR in humans ($\text{HOMA-IR} = \text{fasting serum insulin (mU/L)} \times \text{fasting glucose (mmol/L)} / 22.5$),[41] as our primary outcome. We

also measured HbA1c, fasting blood glucose (FBG), fasting insulin, 2h-BG, BMI, waist circumference, adiponectin, leptin, high-sensitive C-reactive protein (hsCRP), total IgE, and prevalence of soil-transmitted helminths (STH) as our secondary outcomes.

Intervention Study

Subjects were examined before and after a 5-day HFD intervention, consisting of the subject's regular diet supplemented with 375 mL cream (Greenfields™ Whipping Cream, Greenfields Indonesia Ltd, Jakarta, Indonesia) per day [1,500 kcal/day, 83% fat (60% saturated fat)]. After baseline measurements, each subject received three bottles of 125 mL cream per day for five consecutive days. Subjects were instructed to continue their regular diet, and to consume one bottle of cream after each meal (3 meals per day) to make sure they could adhere to their regular dietary habits.

Subjects were asked to keep a food diary before and during the HFD intervention to estimate normal dietary intake and to check for compliance and compensatory behavior. Dietary assessment, using a 24 hours food recall, was performed by a trained dietician. Compliance was further assessed by interviewing the subject and collecting the bottles every day. During the study, subjects were asked not to change lifestyle habits. Measurements of clinical parameters and blood drawing were done on the day before starting the HFD intervention (D-0) and one day after the fifth day of the HFD intervention (D-6).

In this intervention study, we had HOMA-IR as our primary outcome. As our secondary outcomes, we measured adipose-IR index, a measure of adipose tissue IR, which was calculated as the product of the fasting serum free fatty acid (FFA) and insulin (Adipose-IR index = FFA[mM] x Insulin [pM]).[42,43] In addition, we also measured hsCRP, CETP, and lipid levels (TC, HDL-C, TG, LDL-C). Due to limited amount of sera after intervention, adiponectin and leptin level were measured only at baseline. All others measurements for the interventional study were performed pairwise (before and after intervention).

Laboratory measurements

Fasting blood glucose and 2h-post-load glucose were measured in capillary blood using Breeze®2 glucose meters (Bayer Health Care LLC, Basel, Switzerland) in the FSC. All sera, plasma and whole blood samples from rural area were frozen at -20°C in the FSC, and subsequently shipped and stored at -80°C in Faculty of Medicine Universitas Indonesia (FKUI), Jakarta, Indonesia and Leiden University Medical Centre (LUMC), Leiden, The Netherlands. All sera, plasma and whole blood samples from urban area were directly transported from FSC (Jakarta) to be stored at -80°C in FKUI, and subsequently shipped and stored at -80°C at LUMC.

Serum insulin concentrations were determined by a solid-phase, enzyme-labelled chemiluminescent immunometric assay, while HbA1c was measured using a cation-exchange chromatography (IC)-based high performance liquid chromatography (HPLC) assay. A latex-enhanced immunoturbidimetric method was used to measure hsCRP. Assays of TC, HDL-C, and TG were based on enzymatic colorimetric methods. These measurements have been described previously.[16]

Plasma CETP levels were measured with enzyme-linked immunosorbent assays (ELISA) kits according to the manufacturer's instructions (DAIICHI CETP ELISA, Daiichi, Tokyo, Japan). FFA were measured using ELISA kits according to the manufacturer's instructions (abcam ab 65341 FFA Quantification Assay Kit, Cambridge, UK). Adiponectin and leptin were also measured by using ELISA commercial reagents (DuoSet ELISA R&D System Europe Ltd, Abingdon, UK). The inter- and intra-assay coefficients of variance (CV) of adiponectin were 3.1% and 7.0% respectively. While for leptin, the inter- and intra-assay CV were 2.2% and 3.2%, respectively. The levels of total IgE, an important determinant of helminth infection,[31] were measured using ELISA as described previously.[44] The presence of STH [hookworm (*Necator americanus*, *Ancylostoma duodenale*), *Ascaris lumbricoides*, *Trichuris trichiura*, *Strongyloides stercoralis*] was assessed using PCR as described in detail elsewhere.[44,45]

Statistical Analysis

Normally distributed continuous variables were summarized as mean and standard deviation [mean (SD)], while non-normally distributed data were summarized

as geometric mean and its 95% confidence interval [geomean (95% CI)]. Based on previous studies,[14,25] we aimed to recruit 45 subjects from each urban and rural area for the cross-sectional study, while for the interventional study we aimed to recruit 15 subjects from each group (see **Supplementary Material**).

The original plan for the linear regressions was based on a conceptual framework (**Figure 3**) of the proposed causal pathways. In the cross-sectional study (A), we assessed whether the difference between urban and rural subjects, in term of past or current exposure to STH, by using total IgE level as a proxy, contributes to the difference in insulin resistance (IR) between subjects living in urban and rural area, and whether this difference in IR is independent from adiposity, by performing mediation analysis. Next, we further stratified the urban and rural group based on their STH infection status (see **Supplementary Material**). In addition, we also assessed the association between length of stay in urban area and metabolic profiles (IR, adiposity, and leptin) among subjects living in urban area using age-adjusted linear regression model.

In the HDF intervention study (B), first, we assessed whether the difference between urban and rural in term of past or current exposure to STH infections affect the HFD-associated increase in IR, independent of adiposity, by matching both groups for BMI. Next, among subjects living in rural area, similar model was used to further assess whether the presence of current STH infections protect against the HFD-associated increase in IR. The mixed model analysis was performed using R software (lme4).




Figure 3. Conceptual framework. In the cross-sectional study (A), we assessed whether the differences in past or current exposure to helminths contribute to the difference in insulin resistance (IR) between subjects living in urban and rural area, and whether the observed difference in IR is independent from adiposity. In the high-fat diet (HFD) study (B), first, we assessed whether past or current exposure to helminths protect against the HFD-associated increase in IR, independent of adiposity. Next, we also assessed whether the presence of current helminth infection protect against the HFD-associated increase in IR. *Past and current exposure to helminths was assessed by measuring serum total IgE level, a general marker for Th2 responses, commonly induced by soil-transmitted helminth (STH). **Current exposure to helminths was assessed using stool PCR. #Other factors that were not specifically assessed in this study.

Acknowledgements

This study was funded by Universitas Indonesia (1561/UN2.R12/HKP.05.00/2015), Ministry of Research, Technology and Higher Education Republic of Indonesia (0499/UN2.R12/HKP.05.00/2015 and 1100/UN2.R12/HKP.05.00/2016), and The Royal Netherlands Academy of Arts and Science (KNAW), Ref 57-SPIN3-JRP. The authors would like first to thank all study participants in Nangapanda and Jakarta, Indonesia. The authors would like to thank all local government and health officers in Nangapanda who supported this project, and also all field workers from Universitas Indonesia, Nangapanda and Jakarta. The authors would also like to thank Abdurrahman Hadi and Mohammad Rizki for their technical supports.

Author Contributions Statement

Conception or study design: D.L.T., K.R., P.C.N.R., E.S., P.S., D.S.H., J.W.A.S., M.Y. Sample collection, data analysis: D.L.T., K.R., F.K., Y.D., Y.W., S.M.E.N., E.I., D.M., E.Y., P.C.N.R., E.S., J.W.A.S., M.Y. Interpretation of data: D.L.T., K.R., F.K., Y.D., Y.W., S.M.E.N., E.I., T.S., E.Y., B.G., P.C.N.R., E.S., P.S., D.S.H., J.W.A.S., M.Y. Drafting the manuscript: D.L.T., E.S., M.Y., K.R. Revising the manuscript critically for important intellectual content; D.L.T., K.R., F.K., Y.D., Y.W., S.M.E.N., E.I., D.M., E.Y., B.G., T.S., P.C.N.R., E.S., P.S., D.S.H., J.W.A.S., M.Y. Principal investigator of this study: D.S.H. Scientific coordinator of this study: P.S., J.W.A.S., M.Y. All authors have given approval for publication.

Competing Interests

All authors declare no competing interests. The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Author-details

¹*Division of Endocrinology, Metabolism, and Diabetes, Department of Internal Medicine, Dr. Cipto Mangunkusumo National General Hospital, Faculty of Medicine Universitas Indonesia, Jakarta, Indonesia.*

²*Department of Parasitology, Leiden University Medical Center, Leiden, The Netherlands.*

³*Nangapanda Community Research Cluster, The Indonesian Medical Education and Research Institute, Jakarta, Universitas Indonesia, Indonesia*

⁴*Metabolic, Cardiovascular and Aging Research Cluster, The Indonesian Medical Education and Research Institute, Universitas Indonesia, Jakarta, Indonesia*

⁵*Department of Parasitology, Faculty of Medicine Universitas Indonesia, Jakarta, Indonesia.*

⁶*Department of Medicine, Division of Endocrinology, Leiden University Medical Center, Leiden, The Netherlands.*

⁷*Laboratory Unit, South East Asian Minister of Education Organization Regional Centre For Food And Nutrition, Jakarta, Indonesia*

⁸*Dr. W.Z. Johannes Hospital, Kupang, Indonesia*

⁹*Department of Internal Medicine, Radboud University Medical Centre, Nijmegen, The Netherlands.*

REFERENCES

1. Collaboration NCDRF. Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* **2016**; 387(10027): 1513-30.
2. IDF. IDF Diabetes Atlas, 7th edn. 7 ed. Brussels, Belgium: International Diabetes Federation, **2015**.
3. Ebrahim S, Kinra S, Bowen L, et al. The effect of rural-to-urban migration on obesity and diabetes in India: a cross-sectional study. *PLoS Med* **2010**; 7(4): e1000268.
4. Carrillo-Larco RM, Bernabe-Ortiz A, Pillay TD, et al. Obesity risk in rural, urban and rural-to-urban migrants: prospective results of the PERU MIGRANT study. *Int J Obes (Lond)* **2016**; 40(1): 181-5.
5. Lyngdoh T, Kinra S, Shlomo YB, et al. Sib-recruitment for studying migration and its impact on obesity and diabetes. *Emerg Themes Epidemiol* **2006**; 3: 2.
6. Unwin N, McLarty D, Machibya H, et al. Changes in blood pressure and lipids associated with rural to urban migration in Tanzania. *J Hum Hypertens* **2006**; 20(9): 704-6.
7. Unwin N, James P, McLarty D, et al. Rural to urban migration and changes in cardiovascular risk factors in Tanzania: a prospective cohort study. *BMC Public Health* **2010**; 10: 272.
8. Lindgarde F, Ercilla MB, Correa LR, Ahren B. Body adiposity, insulin, and leptin in subgroups of Peruvian Amerindians. *High Alt Med Biol* **2004**; 5(1): 27-31.
9. Miranda JJ, Gilman RH, Smeeth L. Differences in cardiovascular risk factors in rural, urban and rural-to-urban migrants in Peru. *Heart* **2011**; 97(10): 787-96.
10. Hernandez AV, Pasupuleti V, Deshpande A, Bernabe-Ortiz A, Miranda JJ. Effect of rural-to-urban within-country migration on cardiovascular risk factors in low- and middle-income countries: a systematic review. *Heart* **2012**; 98(3): 185-94.
11. Delavari M, Sonderlund AL, Swinburn B, Mellor D, Renzaho A. Acculturation and obesity among migrant populations in high income countries--a systematic review. *BMC Public Health* **2013**; 13: 458.
12. Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the great neglected tropical diseases. *J Clin Invest* **2008**; 118(4): 1311-21.
13. Tracey EF, McDermott RA, McDonald MI. Do worms protect against the metabolic syndrome? A systematic review and meta-analysis. *Diabetes Res Clin Pract* **2016**; 120: 209-20.
14. Wiria AE, Hamid F, Wammes LJ, et al. Infection with Soil-Transmitted Helminths Is Associated with Increased Insulin Sensitivity. *PLoS One* **2015**; 10(6): e0127746.
15. Hays R, Esterman A, Giacomini P, Loukas A, McDermott R. Does Strongyloides stercoralis infection protect against type 2 diabetes in humans? Evidence from Australian Aboriginal adults. *Diabetes Res Clin Pract* **2015**; 107(3): 355-61.
16. Tahapary DL, de Ruiter K, Martin I, et al. Effect of Anthelmintic Treatment on Insulin Resistance: A Cluster-Randomized Placebo-Controlled Trial in Indonesia. *Clinical Infectious Diseases* **2017**; 65:764-71
17. Wammes LJ, Mpairwe H, Elliott AM, Yazdanbakhsh M. Helminth therapy or elimination: epidemiological, immunological, and clinical considerations. *Lancet Infectious Diseases* **2014**; 14(11): 1150-62.
18. Wiria AE, Sartono E, Supali T, Yazdanbakhsh M. Helminth infections, type-2 immune response, and metabolic syndrome. *PLoS Pathog* **2014**; 10(7): e1004140.
19. de Ruiter K, Tahapary DL, Sartono E, et al. Helminths, hygiene hypothesis and type 2 diabetes. *Parasite Immunol* **2016**.
20. Yamauchi T, Umezaki M, Ohtsuka R. Influence of urbanisation on physical activity and dietary changes in Huli-speaking population: a comparative study of village dwellers and migrants in urban settlements. *Br J Nutr* **2001**; 85(1): 65-73.
21. Marshall JA, Bessesen DH. Dietary fat and the

- development of type 2 diabetes. *Diabetes Care* **2002**; 25(3): 620-2.
22. Winzell MS, Ahren B. The high-fat diet-fed mouse: a model for studying mechanisms and treatment of impaired glucose tolerance and type 2 diabetes. *Diabetes* **2004**; 53 Suppl 3: S215-9.
 23. Lee YS, Li P, Huh JY, et al. Inflammation is necessary for long-term but not short-term high-fat diet-induced insulin resistance. *Diabetes* **2011**; 60(10): 2474-83.
 24. Boon MR, Bakker LE, Haks MC, et al. Short-term high-fat diet increases macrophage markers in skeletal muscle accompanied by impaired insulin signalling in healthy male subjects. *Clin Sci (Lond)* **2015**; 128(2): 143-51.
 25. Bakker LE, van Schinkel LD, Guigas B, et al. A 5-day high-fat, high-calorie diet impairs insulin sensitivity in healthy, young South Asian men but not in Caucasian men. *Diabetes* **2014**; 63(1): 248-58.
 26. Brons C, Jensen CB, Storgaard H, et al. Impact of short-term high-fat feeding on glucose and insulin metabolism in young healthy men. *J Physiol* **2009**; 587(Pt 10): 2387-97.
 27. Thamer C, Haap M, Bachmann O, et al. Serum adiponectin levels predict the effect of short-term dietary interventions on insulin sensitivity in humans. *Diabetologia* **2004**; 47(7): 1303-5.
 28. Wulan SN, Westerterp KR, Plasqui G. Metabolic profile before and after short-term overfeeding with a high-fat diet: a comparison between South Asian and White men. *Br J Nutr* **2014**; 111(10): 1853-61.
 29. Widya RL, Hammer S, Boon MR, et al. Effects of short-term nutritional interventions on right ventricular function in healthy men. *PLoS One* **2013**; 8(9): e76406.
 30. Wang Y, van der Tuin S, Tjeerdema N, et al. Plasma cholesteryl ester transfer protein is predominantly derived from Kupffer cells. *Hepatology* **2015**; 62(6): 1710-22.
 31. Cooper PJ, Alexander N, Moncayo AL, et al. Environmental determinants of total IgE among school children living in the rural Tropics: importance of geohelminth infections and effect of anthelmintic treatment. *BMC Immunol* **2008**; 9: 33.
 32. Finucane FM, Luan J, Wareham NJ, et al. Correlation of the leptin:adiponectin ratio with measures of insulin resistance in non-diabetic individuals. *Diabetologia* **2009**; 52(11): 2345-9. Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care* **2004**; 27(6): 1487-95.
 33. Satoh K, Nagano T, Seki N, et al. High level of serum cholesteryl ester transfer protein in active hepatitis C virus infection. *World J Hepatol* **2016**; 8(5): 291-300.
 34. National Institute for Health Research and Development, Ministry of Health Republic of Indonesia. Riset Kesehatan Dasar (Riskesdas) 2013. Jakarta. **2013**.
 35. Haas JT, Staels B. Cholesteryl-ester transfer protein (CETP): A Kupffer cell marker linking hepatic inflammation with atherogenic dyslipidemia? *Hepatology* **2015**; 62(6): 1659-61.
 36. Wan Z, Durrer C, Mah D, Simtchouk S, Robinson E, Little JP. Reduction of AMPK activity and altered MAPKs signalling in peripheral blood mononuclear cells in response to acute glucose ingestion following a short-term high fat diet in young healthy men. *Metabolism* **2014**; 63(9): 1209-16.
 37. Guay V, Lamarche B, Charest A, Tremblay AJ, Couture P. Effect of short-term low- and high-fat diets on low-density lipoprotein particle size in normolipidemic subjects. *Metabolism* **2012**; 61(1): 76-83.
 38. Samaha FF. Effect of very high-fat diets on body weight, lipoproteins, and glycemic status in the obese. *Curr Atheroscler Rep* **2005**; 7(6): 412-20.
 39. Hooper L, Summerbell CD, Thompson R, et al. Reduced or modified dietary fat for preventing cardiovascular disease. *The Cochrane* **2012**; (5): CD002137.
 40. Tahapary DL, de Ruiter K, Martin I, et al. Helminth infections and type 2 diabetes: a cluster-randomized placebo controlled SUGARSPIN trial in Nangapanda, Flores, Indonesia. *BMC Infect Dis* **2015**; 15: 133.
 41. Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care* **2004**; 27(6): 1487-95.
 42. Gastaldelli A, Cusi K, Pettiti M, et al. Relationship

- between hepatic/visceral fat and hepatic insulin resistance in nondiabetic and type 2 diabetic subjects. *Gastroenterology* 2007; 133(2): 496-506.
43. Groop LC, Bonadonna RC, DelPrato S, et al. Glucose and free fatty acid metabolism in non-insulin-dependent diabetes mellitus. Evidence for multiple sites of insulin resistance. *J Clin Invest* **1989**; 84(1): 205-13.
44. Wiria AE, Prasetyani MA, Hamid F, et al. Does treatment of intestinal helminth infections influence malaria? Background and methodology of a longitudinal study of clinical, parasitological and immunological parameters in Nangapanda, Flores, Indonesia (ImmunoSPIN Study). *BMC Infect Dis* **2010**; 10: 77.
45. Kaiser MM, Brienen EA, Djuardi Y, et al. Improved diagnosis of *Trichuris trichiura* by using a bead-beating procedure on ethanol preserved stool samples prior to DNA isolation and the performance of multiplex real-time PCR for intestinal parasites. *Parasitology* **2017**: 1-10.

SUPPLEMENTARY MATERIALS




Figure S1. Comparison of metabolic profile between urban participants and rural participants (rural participants stratified by helminth infection status). The levels of total IgE, HOMA-IR, waist circumference, and leptin on different group of living area and soil-transmitted helminth (STH) infection status are presented as geometric mean and its 95% confidence interval, except for waist circumference which are presented as mean and its standard deviation. The number of urban subjects with helminth infections was very low (2/42) and was not included in this graph. Trend analysis was performed between three groups, namely: (1) rural subjects with STH infections [Rural (+)], (2) rural subjects without STH infections [Rural (-)], and (3) urban subjects without STH infections [Urban (-)]. Total IgE level was the lowest in Urban (-) group and progressively become higher in Rural (-) and Rural (+) groups (A). The contrary was observed for HOMA-IR (B), waist circumference (C), leptin level (D). * $p < 0.05$ in unadjusted model, ** $p < 0.05$ in age-adjusted model, *** $p < 0.05$ in age-waist circumference-adjusted model.

Table S1. Comparison of dietary compositions between urban and rural participants.

	Urban		Rural	
	Pre-HFD*	HFD	Pre-HFD*	HFD
Energy (kcal)	1617.6 (78.2)	2533.0 (124.0)	1890.3 (433.7)	2822.6 (248.6)
Fat (g)	51.8 (23.4)	161.2 (7.5)	50.7 (25.1)	173.9 (13.9)
Carbohydrate (g)	222.1 (69.3)	200.1 (49.6)	268.9 (66.4)	220.2 (32.3)
Protein (g)	67.2 (9.1)	68.8 (5.6)	78.0 (24.3)	81.5 (16.8)
Fiber (g)	9.4 (3.9)	9.5 (1.5)	10.0 (3.1)	11.3 (2.3)

*At baseline, the proportion of energy derived from fat, carbohydrate, and protein for subjects living in urban area were 29%, 55%, and 16% respectively, while for subjects living in rural area the percentages were 24%, 58%, and 18% respectively.

Table S2. Comparison of metabolic responses towards a short-term HFD between subjects living in an urban and rural area.

Variables	Urban n=17			Rural n=17			Comparison of the magnitude of changes between urban and rural subjects**	
	Pre HFD	Post HFD	p-value*	Pre HFD	Post HFD	p-value*	Estimated differences	p-value
Age (years)	30.1 (6.4)	-	-	29.5 (8.0)	-	-	-	-
Body Mass Index (kg/m ²)	23.1 (4.7)	-	-	21.6 (3.6)	-	-	-	-
HOMA-IR	0.78 (0.51 - 1.09)	1.13 (0.78 - 1.57)	0.03	0.87 (0.59 - 1.21)	1.69 (1.01 - 2.45)	0.001	-0.77 (-1.95 - 0.41)	0.21
Fasting Blood Glucose (mmol/L)	5.15 (0.44)	5.23 (0.50)	0.59	4.96 (0.19)	5.46 (0.73)	0.005	-0.42 (-0.82 - 0.03)	0.04
Fasting Insulin (mU/L)	4.05 (2.98 - 5.52)	5.59 (4.18 - 7.47)	0.02	4.63 (3.42 - 6.26)	7.68 (5.70 - 10.34)	0.001	-2.35 (-6.55 - 1.84)	0.28
Adipose-IR Index	51.6 (28.5 - 93.3)	71.9 (45.0 - 114.7)	0.23	40.5 (24.0 - 68.4)	72.0 (44.8 - 115.7)	0.006	-41.2 (-115.1 - 32.7)	0.28
Free Fatty Acid (mmol/L)	3.93 (3.37 - 4.59)	3.43 (2.91 - 4.04)	0.29	2.83 (2.42 - 3.30)	2.65 (2.27 - 3.09)	0.39	-0.32 (-1.24 - 0.59)	0.49
CETP (µg/mL) [#]	1.96 (0.58)	2.28 (0.63)	0.004	2.59 (0.64)	2.58 (0.72)	0.93	0.33 (0.06 - 0.60)	0.02
CRP (mg/L)	2.04 (0.94 - 4.74)	2.24 (1.21 - 3.75)	0.82	1.14 (0.59 - 1.89)	1.06 (0.58 - 1.71)	0.81	-0.31 (-3.61 - 2.98)	0.85
Total Cholesterol (mg/dL)	4.11 (0.61)	4.13 (0.59)	0.84	4.38 (0.61)	4.42 (0.71)	0.65	-0.02 (-0.29 - 0.24)	0.87
Triglyceride (mg/dL)	1.29 (0.53)	1.31 (0.54)	0.83	1.35 (0.36)	1.24 (0.43)	0.37	0.14 (-0.17 - 0.44)	0.38
HDL-C (mg/dL)	1.08 (0.25)	1.18 (0.23)	0.01	1.19 (0.25)	1.20 (0.24)	0.60	0.09 (0.01 - 0.17)	0.04
LDL-C (mg/dL)	2.44 (0.59)	2.35 (0.570)	0.251	2.57 (0.61)	2.65 (0.59)	0.21	-0.17 (-0.36 - 0.01)	0.08

All variables are presented as mean and its standard deviation, however, HOMA-IR, Fasting Insulin, Adipose-IR Index, Free Fatty Acid, and CRP levels are presented as geometric (95%CI). *The differences between before and after HFD diet intervention were analysed using paired t-test. **The difference in changes (before and after HFD) of different parameters between urban and rural were analysed using linear mixed model and are presented as [Estimated Differences in Changes (95%CI), p-value]. #CETP measurements were only available for 33 subjects. Abbreviation: HOMA-IR= homeostatic model assessment of insulin resistance, CETP= cholesteryl ester transfer protein, CRP= C-reactive protein, HDL-C= high-density lipoprotein cholesterol, LDL-C= low-density lipoprotein cholesterol.

Table S3. Comparison of metabolic responses towards a short-term HFHC diet between STH-infected and uninfected subjects living in rural area.

Variables	STH-infected n=8				STH-uninfected n=8				Comparison of the magnitude of changes between STH-infected and STH-uninfected subjects**
	Pre HFD	Post HFD	p-value*	Pre HFD	Post HFD	p-value*	Estimated differences	p-value*	
Age (years)	27.0 (9.6)	-	-	32.0 (6.3)	-	-	-	-	-
Body Mass Index (kg/m ²)	20.1 (3.5)	-	-	23.1 (2.4)	-	-	-	-	-
HOMA-IR	0.73 (0.37 - 1.19)	1.47 (1.08 - 1.93)	0.002	1.00 (0.45 - 1.75)	2.03 (0.72 - 4.34)	0.06	-1.08 (-3.38 - 1.22)	0.36	
Fasting Blood Glucose (mmol/L)	5.03 (0.22)	5.69 (0.88)	0.04	4.90 (0.17)	5.26 (0.57)	0.13	0.30 (-0.30 - 0.90)	0.34	
Fasting Insulin (mIU/L)	4.04 (2.60 - 6.29)	6.80 (5.17 - 8.95)	0.008	5.12 (2.89 - 9.08)	8.97 (4.69 - 17.17)	0.04	-3.74 (-11.67 - 4.18)	0.36	
Adipose-IR Index	36.6 (16.9 - 79.6)	55.8 (34.9 - 89.0)	0.22	51.7 (20.4 - 73.4)	103.2 (39.6 - 268.8)	0.02	-87.8 (-222.1 - 46.4)	0.21	
Free Fatty Acid (mmol/L)	2.88 (2.41 - 3.45)	2.44 (1.94 - 3.09)	0.21	3.03 (2.42 - 3.83)	2.99 (2.34 - 3.81)	0.87	-0.37 (-1.18 - 0.44)	0.37	
CETP (µg/mL)	2.49 (0.82)	2.38 (0.72)	0.46	2.82 (0.23)	2.91 (0.61)	0.59	-0.21 (-0.62 - 0.20)	0.32	
CRP (mg/L)	0.96 (0.28 - 2.00)	1.60 (1.48 - 3.57)	0.13	1.32 (0.29 - 3.17)	0.71 (0.39 - 1.11)	0.25	2.27 (-0.63 - 5.18)	0.13	
Total Cholesterol (mg/dL)	4.07 (0.17)	4.23 (0.33)	0.20	4.77 (0.68)	4.70 (0.90)	0.69	0.23 (0.12 - 0.58)	0.21	
Triglyceride (mg/dL)	1.32 (0.28)	1.14 (0.46)	0.40	1.40 (0.46)	1.38 (0.41)	0.92	-0.16 (-0.67 - 0.35)	0.54	
HDL-C (mg/dL)	1.24 (0.25)	1.27 (0.28)	0.50	1.17 (0.26)	1.16 (0.20)	0.88	0.03 (0.07 - 0.13)	0.54	
LDL-C (mg/dL)	2.22 (0.31)	2.44 (0.28)	0.01	2.97 (0.63)	2.92 (0.74)	0.65	0.27 (0.05 - 0.49)	0.03	

All variables are presented as mean and its standard deviation, however, HOMA-IR, Fasting Insulin, Adipose-IR Index, Free Fatty Acid, and CRP levels are presented as geometric (95%CI). *The differences between before and after HFD intervention were analysed using paired t-test. **The difference in changes (before and after HFD) of different parameters between STH-infected and STH-uninfected were analysed using linear mixed model and are presented as [Estimated Differences in Changes (95%CI), p-value]. Abbreviation: HOMA-IR= homeostatic model assessment of insulin resistance, CETP= cholesteryl ester transfer protein, CRP= C-reactive protein, HDL-C= high-density lipoprotein cholesterol, LDL-C= low-density lipoprotein cholesterol.

Supplementary Methods

Statistical Analysis

In the cross-sectional study, sample size was calculated to aim at a difference in HOMA-IR between urban and rural group of 0.5. The SD of HOMA-IR from previous study was 0.84. We used a significance level of 5% and a power of 80%, thus we needed at least 45 subjects for each group. For the interventional study, sample size was calculated to aim at a difference in changes of HOMA-IR between urban and rural group of 0.70. The SD of the HOMA-IR changes after HFD intervention from previous study was 0.68.2 We used a significance level of 5% and a power of 80%, thus we needed at least 15 subjects per group or 30 subjects in total. Next, to assess STH effect on the metabolic response upon HFD intervention we used similar calculation, aiming at having at least 15 subjects per group.

For the cross-sectional study, we further stratified the urban and rural group based on their STH infection status. However, as the number of urban subjects with STH infections was very low and therefore was excluded from analysis, eventually we had three groups: rural subjects with STH infections, rural subjects without STH infections, and urban subjects without STH infections. We calculated variance inflation factors (VIFs) to check multicollinearity in our regression models and VIF values below 4 were considered appropriate. Due to multicollinearity between BMI and WC, we used WC as clinical marker for adiposity. Analyses were performed using IBM Statistics 23.

For the HFD intervention study, to compare the parameter before and after the HFD intervention for each group, whenever appropriate, paired t-test or Wilcoxon-signed ranked test was performed. A mixed model was applied to assess mean differences before and after intervention between group. Groups were modelled as fixed effects, and to model correlation within subjects, random-specific intercept was used.