



Universiteit  
Leiden  
The Netherlands

## **Immunity in atherosclerosis: novel assays, biomarkers and therapeutic approaches**

Grievink, H.W.

### **Citation**

Grievink, H. W. (2023, October 26). *Immunity in atherosclerosis: novel assays, biomarkers and therapeutic approaches*. Retrieved from <https://hdl.handle.net/1887/3655349>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3655349>

**Note:** To cite this publication please use the final published version (if applicable).



CHAPTER 7

**Immunosuppression  
by hydroxychloroquine:  
mechanistic proof in *in vitro*  
experiments but limited systemic  
activity in a randomized placebo**

*Published in Immunol. Res. 2023 Feb 22.*

*Doi: 10.1007/s12026-023-09367-3*

Aliede E. in 't Veld<sup>1,2\*</sup>, Hendrika W. Grievink<sup>1,3\*</sup>,  
Johan L. van der Plas<sup>1,2</sup>, Boukje C. Eveleens Maarse<sup>1,2</sup>,  
Sebastiaan J.W. van Kraaij<sup>1,2</sup>, Tess D. Woutman<sup>1</sup>,  
Mascha Schoonakker<sup>1</sup>, Naomi B. Klarenbeek<sup>1</sup>,  
Marieke L. de Kam<sup>1</sup>, Ingrid M.C. Kamerling<sup>1,2</sup>,  
Manon A.A. Jansen<sup>1</sup>, Matthijs Moerland<sup>1,2</sup>

*\* Equal contribution*

1. Centre for Human Drug Research, Leiden, NL

2. Leiden University Medical Centre, Leiden, NL

3. Division of BioTherapeutics, Leiden Academic Center  
for Drug Research, Leiden University, Leiden, NL

## Graphical abstract




Figure is created using biorender.com.

## Abstract

Based on its wide range of immunosuppressive properties, hydroxychloroquine (HCQ) is used for the treatment of several autoimmune diseases. Limited literature is available on the relationship between HCQ concentration and its immunosuppressive effect. To gain insight in this relationship we performed *in vitro* experiments in human PBMCs and explored the effect of HCQ on T and B cell proliferation and Toll like receptor (TLR)<sub>3</sub>/TLR<sub>7</sub>/TLR<sub>9</sub>/RIG-I-induced cytokine production. In a placebo-controlled clinical study these same endpoints were evaluated in healthy volunteers that were treated with a cumulative dose of 2400 mg HCQ over 5 days. *In vitro*, HCQ inhibited TLR responses with IC<sub>50</sub>s >100 ng/mL and reaching 100% inhibition. In the clinical study, maximal HCQ plasma concentrations ranged from 75 to 200 ng/mL. No *ex vivo* HCQ effects were found on RIG-I-mediated cytokine release, but there was significant suppression of TLR7 responses and mild suppression of TLR3 and TLR9 responses. Moreover, HCQ treatment did not affect B cell and T cell proliferation. These investigations show that HCQ has clear immunosuppressive effects on human PBMCs, but the effective concentrations exceed the circulating HCQ concentrations under conventional clinical use. Of note, based on HCQ's physico-chemical properties, tissue drug concentrations may be higher, potentially resulting in significant local immunosuppression.

This trial is registered in the International Clinical Trials Registry Platform (ICTRP) under study number NL8726



## Introduction

Hydroxychloroquine (HCQ) is a broad immunosuppressive drug, initially developed as an anti-malarial drug. However, due to its anti-inflammatory properties, HCQ is now widely used in the treatment of autoimmune diseases such as rheumatoid arthritis (RA),<sup>1</sup> systemic lupus erythematosus (SLE)<sup>2</sup> and Sjögren's syndrome.<sup>3</sup> The use of HCQ in other diseases has been under investigation, a pilot trial investigating the use of HCQ in patients after myocardial infarction showed a decrease in plasma IL-6 levels compared to placebo, and a larger trial studying the effect on recurrent cardiovascular events is currently ongoing.<sup>4</sup> Furthermore, HCQ was under investigation for use in moderate to severe COVID-19 patients during the COVID-19 pandemic.<sup>5</sup>

The exact mechanisms behind HCQ's immunosuppressive functions remain unclear. HCQ accumulates in the lysosomes and inhibits lysosomal function by autophagosome fusion with lysosomes,<sup>6</sup> thereby inhibiting antigen presentation.<sup>7,8</sup> In addition, HCQ inhibits pro-inflammatory cytokine production by myeloid cells, possibly via the inhibition of endosomal Toll-like receptor (TLR) signalling.<sup>9</sup> It has been shown that HCQ treatment is associated with decreased interferon (IFN) $\alpha$  serum levels in SLE patients.<sup>10</sup> Furthermore, several studies investigating the effect of HCQ on peripheral blood mononuclear cells (PBMCs) or cell lines show that HCQ treatment reduces phorbol 12-myristate 13-acetate (PMA) and ionomycin or lipopolysaccharide induced cytokine production.<sup>11-13</sup>

Besides effects on the innate immune system, HCQ affects the adaptive immune response as well. It has been shown that HCQ inhibits differentiation of class-switched memory B cells into plasmablasts and thereby decreases IgG production in response to TLR9 stimulation or inoculation with inactivated virus.<sup>14,15</sup> HCQ inhibits T cell activation as well, via the inhibition of T cell receptor induced calcium mobilization and dysregulation of mitochondrial superoxide production.<sup>16-18</sup>

However, the concentrations used in such *in vitro* experiments studying the immunomodulatory effects of HCQ largely exceeded obtainable clinical concentrations in patients. A study in cutaneous lupus erythematosus patients receiving HCQ in clinical doses showed that higher HCQ blood levels corresponded with lower *ex vivo* IFN $\alpha$  responses after TLR9

stimulation, but not after TLR7/8 stimulation.<sup>13</sup> Moreover, influenza antibody titers after vaccination in Sjögren's syndrome patients receiving HCQ were lower compared to HCQ naïve patients.<sup>15</sup> Unfortunately, little additional literature is available on the *in vivo* immunomodulatory effects of HCQ and comparing it to *in vitro* experiments.

We aimed to assess and quantify the immunomodulatory effects of HCQ on primary human immune cells, both *in vitro* and *ex vivo* in a randomized clinical trial. We assessed the effect of HCQ on cytokine production after endosomal TLR stimulation in isolated PBMCs and on T and B cell proliferation (*in vitro* as well as *ex vivo*). In the clinical trial, healthy subjects were dosed with HCQ in the standard dosing regimen for moderate-to-severe COVID-19 that was advised in the Netherlands when the study was conceived. In the study design, we accounted for a potential age effect on the study outcomes, since general immunocompetence and drug metabolism has been reported to be age-dependent.<sup>19,20</sup> Here we present the outcomes of the *in vitro* experiment and the randomized clinical trial.

## Methods

### IN VITRO EXPERIMENTS

Blood was collected by venipuncture using Sodium Heparin vacutainer tubes or Cell Preparation Tubes (CPT, Becton Dickinson, Franklin Lakes, NJ, USA) from healthy volunteers after written informed consent, in accordance with Good Clinical Practice guidelines and the Declaration of Helsinki. Blood was used for the evaluation of the *in vitro* immunomodulatory activity of hydroxychloroquine (10 - 10,000 ng/mL, Sigma-Aldrich, Deisenhofen, Germany). All experiments were started within one hour after blood withdrawal, and incubations were performed in duplicate. Hydroxychloroquine and stimulant were added simultaneously. Per experiment, blood of 6 donors was used.

### CLINICAL STUDY

We conducted a single-blind, randomized, placebo-controlled multiple dose study in forty healthy male volunteers, comprising twenty young (18-30 years) and twenty elderly (65-75 years) subjects. The study was conducted at the Centre for Human Drug Research in Leiden, The Netherlands,

between June and September 2020, during the COVID-19 pandemic. All subjects in the clinical trial gave written informed consent according to Declaration of Helsinki recommendations, prior to any study-related activity. The study was approved by the Independent Ethics Committee of the Foundation 'Evaluation of Ethics in Biomedical Research' (Stichting Beoordeling Ethiek Biomedisch Onderzoek, Assen, The Netherlands) and registered in the Toetsingonline Registry (study number NL73816.056.20), and in the International Clinical Trials Registry Platform (NL8726).

### **VOLUNTEER SELECTION**

To avoid sex-related inter-individual variability in immune responses, only male subjects were included.<sup>21</sup> Subjects were included if they were overtly healthy. The health status of subjects was assessed by medical screening, including medical history, physical examination, vital signs measurements, 12-lead electrocardiography (ECG), urine analysis, drug screen and safety chemistry, coagulation, and hematology blood sampling. BMI of study participants had to be between 18 and 32 kg/m<sup>2</sup>. Subjects with a known hypersensitivity reaction to chloroquine, HCQ or other 4-aminoquinolines, abnormalities in the resting ECG (including QTcF-interval >450ms), evidence of any active or chronic disease or condition (including long QT syndrome, retinal disease, G6PD deficiency, autoimmune diseases, diabetes mellitus type I or II, psychiatric disorders) or a positive SARS-COV-2 PCR test were excluded from study participation. Use of concomitant medication was not permitted during the study, and 14 days (or 5 half-lives) prior to the study drug administration, except for paracetamol.

### **STUDY DESIGN**

Subjects were randomized to receive either hydroxychloroquine sulphate (plaquenil®) or placebo tablets, in a 1:1 ratio. Tablets were dispensed by the pharmacy, according to a randomization list generated by a study-independent statistician. Plaquenil® and placebo tablets were packaged in the same way but the tablets were not indistinguishable, study drug administration was therefore performed by dedicated unblinded personnel not involved in any other study tasks. Subjects received HCQ or placebo by a loading dose of 400 mg twice daily (t=0h and t=12h) followed by a 400 mg once daily dose regimen (t=24h, t=48h, t=72h, and t=96h), giving a cumulative dose of

2400 mg. This reflected the standard dosing regimen for moderate-to-severe COVID-19 patients in the Netherlands when the study was conceived (total dose between 2000 and 3800 mg).

### **PHARMACOKINETIC EVALUATION**

For pharmacokinetic (PK) assessments, blood was collected in 3 mL Vacutainer® K<sub>2</sub>EDTA tubes (Becton Dickinson) on study day 0 (baseline and 3 hours post-dosing), and day 1, 4 and 9 (3 hours post-dosing). Hydroxychloroquine plasma concentrations were measured by Ardena Bioanalytical Laboratory (Assen, the Netherlands) using a validated LC-MS/MS method. The lower limit of quantification (LLOQ) of the analysis was 5 ng/mL.

### **WHOLE BLOOD STIMULATION**

Whole blood was stimulated with 10 µg/mL phytohemagglutinin (PHA, Sigma-Aldrich) for 6 hours and 24 hours. After 6 hours, activation markers on T-cells were measured using CD69-APC (clone: REA824), CD71-FITC (clone: REA902), CD154-VIOBLUE (REA238) and CD25-PE (clone: 3G10), CD3-VioGreen (REA613), CD4-APC-VIO770 (REA623) and CD8-PE-VIO770 (REA734) antibodies and propidium iodide as viability dye (all Miltenyi Biotec, Bergisch-Gladbach, Germany) using a MACSQuant 16 analyzer (Miltenyi Biotec). After 24 hours, culture supernatants were collected for cytokine analysis.

### **PBMC ISOLATION AND TLR STIMULATION**

PBMCs were isolated from CPT after centrifugation at 1800 x g for 30 minutes, and washed 2x using phosphate buffered saline (PBS, pH 7.2, Gibco, Thermo Fisher, Waltham, MA, USA). PBMCs were stimulated with endosomal TLR ligands PolyI:C (TLR3, 50 µg/mL), imiquimod (TLR7, 1 µg/mL), CpG class A (TLR9, oligodeoxynucleotides [ODN] 2.5 µM) and POLYI:C/Iyovec (RIG-I, 1 µg/mL; all Invivogen, Toulouse, France). Supernatants were collected after 24 hours for cytokine quantification.

### **PROLIFERATION ASSAY**

PBMCs were stained with 2.5µM cell trace violet (CTV, Thermo Fisher) according to user's manual. T cells were stimulated with 5 µg/mL phytohemagglutinin (PHA), and B cells with a monoclonal CD40 antibody (5 µg/mL; clone: G28.5, BioXCell) and cpG class B (2.5 µM; ODN Invivogen). After 5 days

of stimulation PBMCs were stained using, CD4-PE (clone: OKT4), CD8-APC (clone: HIT8A), CD19-PE (clone: HIB19, all Biolegend, San Diego, CA, USA) and fixable viability dye eFluor780 (Thermo Fisher) and proliferation was quantified by flow cytometry, using the MACSQuant 16 analyzer.

### FLOW CYTOMETRY

Circulating leukocyte subsets were analyzed using flow cytometry. Red blood cell lysis was performed on sodium heparinized blood using RBC lysis buffer (Thermo Fisher Scientific). After washing with PBS (PH 7.2), leukocytes were incubated with fluorochrome-labeled antibodies for 30 minutes on ice. After a final washing step, leukocytes were measured on a MACSQuant 16 analyzer (Miltenyi Biotec). See supplemental table I for a full list of antibodies used.

### CYTOKINE MEASUREMENTS

IFN $\gamma$  and IL-2 were quantified using the Vplex-2 kit (Meso Scale Discovery). IFN $\alpha$  and IL-6 were quantified using the pan-specific IFN $\alpha$  ELISAPRO HRP kit and the IL-6 ELISAPRO HRP kit (both Mabtech, Nacka Strand, Sweden).

### STATISTICAL ANALYSIS

*In vitro* data are reported as mean  $\pm$  standard deviation (SD). The IC<sub>50</sub> was calculated using a inhibitory sigmoid Emax function where applicable. Analyses were performed using Graphpad Prism version 6.05 (Graphpad, San Diego, CA, USA).

Repeatedly measured pharmacodynamic data were evaluated with a mixed model analysis of variance with fixed factors treatment, age group, time, treatment by time, age group by time, treatment by age group and treatment by age group by time and a random factor subject and the average pre-value as covariate. If needed, variables were log transformed before analysis. Contrasts between the placebo and HCQ treatment groups were calculated per endpoint. In addition, a potential age-specific HCQ effect was evaluated by comparing the 18-30 years with the 65-75 years age group. For the contrasts, an estimate of the difference (back-transformed in percentage for log transformed parameters), a 95% confidence interval (in percentage for log-transformed parameters), Least Square Means (geometric means for log transformed parameters), and the p-value were calculated. A P-value  $\leq$  0.05 was considered to be statistically significant.

All calculations were performed using SAS for windows V9.4 (SAS Institute, Inc., Cary, NC, USA).

## Results

### HYDROXYCHLOROQUINE SUPPRESSED ENDOSOMAL TLR-INDUCED IFN $\alpha$ AND IL-6 RELEASE IN VITRO

PBMCs were stimulated with endosomal TLR ligands in the presence of a dose range of HCQ for 24 hours, and supernatants were analyzed for IRF-mediated IFN $\alpha$  and for NF $\kappa$ B-mediated IL-6 secretion. PBMCs were stimulated with different endosomal TLR ligands: POLY I:C (TLR3), imiquimod (TLR7), CpG class A (TLR9) and POLY I:C lyovec (RIG-I). HCQ dose-dependently inhibited endosomal TLR-induced IFN $\alpha$  and IL-6 secretion (Figure 1). POLY I:C-induced IFN $\alpha$  and IL-6 release was strongly suppressed at 10.000 ng/mL (IFN $\alpha$ : -83.9%, IL-6: -96.6%, IC<sub>50</sub> IL-6=637.2 ng/mL). Imiquimod (IMQ)-induced cytokine release was completely suppressed at the highest concentration (IFN $\alpha$ : -96.3%, IL-6: -96.3%, IC<sub>50</sub> IFN $\alpha$ : 695.8 ng/mL, IL-6: 237.9 ng/mL). The same was observed for stimulation with cpG class A, IFN $\alpha$  was suppressed by 99.6% with an IC<sub>50</sub> of 145.3 ng/mL, and IL-6 was suppressed by 96.4%, with an IC<sub>50</sub> of 86.9 ng/mL. The RIG-I response to POLY I:C/lyovec was less affected by HCQ, while IFN $\alpha$  release was suppressed by 66.1% at 10,000 ng/mL HCQ, IL-6 release was not significantly altered.

### HCQ INHIBITED B CELL PROLIFERATION BUT NOT T CELL PROLIFERATION IN VITRO

PBMCs were stimulated with phytohaemagglutinin (PHA) or monoclonal anti-CD40 with cpG-B to induce T cell and B cell proliferation respectively, in the presence of a dose range of HCQ. No effect of HCQ was seen on T cell proliferation (Figure 2A). Also, no effects were observed on T cell activation markers after PHA stimulation for 6 hours (Figure S1). At HCQ concentrations  $>$ 100 ng/mL, a decrease in B cell proliferation was observed, with an IC<sub>50</sub> of 1138 ng/mL (Figure 2B).

### CLINICAL STUDY

#### *Demographics and safety*

Of the 40 enrolled and randomized healthy subjects, 20 received a cumulative dose of 2400 mg HCQ in 5 days and 20 received placebo (Figure 3). The

different age groups (18–30 and 65–75 years) were of equal size. Baseline characteristics are described in Table 1. All subjects completed their study treatment. One subject in the 65–75 years group erroneously took an additional 400 mg dose of HCQ on study day 2, after which the subject received 400 mg doses (once daily) for two consecutive days to not exceed the cumulative dose of 2400 mg.

Treatment-emergent adverse events were transient, of mild severity and did not lead to study discontinuation. Adverse events were reported more often by subjects in the active treatment arm (50%) compared to placebo (35%). Gastrointestinal complaints (20%) and dizziness (15%) were the most frequently reported adverse events in the active group. There were no findings of clinical concern following assessments of urinalysis, hematology and chemistry laboratory tests, vital signs, physical examination and ECGs.<sup>22</sup>

### Pharmacokinetics

Mean HCQ concentration time profiles in plasma are depicted in Figure 4A. Individual concentration profiles have been published previously.<sup>22</sup> There were no significant differences in HCQ exposures between age groups (Figure 4B). Mean concentrations measured 27 hours after starting the treatment course (day 1, 121.0 ± 40.54 ng/mL) were in a similar range to those measured on the last day of the treatment course (day 4, 109.2 ± 35.59 ng/mL).

## PHARMACODYNAMICS

### Hydroxychloroquine did not affect circulating immune cells

The effects of HCQ on different circulating cell populations, both absolute as relative, were evaluated using flow cytometry. No apparent effects were seen on absolute values of total leukocytes, lymphocytes, monocytes or neutrophils (Table S2), as well as CD14<sup>+</sup> monocytes, CD19<sup>+</sup> B cells, CD3<sup>+</sup> T cells, CD4<sup>+</sup> T cells and CD8<sup>+</sup> T cells (Table S3). Furthermore, no effects were seen on relative T cell populations (CD3<sup>+</sup>) in general, nor on subpopulations of T helper cells (CD4<sup>+</sup>), cytotoxic T cells (CD8<sup>+</sup>), and regulatory T cells (CD4<sup>+</sup>CD25<sup>+</sup>CD127<sup>-</sup>). Similarly, no apparent treatment effects were observed in natural killer cells (CD56<sup>+</sup>), B cells (CD19<sup>+</sup>) and subpopulations of regulatory (CD5<sup>+</sup>CD1d<sup>hi</sup>), transitional (CD24<sup>hi</sup>CD38<sup>hi</sup>) and antibody secreting B cells (CD27<sup>+</sup>CD38<sup>+</sup>). Moreover, also in classical (CD14<sup>+</sup>), non-classical (CD16<sup>+</sup>) and intermediate (CD14<sup>+</sup>CD16<sup>+</sup>) monocytes and plasmacytoid dendritic cells

(pDCs, HLA-DR<sup>+</sup>CD14<sup>+</sup>CD16<sup>+</sup>CD123<sup>+</sup>) no differences were found between treatment groups. Also, between both age groups, no evident HCQ effects were observed (Table S3).

### In vivo hydroxychloroquine suppressed IFN $\alpha$ secretion following TLR7 stimulation, but not after TLR3, TLR9 or RIG-I-like receptor stimulation

To study the effects of HCQ on TLR/RIG-I-mediated IRF activation, PBMCs were stimulated with different endosomal TLR ligands: POLY I:C (TLR3), imiquimod (TLR7), CpG class A (TLR9) and POLY I:C Iyovec (RIG-I). Overall, no HCQ effect was observed on IFN $\alpha$  responses (Figure 5), except for a significant suppression of IMQ-driven IFN $\alpha$  production (inhibition of -48.2%, CI95 -72.1%– -4.0%,  $p=0.038$ ). POLY I:C-driven IFN $\alpha$  release also appeared to be suppressed by HCQ, but not significantly (inhibition -34.2%, CI95 -57.7%– -7.5%,  $p=0.091$ ). No differences in HCQ effect on IFN $\alpha$  responses were observed between the young and elderly population (Figure S3).

### In vivo hydroxychloroquine significantly suppressed IL-6 secretion after TLR7 stimulation, but not following TLR3, TLR9 or RIG-I-like receptor stimulation

Activation of NF $\kappa$ B signaling via endosomal TLR and RIG-I-like ligands was assessed by measuring downstream IL-6 production (Figure 6). HCQ significantly suppressed IMQ-driven IL-6 production (inhibition of -71.3%, CI95 -84.7%– -46.1%,  $p=0.0005$ ). No significant HCQ effects were observed on IL-6 production driven by CpG A (TLR9) and POLY I:C (TLR3) stimulations (inhibition of -35.9%, CI95 -60.3%– -3.6%,  $P=0.068$  and -37.7%, CI95 -62.6%– -3.7%,  $P=0.067$ , respectively). No differences in HCQ effect on IL-6 responses were observed between the young and elderly population (Figure S3).

### In vivo hydroxychloroquine did not alter T cell activation

To further investigate the potential immunomodulatory effect of HCQ on T cell activation, whole blood samples were incubated with PHA, which is known to induce a general T cell response.<sup>23</sup> HCQ treatment did not modulate expression of T cell activation markers (CD25, CD69, CD71, CD154) following PHA-stimulation (Figure S3). In addition, PHA-induced secretion of IL-2 and IFN $\gamma$  was assessed, no apparent differences were observed between HCQ and placebo (Figure S4).



### Hydroxychloroquine did not alter *ex vivo* B and T cell proliferation after *in vivo* administration

Proliferative capability of B cells was assessed by stimulating PBMCs *ex vivo* with anti-CD40 mAb + CpG B ODN, a known stimulus for human B cell activation.<sup>24</sup> Following stimulation of PBMCs, the percentage of proliferative B cells in the HCQ-treated group was similar to that of the placebo group (70.47% at day 4 for placebo, 70.03% for HCQ) (Figure 7). In addition, PBMCs were stimulated with PHA to induce T helper cells (CD4<sup>+</sup>) and cytotoxic T cells (CD8<sup>+</sup>) proliferation. Proliferation of both CD4<sup>+</sup> and CD8<sup>+</sup> cells was comparable between the HCQ- and placebo-treated group (>95% for both groups for all time points for CD4, >92% for both groups for all time points for CD8). No differences were observed for B and T cell proliferation in the separate age groups (Figure S5).

## Discussion

Although HCQ is widely used for the treatment of autoimmune diseases, the exact mechanism behind its immunomodulatory properties remains unclear. In this study we therefore aimed to quantify the immunosuppressive effect of HCQ by studying the endosomal TLR response and lymphocyte proliferation and activation both in *in vitro* experiments and *in vivo* in a randomized placebo-controlled trial in healthy volunteers.

In our *in vitro* experiments, HCQ dose-dependently inhibited TLR3-, 7- and 9-driven IL-6 and IFN $\alpha$  production, with profound effects at concentrations >100 ng/mL. These findings are in line with literature on TLR signaling modulation by chloroquine.<sup>9,25</sup> Limited data are available on the immunomodulatory effect of HCQ/chloroquine on RIG-I signaling.<sup>26</sup> RIG-I functions as a cytosolic sensor of nucleic acids, inducing a type I IFN response after activation. HCQ inhibited the IFN responses in THP-1 cells transfected with RIG-I ligands,<sup>27</sup> but this effect was not confirmed in cultures of human bronchial smooth muscle and epithelial cells.<sup>28,29</sup> This is in line with the observations in the current study, which shows that HCQ only mildly modulated RIG-I-mediated IFN $\alpha$  production in PBMCs, without affecting IL-6 release. Our results suggest that HCQ has a profound effect on endo-lysosomal TLR functioning *in vitro* but affects the cytosolic RIG-I-mediated pathway to a lesser degree. This could be explained by HCQ's excessive affinity to the lysosomal intracellular compartment (expected to be 56,000-fold higher than cytosol).<sup>30</sup>

HCQ did not affect T cell activation *in vitro*. Although a dose-dependent inhibition of T cell proliferation by chloroquine following stimulation with anti-CD3/CD28 has been described,<sup>31-33</sup> we did not see any inhibitory effect of HCQ on T cell proliferation or expression of activation markers in our *in vitro* experiments. This may be explained by the fact that a different and more potent stimulus was used in this study (PHA), which might be more difficult to suppress. For B cell proliferation, on the other hand, a dose-dependent HCQ-mediated inhibition was observed *in vitro*, confirming previous research.<sup>34</sup> Although the HCQ-mediated inhibition was not as strong as the inhibition of cytokine production (IC<sub>50</sub> of 1138 ng/mL for B cell proliferation vs 145-696 ng/mL for cytokine production), at concentrations > 100 ng/mL a clear HCQ-mediated decrease in B cell proliferation was found.

While HCQ had strong immunosuppressive effects *in vitro*, especially at high concentrations, less pronounced *ex vivo* effects of the compound were observed in our clinical study. Compared to placebo, 5-day HCQ treatment did not significantly suppress B cell proliferation or *ex vivo* TLR-driven IFN $\alpha$  and IL-6 secretion in PBMC cultures, except for a suppressive effect on TLR7-driven responses. The most likely explanation for this discrepancy between *in vitro* and *ex vivo* is that there was insufficient drug exposure at the evaluated HCQ dose and regimen in the clinical study. By using a 5-day dose regimen of HCQ (the recommended off-label dose for COVID-19 at the time of study conduct), an average maximum plasma concentration of 121 ng/mL was reached. This concentration is considerably lower than plasma levels found in RA patients receiving HCQ treatment of 200 mg daily for a longer time period, which ranges from 200–500 ng/mL.<sup>35-37</sup> Peak exposures of 100-150 ng/mL from the clinical study translate into a maximal inhibitory effect of 20 to 50% in most cellular assays. In combination with the observed variability of the endpoints, such effects remain easily undetected. However, whole blood concentrations are expected to be approximately 2-to-7-fold higher than plasma concentrations due to intracellular uptake in blood components,<sup>38-40</sup> which would make the concentrations more in range with the *in vitro* experiments. Also, due to the large volume of distribution,<sup>39</sup> and the high HCQ tissue concentrations as compared to plasma,<sup>41,42</sup> immunosuppressive effects in specific tissues may be significant. Moreover, HCQ has a gradual onset of action for HCQ, and is biologically active even after drug discontinuation.<sup>8</sup> This would mean that the five-day treatment that was used in the current study is insufficient to detect *ex vivo* drug effects. Other studies, for



example investigating HCQ effect in HIV patients,<sup>43</sup> showed a discrepancy between plasma levels and drug efficacy.

The widespread use of hydroxychloroquine following the onset of the COVID-19 pandemic was the reason to initiate our experiments. The initial off-label use of HCQ was primarily based on studies that assessed *in vitro* antiviral activity against SARS-COV-2.<sup>44</sup> However, there is also a long-standing hypothesis that the immunomodulatory properties of chloroquine and HCQ could dampen immunopathology caused by viral infections such as influenza, Severe Acute Respiratory Syndrome (SARS), Middle East Respiratory Syndrome (MERS) and COVID-19 by suppressing the host immune response.<sup>45-47</sup> Use of HCQ in COVID-19 patients did not show evident favorable effects for clinical endpoints such as mortality and mechanical ventilation for both prophylaxis and treatment.<sup>48</sup> Our study provides mechanistic insight in the immuno-modulatory effects of a HCQ dosing regimen that was used to treat COVID-19. We found that a 5-day treatment course of HCQ did not have extensive immuno-modulatory effect in healthy individuals. HCQ treatment only significantly inhibited TLR7 responses. In theory, inhibition of the TLR7-mediated innate response to viral agents may be disadvantageous during the initial stages of viral infection.<sup>49,50</sup> However, recent COVID-19 trials did not show an effect of HCQ treatment on disease incidence, and long-term HCQ use in rheumatoid arthritis is not associated with higher incidence of upper respiratory tract infections.<sup>51,52</sup>


In conclusion, we showed extensive and profound immunomodulation by HCQ *in vitro*, however in a clinical study in healthy volunteers, the overall immunomodulatory effects of a 5-day HCQ treatment regimen of 2400 mg were limited. The pharmacological activity of HCQ in autoimmunity remains to be studied in greater detail, based on the assays as presented in our studies and at a therapeutic dose and regimen relevant for the condition of interest.

**Table 1 Baseline characteristics.**


	Hydroxychloroquine		Placebo	
	Age group 18-30 yrs (n=10)	Age group 65-75 yrs (n=10)	Age group 18-30 yrs (n=10)	Age group 65-75 yrs (n=10)
Age, median (range)	23 (20-26)	68 (65-70)	23 (18-25)	68 (65-71)
BMI, mean (SD)	21.8 (1.5)	25.8 (2.0)	24.4 (1.9)	24.2 (3.0)
Race or ethnicity,* n (%)				
White	10 (100)	10 (100)	10 (100)	10 (100)
Other	0 (0)	0 (0)	0 (0)	0 (0)

\*Self-reported race or ethnicity of subjects. BMI=body mass index; SD=standard deviation.


**Figure 1** HCQ dose-dependently inhibited endosomal TLR induced IFN $\alpha$  and IL-6 release in vitro. PBMCs were stimulated with 50  $\mu$ g/mL POLY I:C (TLR3), 1  $\mu$ g/mL IMQ (TLR7), 2.5  $\mu$ M CpG-A (TLR9) or 1  $\mu$ g/mL POLY I:C/Iyovec (RIG-I) for 24 hours in the presence of a dose range of HCQ. IFN $\alpha$  and IL-6 release were measured by ELISA. The mean  $\pm$  SD of the change from baseline of 6 subjects is shown. The IC<sub>50</sub> was calculated using a four-parameter non-linear regression fit where applicable.



**Figure 2** HCQ dose-dependently inhibited B cell, but not T cell proliferation in vitro. PBMCs from 6 healthy donors were stained with CFV and stimulated for 5 days with 5  $\mu$ g/ml PHA for T cell proliferation (A), or 5  $\mu$ g/mL anti-CD40 MAB + 2.5  $\mu$ M CpG B for B cell proliferation (B). Proliferation was measured by flow cytometry. The mean  $\pm$  SD of the change from baseline are shown. The IC<sub>50</sub> was calculated using a four-parameter non-linear regression fit where applicable.




**Figure 3 Trial flow chart (CONSORT diagram).**




\* Drug concentrations were only analyzed in the active treatment group.


**Figure 4 Pharmacokinetic profile of HCQ.** Mean and standard deviation of hydroxychloroquine plasma concentrations for HCQ treatment group (A), and split for young and elderly volunteers (B). Dotted vertical lines indicate timing of HCQ dosing (0, 12, 24, 48, 72, 96 hrs).



**Figure 5** In vivo HCQ inhibited IMQ-induced IFN $\alpha$  release, but not TLR3, TLR9 and RIG-I. PBMCs were stimulated with 50  $\mu$ g/mL POLY I:C (TLR3), 1  $\mu$ g/mL IMQ (TLR7), 2.5 $\mu$ M CpG A (TLR9) or 1  $\mu$ g/mL POLY I:C/lyovec (RIG-I) at 0, 12, 24, 48, 72 and 92 hours after primary HCQ dosing. IFN $\alpha$  release was measured by ELISA. Data is shown as mean + SD as one-sided error bars. Dotted vertical lines indicate HCQ dosing times.




**Figure 6** In vivo HCQ inhibited IMQ-induced IL-6 release, but not TLR3, TLR9 and RIG-I. PBMCs were stimulated with 50  $\mu$ g/mL POLY I:C (TLR3), 1  $\mu$ g/mL IMQ (TLR7), 2.5 $\mu$ M CpG A (TLR9) or 1  $\mu$ g/mL POLY I:C/lyovec (RIG-I) at 0, 12, 24, 48, 72 and 92 hours after primary HCQ dosing. IFN $\alpha$  release was measured by ELISA. Data is shown as mean + SD as one-sided error bars. Dotted vertical lines indicate HCQ dosing times.





**Figure 7 In vivo HCQ did not affect T and B cell proliferation.** PBMCs were stained with CFV and stimulated for 5 days with 5µg/ml PHA for T cell proliferation (A), or 5 µg/mL anti-CD40 mAb + 2.5 µM CpG B for B cell proliferation (B). Proliferation was measured by flow cytometry. The mean ± SD are shown. Dotted vertical lines indicate HCQ dosing times.



All mentioned supplementary figures and tables in this chapter can be found on the corresponding website by scanning this QR code.

## REFERENCES

- Rempnault, C., et al., Clinical and Structural Efficacy of Hydroxychloroquine in Rheumatoid Arthritis: A Systematic Review. *Arthritis Care & Research*, 2020. 72(1): p. 36-40.
- Ponticelli, C. and G. Moroni, Hydroxychloroquine in systemic lupus erythematosus (SLE). *Expert Opin Drug Saf*, 2017. 16(3): p. 411-419.
- Wang, X., et al., The Efficiency of Hydroxychloroquine for the Treatment of Primary Sjögren's Syndrome: A Systematic Review and Meta-Analysis. *Frontiers in Pharmacology*, 2021. 12.
- Ulander, L., et al., Hydroxychloroquine reduces interleukin-6 levels after myocardial infarction: The randomized, double-blind, placebo-controlled OXI pilot trial. *Int J Cardiol*, 2021. 337: p. 21-27.
- Axfors, C., et al., Mortality outcomes with hydroxychloroquine and chloroquine in COVID-19 from an international collaborative meta-analysis of randomized trials. *Nat Commun*, 2021. 12(1): p. 2349.
- Mauthe, M., et al., Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome fusion. *Autophagy*, 2018. 14(8): p. 1435-1455.
- Lotteau, V., et al., Intracellular transport of class II MHC molecules directed by invariant chain. *Nature*, 1990. 348(6302): p. 600-5.
- Schrezenmeier, E. and T. Dörner, Mechanisms of action of hydroxychloroquine and chloroquine: implications for rheumatology. *Nature Reviews Rheumatology*, 2020. 16(3): p. 155-166.
- Kuznik, A., et al., Mechanism of endosomal TLR inhibition by antimalarial drugs and imidazoquinolines. *J Immunol*, 2011. 186(8): p. 4794-804.
- Willis, R., et al., Effect of hydroxychloroquine treatment on pro-inflammatory cytokines and disease activity in SLE patients: data from LUMINA (LXXV), a multiethnic US cohort. *Lupus*, 2012. 21(8): p. 830-5.
- Jang, C.-H., et al., Chloroquine inhibits production of TNF- $\alpha$ , IL-1 $\beta$  and IL-6 from lipopolysaccharide-stimulated human monocytes/macrophages by different modes. *Rheumatology*, 2006. 45(6): p. 703-710.
- Silva, J.C.d., et al., Hydroxychloroquine decreases Th17-related cytokines in systemic lupus erythematosus and rheumatoid arthritis patients. *Clinics (Sao Paulo, Brazil)*, 2013. 68(6): p. 766-771.
- Gardet, A., et al., Effect of in vivo Hydroxychloroquine and ex vivo Anti-BDCA2 mAb Treatment on pDC IFN $\alpha$  Production From Patients Affected With Cutaneous Lupus Erythematosus. *Front Immunol*, 2019. 10: p. 275.
- Torigoe, M., et al., Hydroxychloroquine efficiently suppresses inflammatory responses of human class-switched memory B cells via Toll-like receptor 9 inhibition. *Clin Immunol*, 2018. 195: p. 1-7.
- Brauner, S., et al., H1N1 vaccination in Sjögren's syndrome triggers polyclonal B cell activation and promotes autoantibody production. *Ann Rheum Dis*, 2017. 76(10): p. 1755-1763.
- Goldman, F.D., et al., Hydroxychloroquine inhibits calcium signals in T cells: a new mechanism to explain its immunomodulatory properties. *Blood*, 2000. 95(11): p. 3460-6.
- Wu, S.F., et al., Hydroxychloroquine inhibits CD154 expression in CD4(+) T lymphocytes of systemic lupus erythematosus through NFAT, but not STAT5, signaling. *Arthritis Res Ther*, 2017. 19(1): p. 183.
- Kim, M.L., et al., Hydroxychloroquine inhibits the mitochondrial antioxidant system in activated T cells. *iScience*, 2021. 24(12): p. 103509.
- Le Couteur, D.G., A.J. McLachlan, and R. de Cabo, Aging, Drugs, and Drug Metabolism. *The Journals of Gerontology: Series A*, 2011. 67A(2): p. 137-139.
- Mangoni, A.A. and S.H.D. Jackson, Age-related changes in pharmacokinetics and pharmacodynamics: basic principles and practical applications. *British journal of clinical pharmacology*, 2004. 57(1): p. 6-14.
- Klein, S.L. and K.L. Flanagan, Sex differences in immune responses. *Nature Reviews Immunology*, 2016. 16(10): p. 626-638.
- Eveleens Maarse, B.C., et al., Effect of hydroxychloroquine on the cardiac ventricular repolarization: A randomized clinical trial. *Br J Clin Pharmacol*, 2021.
- Nowell, P.C., Phytohemagglutinin: an initiator of mitosis in cultures of normal human leukocytes. *Cancer Res*, 1960. 20: p. 462-6.
- Carpenter, E.L., et al., Activation of human B cells by the agonist CD40 antibody CP-870,893 and augmentation with simultaneous toll-like receptor 9 stimulation. *J Transl Med*, 2009. 7: p. 93.
- Macfarlane, D.E. and L. Manzel, Antagonism of Immunostimulatory CpG-Oligodeoxynucleotides by Quinacrine, Chloroquine, and Structurally Related Compounds. *The Journal of Immunology*, 1998. 160(3): p. 1122-1131.
- Gies, V., et al., Beyond Anti-viral Effects of Chloroquine/Hydroxychloroquine. *Front Immunol*, 2020. 11: p. 1409.
- An, J., et al., Cutting edge: Antimalarial drugs inhibit IFN-beta production through blockade of cyclic GMP-AMP synthase-DNA interaction. *J Immunol*, 2015. 194(9): p. 4089-93.
- Calvén, J., et al., Viral stimuli trigger exaggerated thymic stromal lymphopoietin expression by chronic obstructive pulmonary disease epithelium: role of endosomal TLR3 and cytosolic RIG-I-like helicases. *J Innate Immun*, 2012. 4(1): p. 86-99.
- Calvén, J., Y. Yudina, and L. Uller, Rhinovirus and dsRNA induce RIG-I-like receptors and expression of interferon  $\beta$  and  $\lambda$ 1 in human bronchial smooth muscle cells. *PLoS One*, 2013. 8(4): p. e62718.
- Derendorf, H., Excessive lysosomal ion-trapping of hydroxychloroquine and azithromycin. *International journal of antimicrobial agents*, 2020. 55(6): p. 106007-106007.
- Schmidt, R.L., et al., Chloroquine inhibits human CD4(+) T-cell activation by AP-1 signaling modulation. *Sci Rep*, 2017. 7: p. 42191.

- 32 Landewe, R.B., et al., Chloroquine inhibits T cell proliferation by interfering with IL-2 production and responsiveness. *Clin Exp Immunol*, 1995. 102(1): p. 144-51.
- 33 Muschalek R, J.P., Kerman R, Kueht M, Etheridge W, Cusick M. Chloroquine's Effect on T cell Proliferation [abstract]. 2019 Accessed January 12, 2022.; 19 (suppl 3):[Available from: <https://atcmeetingabstracts.com/abstract/chloroquines-effect-on-t-cell-proliferation/>].
- 34 Ma, X., et al., Chloroquine Suppresses Effector B-Cell Functions and Has Differential Impact on Regulatory B-Cell Subsets. *Front Immunol*, 2022. 13: p. 818704.
- 35 Walbi, I.A., et al., The relationship between hydroxychloroquine plasma concentration and COVID-19 outcomes in rheumatoid arthritis patients in Saudi Arabia. *Saudi Pharmaceutical Journal*, 2022.
- 36 Carmichael, S.J., B. Charles, and S.E. Tett, Population Pharmacokinetics of Hydroxychloroquine in Patients With Rheumatoid Arthritis. *Therapeutic Drug Monitoring*, 2003. 25(6): p. 671-681.
- 37 Tett, S.E., R.O. Day, and D.J. Cutler, Concentration-effect relationship of hydroxychloroquine in rheumatoid arthritis--a cross sectional study. *J Rheumatol*, 1993. 20(11): p. 1874-9.
- 38 Carlsson, H., et al., Measurement of hydroxychloroquine in blood from SLE patients using LC-HRMS-evaluation of whole blood, plasma, and serum as sample matrices. *Arthritis Res Ther*, 2020. 22(1): p. 125.
- 39 Tett, S.E., et al., A dose-ranging study of the pharmacokinetics of hydroxy-chloroquine following intravenous administration to healthy volunteers. *Br J Clin Pharmacol*, 1988. 26(3): p. 303-13.
- 40 Chakrabarti, K. and W.J. McCune, Advances in the clinical use of hydroxychloroquine levels. *Current Opinion in Rheumatology*, 2022. 34(3).
- 41 Popert, A.J., Chloroquine: a review. *Rheumatol Rehabil*, 1976. 15(3): p. 235-8.
- 42 Liu, Q., et al., Time-Dependent Distribution of Hydroxychloroquine in Cynomolgus Macaques Using Population Pharmacokinetic Modeling Method. *Front Pharmacol*, 2020. 11: p. 602880.
- 43 Savarino, A. and I.L. Shtaj, Chloroquine and beyond: exploring anti-rheumatic drugs to reduce immune hyperactivation in HIV/AIDS. *Retrovirology*, 2015. 12: p. 51.
- 44 Liu, J., et al., Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov*, 2020. 6: p. 16.
- 45 Paton, N.I., et al., Chloroquine for influenza prevention: a randomised, double-blind, placebo controlled trial. *Lancet Infect Dis*, 2011. 11(9): p. 677-83.
- 46 Al-Bari, M.A.A., Targeting endosomal acidification by chloroquine analogs as a promising strategy for the treatment of emerging viral diseases. *Pharmacol Res Perspect*, 2017. 5(1): p. e00293.
- 47 Sinha, N. and G. Balayla, Hydroxychloroquine and COVID-19. *Postgrad Med J*, 2020. 96(1139): p. 550-555.
- 48 Singh, B., et al., Chloroquine or hydroxychloroquine for prevention and treatment of COVID-19. *Cochrane Database of systematic Reviews*, 2021(2).
- 49 Lund, J.M., et al., Recognition of single-stranded RNA viruses by Toll-like receptor 7. *Proc Natl Acad Sci U S A*, 2004. 101(15): p. 5598-603.
- 50 In 't Veld, A.E., et al., Hydroxychloroquine Effects on TLR Signalling: Underexposed but Unneglectable in COVID-19. *J Immunol Res*, 2021. 2021: p. 6659410.
- 51 Lamontagne, F., et al., A living WHO guideline on drugs to prevent covid-19. *BMJ*, 2021. 372: p. n526.
- 52 Kremer, J.M., et al., Hydroxychloroquine and the risk of respiratory infections among RA patients. *RMD Open*, 2020. 6(3).