



Universiteit
Leiden
The Netherlands

Thromboinflammation in high-risk human populations

Yuan, L.

Citation

Yuan, L. (2023, November 7). *Thromboinflammation in high-risk human populations*. Retrieved from <https://hdl.handle.net/1887/3656071>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3656071>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER

2

Sex-specific association between microvascular health and coagulation parameters: the Netherlands Epidemiology of Obesity (NEO) Study

Lushun Yuan¹, Jihee Han², Anouk I.M. van der Velden¹, Hans Vink^{3,4}, Renée de Mutsert², Frits R. Rosendaal², Astrid van Hylckama Vlieg², Ruifang Li-Gao^{2,5}, T.J Rabelink¹, Bernard M. van den Berg¹

¹*Eindhoven Laboratory for Vascular and Regenerative Medicine, department of Internal Medicine, Nephrology, Leiden University Medical Center, Leiden, the Netherlands.*

²*Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands.*

³*Department of Physiology, Cardiovascular Research Institute Maastricht, Maastricht, the Netherlands.*

⁴*MicroVascular Health Solutions LLC, Alpine, Utah, USA.*

⁵*Metabolon Inc., Morrisville, North Carolina, USA*

Abstract

Background: Microvascular dysfunction is a growing determinant of sex difference in coronary heart disease (CHD). The dysregulation of the coagulation system is involved in CHD pathogenesis and can be induced by endothelial glycocalyx (EG) perturbation. However, little is known about the link between EG function and coagulation parameters in population-based studies on sex specificity.

Objectives: We sought to examine sex differences in the relationship between EG function and coagulation parameters in a middle-aged Dutch population.

Methods: Using baseline measurements of 771 participants from the Netherlands Epidemiology of Obesity (NEO) study (age: 56 [IQR: 51-61] years, 53% women, BMI: 27.9 [IQR: 25.1-30.9] kg/m²), associations between glycocalyx-related perfused boundary region (PBR) derived from sidestream dark-field imaging and coagulation parameters (factor VIII/IX/XI, thrombin generation parameters, and fibrinogen) were investigated using linear regression analyses, adjusting for possible confounders (including CRP, leptin, and GlycA), followed by sex-stratified analyses.

Results: There was a sex difference in the associations between PBR and coagulation parameters. Particularly in women, one SD PBR (both total and feed vessel, indicating poorer glycocalyx status) was associated with higher FIX activity (1.8%, 95% CI: 0.3-3.3, and 2.0%, 95% CI: 0.5-3.4) and plasma fibrinogen levels (5.1 mg/dL, 95% CI: 0.4-9.9, and 5.8 mg/dL, 95% CI: 1.1-10.6). Furthermore, one SD PBR_{capillary} was associated with higher FVIII activity (3.5%, 95% CI: 0.4-6.5) and plasma fibrinogen levels (5.3 mg/dL, 95% CI: 0.6-10.0).

Conclusion: We revealed a sex-specific association between microcirculatory health and procoagulant status, which suggests considering microvascular health in the early development of CHD in women.

Keywords: coagulation factors, thrombin generation, fibrinogen, endothelial glycocalyx (EG), perfused boundary region (PBR)

Essentials

- Little is known about the link between microvascular health and coagulation parameters in population-based studies on sex specificity.
- We studied sex differences in relation between endothelial glycocalyx function and coagulation parameters in a middle-aged Dutch population.
- Particularly in women, associations between endothelial glycocalyx health and coagulation parameters were observed.
- Microcirculatory changes together with coagulation factors provide possible cues to prevent development of CHD in women.

Introduction

In previous decades, men were thought to be more susceptible to coronary heart disease (CHD) than women [1]. However, the risk of CHD in women is frequently underestimated due to the under-recognition of CHD and distinct clinical presentations, which resulted in a poor prognosis [2]. Nowadays, the systemic non-obstructive microvascular dysfunction, causing problems in small blood vessels feeding the muscular tissue, which can result in CHD, is believed to be more common in women than men. Therefore, this type of dysfunction is becoming a growing determinant of sex difference in CHD patients with normal or near normal coronary angiographic assessment [3, 4].

Previously, we examined endothelial surface perturbation using the non-invasive sidestream dark-field (SDF) imaging technique in a subpopulation of the Netherlands Epidemiology of Obesity (NEO) study [5-7]. While SDF imaging provides high contrast images of the sublingual microvasculature allowing to determine lateral red blood cell (RBC) movement within the vessels of interest, with the newly developed software automatic analysis of RBC velocity allows to include flow changes between feed vessels (10-25 μ m) and capillaries (< 10 μ m) to be coupled to the perfused boundary region (PBR), vessel density, and capillary blood volume [8, 9]. Already in various studies, it was previously shown that the detected changes in PBR, inversely related to the endothelial glycocalyx thickness, correlated with glycocalyx degradation products including circulating Syndecan-1, heparan sulfate (HS), hyaluronan, and soluble thrombomodulin and endothelial activation markers such as E-selection, soluble angiotensin-2, and soluble Tie-2 [10-13]. It was also found to be negatively correlated with coronary flow reserve (CFR), an assessment to evaluate coronary microvascular dysfunction (CMD). Besides, according to miCROvascular rarefaction in vasCUlar Cognitive Impairment and heArt failUre (CRUCIAL) study protocol [14], SDF imaging technique will be used to quantify microvascular health of cerebral and cardiac microvasculature, suggesting that this technique is a valid surrogate for cardiac microvasculature health examination [15].

The endothelial glycocalyx (EG) is a negatively charged gel-like surface matrix of proteoglycans and covalently bound glycosaminoglycans, glycoproteins, and glycolipids [16], and exerts an anti-inflammatory and antithrombotic role by covering various glycoprotein adhesion receptors for leukocytes [17], and platelets [18], or aiding in the sequestration of anti-adhesion factors [19]. EG perturbation acts as a representative of impaired microvascular function [6, 9, 20, 21]. In particular, several EG-bound proteins including specific sulfation patterns of HS proteoglycans or HS binding proteins such as anti-thrombin III (ATIII) and tissue factor pathway inhibitor (TFPI) could prevent the coagulation cascade *in vitro* and *in vivo* [22-24]. In disease cases such as nephrotic syndrome, sepsis, brain injury, and severe COVID-19, derangement of EG induced by microvascular dysfunction could accelerate hypercoagulation [8, 25-28].

Specifically, women were found to have already higher fibrinogen plasma levels than men of the same age and ethnic group [29]. Increased fibrinogen levels could push the balance from fibrinolysis to clotting [30] and have been reported as a potential risk factor for CHD [31-34]. Furthermore, other coagulation factors were also associated with increased risk of CHD [33, 35]. However, little is known about the link between EG function and coagulation parameters in population-based studies as well as whether a sex-specific determinant is present. Based on *in vitro* and *in vivo* evidence, we hypothesized that poorer endothelial glycocalyx function could lead to a procoagulant state in a population-based study, which might demonstrate sex specificity and be involved in the risk of CHD. The aim of our study was therefore to investigate the association between microvascular health, assessed by the new SDF imaging parameters, and levels of procoagulant factor activities (Factor VIII [FVIII], FIX, and FXI), fibrinogen concentration, and thrombin generation parameters in the general Dutch population and to further perform analyses stratified by sex, revealing sex differences in vascular vulnerability and its potential roles in the risk of CHD.

Materials and methods

Study population and study design

This study was performed in a population-based prospective cohort, the Netherlands Epidemiology of Obesity (NEO) study [7]. All 6,671 participants gave written informed consent, and the Medical Ethical Committee of the Leiden University Medical Center (LUMC) approved the study design. Initiated in 2008, the NEO study was designed to study pathways that lead to obesity-related diseases. Detailed information about the study design and data collection have been described elsewhere [7]. Briefly, men and women aged between 45 and 65 years with a self-reported body mass index (BMI) of 27 kg/m² or higher living in the greater area of Leiden (in the west of the Netherlands) were eligible to participate in the NEO study. In addition, all inhabitants aged between 45 and 65 years from one municipality (Leiderdorp) were invited irrespective of their BMI, representing the BMI distribution of the Dutch general population. Prior to their visits, participants completed questionnaires at home with demographic, lifestyle, and clinical information. At the baseline visit, fasting blood samples were drawn from the antecubital vein after the NEO participants had rested for 5 minutes. The present study is a cross-sectional analysis using a subpopulation of 918 NEO participants in whom SDF imaging was performed between January and October 2012, as part of the baseline visit at the Leiden University Medical Center NEO study center. Individuals were excluded from the analyses (Figure 1), when: 1) with missing SDF imaging parameters (n = 110); 2) with missing values of confounding factors (n = 8); 3) with current use of anticoagulant therapy (using vitamin K antagonists or heparin) (n = 7); 4) with missing values of coagulation parameters (n = 18); 5) with outliers values (z score > 5) in outcomes (i.e., coagulation factors and thrombin generation parameters) (n = 4), and 771 participants were included for all analyses using coagulation parameters as outcomes.




Figure 1. Study Flow chart.

SDF microcirculation imaging

Intravital microscopy was performed earlier on individuals in a supine position using an SDF camera (MicroVision Medical Inc., Wallingford, Pennsylvania) and acquired using Glycocheck software (Microvascular Health Solutions Inc., Salt Lake City, Utah) as described elsewhere [6, 9]. The software automatically identifies all available measurable microvessels distributed at a $1\mu\text{m}$ interval between $4\text{-}25\mu\text{m}$ and RBC velocity was included as a new parameter using the new software. After reanalyzing, the following validated parameters [8] were included in this study, divided into two categories: glycocalyx-related parameters, i.e., total PBR ($\text{PBR}_{\text{Total}}$, $4\text{-}25\mu\text{m}$), PBR feed vessel ($\text{PBR}_{\text{feed vessel}}$, $10\text{-}19\mu\text{m}$), and PBR capillary ($\text{PBR}_{\text{capillary}}$, $4\text{-}9\mu\text{m}$); microcirculatory-perfusion-related parameters, i.e., feed vessel RBC velocity ($\text{RBCV}_{\text{feed vessel}}$), capillary RBC velocity ($\text{RBCV}_{\text{capillary}}$), total valid vessel density (with measurable RBC velocity; D_{Total}), perfused feed vessel density ($D_{\text{feed vessel}}$), perfused capillary density ($D_{\text{capillary}}$), and capillary blood volume ($\text{CBV}_{\text{static}}$). Detailed

information about the new software used in the NEO study is described previously [8]. PBR_{Total} , $PBR_{feed\ vessel}$, and $PBR_{capillary}$ were expressed in μm . Levels of $RBCV_{feed\ vessel}$ and $RBCV_{capillary}$ were expressed in $\mu m/s$. Levels of D_{Total} , $D_{feed\ vessel}$, and $D_{capillary}$ were expressed in $\mu m/mm^2$. CBV_{static} was expressed in pL/mm^2 .

Coagulation factor activity

Blood samples for measurements of coagulation factors activity were collected and processed within 4 hours and drawn into tubes containing 0.106M trisodium citrate (Sarstedt, Etten Leur, the Netherlands) without the measurement of residual platelets, as discussed earlier [7]. Fasting fibrinogen levels were measured according to the method of Claus, as described earlier [36]. Fasting FVIII, FIX, and FXI activity were measured with a coagulometric clot detection method on an ACL TOP 700 analyzer (Werfen, Barcelona, Spain), and the Werfen plasma was used as reference plasma to calibrate factor measurements (Werfen). Levels of FVIII, FIX, and FXI activity were expressed in percentages (%). Levels of fibrinogen were expressed in mg/dL.

Thrombin generation

All blood samples were collected and processed within 4hrs. Tubes were centrifuged for 10 minutes at 2500g at 18°C and aliquoted plasma was stored at -80°C until further use, only after first time thawing for 3 min 37°C (waterbath), as discussed earlier [7]. Thrombin generation was measured according to the protocols earlier described by Hemker *et al.* [37]: calibrated automated thrombogram (CAT; Thrombinoscope BV, Maastricht, the Netherlands). Briefly, 20 μL of PPP-Reagent LOW (86194,TS31.00, STAGO, France) and thrombin calibrator (86192, TS30.00, STAGO, France) were dispensed into the wells of a round-bottom 96-well plate (#3655, Thermo Scientific, Uden, the Netherlands).

Thermostable inhibitor of contact activation (TICA, PS-0177-oxoxox, Maastricht, the Netherlands) was added to plasma samples from participants in the NEO study and normal pooled plasma (NPP, as an internal control for each plate). Then 80 μ L of mixed plasma was added to the plate and the plate was placed in the Fluorometer for a 10-minute, 37°C incubation. Thrombin formation was initiated by adding 20 μ L of the fluorogenic substrate with calcium (FluCa-kit, 86197, TS 50.00, STAGO, France). The final reaction volume was 120 μ L. Thrombin formation was determined every 10 seconds for 50 minutes and corrected for the calibrator using Thrombinscope software. Thrombin generation parameters included lag time, time to Peak (ttPeak), peak thrombin generation (Peak), endogenous thrombin potential (ETP), and thrombin generation velocity (VelIndex). Levels of lag time and ttPeak were expressed in minutes (min); levels of Peak height were expressed in nM; levels of ETP were expressed in nM \times min; levels of VelIndex were expressed in nM/min.

Serum inflammatory markers

Serum concentrations of C-reactive protein (CRP) were determined using a high sensitivity CRP assay (hs-CRP, TINA-Quant CRP HS system, Roche, Germany and Modular P800, Roche, Germany) [36]. Serum leptin concentration was measured using a human leptin competitive RadioImmunoAssay (RIA) (HL-81HK, Merck Millipore, Darmstadt, Germany). The leptin concentration was counted using a gamma counter, as described elsewhere [38]. Glycoprotein acetyls (GlycA) concentrations were measured in plasma that had undergone one previous freeze-thaw cycle, using a high-throughput proton nuclear magnetic resonance (NMR) spectroscopy (Nightingale Health Ltd., Helsinki, Finland) [39].

Statistical analyses

Descriptive baseline characteristics of the study population were expressed as median with interquartile range (IQR) for non-normally distributed variables, mean with standard deviation (SD) for normally distributed variables, or percentages (%) for dichotomous variables.

First, the distributions of confounding factors and SDF imaging parameters were evaluated, and z-transformation (i.e., with mean=0 and SD=1) was performed on age, BMI, and SDF imaging parameters. After checking the distributions of outcome variables, thrombin generation velocity (VelIndex) showed a right-skewed distribution and, \log_2 transformation was performed on VelIndex. Afterwards, linear regression analysis was used to investigate the associations between SDF imaging parameters (exposures) and pro-coagulation factors (i.e., FVIII, FIX, and FXI), fibrinogen, and thrombin generation parameters (outcomes). The analyses were adjusted for several potential confounding factors. First, crude analyses were performed (model 1). Second, we adjusted the models for the confounding factors age (unit: years, continuous variable) and sex (categories: women and men, dichotomous variable) (model 2). Third, models were adjusted for other confounding factors: BMI (unit: kg/m^2 , continuous variable), current smoking status (categories: current users and non-current users, dichotomous variable), menopausal status (categories: premenopausal and postmenopausal, dichotomous variable), current use of the oral contraceptive pill (categories: current users and non-current users, dichotomous variable), and current use of hormone replacement therapy (categories: current users and non-current users, dichotomous variable) (model 3). Fourth, we adjusted for serum C-reactive protein (CRP) (unit: mg/L , continuous variable), serum leptin concentration (unit: $\mu\text{g}/\text{L}$, continuous variable) and glycoprotein acetyls (GlycA, unit: mmol/L , continuous variable) in a separate model as these systemic inflammation markers might be potential confounders of the association between microvascular health and coagulation parameters (model 4). We calculated differences in the mean levels of fibrinogen, FVIII, FIX, FXI, and thrombin generation parameters, with 95% confidence intervals (CI), associated with SDF imaging

parameters. Furthermore, the association between SDF imaging parameters and coagulation factor levels was examined through sex-stratified analysis. Mean coagulation factor levels were calculated for sex-stratified SDF imaging parameters. As a sensitivity analysis, all analyses were performed separately in premenopausal and postmenopausal women.

To our knowledge, the present study is the first to investigate the association between the microvascular health index derived from SDF imaging and the levels of coagulation factors as well as the parameters of thrombin generation on a population level; we have no prior information on effect sizes to perform a power calculation. Noteworthy, most of previous studies with relatively small sample sizes have identified associations between the microvascular health index and disease outcomes, e.g., COVID-19 in 38 participants [13], coronary artery disease in 115 participants [40], and sepsis in 51 participants [8]. To address the power that we could achieve using 771 individuals, we used the following setting: significant level 0.05, power 80%, with the linear regression $PBR_{Total} \sim \text{fibrinogen}$ as an example; we may need 394, 54, and 24 samples for a small, medium and large effect size, respectively [41]. Therefore, a sample size of 771 individuals may have the statistical power to capture even small effect sizes.

Results

Participant characteristics

Table 1 shows the characteristics of individuals included in the present study. Of 771 included participants, 53% were women, with a median age of 56 years (interquartile range, 51–61 years), and 12.5% were current smokers. Participants had a median BMI of 27.9 kg/m² (interquartile range, 25.1–30.9). Among women, 84% had a postmenopausal status, 7.6% used oral contraceptives, and 3.4 % hormone replacement therapy. Apart from this, we observed a higher level of inflammatory markers and procoagulant status. While

capillary glycocalyx is very important for proper endothelial function, the PBR dimensions are measured in vessels with diameters ranging from 4 – 25 microns. Therefore the contribution of capillary glycocalyx to PBR_{total} is very limited. PBR_{total} is comparable to $PBR_{feed\ vessel}$ and both are the main parameters presenting possible disturbances of the endothelial surface layer, which in women are both significantly increased (median, women vs. men: PBR_{total} , 2.35 vs. 2.3; $PBR_{feed\ vessel}$, 2.31 vs. 2.24). The density of perfused capillaries is higher in women vs. men, as measured by both a higher capillary density ($D_{capillary}$) and higher capillary blood volume (CBV_{static}), which is associated with a lower capillary RBC velocity ($RBCV_{capillary}$), suggests a more advantages capillary presence in women.

Association between microvascular health and levels of coagulation parameters in the total population

Linear regression analyses were used to investigate the association between one standard deviation (SD) difference in microvascular parameters and the coagulation factors. In model 3 of the total population, both glycocalyx-related parameters, PBR_{Total} and $PBR_{feed\ vessel}$ were associated with higher fibrinogen levels (Table S1). Nevertheless, with an additional adjustment for systemic inflammatory markers (model 4), only the association of the $PBR_{feed\ vessel}$ remained (Table S1 and Figure S1) and we observed that one SD (i.e., $0.3\mu m$) in the $PBR_{feed\ vessel}$ was associated with an higher fibrinogen concentration of 3.2 mg/dL (95% CI: 0.02-6.4). Interestingly, the microcirculatory perfusion parameters, in particular all three vessel density parameters, were positively associated with the endogenous thrombin potential (ETP) in all models with very similar effect size estimations between model 3 and model 4 (Table S2 and Figure S2). In model 4 of the total population, one SD in vessel density (D_{Total} , $D_{feed\ vessel}$, $D_{capillary}$; 65.84, 42.17, 41.39 $\mu m/mm^2$, respectively), corresponded to higher ETP levels (31.9 nM \times min, 95% CI: 9.6-54.2; 23.7 nM \times min, 95% CI: 1.2-46.2; 25.2 nM \times min, 95% CI: 2.9-47.5, respectively).

Table 1. Characteristics of the study population

	Total (100%)	Women (53%)	Men (47%)
Demographic			
Age, y	56 (51-61)	56 (50.75-61)	57 (51-61)
BMI, kg/m ²	27.93 (25.08-30.91)	27.75 (24.52-31.25)	27.99 (25.73-30.59)
Tobacco smoking (% current user)	12.5	8.1	17.4
Menopause status (% of female postmenopausal/perimenopausal)	NA	83.8	NA
Medication (current use)			
Oral contraceptive pill (% of females)	NA	7.6	NA
HRT (% of female)	NA	3.4	NA
Systemic inflammatory markers			
C-reactive protein, mg/L	1.37 (0.74-3.06)	1.61 (0.8-3.5) ***	1.25 (0.64-2.26)
Leptin, µg/L	14.7 (8-27.15)	25.2 (16-37) ****	8.4 (5.55-12.9)
Glycoprotein acetyls, mmol/L	1.23 (1.13-1.34)	1.21 (1.11-1.32) *	1.24 (1.14-1.35)
Coagulation parameters			
Fibrinogen, mg/dL	289.32 (257.77-330.88)	300.55 (265.66-346.02) ****	278.98 (247.19-312.78)
Factor VIII (%)	122.98 (101.39-147.01)	123.53 (102.05-148.34)	122.98 (101.39-145.04)
Factor IX (%)	119.31 (108.41-134.28)	119.31 (107.27-136.11)	119.31 (110.74-131.41)
Factor XI (%)	116.77 (103.75-129.16)	120.02 (107.58-134.01) ****	111.56 (100.06-123.93)
Lag time, min	6.72 (6-7.75)	6.5 (5.75-7.58) ****	7.08 (6.17-7.92)
ttPeak, min	14.75 (13.46-16.08)	14.36 (13.19-15.75) ****	15.04 (13.83-16.33)
Peak, nM	83.27 (62.1-106.28)	84.77 (61.11-111.09)	81.69 (63.92-102.77)
ETP, nM-min	1149.22 (915.56-1396.66)	1175.06 (916.94-1439.8) *	1124.76 (914.49-1329.16)
VelIndex, nM/min	10.42 (7.47-14.63)	10.79 (7.36-15.32)	10.18 (7.76-13.8)
SDF imaging parameters			
PBR of total vessels, µm	2.33 (2.19-2.5)	2.35 (2.2-2.54) **	2.3 (2.17-2.44)
PBR of feed vessels, µm	2.28 (2.09-2.48)	2.31 (2.11-2.52) **	2.24 (2.06-2.44)
PBR of capillaries, µm	1.2 (1.13-1.27)	1.2 (1.14-1.27)	1.2 (1.13-1.26)
RBC velocity in feed vessels, µm/s	55.85 (43.93-68.91)	51.24 (39.7-64.4) ****	60.26 (49.5-73.13)
RBC velocity in capillaries, µm/s	55.74 (41.94-70.55)	50.77 (37.78-66.86) ****	60.45 (47.77-72.65)
Total vessel density, µm/mm ²	257.02 (209.81-309.06)	255.58 (206.18-309.23)	259.16 (213.37-306.45)
Feed vessel density, µm/mm ²	131.57 (107.1-160.52)	129.5 (103.51-158.26) *	134.83 (110.94-165.09)
Capillary density, µm/mm ²	100.71 (79.69-131.2)	104.25 (81.76-137.28) *	96.82 (74.81-121.83)
Capillary blood volume, pL/mm ²	2.52 (1.4-4.35)	2.81 (1.41-4.76) *	2.29 (1.38-3.92)

Data are shown as mean (± SD), median (25th percentile–75th percentile), or percentage.

Abbreviations: BMI, body mass index; ETP, endogenous thrombin potential; HRT, hormonal replacement therapy; NA, not applicable; PBR, perfused boundary region; Peak, peak thrombin generation; ttPeak, time to peak; VelIndex, thrombin generation velocity.

Non-paired t-test or Wilcoxon test were performed between women and men; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Effect modification by sex of the association between microvascular health and coagulation factors

After sex stratification, we observed differences between men and women between the glycocalyx-related parameters and coagulation factor levels (Figure 2 and Figure 3). Consistent associations were found between PBR_{Total} and $PBR_{feed\ vessel}$ and FIX activity as well as plasma fibrinogen concentration specifically in women, which remained after adjusting for inflammatory markers: one SD in PBR_{Total} or $PBR_{feed\ vessel}$ (0.25 μm and 0.3 μm , respectively) was associated with higher fibrinogen levels (5.1 mg/dL, 95% CI: 0.4-9.9 and 5.8 mg/dL, 95% CI: 1.1-10.6; respectively) and higher FIX activity (1.8%, 95% CI: 0.3-3.3 and 2.0%, 95% CI: 0.5-3.4; respectively) (Figure 2 and Table S3); while no associations were observed in men (Figure 3 and Table S4). For $PBR_{capillary}$, in addition to the association with fibrinogen concentration, $PBR_{capillary}$ was also associated with FVIII activity, which remained after further adjustment for systemic inflammatory markers (model 4): one SD in $PBR_{capillary}$ (0.11 μm) was associated with higher fibrinogen levels (5.3 mg/dL, 95% CI: 0.6-10.0) and FVIII activity (3.5%, 95% CI: 0.4-6.5) (Figure 2 and Table S3).

There were differences in the associations between microcirculatory-perfusion-related parameters and coagulation factor levels stratified by sex. For women, the associations between vessel density and ETP were observed in the crude and age- and sex-adjusted model (Figure S3 and Table S5); however, only $D_{capillary}$ was positively associated with ETP after further adjusting for demographic and lifestyle factors (model 3) and systemic inflammatory markers (model 4), in which one SD of $D_{capillary}$ (42.26 $\mu m/mm^2$) was associated with higher levels of ETP (32.0 nM \times min, 95% CI: 0.1-63.8). In men, only one consistent

negative association was observed between the $RBCV_{\text{feed vessel}}$ and thrombin generation parameter time to peak (ttPeak) (Figure S4 and Table S6), in which one SD of the $RBCV_{\text{feed vessel}}$ ($61.52 \mu\text{m/s}$) was associated with lower levels of ttPeak (-0.2 , 95% CI: -0.4 - -0.02). In addition, one SD of D_{Total} ($68.08 \mu\text{m/mm}^2$) was associated with higher ETP levels ($31.9 \text{ nM} \times \text{min}$, 95% CI: 1.3 - 62.6).

Sensitivity analyses were performed to test the robustness. Using premenopausal women yielded even more notable results than the main analysis for total women that PBR difference was associated with higher fibrinogen concentration, higher FXI and FVIII activity, and higher thrombin generation parameters (ETP, peak, and VelIndex) (Figure S5, Table S7, Figure S7, and Table S9). However, the results in postmenopausal women were less significant but comparable to the main analysis for all women (Figure S6, Table S8, Figure S8, and Table S10).




Figure 2. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in women. After sex stratification, differences in women between the glyocalyx-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2 = model 1 + age; Model 3: model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: PBR_{Total} : PBR of total vessels from 4 - $25\mu\text{m}$; $PBR_{\text{feed vessel}}$: PBR of feed vessels from 10 - $19\mu\text{m}$; $PBR_{\text{capillary}}$: PBR of capillaries from 4 - $9\mu\text{m}$.




Figure 3. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in men. After sex stratification, differences in men between the glyocalyx-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI + current smoking status; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: PBR_{Total}: PBR of total vessels from 4-25 μ m; PBR_{feed vessel}: PBR of feed vessels from 10-19 μ m; PBR_{capillary}: PBR of capillaries from 4-9 μ m.

Discussion

In this population-based cross-sectional study the PBR in feed vessels was positively associated with plasma fibrinogen levels in the total population. Remarkably, we discovered a sex difference in the associations between PBR and coagulation parameters, in which higher PBR (both total and feed vessel, indicating perturbed glyocalyx) was associated with higher FIX activity and plasma fibrinogen levels in women. Furthermore, in women, higher PBR_{capillary} (i.e., poorer glyocalyx status in capillaries) was associated with higher FVIII activity and fibrinogen levels and higher D_{capillary} was associated with higher levels of ETP, while none of these associations were present in men.

SDF imaging technique could provide two types of parameters concerning endothelial glyocalyx and microcirculatory perfusion function. We proposed that the change in the endothelial surface properties, i.e., the endothelial glyocalyx layer (PBR), is a likely functional unit that could present as a marker for microvascular dysfunction. In the present

study, we observed an association between early (pre-clinical) microvascular health changes and coagulation factor activation and discovered a striking sex difference in microvascular health, in which women showed a perturbed EG concomitant with a more pro-coagulable endothelial surface. This association was not observed in men. These findings highlight the importance of sex differences in microcirculatory perturbation in CHD.

Previously, studies reported the association between CHD and the underlying systemic presence of a hypercoagulable state [42-44]. Our results showed that in women, one SD in PBR_{Total} and $PBR_{feed\ vessel}$ was associated with a 5.1 and 5.8 mg/dL increase in fibrinogen concentration. Based on a large individual participant meta-analysis study that assessed the association between fibrinogen concentration and CHD risk [45], such increase in PBR would suggest a 12-14% increase in CHD risk in women, which could be clinically relevant. In line with our observations, Brands *et al.* reported that reduced endothelial glycocalyx barrier properties were only found in women with CHD, while not present in men [40]. In addition, in the REasons for Geographic and Racial Differences in Stroke Study (REGARDS), a large population-based observational study, higher levels of FVIII and FIX were associated with increased risk of CHD [35, 46]. According to these associations, per SD difference in PBR could correspond to a 2-5% increase in the risk of CHD in women. FVIII and vWF are two distinct but related glycoproteins that circulate in plasma as a tightly bound complex (FVIII/VWF) [47]. As one of the endothelial activation markers, vWF can bind to HS at the endothelial cell surface [26, 48]. Based on the observed tight correlation between FVIII and vWF in the Multiple Environmental and Genetic Assessment of risk factors for venous thrombosis (MEGA) study [49], we deduced an association between EG health and vWF in our present study, to further indicate the link between EG dysfunction and endothelial activation. Our current findings imply a role of microvascular health in CHD risk in women through the association between higher microvascular PBR (perturbed glycocalyx) and differences in both FIX and FVIII activity, together with the already high plasma fibrinogen levels (Figure 4).




Figure 4. Proposed pathophysiologic concept on the association between microvascular health and endothelial response to coagulation. Here a healthy glycocalyx (i.e., PBR_{low}, marked by the arbitrary dotted lines) provides a robust anti-coagulant barrier through binding of AT-III [22], presence of thrombomodulin [26], PAI-1 [54] and TFPI [23] to HS proteoglycans [19]. The thrombin-TM complex activates protein C to produce APC, which inactivates factors VIIIa and Va in the presence of protein S, thereby inhibiting further thrombin (F-II) formation [55]. The pro-coagulant endothelial surface is a result of multiple mechanisms involving effects such as, gene transcription, protein expression and release. We hypothesized that in the event of a perturbed glycocalyx (i.e., PBR_{high}, marked by arbitrary dotted lines further apart), a pro-coagulant surface appears, with increasing binding possibilities of fibrinogen (interacting with ICAM1) [56, 57], FXI [58, 59] and F-VIII (to surface expressed von Willebrand factor, vWF) [47] and reduced presence of the anti-coagulant factors. While F-IX can bind directly to the cell surface or abluminal collagen, fibrinogen and F-VIII binding to the surface depends more on the activation state of the endothelial cells. This increased interaction of coagulation factor IX and VIII activity and fibrinogen concentration, as observed in the present study, together with possible diminished surface anti-coagulation pathways, would especially in women play a critical role in increased systemic microvascular dysfunction and development of coronary heart disease (CHD).

The present study also included various thrombin generation parameters which represent the global coagulation cascade in addition to coagulation factor activity. Interestingly, no associations were observed between the microvascular health parameters measured and

the dynamic parameters of thrombin generation (such as lag time and time to peak); except for an association between vessel density and ETP. Thrombin was found to be involved in both pro-thrombotic and inflammatory endothelial processes [50]. While previous studies reported an association between *in vivo* thrombin generation potential and severity of coronary vessel disease [51], increased thrombin generation potential was a characteristic in patients with clinically stable CHD [52]. Although we found sex differences in the association between vessel density and ETP, when we compared the findings in the total population, men, and women, we found that the associations in men and women were close to significance; therefore, this observation might reflect a general feature of microvascular health and thrombin formation rather than a sex specific feature. Furthermore, comparing with previous studies using thrombin generation assay, the participants in the present study were preclinical and did not have CHD yet, which might limit the effect of thrombin and findings related to thrombin generation. In the context with the associations between endothelial glycocalyx (PBR) and coagulation factors, the associations between vessel densities and ETP were very marginal.

To the best of our knowledge, this is the first study to show the association between microvascular health derived from SDF imaging and the levels of coagulation factors and thrombin generation parameters in a large population-based study. In addition, adjustment for extensive potential confounding factors including systemic inflammatory markers strongly suggested a sex-specific role in monitoring microvascular health change in women. Additionally, the present study used the new software with red blood cell (RBC) velocity being included as a new parameter to better quantify the microcirculatory difference.

Limitations of the present study should also be addressed. First, it is an observational, cross-sectional study and residual confounding may still be present. Second, the present study population is mainly comprised of European white participants in middle age (i.e., between 45 and 65 years). Therefore, it is not clear whether the results of the present study can be generalizable to other ethnicities or age groups. Third, the majority of women are postmenopausal; although sensitivity analysis revealed similar findings in premenopausal

women, future studies including more premenopausal women need to be done. Fourth, a recent study reported the variability of microcirculatory measurements in healthy volunteers [53] showing that when three consecutive measurements are averaged, SDF imaging with GlycoCheck software can be used with acceptable reliability and reproducibility for microcirculation measurements on a population level. In our current study, repeated measurement of SDF imaging was not performed for each participant and limited information was extracted from the smallest capillaries (diameter 4 μm). However, the measurement by itself is already an average from at least ten recordings on one individual at the same time point rather than averaging three consecutive measurements on the same individual, which could only lead to random error and subsequent underestimation of the effect size. Fifth, due to the technical limitation of the *ex vivo* thrombin generation assay, we could not avoid batch effects and this assay could not represent the real circumstances of the coagulation cascade in humans. Sixth, as a result of sample availability to measure the circulating EG disruption and EC activation markers, our study lacks possible mechanistic correlations between EG dysfunction, increased procoagulant state and increased CHD risk in females. Seventhly, although we focussed on the role of EG perturbation in coagulation in the present study, it is important to note that the formation of a pro-coagulant surface is a result of multiple mechanisms, among others, such as gene transcription changes and protein expression and release. Finally, in model 4 we adjusted for systemic inflammatory markers (such as CRP, leptin, and GlycA) in the association between microvascular health and coagulation factor levels. It should be noted that these systemic inflammatory markers could act as mediators instead of confounders in the estimated associations. If so, adjustment for systemic inflammatory markers could lead to an underestimation of the association between microvascular health and coagulation factor levels.

In conclusion, our data reveal a hitherto unreported sex-specific association between microcirculatory health and procoagulant status. Microcirculatory differences between men and women identified in our study implied that microvascular health changes might be

the earliest detectable clue prior to the general higher procoagulant status in women ultimately developing CHD, which is also independent of increased systemic inflammatory state (Figure 4). Our study suggested the potential clinical utility of monitoring microcirculatory change specifically in women to prevent the development of CHD.

Authors' contributions

The inclusion of participants for the NEO study, the study design of the NEO study, and the acquisition of data of the NEO study have been performed by Renée de Mutsert and Frits R. Rosendaal. Of the present study, the study concept and design was performed by Lushun Yuan, Jihee Han, Ruifang Li-Gao, T.J Rabelink and Bernard M. van den Berg. The analyses and interpretation of the data were performed by Lushun Yuan, Jihee Han, Anouk I.M. van der Velden, Hans Vink, Astrid van Hylckama Vlieg, Renée de Mutsert, Ruifang Li-Gao, T.J Rabelink and Bernard M. van den Berg. The draft of the manuscript and design of figures are performed by Lushun Yuan, Jihee Han, Ruifang Li-Gao, T.J Rabelink and Bernard M. van den Berg. All authors read and approved the final manuscript.

Acknowledgments

We express our gratitude toward all participants of the NEO study. Furthermore, we thank P.R. van Beelen and all research nurses for the data collection, P.J. Noordijk and her team for the handling and storage of the blood samples, and I. de Jonge for the data management.

Conflict of interest

Hans Vink works for MicroVascular Health Solutions LLC and Ruifang Li-Gao works for Metabolon Inc. All of the authors, including Lushun Yuan, Jihee Han, Anouk I.M. van der Velden, Hans Vink, Renée de Mutsert, Frits R. Rosendaal, Astrid van Hylckama Vlieg,

Ruifang Li-Gao, T.J Rabelink, Bernard M. van den Berg, hereby confirm that they have no conflict of interest and have nothing to disclose.

Funding

The NEO study is supported by the participating departments, the Division and the Board of Directors of the Leiden University Medical Centre, and by the Leiden University, Research Profile Area 'Vascular and Regenerative Medicine'. Coagulation factor analyses were funded by 'Stichting De Merel'. Stichting De Merel had no role in the study design, data collection, and analysis, decision to publish, or preparation, review, or approval of the manuscript. This work was supported by the China Scholarship Council grant to Lushun Yuan (CSC no. 201806270262).

Reference

- 1 Bots SH, Peters SAE, Woodward M. Sex differences in coronary heart disease and stroke mortality: a global assessment of the effect of ageing between 1980 and 2010. *BMJ Glob Health*. 2017; **2**: e000298. 10.1136/bmjgh-2017-000298.
- 2 Maas AH, Appelman YE. Gender differences in coronary heart disease. *Neth Heart J*. 2010; **18**: 598-602. 10.1007/s12471-010-0841-y.
- 3 Koller A. Perspectives: Microvascular endothelial dysfunction and gender. *Eur Heart J Suppl*. 2014; **16**: A16-A9. 10.1093/eurheartj/sut005.
- 4 Padro T, Manfrini O, Bugiardini R, Canty J, Cenko E, De Luca G, Duncker DJ, Eringa EC, Koller A, Tousoulis D, Trifunovic D, Vavlukis M, de Wit C, Badimon L. ESC Working Group on Coronary Pathophysiology and Microcirculation position paper on 'coronary microvascular dysfunction in cardiovascular disease'. *Cardiovasc Res*. 2020; **116**: 741-55. 10.1093/cvr/cvaa003.
- 5 Lekakis J, Abraham P, Balbarini A, Blann A, Boulanger CM, Cockcroft J, Cosentino F, Deanfield J, Gallino A, Ikonomidis I, Kremastinos D, Landmesser U, Protogerou A, Stefanadis C, Tousoulis D, Vassalli G, Vink H, Werner N, Wilkinson I, Vlachopoulos C. Methods for evaluating endothelial function: a position statement from the European Society of Cardiology Working Group on Peripheral Circulation. *Eur J Cardiovasc Prev Rehabil*. 2011; **18**: 775-89. 10.1177/1741826711398179.

- 6 Lee DH, Dane MJ, van den Berg BM, Boels MG, van Teeffelen JW, de Mutsert R, den Heijer M, Rosendaal FR, van der Vlag J, van Zonneveld AJ, Vink H, Rabelink TJ, group NEOs. Deeper penetration of erythrocytes into the endothelial glycocalyx is associated with impaired microvascular perfusion. *PLoS one*. 2014; **9**: e96477. 10.1371/journal.pone.0096477.
- 7 de Mutsert R, den Heijer M, Rabelink TJ, Smit JW, Romijn JA, Jukema JW, de Roos A, Cobbaert CM, Kloppenburg M, le Cessie S, Middeldorp S, Rosendaal FR. The Netherlands Epidemiology of Obesity (NEO) study: study design and data collection. *European journal of epidemiology*. 2013; **28**: 513-23. 10.1007/s10654-013-9801-3.
- 8 Rovas A, Sackarnd J, Rossaint J, Kampmeier S, Pavenstadt H, Vink H, Kumpers P. Identification of novel sublingual parameters to analyze and diagnose microvascular dysfunction in sepsis: the NOSTRADAMUS study. *Crit Care*. 2021; **25**: 112. 10.1186/s13054-021-03520-w.
- 9 van der Velden AIM, van den Berg BM, de Mutsert R, van der Vlag J, Jukema JW, Rosendaal FR, Rabelink TJ, Vink H. Microvascular differences in individuals with obesity at risk of developing cardiovascular disease. *Obesity*. 2021; **29**: 1439-44. 10.1002/oby.23222.
- 10 Drost CC, Rovas A, Kusche-Vihrog K, Van Slyke P, Kim H, Hoang VC, Maynes JT, Wennmann DO, Pavenstadt H, Linke W, Lukasz A, Hesse B, Kumpers P. Tie2 Activation Promotes Protection and Reconstitution of the Endothelial Glycocalyx in Human Sepsis. *Thromb Haemost*. 2019; **119**: 1827-38. 10.1055/s-0039-1695768.
- 11 Rovas A, Seidel LM, Vink H, Pohlkötter T, Pavenstadt H, Ertmer C, Hessler M, Kumpers P. Association of sublingual microcirculation parameters and endothelial glycocalyx dimensions in resuscitated sepsis. *Crit Care*. 2019; **23**: 260. 10.1186/s13054-019-2542-2.
- 12 Bol ME, Huckriede JB, van de Pas KGH, Delhaas T, Lorusso R, Nicolaes GAF, Sels JEM, van de Poll MCG. Multimodal measurement of glycocalyx degradation during coronary artery bypass grafting. *Front Med (Lausanne)*. 2022; **9**: 1045728. 10.3389/fmed.2022.1045728.
- 13 Rovas A, Osiaevi I, Buscher K, Sackarnd J, Tepasse PR, Fobker M, Kuhn J, Braune S, Gobel U, Tholking G, Groschel A, Pavenstadt H, Vink H, Kumpers P. Microvascular dysfunction in COVID-19: the MYSTIC study. *Angiogenesis*. 2021; **24**: 145-57. 10.1007/s10456-020-09753-7.
- 14 van Dinther M, Bennett J, Thornton GD, Voorter PHM, Ezponda Casajus A, Hughes A, Captur G, Holtackers RJ, Staals J, Backes WH, Bastarika G, Jones EAV, Gonzalez A, van Oostenbrugge R, Treibel TA. Evaluation of microvascular rarefaction in vascular Cognitive Impairment and heart failure (CRUCIAL): Study protocol for an observational study. *Cerebrovasc Dis Extra*. 2023; **13**: 18-32. 10.1159/000529067.
- 15 Ikonomidis I, Pavlidis G, Lambadiari V, Rafouli-Stergiou P, Makavos G, Thymis J, Kostelli G, Varoudi M, Katogiannis K, Theodoropoulos K, Katsimbri P, Parissis J, Papadavid

E. Endothelial glycocalyx and microvascular perfusion are associated with carotid intima-media thickness and impaired myocardial deformation in psoriatic disease. *J Hum Hypertens*. 2022; **36**: 1113-20. 10.1038/s41371-021-00640-2.

16 Dane MJ, van den Berg BM, Lee DH, Boels MG, Tiemeier GL, Avramut MC, van Zonneveld AJ, van der Vlag J, Vink H, Rabelink TJ. A microscopic view on the renal endothelial glycocalyx. *Am J Physiol Renal Physiol*. 2015; **308**: F956-F66. 10.1152/ajprenal.00532.2014.

17 Schmidt EP, Yang Y, Janssen WJ, Gandjeva A, Perez MJ, Barthel L, Zemans RL, Bowman JC, Koyanagi DE, Yunt ZX, Smith LP, Cheng SS, Overdier KH, Thompson KR, Geraci MW, Douglas IS, Pearse DB, Tudor RM. The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. *Nature medicine*. 2012; **18**: 1217-23. 10.1038/nm.2843.

18 Reitsma S, Oude Egbrink MG, Heijnen VV, Megens RT, Engels W, Vink H, Slaaf DW, van Zandvoort MA. Endothelial glycocalyx thickness and platelet-vessel wall interactions during atherogenesis. *Thromb Haemost*. 2011; **106**: 939-46. 10.1160/TH11-02-0133.

19 Boels MG, Lee DH, van den Berg BM, Dane MJ, van der Vlag J, Rabelink TJ. The endothelial glycocalyx as a potential modifier of the hemolytic uremic syndrome. *Eur J Intern Med*. 2013; **24**: 503-9. 10.1016/j.ejim.2012.12.016.

20 de Jongh RT, Serne EH, RG IJ, de Vries G, Stehouwer CD. Impaired microvascular function in obesity: implications for obesity-associated microangiopathy, hypertension, and insulin resistance. *Circulation*. 2004; **109**: 2529-35. 10.1161/01.CIR.0000129772.26647.6F.

21 Vlahu CA, Lemkes BA, Struijk DG, Koopman MG, Krediet RT, Vink H. Damage of the endothelial glycocalyx in dialysis patients. *J Am Soc Nephrol*. 2012; **23**: 1900-8. 10.1681/ASN.2011121181.

22 Chopra P, Joshi A, Wu J, Lu W, Yadavalli T, Wolfert MA, Shukla D, Zaia J, Boons GJ. The 3-O-sulfation of heparan sulfate modulates protein binding and lyase degradation. *Proc Natl Acad Sci U S A*. 2021; **118**. 10.1073/pnas.2012935118.

23 Tinholt M, Stavik B, Louch W, Carlson CR, Sletten M, Ruf W, Skretting G, Sandset PM, Iversen N. Syndecan-3 and TFPI colocalize on the surface of endothelial-, smooth muscle-, and cancer cells. *PLoS one*. 2015; **10**: e0117404. 10.1371/journal.pone.0117404.

24 Rezaie AR, Giri H. Anticoagulant and signaling functions of antithrombin. *Journal of thrombosis and haemostasis : JTH*. 2020; **18**: 3142-53. 10.1111/jth.15052.

25 Uchimido R, Schmidt EP, Shapiro NI. The glycocalyx: a novel diagnostic and therapeutic target in sepsis. *Crit Care*. 2019; **23**: 16. 10.1186/s13054-018-2292-6.

26 Yuan L, Cheng S, Sol W, van der Velden AIM, Vink H, Rabelink TJ, van den Berg BM. Heparan sulfate mimetic fucoidan restores the endothelial glycocalyx and protects against

dysfunction induced by serum of COVID-19 patients in the intensive care unit. *ERJ Open Res.* 2022; **8**. 10.1183/23120541.00652-2021.

27 Rabelink TJ, van den Berg BM, Garsen M, Wang G, Elkin M, van der Vlag J. Heparanase: roles in cell survival, extracellular matrix remodelling and the development of kidney disease. *Nat Rev Nephrol.* 2017; **13**: 201-12. 10.1038/nrneph.2017.6.

28 Haeren RH, Vink H, Staals J, van Zandvoort MA, Dings J, van Overbeeke JJ, Hoogland G, Rijkers K, Schijns OE. Protocol for intraoperative assessment of the human cerebrovascular glycocalyx. *BMJ open.* 2017; **7**: e013954. 10.1136/bmjopen-2016-013954.

29 Vorster HH. Fibrinogen and women's health. *Thromb Res.* 1999; **95**: 137-54. 10.1016/s0049-3848(99)00033-x.

30 Kim PY, Stewart RJ, Lipson SM, Nesheim ME. The relative kinetics of clotting and lysis provide a biochemical rationale for the correlation between elevated fibrinogen and cardiovascular disease. *J Thromb Haemost.* 2007; **5**: 1250-6. 10.1111/j.1538-7836.2007.02426.x.

31 Hsieh CT, Chien KL, Hsu HC, Lin HJ, Su TC, Chen MF, Lee YT. Associations between fibrinogen levels and the risk of cardiovascular disease and all-cause death: a cohort study from the Chin-Shan community in Taiwan. *BMJ Open.* 2022; **12**: e054638. 10.1136/bmjopen-2021-054638.

32 Wilhelmsen L, Svardsudd K, Korsan-Bengtson K, Larsson B, Welin L, Tibblin G. Fibrinogen as a risk factor for stroke and myocardial infarction. *N Engl J Med.* 1984; **311**: 501-5. 10.1056/NEJM198408233110804.

33 Rudnicka AR, Mt-Isa S, Meade TW. Associations of plasma fibrinogen and factor VII clotting activity with coronary heart disease and stroke: prospective cohort study from the screening phase of the Thrombosis Prevention Trial. *J Thromb Haemost.* 2006; **4**: 2405-10. 10.1111/j.1538-7836.2006.02221.x.

34 Lee AJ, Lowe GD, Smith WC, Tunstall-Pedoe H. Plasma fibrinogen in women: relationships with oral contraception, the menopause and hormone replacement therapy. *Br J Haematol.* 1993; **83**: 616-21. 10.1111/j.1365-2141.1993.tb04699.x.

35 Olson NC, Cushman M, Judd SE, Kissela BM, Safford MM, Howard G, Zakai NA. Associations of coagulation factors IX and XI levels with incident coronary heart disease and ischemic stroke: the REGARDS study. *J Thromb Haemost.* 2017; **15**: 1086-94. 10.1111/jth.13698.

36 Dekkers IA, de Mutsert R, de Vries APJ, Rosendaal FR, Cannegieter SC, Jukema JW, le Cessie S, Rabelink TJ, Lamb HJ, Lijfering WM. Determinants of impaired renal and vascular function are associated with elevated levels of procoagulant factors in the general population. *Journal of thrombosis and haemostasis : JTH.* 2018; **16**: 519-28. 10.1111/jth.13935.

- 37 Hemker HC, Giesen P, Al Dieri R, Regnault V, de Smedt E, Wagenvoord R, Lecompte T, Beguin S. Calibrated automated thrombin generation measurement in clotting plasma. *Pathophysiol Haemost Thromb*. 2003; **33**: 4-15. 10.1159/000071636.
- 38 Buis DTP, Christen T, Smit RAJ, de Mutsert R, Jukema JW, Cannegieter SC, Lijfering WM, Rosendaal FR. The association between leptin concentration and blood coagulation: Results from the NEO study. *Thrombosis research*. 2020; **188**: 44-8. 10.1016/j.thromres.2020.01.021.
- 39 Huxley VH, Kemp SS. Sex-Specific Characteristics of the Microcirculation. *Adv Exp Med Biol*. 2018; **1065**: 307-28. 10.1007/978-3-319-77932-4_20.
- 40 Brands J, Hubel CA, Althouse A, Reis SE, Pacella JJ. Noninvasive sublingual microvascular imaging reveals sex-specific reduction in glycocalyx barrier properties in patients with coronary artery disease. *Physiol Rep*. 2020; **8**: e14351. 10.14814/phy2.14351.
- 41 Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. New York: Routledge, 1988.
- 42 Song CJ, Nakagomi A, Chandar S, Cai H, Lim IG, McNeil HP, Freedman SB, Geczy CL. C-reactive protein contributes to the hypercoagulable state in coronary artery disease. *J Thromb Haemost*. 2006; **4**: 98-106. 10.1111/j.1538-7836.2005.01705.x.
- 43 Tantry US, Bliden KP, Suarez TA, Kreutz RP, Dichiara J, Gurbel PA. Hypercoagulability, platelet function, inflammation and coronary artery disease acuity: results of the Thrombotic Risk Progression (TRIP) study. *Platelets*. 2010; **21**: 360-7. 10.3109/09537100903548903.
- 44 Bratseth V, Pettersen AA, Opstad TB, Arnesen H, Seljeflot I. Markers of hypercoagulability in CAD patients. Effects of single aspirin and clopidogrel treatment. *Thromb J*. 2012; **10**: 12. 10.1186/1477-9560-10-12.
- 45 Fibrinogen Studies C, Danesh J, Lewington S, Thompson SG, Lowe GD, Collins R, Kostis JB, Wilson AC, Folsom AR, Wu K, Benderly M, Goldbourt U, Willeit J, Kiechl S, Yarnell JW, Sweetnam PM, Elwood PC, Cushman M, Psaty BM, Tracy RP, Tybjaerg-Hansen A, Haverkate F, de Maat MP, Fowkes FG, Lee AJ, Smith FB, Salomaa V, Harald K, Rasi R, Vahtera E, Jousilahti P, Pekkanen J, D'Agostino R, Kannel WB, Wilson PW, Tofler G, Arocha-Pinango CL, Rodriguez-Larralde A, Nagy E, Mijares M, Espinosa R, Rodriguez-Roa E, Ryder E, Diez-Ewald MP, Campos G, Fernandez V, Torres E, Marchioli R, Valagussa F, Rosengren A, Wilhelmsen L, Lappas G, Eriksson H, Cremer P, Nagel D, Curb JD, Rodriguez B, Yano K, Salonen JT, Nyyssonen K, Tuomainen TP, Hedblad B, Lind P, Loewel H, Koenig W, Meade TW, Cooper JA, De Stavola B, Knottenbelt C, Miller GJ, Cooper JA, Bauer KA, Rosenberg RD, Sato S, Kitamura A, Naito Y, Palosuo T, Ducimetiere P, Amouyel P, Arveiler D, Evans AE, Ferrieres J, Juhan-Vague I, Bingham A, Schulte H, Assmann G, Cantin B, Lamarche B, Despres JP, Dagenais GR, Tunstall-Pedoe H, Woodward M, Ben-Shlomo Y, Davey Smith G, Palmieri V, Yeh JL, Rudnicka A, Ridker P, Rodeghiero F, Tosetto A, Shepherd J, Ford I,

Robertson M, Brunner E, Shipley M, Feskens EJ, Kromhout D, Dickinson A, Ireland B, Juzwishin K, Kaptoge S, Lewington S, Memon A, Sarwar N, Walker M, Wheeler J, White I, Wood A. Plasma fibrinogen level and the risk of major cardiovascular diseases and nonvascular mortality: an individual participant meta-analysis. *JAMA*. 2005; **294**: 1799-809. 10.1001/jama.294.14.1799.

46 Zakai NA, Judd SE, Kissela B, Howard G, Safford MM, Cushman M. Factor VIII, Protein C and Cardiovascular Disease Risk: The REasons for Geographic and Racial Differences in Stroke Study (REGARDS). *Thromb Haemost*. 2018; **118**: 1305-15. 10.1055/s-0038-1655766.

47 Federici AB. The factor VIII/von Willebrand factor complex: basic and clinical issues. *Haematologica*. 2003; **88**: EREPO2.

48 Kalagara T, Moutsis T, Yang Y, Pappelbaum KI, Farken A, Cladder-Micus L, Vidal YSS, John A, Bauer AT, Moerschbacher BM, Schneider SW, Gorzelanny C. The endothelial glycocalyx anchors von Willebrand factor fibers to the vascular endothelium. *Blood Adv*. 2018; **2**: 2347-57. 10.1182/bloodadvances.2017013995.

49 Rietveld IM, Lijfering WM, le Cessie S, Bos MHA, Rosendaal FR, Reitsma PH, Cannegieter SC. High levels of coagulation factors and venous thrombosis risk: strongest association for factor VIII and von Willebrand factor. *J Thromb Haemost*. 2019; **17**: 99-109. 10.1111/jth.14343.

50 Foley JH, Conway EM. Cross Talk Pathways Between Coagulation and Inflammation. *Circ Res*. 2016; **118**: 1392-408. 10.1161/CIRCRESAHA.116.306853.

51 Rho R, Tracy RP, Bovill EG, Ball SP, Becker RC. Plasma Markers of Procoagulant Activity Among Individuals with Coronary Artery Disease. *J Thromb Thrombolysis*. 1995; **2**: 239-43. 10.1007/BF01062716.

52 Tosi F, Micaglio R, Sandri M, Castagna A, Minguzzi D, Stefanoni F, Chiariello C, Franzese I, Luciani GB, Faggian G, Girelli D, Olivieri O, Martinelli N. Increased plasma thrombin potential is associated with stable coronary artery disease: An angiographically-controlled study. *Thromb Res*. 2017; **155**: 16-22. 10.1016/j.thromres.2017.04.021.

53 Bol ME, Broddin BEK, Delhaas T, Sels JEM, van de Poll MCG. Variability of microcirculatory measurements in healthy volunteers. *Sci Rep*. 2022; **12**: 19887. 10.1038/s41598-022-22947-x.

54 Dolleman SC, Agten SM, Spronk HMH, Hackeng TM, Bos MHA, Versteeg HH, van Zonneveld AJ, de Boer HC. Thrombin in complex with dabigatran can still interact with PAR-1 via exosite-I and instigate loss of vascular integrity. *J Thromb Haemost*. 2022; **20**: 996-1007. 10.1111/jth.15642.

55 Ikezoe T. Thrombomodulin/activated protein C system in septic disseminated intravascular coagulation. *J Intensive Care*. 2015; **3**: 1. 10.1186/s40560-014-0050-7.

- 56 Pluskota E, D'Souza SE. Fibrinogen interactions with ICAM-1 (CD54) regulate endothelial cell survival. *Eur J Biochem.* 2000; **267**: 4693-704. 10.1046/j.1432-1327.2000.01520.x.
- 57 Tsakadze NL, Zhao Z, D'Souza SE. Interactions of intercellular adhesion molecule-1 with fibrinogen. *Trends Cardiovasc Med.* 2002; **12**: 101-8. 10.1016/s1050-1738(01)00157-8.
- 58 Kossmann S, Lagrange J, Jackel S, Jurk K, Ehlken M, Schonfelder T, Weihert Y, Knorr M, Brandt M, Xia N, Li H, Daiber A, Oelze M, Reinhardt C, Lackner K, Gruber A, Monia B, Karbach SH, Walter U, Ruggeri ZM, Renne T, Ruf W, Munzel T, Wenzel P. Platelet-localized FXI promotes a vascular coagulation-inflammatory circuit in arterial hypertension. *Sci Transl Med.* 2017; **9**. 10.1126/scitranslmed.aah4923.
- 59 Shariat-Madar Z, Mahdi F, Schmaier AH. Factor XI assembly and activation on human umbilical vein endothelial cells in culture. *Thromb Haemost.* 2001; **85**: 544-51.

Supporting information




Figure S1. Association between glycoalyx-related SDF imaging parameters and levels of coagulation parameters in total population. Differences in the total population between the glycoalyx-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + sex + age; Model 3: model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: PBR_{Total}: PBR of total vessels from 4-25 μ m; PBR_{feed vessel}: PBR of feed vessels from 10-19 μ m; PBR_{capillary}: PBR of capillaries from 4-9 μ m.

Table S1. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in total population.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VellIndex	Fibrinogen
PBR _{Total}	Model 1	0.3 (-1.2-1.7)	0.3 (-1.1-1.7)	0 (-2.3-2.4)	-0.1 (-0.2-0)	-0.1 (-0.3-0)	1.4 (-1.2-3.9)	8.2 (-16.7-33.1)	0 (0-0.1)	4 (-0.1-8)
	Model 2	-0.2 (-1.7-1.2)	0.3 (-1.1-1.7)	0.2 (-2.2-2.5)	-0.1 (-0.2-0)	-0.1 (-0.2-0)	0.8 (-1.7-3.3)	1.7 (-22.9-26.3)	0 (0-0.1)	3 (-0.9-7)
	Model 3	0.1 (-1.3-1.6)	0.9 (-0.4-2.1)	0 (-2.3-2.4)	0 (-0.1-0.1)	-0.1 (-0.2-0.1)	1 (-1.3-3.2)	3.4 (-19.5-26.4)	0 (0-0.1)	4.4 (0.7-8.1)
	Model 4	0 (-1.4-1.4)	0.5 (-0.5-1.6)	-0.3 (-2.6-1.9)	-0.1 (-0.1-0)	-0.1 (-0.2-0.1)	0.5 (-1.7-2.8)	-0.8 (-23.2-21.6)	0 (0-0.1)	3 (-0.3-6.2)
PBR _{feed vessel}	Model 1	0.9 (-0.6-2.3)	0.2 (-1.2-1.6)	0.2 (-2.2-2.5)	-0.1 (-0.2-0)	-0.1 (-0.3-0)	1.1 (-1.4-3.7)	6.1 (-18.8-31)	0 (0-0.1)	3.7 (-0.4-7.7)
	Model 2	0.4 (-1.1-1.8)	0.2 (-1.2-1.5)	0.2 (-2.1-2.6)	-0.1 (-0.2-0)	-0.1 (-0.2-0)	0.6 (-1.9-3.1)	0.2 (-24.4-24.9)	0 (0-0.1)	2.6 (-1.4-6.6)
	Model 3	0.7 (-0.7-2.1)	0.7 (-0.5-1.9)	0.2 (-2.1-2.6)	0 (-0.1-0.1)	-0.1 (-0.2-0)	0.9 (-1.4-3.2)	3 (-19.9-25.9)	0 (0-0.1)	3.9 (0.2-7.6)
	Model 4	0.7 (-0.7-2)	0.6 (-0.5-1.7)	0 (-2.3-2.3)	0 (-0.1-0.1)	-0.1 (-0.2-0.1)	0.6 (-1.6-2.8)	0 (-22.4-22.3)	0 (0-0.1)	3.2 (0-6.4)
PBR _{capillary}	Model 1	-0.2 (-1.6-1.3)	0.1 (-1.3-1.4)	1.1 (-1.2-3.5)	0 (-0.1-0.1)	0 (-0.1-0.1)	-1.5 (-4-1)	-12 (-36.9-12.9)	0 (-0.1-0)	0.7 (-3.3-4.8)
	Model 2	0 (-1.5-1.4)	0.2 (-1.2-1.5)	1.5 (-0.8-3.8)	0 (-0.1-0.1)	0 (-0.1-0.2)	-1.9 (-4.4-0.6)	-16 (-40.5-8.5)	0 (-0.1-0)	1.5 (-2.5-5.4)
	Model 3	0.1 (-1.3-1.5)	0.5 (-0.7-1.7)	1.4 (-0.9-3.7)	0 (-0.1-0.1)	0 (-0.1-0.1)	-1.4 (-3.7-0.9)	-10.9 (-33.7-11.9)	0 (-0.1-0)	2.3 (-1.4-6)
	Model 4	0.1 (-1.2-1.5)	0.5 (-0.6-1.6)	1.4 (-0.8-3.7)	0 (-0.1-0.1)	0 (-0.1-0.1)	-1.4 (-3.6-0.8)	-10.6 (-32.8-11.6)	0 (-0.1-0)	2.1 (-1.1-5.3)

Results are based on linear regression analyses of the general Dutch population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + sex + age.

Model 3 = model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: PBR_{Total}: PBR of total vessels from 4-25µm; PBR_{feed vessel}: PBR of feed vessels from 10-19µm; PBR_{capillary}: PBR of capillaries from 4-9µm.




Figure S2. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in total population. Differences in the total population between the microcirculatory-perfusion-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + sex + age; Model 3: model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.

Table S2. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in total population.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
RBCV _{feed vessel}	Model 1	0 (-1.5-1.4)	1.6 (0.2-2.9)	2 (-0.3-4.3)	0.1 (0-0.2)	0 (-0.1-0.2)	1.8 (-0.7-4.4)	20.6 (-4.3-45.4)	0 (0-0.1)	-0.1 (-4.1-4)
	Model 2	1.1 (-0.4-2.5)	1.8 (0.4-3.2)	2.1 (-0.3-4.5)	0 (-0.1-0.1)	-0.1 (-0.2-0.1)	2.9 (0.4-5.5)	33.4 (8.3-58.5)	0.1 (0-0.1)	2.6 (-1.5-6.6)
	Model 3	0.5 (-0.9-2)	0.5 (-0.8-1.8)	2 (-0.4-4.4)	0 (-0.1-0.1)	-0.1 (-0.2-0)	1.8 (-0.5-4.2)	21.6 (-2-45.2)	0 (0-0.1)	-0.3 (-4.2-3.5)
	Model 4	0.3 (-1.1-1.7)	0.1 (-1-1.2)	1.9 (-0.5-4.2)	-0.1 (-0.2-0)	-0.1 (-0.3-0)	1.8 (-0.4-4.1)	22 (-1.1-45)	0 (0-0.1)	-1.3 (-4.7-2)
RBCV _{capillary}	Model 1	-0.1 (-1.6-1.3)	1.2 (-0.1-2.6)	1.8 (-0.5-4.2)	0.1 (0-0.2)	0 (-0.1-0.2)	1.4 (-1.1-4)	16.1 (-8.7-41)	0 (0-0.1)	0.1 (-4-4.1)
	Model 2	0.9 (-0.6-2.4)	1.4 (0-2.8)	2 (-0.4-4.4)	0 (-0.1-0.1)	-0.1 (-0.2-0.1)	2.3 (-0.2-4.9)	27 (1.9-52)	0 (0-0.1)	2.7 (-1.3-6.7)
	Model 3	0.5 (-1.1-1.9)	0.4 (-0.8-1.7)	1.9 (-0.5-4.3)	0 (-0.1-0.1)	-0.1 (-0.2-0.1)	1.3 (-1-3.7)	17.1 (-6.3-40.5)	0 (0-0.1)	0.7 (-3.1-4.6)
	Model 4	0.4 (-1-1.8)	0.3 (-0.8-1.4)	1.9 (-0.4-4.2)	0 (-0.1-0.1)	-0.1 (-0.2-0.1)	1.4 (-0.9-3.6)	17.5 (-5.3-40.4)	0 (0-0.1)	0.4 (-2.9-3.6)
D _{Total}	Model 1	0.7 (-0.7-2.2)	1.1 (-0.3-2.5)	2.2 (-0.1-4.5)	0 (-0.1-0.1)	0 (-0.1-0.2)	2.7 (0.2-5.3)	39.5 (14.7-64.2)	0 (0-0.1)	3.2 (-0.8-7.3)
	Model 2	0.7 (-0.7-2.2)	1 (-0.3-2.4)	1.8 (-0.5-4.1)	0 (-0.1-0.1)	0 (-0.1-0.1)	3.3 (0.8-5.8)	45.3 (21-69.7)	0 (0-0.1)	2.7 (-1.2-6.7)
	Model 3	0.3 (-1.1-1.7)	0 (-1.2-1.2)	1.6 (-0.7-4)	0 (-0.1-0.1)	0 (-0.1-0.1)	1.9 (-0.3-4.2)	32.8 (10-55.7)	0 (0-0.1)	0.8 (-3-4.5)
	Model 4	0.3 (-1.1-1.7)	0 (-1.1-1.1)	1.6 (-0.7-3.9)	0 (-0.1-0.1)	0 (-0.1-0.1)	1.9 (-0.4-4.1)	31.9 (9.6-54.2)	0 (0-0.1)	0.7 (-2.6-3.9)
D _{feed vessel}	Model 1	0.5 (-0.9-2)	1.6 (0.3-3)	2.4 (0-4.7)	0.1 (0-0.2)	0.1 (0-0.2)	1.7 (-0.8-4.2)	28.9 (4.1-53.7)	0 (0-0.1)	2.9 (-1.1-7)
	Model 2	0.8 (-0.7-2.2)	1.6 (0.2-2.9)	1.9 (-0.5-4.2)	0 (-0.1-0.1)	0.1 (-0.1-0.2)	2.6 (0.1-5.1)	38.6 (14.1-63.1)	0 (0-0.1)	2.9 (-1-6.9)
	Model 3	0.4 (-1.1-1.8)	0.5 (-0.8-1.7)	1.7 (-0.7-4)	0 (-0.1-0.1)	0.1 (-0.1-0.2)	1.1 (-1.2-3.4)	25.4 (2.4-48.4)	0 (0-0.1)	1 (-2.7-4.8)
	Model 4	0.2 (-1.2-1.6)	0.2 (-0.9-1.2)	1.4 (-0.9-3.7)	0 (-0.1-0.1)	0.1 (-0.1-0.2)	0.9 (-1.3-3.2)	23.7 (1.2-46.2)	0 (0-0.1)	0.7 (-2.6-3.9)
D _{capillary}	Model 1	0.6 (-0.8-2.1)	-0.1 (-1.5-1.2)	0.8 (-1.6-3.1)	-0.1 (-0.2-0)	-0.1 (-0.2-0.1)	2.5 (0-5)	32.6 (7.8-57.4)	0 (0-0.1)	2.2 (-1.9-6.2)
	Model 2	0.3 (-1.2-1.7)	-0.2 (-1.6-1.2)	0.7 (-1.6-3)	0 (-0.1-0)	0 (-0.2-0.1)	2.3 (-0.2-4.8)	30.2 (5.8-54.6)	0 (0-0.1)	1.2 (-2.7-5.2)
	Model 3	0.1 (-1.3-1.5)	-0.6 (-1.8-0.6)	0.6 (-1.7-3)	-0.1 (-0.2-0)	0 (-0.2-0.1)	1.8 (-0.5-4)	24.5 (1.7-47.3)	0 (0-0.1)	0.3 (-3-4.4)
	Model 4	0.3 (-1.1-1.7)	-0.2 (-1.3-0.8)	0.8 (-1.4-3.1)	0 (-0.1-0.1)	0 (-0.2-0.1)	1.8 (-0.4-4.1)	25.2 (2.9-47.5)	0 (0-0.1)	0.5 (-2.7-3.7)
CBV _{static}	Model 1	-0.3 (-1.7-1.2)	-1.5 (-2.9-0.1)	-0.2 (-2.5-2.1)	0 (-0.1-0.1)	0 (-0.2-0.1)	0.6 (-1.9-3.1)	2.6 (-22.3-27.5)	0 (0-0.1)	0.7 (-3-4-7)
	Model 2	-0.7 (-2.1-0.7)	-1.6 (-3-0.2)	-0.5 (-2.8-1.9)	0 (-0.1-0.1)	0 (-0.2-0.1)	0.6 (-1.9-3.1)	2 (-22.5-26.6)	0 (0-0.1)	-0.6 (-4.6-3-3)
	Model 3	-0.6 (-2-0.8)	-1 (-2.2-0.2)	-0.4 (-2.7-2)	0 (-0.1-0.1)	0 (-0.2-0.1)	1.5 (-0.8-3.8)	10.5 (-12.3-33.4)	0 (0-0.1)	0.5 (-3-3-4.2)
	Model 4	-0.4 (-1.7-1)	-0.6 (-1.7-0.4)	-0.2 (-2.5-2.1)	0 (-0.1-0.1)	0 (-0.1-0.1)	1.5 (-0.7-3.7)	10.5 (-11.8-32.9)	0 (0-0.1)	0.8 (-2-4-4)

Results are based on linear regression analyses of the general Dutch population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + sex + age.

Model 3 = model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.

Table S3. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in women.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
PBR _{Total}	Model 1	0.4 (-1.7-2.6)	1.2 (-0.9-3.2)	0.8 (-2.3-4)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	3 (-0.9-6.9)	29 (-8.6-66.7)	0 (0-0.1)	5 (-0.8-10.7)
	Model 2	0.7 (-1.5-2.8)	1.3 (-0.8-3.4)	1.4 (-1.7-4.6)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	1.9 (-1.9-5.7)	16.3 (-20-52.6)	0 (0-0.1)	5.5 (-0.3-11.3)
	Model 3	1.2 (-0.8-3.3)	2.3 (0.5-4)	1.3 (-1.9-4.5)	0 (-0.2-0.1)	-0.1 (-0.2-0.1)	2.6 (-0.7-5.9)	22.8 (-10-55.6)	0 (0-0.1)	7.5 (2.2-12.9)
	Model 4	1.2 (-0.9-3.2)	1.8 (0.3-3.3)	0.5 (-2.5-3.6)	0 (-0.2-0.1)	0 (-0.2-0.1)	1.8 (-1.5-5)	15.5 (-16.8-47.8)	0 (0-0.1)	5.1 (0.4-9.9)
PBR _{feed vessel}	Model 1	0.9 (-1.2-3.1)	1 (-1.1-3.1)	1.3 (-1.9-4.4)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	2.8 (-1.1-6.7)	25.8 (-11.9-63.5)	0 (0-0.1)	4.5 (-1.2-10.3)
	Model 2	1.1 (-1-3.3)	1.1 (-1-3.2)	1.8 (-1.3-4.9)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	1.8 (-2-5.6)	14.3 (-21.9-50.6)	0 (0-0.1)	5 (-0.8-10.8)
	Model 3	1.7 (-0.4-3.7)	2.2 (0.4-3.9)	1.8 (-1.3-5)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	2.8 (-0.5-6.1)	24 (-8.7-56.7)	0 (0-0.1)	7.1 (1.7-12.4)
	Model 4	1.7 (-0.4-3.7)	2 (0.5-3.4)	1.4 (-1.7-4.5)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	2.2 (-1-5.4)	19 (-13.1-51)	0 (0-0.1)	5.8 (1.1-10.6)
PBR _{capillary}	Model 1	-1.1 (-3.2-1.1)	0.4 (-1.7-2.5)	3 (-0.1-6.2)	0 (-0.2-0.1)	0 (-0.2-0.2)	-0.5 (-4.4-3.4)	0.1 (-37.7-37.8)	0 (-0.1-0.1)	3.7 (-2-9.5)
	Model 2	-1 (-3.1-1.2)	0.5 (-1.6-2.6)	3.4 (0.2-6.5)	0 (-0.1-0.1)	0 (-0.2-0.2)	-1.2 (-4.9-2.6)	-7.5 (-43.6-28.6)	0 (-0.1-0.1)	4 (-1.7-9.8)
	Model 3	-0.7 (-2.7-1.4)	1 (-0.7-2.8)	3.3 (0.2-6.4)	0 (-0.1-0.1)	0 (-0.2-0.2)	-0.4 (-3.7-2.9)	-0.5 (-33.1-32.1)	0 (-0.1-0.1)	5 (-0.4-10.3)
	Model 4	-0.5 (-2.6-1.5)	1.3 (-0.2-2.8)	3.5 (0.4-6.5)	0 (-0.1-0.1)	0 (-0.2-0.2)	-0.3 (-3.5-2.9)	0.3 (-31.6-32.2)	0 (-0.1-0.1)	5.3 (0.6-10)

Results are based on linear regression analyses of the Dutch women population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: PBR_{Total}: PBR of total vessels from 4-25µm; PBR_{feed vessel}: PBR of feed vessels from 10-19µm; PBR_{capillary}: PBR of capillaries from 4-9µm.

Table S4. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in men.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
PBR _{Total}	Model 1	-1.1 (-3-0.7)	-0.9 (-2.7-0.8)	-1.2 (-4.7-2.3)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.5 (-4.5-1.6)	-26.8 (-58.1-4.5)	0 (-0.1-0)	-0.1 (-5.6-5.3)
	Model 2	-1.1 (-2.9-0.8)	-0.9 (-2.7-0.8)	-1.4 (-4.9-2)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.4 (-4.5-1.6)	-27 (-58.4-4.4)	0 (-0.1-0)	-0.6 (-5.9-4.8)
	Model 3	-1 (-2.8-0.9)	-0.7 (-2.4-1)	-1.5 (-5-1.9)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.3 (-4.4-1.8)	-25.1 (-56.3-6.2)	0 (-0.1-0)	0.2 (-4.9-5.3)
	Model 4	-1.1 (-2.8-0.7)	-0.8 (-2.3-0.7)	-1.5 (-4.9-1.9)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.3 (-4.3-1.8)	-24.5 (-55-6)	0 (-0.1-0)	0.2 (-4-4.4)
PBR _{feed vessel}	Model 1	-0.3 (-2.1-1.6)	-0.9 (-2.6-0.9)	-1.3 (-4.8-2.2)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.5 (-4.6-1.5)	-25.8 (-57.1-5.5)	0 (-0.1-0)	-0.1 (-5.6-5.3)
	Model 2	-0.2 (-2-1.6)	-0.9 (-2.7-0.8)	-1.5 (-5-1.9)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.5 (-4.6-1.6)	-26 (-57.4-5.4)	0 (-0.1-0.1)	-0.6 (-6-4.7)
	Model 3	-0.2 (-2-1.7)	-0.8 (-2.5-0.9)	-1.6 (-5-1.9)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	-1.5 (-4.5-1.6)	-24.9 (-56.1-6.3)	0 (-0.1-0.1)	-0.1 (-5-2.5)
	Model 4	-0.1 (-1.9-1.6)	-0.8 (-2.3-0.7)	-1.7 (-5.1-1.8)	0 (-0.2-0.1)	-0.1 (-0.2-0.1)	-1.5 (-4.6-1.5)	-25.6 (-56.1-4.9)	0 (-0.1-0)	-0.3 (-4.5-3.9)
PBR _{capillary}	Model 1	1.2 (-0.7-3)	-0.3 (-2.1-1.4)	-1.1 (-4.6-2.4)	-0.1 (-0.2-0.1)	0 (-0.2-0.2)	-2.6 (-5.7-0.4)	-24.8 (-56.1-6.5)	0 (-0.1-0)	-2.4 (-7.8-3.1)
	Model 2	1 (-0.8-2.9)	-0.2 (-2-1.5)	-0.7 (-4.1-2.8)	0 (-0.2-0.1)	0 (-0.2-0.2)	-2.7 (-5.8-0.4)	-24.7 (-56.1-6.8)	0 (-0.1-0)	-1.5 (-6.8-3.9)
	Model 3	1.1 (-0.7-3)	-0.1 (-1.7-1.6)	-0.8 (-4.3-2.7)	0 (-0.2-0.1)	0 (-0.2-0.2)	-2.5 (-5.6-0.6)	-22.5 (-53.9-8.9)	0 (-0.1-0)	-0.8 (-5.9-4.3)
	Model 4	0.9 (-0.9-2.7)	-0.3 (-1.8-1.2)	-0.8 (-4.3-2.6)	-0.1 (-0.2-0.1)	0 (-0.2-0.2)	-2.6 (-5.6-0.4)	-23.5 (-54.2-7.2)	0 (-0.1-0)	-1.6 (-5.8-2.6)

Results are based on linear regression analyses of the Dutch men population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI + current smoking status.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: PBR_{Total}: PBR of total vessels from 4-25µm; PBR_{feed vessel}: PBR of feed vessels from 10-19µm; PBR_{capillary}: PBR of capillaries from 4-9µm.




Figure S3. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in women. After sex stratification, differences in women between the microcirculatory-perfusion-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.

Table S5. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in women.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	Vellindex	Fibrinogen
RBCV _{feed vessel}	Model 1	1.6 (-0.6-3.7)	2.2 (0.1-4.2)	2.2 (-1.5-3)	0 (-0.1-0.2)	0 (-0.2-0.2)	2.3 (-1.6-6.2)	29.3 (-8.3-67)	0 (0-0.1)	3.1 (-2.7-8.8)
	Model 2	1.6 (-0.6-3.7)	2.2 (0.1-4.2)	2.1 (-1.5-2)	0 (-0.1-0.2)	0 (-0.2-0.2)	2.4 (-1.4-6.1)	30.7 (-5.2-66.7)	0 (0-0.1)	3 (-2.7-8.8)
	Model 3	0.7 (-1.4-2.7)	0.6 (-1.2-2.3)	1.8 (-1.3-5)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.7 (-2.6-4)	14.8 (-18-47.7)	0 (-0.1-0.1)	-0.2 (-5.6-5.3)
	Model 4	0.1 (-1.9-2.2)	-0.6 (-2-0.9)	1.5 (-1.6-4.6)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.7 (-2.5-4)	15.6 (-16.9-48)	0 (-0.1-0.1)	-2.3 (-7.1-2.5)
RBCV _{capillary}	Model 1	1.5 (-0.6-3.7)	1.6 (-0.5-3.6)	2.2 (-1.5-3)	0 (-0.1-0.2)	0 (-0.2-0.2)	1.6 (-2.3-5.6)	24.6 (-13.1-62.3)	0 (-0.1-0.1)	2.7 (-3.1-8.5)
	Model 2	1.5 (-0.6-3.6)	1.5 (-0.5-3.6)	2 (-1.1-5.1)	0 (-0.1-0.2)	0 (-0.2-0.2)	1.9 (-1.8-5.7)	27.9 (-8.1-63.9)	0 (0-0.1)	2.6 (-3.2-8.3)
	Model 3	0.8 (-1.3-2.8)	0.3 (-1.5-2)	1.8 (-1.4-4.9)	0 (-0.1-0.1)	0 (-0.1-0.2)	0.2 (-3.1-3.5)	12.6 (-20.1-45.3)	0 (-0.1-0.1)	0.5 (-4.9-5.9)
	Model 4	0.5 (-1.6-2.5)	-0.4 (-1.9-1.1)	1.5 (-1.6-4.6)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.2 (-3.1-3.4)	12.6 (-19.5-44.7)	0 (-0.1-0.1)	-1 (-5.8-3.7)
D _{Total}	Model 1	1 (-1.1-3.2)	2 (-0.1-4)	1.7 (-1.5-4.9)	0 (-0.1-0.2)	0 (-0.2-0.2)	3.5 (-0.4-7.4)	46.2 (8.7-83.7)	0 (0-0.1)	3.9 (-1.8-9.7)
	Model 2	0.9 (-1.3-3)	1.9 (-0.2-4)	1.3 (-1.8-4.5)	0 (-0.1-0.1)	0 (-0.2-0.2)	4.3 (0.5-8.1)	55.3 (19.5-91.1)	0.1 (0-0.1)	3.7 (-2.1-9.4)
	Model 3	0 (-2.1-2.1)	0.2 (-1.5-2)	0.9 (-2.3-4)	0 (-0.2-0.1)	0 (-0.2-0.2)	1.8 (-1.6-5.1)	33.2 (0.3-66.1)	0 (-0.1-0.1)	1 (-4.5-6.4)
	Model 4	-0.1 (-2.2-1.9)	0 (-1.5-1.5)	0.6 (-2.5-3.7)	0 (-0.2-0.1)	0 (-0.2-0.2)	1.5 (-1.7-4.8)	31 (-1.2-63.3)	0 (-0.1-0.1)	0.6 (-4.2-5.4)
D _{feed vessel}	Model 1	1.4 (-0.8-3.5)	2.5 (0.5-4.6)	2.3 (-0.8-5.5)	0.1 (-0.1-0.2)	0.1 (-0.1-0.3)	2.1 (-1.8-6)	34.4 (-3.2-72)	0 (-0.1-0.1)	3.4 (-2.3-9.2)
	Model 2	1.2 (-1-3.3)	2.5 (0.4-4.5)	1.8 (-1.3-5)	0 (-0.1-0.2)	0.1 (-0.1-0.3)	3.3 (-0.5-7.1)	46.9 (10.9-82.9)	0 (0-0.1)	3 (-2.8-8.8)
	Model 3	0.3 (-1.8-2.4)	0.6 (-1.1-2.4)	1.3 (-1.8-4.5)	0 (-0.1-0.2)	0.1 (-0.1-0.3)	0.5 (-2.9-3.8)	22.8 (-10.3-55.8)	0 (-0.1-0.1)	0.3 (-5.2-5.7)
	Model 4	0.1 (-2-2.1)	0.1 (-1.4-1.6)	0.8 (-2.3-3.9)	0 (-0.1-0.1)	0.1 (-0.1-0.3)	-0.1 (-3.3-3.2)	17.8 (-14.7-50.3)	0 (-0.1-0)	-0.5 (-5.3-4.3)
D _{capillary}	Model 1	0.1 (-2-2.3)	0.4 (-1.7-2.4)	-0.3 (-3.4-2.9)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	3.6 (-0.3-7.5)	40.7 (3.1-78.3)	0.1 (0-0.1)	3.2 (-2.6-8.9)
	Model 2	0.1 (-2-2.3)	0.4 (-1.7-2.4)	-0.2 (-3.4-2.9)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	3.5 (-0.2-7.3)	40 (4.2-75.9)	0.1 (0-0.1)	3.2 (-2.5-9)
	Model 3	-0.4 (-2.4-1.7)	-0.3 (-2.1-1.4)	-0.5 (-3.6-2.7)	-0.1 (-0.2-0)	-0.1 (-0.3-0.1)	2.5 (-0.8-5.7)	30.3 (-2-62.9)	0 (0-0.1)	1.9 (-3.4-7.3)
	Model 4	-0.2 (-2.3-1.8)	0 (-1.5-1.4)	-0.3 (-3.4-2.8)	-0.1 (-0.2-0)	-0.1 (-0.3-0.1)	2.6 (-0.6-5.8)	32 (0.1-63.8)	0 (0-0.1)	2.2 (-2.5-7)
CBV _{static}	Model 1	-0.6 (-2.7-1.6)	-1.6 (-3.7-0.5)	-1 (-4.1-2.2)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.7 (-3.2-4.6)	0.9 (-36.9-38.7)	0 (-0.1-0.1)	0.1 (-5.6-5.9)
	Model 2	-0.7 (-2.9-1.4)	-1.7 (-3.7-0.4)	-1.3 (-4.5-1.8)	0 (-0.2-0.1)	0 (-0.2-0.2)	1.5 (-2.3-5.3)	9.1 (-27-45.3)	0 (-0.1-0.1)	-0.2 (-5.9-5.6)
	Model 3	-0.5 (-2.6-1.5)	-0.8 (-2.5-1)	-1.2 (-4.3-2)	0 (-0.2-0.1)	-0.1 (-0.2-0.1)	2.8 (-0.5-6.1)	21.1 (-11.5-53.8)	0 (0-0.1)	1.2 (-4.2-6.6)
	Model 4	-0.3 (-2.3-1.8)	-0.2 (-1.6-1.3)	-0.8 (-3.9-2.3)	0 (-0.1-0.1)	0 (-0.2-0.1)	3.1 (-0.1-6.2)	23.5 (-8.5-55.5)	0.1 (0-0.1)	2.7 (-2.1-7.4)

Results are based on linear regression analyses of the Dutch men population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI, current smoking status, menopausal status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.




Figure S4. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in men. After sex stratification, differences in men between the microcirculatory-perfusion-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI + current smoking status; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.

Table S6. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in men.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
RBCV _{feed vessel}	Model 1	0.5 (-1.3-2.4)	1.3 (-0.4-3.1)	2.3 (-1.2-5.7)	-0.1 (-0.2-0.1)	-0.2 (-0.3-0)	3 (-0.1-6.1)	30.4 (-0.9-61.6)	0.1 (0-0.1)	2.5 (-2.9-7.9)
	Model 2	0.6 (-1.2-2.5)	1.2 (-0.5-3)	1.9 (-1.5-5.4)	-0.1 (-0.2-0.1)	-0.2 (-0.4-0)	3 (0-6.1)	30.3 (-1.1-61.7)	0.1 (0-0.1)	1.7 (-3.6-7.1)
	Model 3	0.5 (-1.4-2.3)	0.5 (-1.2-2.2)	2.1 (-1.4-5.6)	-0.1 (-0.2-0)	-0.2 (-0.4-0)	2.9 (-0.2-6)	26.5 (-5.1-58)	0.1 (0-0.1)	-0.6 (-5.7-4.6)
	Model 4	0.4 (-1.4-2.2)	0.4 (-1.1-1.9)	2 (-1.5-5.4)	-0.1 (-0.2-0)	-0.2 (-0.4-0)	2.8 (-0.3-5.8)	25.5 (-5.4-56.3)	0.1 (0-0.1)	-1 (-5.2-3.3)
RBCV _{capillary}	Model 1	0.2 (-1.7-2)	1.2 (-0.5-3)	2 (-1.5-5.5)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	2.7 (-0.4-5.8)	25 (-6.3-56.3)	0.1 (0-0.1)	2.9 (-2.5-8.3)
	Model 2	0.2 (-1.6-2.1)	1.2 (-0.6-2.9)	1.9 (-1.6-5.3)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	2.7 (-0.4-5.8)	25 (-6.4-56.3)	0.1 (0-0.1)	2.7 (-2.6-8)
	Model 3	0.1 (-1.7-2)	0.6 (-1-2.3)	2 (-1.5-5.4)	-0.1 (-0.2-0.1)	-0.2 (-0.4-0)	2.6 (-0.5-5.7)	22.7 (-8.7-54.1)	0.1 (0-0.1)	1.1 (-4-6.2)
	Model 4	0.3 (-1.5-2.1)	0.8 (-0.7-2.4)	2 (-1.5-5.4)	0 (-0.2-0.1)	-0.2 (-0.3-0)	2.7 (-0.4-5.7)	23 (-7.7-53.8)	0.1 (0-0.1)	1.6 (-2.7-5.8)
D _{Total}	Model 1	0.4 (-1.4-2.3)	0.1 (-1.7-1.8)	2.8 (-0.7-6.3)	0 (-0.1-0.2)	0 (-0.2-0.2)	1.9 (-1.2-5)	32.2 (0.9-63.4)	0 (0-0.1)	2.6 (-2.9-8)
	Model 2	0.6 (-1.3-2.4)	0 (-1.8-1.8)	2.3 (-1.1-5.8)	0 (-0.1-0.2)	0 (-0.2-0.2)	2 (-1.1-5.1)	32.2 (0.7-63.6)	0 (0-0.1)	1.6 (-3.8-6.9)
	Model 3	0.5 (-1.3-2.4)	-0.3 (-2-1.3)	2.4 (-1.1-5.9)	0 (-0.1-0.1)	0 (-0.2-0.2)	1.9 (-1.2-5)	30.8 (-5.5-62.2)	0 (0-0.1)	0.6 (-4.6-5.7)
	Model 4	0.6 (-1.1-2.4)	-0.1 (-1.7-1.4)	2.7 (-0.7-6.1)	0 (-0.1-0.1)	0 (-0.2-0.2)	2.1 (-1-5.1)	31.9 (1.3-62.6)	0 (0-0.1)	1 (-3.2-5.3)
D _{feed vessel}	Model 1	0.2 (-1.7-2)	0.6 (-1.1-2.4)	2.5 (-1-6)	0 (-0.1-0.2)	0 (-0.1-0.2)	1.6 (-1.5-4.7)	27.4 (-3.9-58.7)	0 (0-0.1)	4 (-1.4-9.4)
	Model 2	0.4 (-1.5-2.2)	0.5 (-1.2-2.3)	1.9 (-1.6-5.4)	0 (-0.1-0.2)	0 (-0.2-0.2)	1.7 (-1.4-4.8)	27.4 (-4.2-59)	0 (0-0.1)	2.7 (-2.7-8.1)
	Model 3	0.3 (-1.5-2.2)	0.2 (-1.5-1.9)	1.9 (-1.6-5.4)	0 (-0.1-0.2)	0 (-0.2-0.2)	1.7 (-1.4-4.8)	26.4 (-5.1-57.9)	0 (0-0.1)	1.8 (-3-3.7)
	Model 4	0.2 (-1.6-2)	0.2 (-1.3-1.8)	2.1 (-1.4-5.5)	0 (-0.1-0.1)	0 (-0.2-0.2)	1.9 (-1.1-5)	29.6 (-1.2-60.4)	0 (0-0.1)	2.5 (-1.7-6.8)
D _{capillary}	Model 1	0.5 (-1.4-2.3)	-0.8 (-2.6-0.9)	1.9 (-1.6-5.3)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.8 (-2.3-3.9)	17.1 (-14.3-48.4)	0 (-0.1-0.1)	-1.1 (-6.5-4.4)
	Model 2	0.5 (-1.4-2.3)	-0.9 (-2.6-0.9)	1.8 (-1.7-5.2)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.8 (-2.3-3.9)	17 (-14.4-48.4)	0 (-0.1-0.1)	-1.2 (-6.5-4.1)
	Model 3	0.4 (-1.4-2.3)	-1 (-2.7-0.7)	1.8 (-1.6-5.3)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.7 (-2.3-3.8)	16.2 (-15-47.5)	0 (-0.1-0.1)	-1.6 (-6.7-3.4)
	Model 4	0.7 (-1-2.5)	-0.6 (-2.1-0.9)	2.2 (-1.2-5.6)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.7 (-2.3-3.8)	15.4 (-15.4-46.2)	0 (-0.1-0.1)	-1.4 (-5.6-2.8)
CBV _{static}	Model 1	-0.8 (-2.7-1)	-1.5 (-3.3-0.2)	0.6 (-2.8-4.1)	0 (-0.1-0.2)	0 (-0.2-0.2)	-0.1 (-3.2-3)	-2.2 (-33.6-29.2)	0 (-0.1-0.1)	-0.9 (-6.3-4.5)
	Model 2	-0.8 (-2.6-1)	-1.5 (-3.3-0.2)	0.6 (-2.8-4.1)	0 (-0.1-0.2)	0 (-0.2-0.2)	-0.1 (-3.2-3)	-2.2 (-33.7-29.2)	0 (-0.1-0.1)	-1 (-6.3-4.4)
	Model 3	-0.8 (-2.6-1.1)	-1.3 (-3-0.4)	0.6 (-2.9-4.1)	0 (-0.1-0.2)	0 (-0.2-0.2)	-0.1 (-3.2-3)	-1.5 (-32.8-29.8)	0 (-0.1-0.1)	-0.4 (-5.4-4.7)
	Model 4	-0.7 (-2.5-1.1)	-1.3 (-2.8-0.3)	0.7 (-2.7-4.1)	0 (-0.1-0.2)	0 (-0.2-0.2)	-0.4 (-3.5-2.7)	-5.9 (-36.8-24.9)	0 (-0.1-0.1)	-1.5 (-5.7-2.7)

Results are based on linear regression analyses of the Dutch men population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI + current smoking status.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.




Figure S5. Association between glycoalyx-related SDF imaging parameters and levels of coagulation parameters in premenopausal women. After stratification for menopausal status, differences in premenopausal women between the glycoalyx-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: PBR_{Total}: PBR of total vessels from 4-25 μ m; PBR_{feed vessel}: PBR of feed vessels from 10-19 μ m; PBR_{capillary}: PBR of capillaries from 4-9 μ m.

Table S7. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in premenopausal women.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
PBR _{Total}	Model 1	3.3 (-1.8-8.4)	5.6 (-0.2-11.3)	6.2 (-2-14.4)	0 (-0.3-0.3)	-0.3 (-0.9-0.3)	12.2 (-0.5-24.9)	108.8 (4.1-213.5)	0.2 (0-0.4)	13.8 (-1.1-28.7)
	Model 2	2.9 (-2.3-8.1)	4.1 (-1.6-9.8)	3.7 (-4.3-11.6)	0 (-0.4-0.3)	-0.2 (-0.8-0.4)	10.9 (-2.1-23.9)	107.4 (-0.7-215.5)	0.2 (0-0.4)	11 (-4-26.1)
	Model 3	3 (-2.1-8.2)	4.1 (-0.4-8.6)	4.9 (-3-12.8)	0 (-0.3-0.3)	-0.2 (-0.7-0.3)	10.1 (-0.4-20.5)	89.3 (-1.1-179.6)	0.2 (0-0.4)	10.1 (-3.6-23.8)
	Model 4	3.3 (-2-8.6)	5.5 (1.9-9.2)	8.4 (1.2-15.7)	0 (-0.2-0.3)	-0.3 (-0.8-0.2)	14.2 (4.2-24.2)	118.9 (31.9-205.8)	0.2 (0.1-0.4)	12.4 (1.2-23.7)
PBR _{feed vessel}	Model 1	3.8 (-1.2-8.9)	6.3 (0.6-12.1)	5.8 (-2.5-14)	0 (-0.3-0.3)	-0.3 (-0.8-0.3)	12 (-0.6-24.7)	99.5 (-5.8-204.7)	0.2 (0-0.4)	13 (-2-27.9)
	Model 2	3.5 (-1.7-8.7)	5 (-0.6-10.6)	3.4 (-4.5-11.3)	0 (-0.3-0.3)	-0.2 (-0.8-0.4)	10.8 (-2.2-23.7)	97.4 (-10.9-205.7)	0.2 (-0.1-0.4)	10.3 (-4.7-25.3)
	Model 3	3.4 (-1.7-8.5)	4.6 (0.2-9)	4.7 (-3.2-12.5)	0 (-0.3-0.3)	-0.2 (-0.7-0.3)	10 (-0.3-20.4)	80.4 (-9.5-170.4)	0.2 (0-0.3)	8.2 (-5.5-21.9)
	Model 4	4 (-1.2-9.1)	5.7 (2.2-9.3)	7 (-0.2-14.1)	0.1 (-0.2-0.3)	-0.2 (-0.7-0.3)	12 (2.1-21.9)	94.5 (7.9-181.1)	0.2 (0-0.4)	8.5 (-2.8-19.7)
PBR _{capillary}	Model 1	0.8 (-4.3-6)	-2.1 (-8-3.8)	0.8 (-7.5-9.2)	0 (-0.3-0.3)	0.2 (-0.4-0.7)	-11.3 (-24-1.4)	-115.6 (-219.9--11.4)	-0.2 (-0.4-0)	4.6 (-10.7-19.8)
	Model 2	0.6 (-4.5-5.8)	-2.7 (-8.3-3)	-0.1 (-8-7.7)	0 (-0.3-0.3)	0.2 (-0.4-0.8)	-12 (-24.7-0.6)	-118.6 (-223.6--13.7)	-0.2 (-0.4-0)	3.4 (-11.6-18.4)
	Model 3	1.7 (-3.3-6.8)	-0.4 (-4.9-4.1)	-1 (-8.8-6.8)	0 (-0.3-0.3)	0.1 (-0.4-0.6)	-8.8 (-19.1-1.4)	-94 (-181.4--6.5)	-0.1 (-0.3-0)	7 (-6.4-20.5)
	Model 4	2.4 (-2.7-7.6)	1 (-2.8-4.8)	1.9 (-5.5-9.2)	0.1 (-0.2-0.4)	0.1 (-0.4-0.6)	-7.1 (-17.3-3)	-82.8 (-169.7-4.1)	-0.1 (-0.3-0.1)	8.1 (-3.1-19.2)

Results are based on linear regression analyses of the Dutch premenopausal women population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: PBR_{Total}: PBR of total vessels from 4-25µm; PBR_{feed vessel}: PBR of feed vessels from 10-19µm; PBR_{capillary}: PBR of capillaries from 4-9µm.




Figure S6. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in premenopausal women. After stratification for menopausal status, differences in premenopausal women between the microcirculatory-perfusion-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.

Table S8. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in premenopausal women.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
RBCV _{feed vessel}	Model 1	1.5 (-3.7-6.6)	3 (-2.9-8.9)	0.2 (-8.2-8.5)	0.1 (-0.2-0.4)	0 (-0.5-0.6)	3.7 (-9.3-16.7)	48.9 (-58.5-156.3)	0.1 (-0.2-0.3)	21.7 (7.4-36)
	Model 2	1.5 (-3.6-6.7)	3.2 (-2.4-8.8)	0.5 (-7.3-8.3)	0.1 (-0.2-0.4)	0 (-0.5-0.6)	3.9 (-9-16.9)	49.7 (-58.4-157.7)	0.1 (-0.2-0.3)	22.1 (8.2-36)
	Model 3	0.1 (-5.2-5.3)	0.2 (-4.5-4.8)	-0.1 (-8.1-8)	0 (-0.3-0.3)	0 (-0.5-0.5)	1.9 (-9-12.7)	17.6 (-76.1-111.2)	0 (-0.2-0.2)	15 (1.6-28.5)
	Model 4	-0.3 (-5.6-5)	-1.2 (-5.1-2.7)	-0.6 (-8.1-7)	0 (-0.3-0.3)	-0.1 (-0.5-0.4)	1.4 (-9.1-12)	6.6 (-84.9-98.1)	0 (-0.2-0.2)	10.3 (-0.9-21.6)
RBCV _{capillary}	Model 1	3.3 (-1.8-8.3)	2.3 (-3.6-8.2)	1.3 (-7.9-7)	0 (-0.3-0.3)	-0.1 (-0.7-0.4)	6.2 (-6.8-19.1)	63.2 (-43.8-170.2)	0.1 (-0.1-0.3)	15.2 (0.4-30)
	Model 2	3.2 (-1.9-8.3)	2 (-3.6-7.7)	0.9 (-6.9-8.7)	0 (-0.3-0.3)	-0.1 (-0.7-0.5)	5.9 (-7-18.8)	62.2 (-45.5-169.9)	0.1 (-0.1-0.3)	14.7 (0.2-29.2)
	Model 3	1.8 (-3.4-7)	-0.8 (-5.4-3.7)	-0.2 (-8.2-7.8)	0 (-0.3-0.3)	0 (-0.5-0.5)	1.9 (-8.8-12.6)	23.1 (-69.6-115.8)	0 (-0.2-0.2)	10.1 (-3.6-23.7)
	Model 4	2.3 (-3-7.5)	-0.4 (-4.3-3.5)	1.7 (-5.8-9.2)	0 (-0.2-0.3)	0 (-0.5-0.5)	3.1 (-7.4-13.6)	28.9 (-62.2-120)	0.1 (-0.1-0.2)	8 (-3.4-19.4)
D _{Total}	Model 1	2.3 (-2.9-7.4)	5.9 (0.2-11.7)	1.2 (-7.1-9.6)	0 (-0.3-0.3)	-0.2 (-0.7-0.4)	12.5 (-0.2-25.1)	126 (22.6-229.5)	0.2 (0-0.4)	22.9 (8.7-37.1)
	Model 2	2.4 (-2.7-7.5)	6.3 (0.8-11.7)	1.8 (-6-9.6)	0 (-0.3-0.3)	-0.2 (-0.8-0.4)	12.9 (0.3-25.4)	127.5 (23.5-231.6)	0.2 (0-0.4)	23.6 (9.8-37.3)
	Model 3	-0.4 (-6-5.2)	1.2 (-3.8-6.1)	1.7 (-6.8-10.3)	0.1 (-0.2-0.4)	0.1 (-0.4-0.6)	3.9 (-7.5-15.4)	42.5 (-56.6-141.7)	0.1 (-0.1-0.3)	17.7 (3.5-31.9)
	Model 4	-1 (-6.5-4.6)	-0.1 (-4.2-4)	0.9 (-7.1-8.8)	0.1 (-0.2-0.4)	0 (-0.5-0.6)	3.5 (-7.6-14.6)	33.2 (-63.1-129.4)	0.1 (-0.1-0.3)	14.5 (2.8-26.1)
D _{feed vessel}	Model 1	3.9 (-1.1-9)	5.2 (-0.6-11)	2.1 (-6.3-10.4)	0.1 (-0.2-0.4)	0 (-0.5-0.6)	7.1 (-5.8-20)	91.4 (-14.4-197.1)	0.1 (-0.1-0.3)	18 (3.4-32.7)
	Model 2	3.8 (-1.2-8.9)	4.9 (-0.6-10.4)	1.5 (-6.3-9.4)	0.1 (-0.2-0.4)	0.1 (-0.5-0.6)	6.8 (-6.1-19.6)	90.3 (-16.2-196.7)	0.1 (-0.1-0.3)	17.4 (3.1-31.7)
	Model 3	1.7 (-3.7-7.1)	0.4 (-4.4-5.2)	1.2 (-7.1-9.5)	0.2 (-0.1-0.5)	0.4 (-0.1-0.9)	-2.8 (-14-8.3)	9.4 (-87.4-106.1)	-0.1 (-0.3-0.1)	12.4 (-1.7-26.6)
	Model 4	1 (-4.5-6.4)	-0.4 (-4.4-3.6)	0.8 (-6.9-8.6)	0.1 (-0.1-0.4)	0.3 (-0.2-0.8)	-2.1 (-12.9-8.7)	11.6 (-82.4-105.7)	0 (-0.2-0.1)	12.8 (1.4-24.2)
D _{capillary}	Model 1	-0.4 (-5.5-4.8)	4.7 (-1.1-10.5)	-0.1 (-8.4-8.3)	-0.1 (-0.4-0.2)	-0.4 (-0.9-0.2)	13.3 (0.7-25.9)	112 (7.6-216.5)	0.2 (0-0.5)	21.2 (6.8-35.6)
	Model 2	-0.2 (-5.3-5)	5.4 (-0.1-11)	1.1 (-6.7-8.9)	-0.1 (-0.4-0.3)	-0.4 (-1-0.2)	14.2 (1.7-26.7)	115.8 (10.5-221.1)	0.2 (0-0.5)	22.8 (8.9-36.7)
	Model 3	-2.6 (-8-2.7)	1.2 (-3.5-6)	1.4 (-6.9-9.6)	-0.1 (-0.4-0.3)	-0.3 (-0.8-0.2)	8.6 (-2.3-19.4)	50.2 (-45.1-145.4)	0.2 (0-0.4)	15.7 (1.9-29.5)
	Model 4	-2.7 (-8.1-2.8)	0.1 (-3.9-4.1)	0.7 (-7.1-8.4)	0 (-0.3-0.2)	-0.3 (-0.8-0.2)	7.3 (-3.4-18)	33.1 (-60.9-127.1)	0.1 (-0.1-0.3)	9.6 (-2.1-21.2)
CBV _{static}	Model 1	-2.1 (-7.3-3)	1.4 (-4.5-7.3)	-1.6 (-10-6.7)	-0.1 (-0.4-0.2)	-0.3 (-0.9-0.3)	8.3 (-4.5-21.2)	37.4 (-70.3-145.2)	0.2 (-0.1-0.4)	6.3 (-8.9-21.6)
	Model 2	-1.8 (-7-3.4)	2.6 (-3.1-8.3)	0.3 (-7.6-8.2)	-0.1 (-0.4-0.2)	-0.4 (-0.9-0.2)	9.9 (-3-22.7)	42.9 (-66.6-152.4)	0.2 (0-0.4)	8.8 (-6.1-23.8)
	Model 3	-2.2 (-7.3-2.9)	1.8 (-2.7-6.3)	1.4 (-6.5-9.3)	-0.1 (-0.4-0.2)	-0.4 (-0.9-0.1)	9.7 (-0.6-20)	45.8 (-45.4-136.9)	0.2 (0-0.4)	8 (-5.6-21.6)
	Model 4	-2.4 (-7.5-2.7)	0.7 (-3.1-4.4)	-0.2 (-7.5-7.2)	-0.1 (-0.4-0.1)	-0.4 (-0.9-0.1)	8 (-2-18.1)	29.2 (-59.6-118)	0.2 (0-0.4)	4.1 (-7.1-15.4)

Results are based on linear regression analyses of the Dutch premenopausal women population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.




Figure S7. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in postmenopausal women. After stratification for menopausal status, differences in postmenopausal women between the glyocalyx-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: PBR_{Total}: PBR of total vessels from 4-25 μ m; PBR_{feed vessel}: PBR of feed vessels from 10-19 μ m; PBR_{capillary}: PBR of capillaries from 4-9 μ m.

Table S9. Association between glyocalyx-related SDF imaging parameters and levels of coagulation parameters in postmenopausal women.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
PBR _{Total}	Model 1	0.2 (-2.1-2.6)	0.5 (-1.7-2.7)	0.2 (-3.2-3.6)	-0.1 (-0.2-0.1)	0 (-0.3-0.2)	0.3 (-3.4-4.1)	3.9 (-34.4-42.2)	0 (-0.1-0.1)	4 (-2.3-10.2)
	Model 2	0.3 (-2-2.7)	0.5 (-1.7-2.7)	0.6 (-2.8-4)	0 (-0.2-0.1)	0 (-0.2-0.2)	-0.4 (-4.1-3.3)	-4.1 (-41.7-33.4)	0 (-0.1-0.1)	4.1 (-2.2-10.4)
	Model 3	1 (-1.4-3.3)	1.9 (0-3.8)	0.8 (-2.7-4.2)	0 (-0.2-0.1)	0 (-0.2-0.2)	1.1 (-2.4-4.6)	10.5 (-25.1-46.1)	0 (-0.1-0.1)	6.5 (0.6-12.4)
	Model 4	0.8 (-1.5-3.1)	1.3 (-0.3-3)	-0.2 (-3.6-3.2)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.2 (-3.2-3.6)	2 (-33.3-37.3)	0 (-0.1-0.1)	3.6 (-1.8-9)
PBR _{feed vessel}	Model 1	0.6 (-1.7-3)	0.2 (-2-2.4)	0.7 (-2.7-4.1)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	0.6 (-3.2-4.3)	6.3 (-31.9-44.6)	0 (-0.1-0.1)	3.5 (-2.8-9.7)
	Model 2	0.7 (-1.6-3.1)	0.2 (-2-2.4)	1.1 (-2.3-4.6)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	-0.2 (-3.9-3.6)	-1.9 (-39.5-35.7)	0 (-0.1-0.1)	3.6 (-2.6-9.9)
	Model 3	1.4 (-0.9-3.7)	1.7 (-0.2-3.6)	1.5 (-1.9-4.9)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	1.4 (-2-4.9)	13.2 (-22.3-48.8)	0 (-0.1-0.1)	6.2 (0.3-12.1)
	Model 4	1.3 (-1-3.6)	1.4 (-0.2-3)	1 (-2.3-4.4)	0 (-0.2-0.1)	-0.1 (-0.3-0.1)	0.9 (-2.5-4.3)	7.9 (-27.1-43)	0 (-0.1-0.1)	5.1 (-0.2-10.4)
PBR _{capillary}	Model 1	-1.3 (-3.7-1)	0.8 (-1.5-3)	3.4 (-0.1-6.8)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.9 (-2.9-4.7)	14.6 (-23.7-52.8)	0 (-0.1-0.1)	3.7 (-2.5-10)
	Model 2	-1.3 (-3.6-1.1)	0.8 (-1.5-3)	3.7 (0.3-7.1)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.3 (-3.4-4)	8.3 (-29.2-45.7)	0 (-0.1-0.1)	3.8 (-2.4-10.1)
	Model 3	-1 (-3.3-1.3)	1.2 (-0.7-3.1)	3.7 (0.3-7)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.6 (-2.8-4.1)	11.6 (-23.6-46.9)	0 (-0.1-0.1)	4.6 (-1.2-10.5)
	Model 4	-0.9 (-3.2-1.3)	1.3 (-0.3-2.9)	3.7 (0.4-7)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.7 (-2.7-4)	11.9 (-22.7-46.5)	0 (-0.1-0.1)	4.8 (-0.4-10.1)

Results are based on linear regression analyses of the Dutch postmenopausal women population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: PBR_{Total}: PBR of total vessels from 4-25µm; PBR_{feed vessel}: PBR of feed vessels from 10-19µm; PBR_{capillary}: PBR of capillaries from 4-9µm.




Figure S8. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in postmenopausal women. After stratification for menopausal status, differences in postmenopausal women between the microcirculatory-perfusion-related parameters and coagulation factor levels can be observed. Model 1: crude model; Model 2: model 1 + age; Model 3: model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy; Model 4: model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration. The effect size and 95% confidence interval were depicted by a horizontal line with a dot. A non-significant association was represented by the color grey, and a substantial positive association was represented by the color red. Abbreviations: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.

Table S10. Association between microcirculatory-perfusion-related SDF imaging parameters and levels of coagulation parameters in postmenopausal women.

		FXI activity	FIX activity	FVIII activity	Lag time	ttPeak	Peak	ETP	VelIndex	Fibrinogen
RBCV _{feed vessel}	Model 1	1.5 (-0.8-3.9)	2 (-0.2-4.2)	2.5 (-0.9-5.9)	0 (-0.1-0.2)	0 (-0.2-0.2)	2.2 (-1.6-6)	28.1 (-10.1-66.2)	0 (0-0.1)	-0.2 (-6.5-6)
	Model 2	1.5 (-0.8-3.9)	2 (-0.2-4.2)	2.5 (-0.9-5.9)	0 (-0.1-0.2)	0 (-0.2-0.2)	2.3 (-1.4-5.9)	28.6 (-8.6-65.8)	0 (0-0.1)	-0.3 (-6.5-6)
	Model 3	0.7 (-1.6-3)	0.5 (-1.4-2.4)	2.1 (-1.4-5.5)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.5 (-2.9-3.9)	12.7 (-22.8-48.2)	0 (-0.1-0.1)	-2.8 (-8.7-3.1)
	Model 4	0.2 (-2.1-2.5)	-0.5 (-2.1-1.1)	1.9 (-1.5-5.3)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.7 (-2.7-4.1)	15.8 (-19.4-50.9)	0 (-0.1-0.1)	-4.4 (-9.7-1)
RBCV _{capillary}	Model 1	1.2 (-1.1-3.6)	1.4 (-0.8-3.6)	2.2 (-1.2-5.7)	0 (-0.1-0.2)	0.1 (-0.2-0.3)	1.1 (-2.7-4.9)	20.4 (-17.8-58.6)	0 (-0.1-0.1)	0.5 (-5.8-6.7)
	Model 2	1.2 (-1.2-3.5)	1.4 (-0.8-3.6)	2.2 (-1.2-5.6)	0 (-0.1-0.2)	0 (-0.2-0.2)	1.3 (-2.4-5)	22.2 (-15.1-59.4)	0 (-0.1-0.1)	0.5 (-5.8-6.7)
	Model 3	0.6 (-1.7-2.8)	0.3 (-1.6-2.2)	1.9 (-1.6-5.3)	0 (-0.1-0.2)	0.1 (-0.1-0.3)	-0.1 (-3.5-3.3)	10 (-25.3-45.3)	0 (-0.1-0.1)	-1.2 (-7.1-4.6)
	Model 4	0.2 (-2.1-2.5)	-0.4 (-2-1.2)	1.7 (-1.7-5)	0 (-0.1-0.1)	0 (-0.2-0.2)	0 (-3.4-3.3)	11.3 (-23.5-46.2)	0 (-0.1-0.1)	-2.6 (-7.9-2.7)
D _{Total}	Model 1	0.8 (-1.5-3.1)	1.3 (-0.9-3.5)	1.8 (-1.7-5.2)	0 (-0.1-0.2)	0.1 (-0.2-0.3)	2.1 (-1.7-5.9)	34.1 (-4-72.2)	0 (-0.1-0.1)	0.8 (-5.5-7)
	Model 2	0.7 (-1.6-3.1)	1.3 (-0.9-3.5)	1.4 (-2-4.8)	0 (-0.2-0.1)	0 (-0.2-0.2)	2.8 (-0.9-6.5)	42 (4.7-79.3)	0 (0-0.1)	0.7 (-5.6-7)
	Model 3	0.1 (-2.2-2.4)	0.2 (-1.7-2.1)	1 (-2.4-4.4)	0 (-0.2-0.1)	0 (-0.2-0.2)	1.7 (-1.8-5.1)	32.2 (-3.2-67.5)	0 (-0.1-0.1)	-1.5 (-7.4-4.4)
	Model 4	0.1 (-2.2-2.3)	0.2 (-1.5-1.8)	0.9 (-2.4-4.3)	0 (-0.2-0.1)	0 (-0.2-0.2)	1.5 (-1.9-4.9)	30.1 (-4.7-64.9)	0 (-0.1-0.1)	-1.3 (-6.7-4)
D _{feed vessel}	Model 1	0.8 (-1.5-3.2)	2.1 (-0.1-4.3)	2.3 (-1.2-5.7)	0.1 (-0.1-0.2)	0.1 (-0.1-0.3)	1.9 (-1.8-7.5)	31.2 (-6.9-69.3)	0 (-0.1-0.1)	0.9 (-5.4-7.1)
	Model 2	0.7 (-1.6-3.1)	2.1 (-0.1-4.3)	1.9 (-1.5-5.3)	0 (-0.1-0.2)	0.1 (-0.1-0.3)	2.7 (-1.1-6.4)	39.5 (2.2-76.8)	0 (-0.1-0.1)	0.8 (-5.5-7.1)
	Model 3	0.1 (-2.2-2.4)	0.7 (-1.2-2.6)	1.3 (-2.1-4.8)	0 (-0.1-0.2)	0.1 (-0.1-0.3)	1 (-2.4-4.5)	24.9 (-10.6-60.5)	0 (-0.1-0.1)	-1.6 (-7.5-4.4)
	Model 4	-0.1 (-2.3-2.2)	0.2 (-1.4-1.9)	0.8 (-2.5-4.2)	0 (-0.1-0.1)	0.1 (-0.1-0.3)	0.4 (-3-3.8)	18.9 (-16.2-54)	0 (-0.1-0.1)	-2.5 (-7.9-2.8)
D _{capillary}	Model 1	0.3 (-2-2.7)	-0.3 (-2.5-1.9)	-0.2 (-3.6-3.3)	-0.1 (-0.2-0.1)	0 (-0.2-0.2)	1.4 (-2.3-5.2)	23.3 (-14.9-61.5)	0 (-0.1-0.1)	0.5 (-5.8-6.7)
	Model 2	0.3 (-2-2.7)	-0.3 (-2.5-1.9)	-0.3 (-3.7-3.1)	-0.1 (-0.2-0.1)	0 (-0.2-0.2)	1.6 (-2-5.3)	25.7 (-11.6-63)	0 (-0.1-0.1)	0.4 (-5.8-6.7)
	Model 3	0 (-2.3-2.3)	-0.4 (-2.3-1.5)	-0.3 (-3.7-3.1)	-0.1 (-0.2-0.1)	-0.1 (-0.3-0.1)	1.8 (-1.6-5.2)	27.1 (-8.1-62.3)	0 (0-0.1)	-0.3 (-6.2-5.6)
	Model 4	0.2 (-2-2.5)	0.1 (-1.5-1.7)	0.1 (-3.2-3.5)	-0.1 (-0.2-0.1)	0 (-0.2-0.2)	2.1 (-1.2-5.5)	30.2 (-4.4-64.7)	0 (0-0.1)	1.1 (-4.2-6.4)
CBV _{static}	Model 1	-0.5 (-2.8-1.9)	-2 (-4.3-0.2)	-1 (-4.4-2.4)	0 (-0.1-0.1)	0 (-0.2-0.2)	0.2 (-3.6-3.9)	0.6 (-37.6-38.9)	0 (-0.1-0.1)	-0.9 (-7.2-5.3)
	Model 2	-0.5 (-2.9-1.8)	-2.1 (-4.3-0.1)	-1.3 (-4.7-2.1)	0 (-0.2-0.1)	0 (-0.2-0.2)	0.7 (-3-4)	6.5 (-31-44)	0 (-0.1-0.1)	-1 (-7.3-5.3)
	Model 3	-0.3 (-2.6-2)	-1 (-2.9-0.9)	-1 (-4.4-2.4)	0 (-0.2-0.1)	0 (-0.2-0.2)	2 (-1.4-5.5)	19.3 (-16.1-54.7)	0 (0-0.1)	0.6 (-5.3-6.5)
	Model 4	0 (-2.2-2.3)	-0.2 (-1.8-1.4)	-0.5 (-3.9-2.9)	0 (-0.1-0.2)	0 (-0.2-0.2)	2.3 (-1-5.7)	21.5 (-13.4-56.4)	0 (0-0.1)	2.4 (-2.9-7.8)

Results are based on linear regression analyses of the Dutch postmenopausal women population. The beta coefficient (95% CI) can be interpreted as differences in coagulation parameter levels per SD change in PBR.

Model 1 = crude model.

Model 2 = model 1 + age.

Model 3 = model 2 + BMI, current smoking status, current use of the oral contraceptive pill, and current use of hormone replacement therapy.

Model 4 = model 3 + serum CRP, serum leptin concentration, and serum GlycA concentration.

Abbreviation: RBCV_{feed vessel}: feed vessel RBC velocity; RBCV_{capillary}: capillary RBC velocity; D_{Total}: total valid vessel density with measurable RBC velocity; D_{feed vessel}: perfused feed vessel density; D_{capillary}: perfused capillary density; CBV_{static}: capillary blood volume.