



Universiteit
Leiden

The Netherlands

Harnessing the immunostimulatory properties of oncolytic reovirus for anticancer immunotherapy

Groeneveldt, P.C.

Citation

Groeneveldt, P. C. (2023, November 23). *Harnessing the immunostimulatory properties of oncolytic reovirus for anticancer immunotherapy*. Retrieved from <https://hdl.handle.net/1887/3663612>


Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3663612>

Note: To cite this publication please use the final published version (if applicable).

ADDENDUM



**TGF- β blockade improves Reo& α PD-L1
therapy in the murine colon MC38
tumor model**

BACKGROUND

In **Chapter 7**, we observed that TGF- β blockade differentially affects the efficacy of reovirus and CD3-bispecific antibody therapy (Reo&CD3-bsAbs). In the murine pancreatic KPC3 tumor model, the efficacy of Reo&CD3-bsAbs was impaired. Tumor regressions were prevented, and survival was shortened by TGF- β blockade. In contrast, in the murine MC38 colon carcinoma tumor model, TGF- β blockade significantly improved the efficacy of Reo&CD3-bsAb therapy, even leading to a 100% complete response rate. Since we also demonstrated that reovirus can enhance the efficacy of PD-L1 blockade in MC38 tumors in **Chapter 5**, we here investigated whether the efficacy of this combination therapy (Reo& α PD-L1) could also be improved by TGF- β blockade.

RESULTS & DISCUSSION

We engrafted immunocompetent C57BL/6j mice with subcutaneous MC38 tumors. Mice received TGF- β neutralizing antibodies (α TGF- β) starting a few days after tumor engraftment, and Reo& α PD-L1 therapy was administered on day 8, 11 and 14 (**Figure 1A**). Anti-TGF- β antibodies alone delayed tumor growth and induced complete tumor clearance in 10% of animals, and Reo& α PD-L1 therapy led to tumor clearance in 50% of animals (**Figure 1B-D**). However, the addition of α TGF- β to Reo& α PD-L1 therapy led to a total tumor clearance in 80% of animals and enhanced survival. Thus, the efficacy of Reo& α PD-L1 therapy can be improved by TGF- β blockade.

It is expected that PD-L1 blockade mediates its efficacy by reinvigorating dysfunctional tumor-specific T cells (1). However, the increased efficacy of Reo& α PD-L1 after α TGF- β could not be attributed to an increase in tumor-specific (Rpl18+) T cells, as similar frequencies were detected in the circulation of both groups receiving Reo& α PD-L1 therapy (**Figure 1E**). In **Chapter 7**, we also observed that Reo treatment does not improve the frequency of Rpl18⁺ CD8⁺ T cells, either in the circulation or in the tumor. Here, we confirmed that the frequency of Rpl18⁺ CD8⁺ T cells was not correlated with clinical outcome, since mice with a complete response (CR; in blue) did not always have the highest frequency of Rpl18⁺ CD8⁺ T cells (**Figure 1E**). Hence, we concluded that the triple combination of Reo, α PD-L1 and α TGF- β blockade demonstrates superior antitumor efficacy compared to all single arms or combinations of two arms, but that this improved efficacy was not associated with an increased frequency of tumor-specific Rpl18⁺ CD8⁺ T cells.

We next investigated whether antitumor immunologic memory was established by the different treatments. We therefore rechallenged the mice that completely cleared the primary tumor on the other flank and monitored tumor outgrowth (**Figure 1F**). Mice that cleared previous tumors had increased presence of Rpl18-specific T cells in the

circulation compared to naive mice, especially the group that received Reo& α PD-L1 therapy together with α TGF- β , suggesting that Rpl18-specific T-cell responses were boosted by the rechallenge with MC38 tumor cells (Figure 1G). While tumors grew out in 100% of naive mice, this was not the case in mice previously treated with Reo& α PD-L1 or Reo& α PD-L1+ α TGF- β (Figure 1H).




Figure 1. TGF- β blockade increases response rate after Reo& α PD-L1 in the MC38 model of colon cancer. (A) Overview of experiment described in (B-E). Immunocompetent C57BL/6J mice were subcutaneously engrafted with MC38 cells (5×10^5 /mouse) and received TGF- β -neutralizing antibodies (α TGF- β , 200 μ g/injection every 3 days) starting directly after tumor engraftment. Mice received Reo (intratumorally, 10^7 plaque-forming units/injection) and α PD-L1 (intraperitoneally, 200 μ g/injection) on day 8, 11 and 14. Tumor growth was measured 3x/week. (B) Individual tumor growth curves of mice receiving indicated treatments. (C) Frequency of Non-Responders (NR), Partial Responders (PR) or Complete Responders (CR) within each treatment group. (D) Kaplan-Meier survival graphs of mice after indicated treatments. (E) Frequency of Rpl18 $^+$ CD8 $^+$ T cells in blood of mice after indicated treatments. (F) Design of rechallenge experiment. All CR mice from (D) were subcutaneously engrafted with MC38 tumor cells (5×10^5 /mouse) in the alternate flank, and tumor outgrowth was measured 3x/week. (G) Frequency of Rpl18 $^+$ CD8 $^+$ T cells in blood of mice after rechallenge. (H) Individual tumor growth curves of mice that were rechallenged with MC38 tumor cells. Data represent mean \pm SEM. Significance between groups in (E) and (G) was determined using an ordinary one-way ANOVA with Tukey's multiple comparisons test. Chi-square test was used to determine statistical differences in response in (C). Log-rank tests were used to compare differences in survival in (D). Significance levels: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$.

These data indicate that tumor-specific immune responses were established by our therapies, but that the efficacy of these therapies could not be explained by the increased frequency of tumor-specific Rpl18⁺ T cells. It would be interesting to investigate whether the improved protection was due to other MC38-specific T-cell responses, for instance those directed towards neo-epitope Adpgk (2). Alternatively, other cell types in the TME may be involved. For instance, tumor-associated M2 macrophages (TAMs) have been shown to hamper the efficacy of checkpoint blockade in the MC38 tumor model (3). Since anti-inflammatory M2 TAMs are induced by TGF- β (4,5), it is possible that neutralization of TGF- β hampered the function or reduced the level of these TAMs and thereby improved the efficacy of Reo& α PD-L1 therapy, without directly affecting the numbers of circulating tumor-specific T cells.

CONCLUSION

Combined, these data show that both Reo&CD3-bsAb and Reo& α PD-L1 therapy can be improved by TGF- β blockade in the MC38 tumor model. This further highlights the necessity to determine which factor(s), mechanism(s) or cell type(s) that are present in MC38 tumors permit or even contribute to this beneficial effect of TGF- β blockade, with the ultimate aim to employ this characteristic for effective viro-immunotherapy in other tumor models that are much harder to treat.

REFERENCES

1. Xu-Monette ZY, Zhang M, Li J, Young KH. PD-1/PD-L1 Blockade: Have We Found the Key to Unleash the Antitumor Immune Response? *Front Immunol* **2017**;8
2. Hos BJ, Camps MGM, van den Bulk J, Tondini E, van den Ende TC, Ruano D, *et al.* Identification of a neo-epitope dominating endogenous CD8 T cell responses to MC-38 colorectal cancer. *Oncoimmunology* **2019**;9:1673125
3. Arlauckas SP, Garriss CS, Kohler RH, Kitaoka M, Cuccarese MF, Yang KS, *et al.* In vivo imaging reveals a tumor-associated macrophage-mediated resistance pathway in anti-PD-1 therapy. *Sci Transl Med* **2017**;9:eaal3604
4. Zhang F, Wang H, Wang X, Jiang G, Liu H, Zhang G, *et al.* TGF- β induces M2-like macrophage polarization via SNAIL-mediated suppression of a pro-inflammatory phenotype. *Oncotarget* **2016**;7:52294-306
5. Batlle E, Massague J. Transforming Growth Factor-beta Signaling in Immunity and Cancer. *Immunity* **2019**;50:924-40

