



**Universiteit
Leiden**
The Netherlands

Characterization of candidate genes in unexplained polyposis and colorectal cancer

Abayzeed Elsayed Osman, F.

Citation

Abayzeed Elsayed Osman, F. (2023, November 28). *Characterization of candidate genes in unexplained polyposis and colorectal cancer*. Retrieved from <https://hdl.handle.net/1887/3665175>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3665175>

Note: To cite this publication please use the final published version (if applicable).

Chapter 1

General introduction and outline of
this thesis

General introduction

Colorectal cancer

Colorectal cancer (CRC; MIM 114500) is the third most commonly diagnosed cancer and the second leading cause of cancer death worldwide ¹⁻³, with almost 1.9 million new cases and 1 million deaths in 2020. CRC accounts for 10% of all cancer diagnoses and 9.4% of all cancer deaths ^{2,4,5}. CRC results from the progressive accumulation of genetic and epigenetic alterations that lead to the transformation of normal colonic epithelium to colon adenocarcinoma ⁶. In the progression from colorectal adenoma to carcinoma, three major pathways are distinguished: chromosomal instability (CIN), microsatellite instability (MSI) and CpG island methylator phenotype (CIMP) ⁷. CIN is the most common type of genomic instability observed in CRC and occurs in 80%-85% of colorectal tumors ⁶. While the majority of CRCs occur sporadically, an estimated 35% of CRCs are due to heritable factors ^{8,9}. Between 5% and 10% of all CRC cases are associated with well-characterized hereditary polyposis and/or CRC syndromes ⁹. The etiology of up to 30% of inherited CRCs is not completely understood, and the underlying genetic factors contributing to the risk of CRC remain undefined ¹⁰. Genome-wide association studies (GWAS) have successfully identified common, low-penetrance single nucleotide polymorphisms (SNPs) associated with the risk of CRC ¹¹⁻¹⁷. In recent years, major efforts have been made to identify the genetic causes, as the identification of germline pathogenic variants substantially facilitates the clinical management of patients and their families.

Hereditary colorectal cancer syndromes

Hereditary CRC syndromes (Table 1), characterized by dramatic increases in the risk of colorectal neoplasia, are phenotypically divided into polyposis and nonpolyposis syndromes, based largely on the number and histology of the colorectal polyps. The polyposis syndromes can be further divided into adenomatous, hamartomatous, serrated and mixed polyposis syndromes according to the predominant type of polyps, e.g., adenomatous polyps, hamartomatous polyps or serrated polyps. Polyposis is defined by the constitutive development of multiple polyps in the colon and rectum. Polyps are benign outgrowths of tissue into the lumen of the colorectum, but they have the potential to evolve into an in situ carcinoma by the accumulation of additional somatic mutations ¹⁸. This phenomenon is known as the adenoma-to-carcinoma sequence, and it is accepted that more than 95% of colorectal cancers arise from adenomas. Syndromic nonpolyposis CRC is subdivided on the basis of molecular tumor phenotype as DNA mismatch repair-deficient (MMRD) or mismatch

repair-proficient (MMRP) CRC¹⁹⁻²¹. The development of polyps in patients with a nonpolyposis CRC predisposition syndrome is rare, but these polyps evolve rapidly into carcinomas since the polyp-to-carcinoma sequence appears to be accelerated in these patients²². Several high-penetrance genes with inherited germline variants, such as *APC* (MIM 611731), *BMPR1A* (MIM 601299), *GREM1* (MIM 603054), *MLH1* (MIM 120436), *MSH2* (MIM 609309), *MSH3* (MIM 600887), *MSH6* (MIM 600678), *MUTYH* (MIM 604933), *NTHL1* (MIM 602656), *PMS2* (MIM 600259), *POLD1* (MIM 174761), *POLE* (MIM 174762), *PTEN* (MIM 601728), *RNF43* (MIM 612482), *SMAD4* (MIM 600993) and *STK11* (MIM 602216), are known to be associated with CRC syndromes²³.

Table 1. CRC predisposition syndromes

| Syndrome | Genes | Pattern of inheritance | Prevalence |
|--|--------------------------------------|------------------------|------------|
| Lynch syndrome | <i>MLH1, MSH2, MSH6, PMS2, EPCAM</i> | Dominant | 2% - 4% |
| Familial adenomatous polyposis | <i>APC</i> | Dominant | < 1% |
| MUTYH-associated polyposis | <i>MUTYH</i> | Recessive | < 1% |
| Polymerase proofreading-associated polyposis | <i>POLE, POLD1</i> | Dominant | Unknown |
| NTHL1-associated polyposis | <i>NTHL1</i> | Recessive | Unknown |
| MSH3-associated polyposis | <i>MSH3</i> | Recessive | Unknown |
| Serrated polyposis syndrome | <i>RNF43</i> | Dominant | Unknown |
| Constitutional MMR deficiency syndrome | <i>MLH1, MSH2, MSH6, PMS2</i> | Recessive | Unknown |
| Hereditary mixed polyposis syndrome | <i>GREM1</i> | Dominant | Unknown |
| Juvenile polyposis | <i>SMAD4, BMPR1A</i> | Dominant | < 1% |
| Peutz-Jeghers Syndrome | <i>STK11</i> | Dominant | < 1% |
| PTEN hamartoma tumor syndrome | <i>PTEN</i> | Dominant | < 1% |

Nonpolyposis syndromes

Lynch syndrome

Lynch syndrome (LS; MIM 120435), previously referred to as hereditary nonpolyposis colorectal cancer (HNPCC), is the most common cause of hereditary CRC, accounting for approximately 2% - 4% of all CRCs^{10, 24, 25}. The lifetime CRC risk is estimated to be 50%-80%^{10, 24}. This syndrome also predisposes patients to extracolonic cancers, such as cancers of the endometrium, small bowel, ureter and renal pelvis, stomach, hepatobiliary tract and ovary²⁶⁻³². LS is inherited in an autosomal dominant pattern and is caused by germline pathogenic variants in one of the MMR genes (*MLH1, MSH2, MSH6, PMS2*) or 3' end deletion of the *EPCAM* gene, leading to transcriptional read-through into and subsequent epigenetic silencing of *MSH2*³³⁻³⁸. For LS, the lifetime risk for CRC is highly variable and dependent on the gene involved. The risk for CRC-associated *MLH1* and *MSH2*

mutations is generally higher than the risks associated with mutations in the other LS-related genes^{32,39}. The MMR system consists of several proteins that repair DNA damage during replication and maintain genome stability mainly by correcting base-base and small insertion–deletion mismatches that are generated during DNA replication. MMR proteins function as heterodimers in two main complexes, MutS heterodimers (MSH2/MSH6 and MSH2/MSH3) and MutL heterodimers (MLH1/PMS2, MLH1/PMS1 and MLH1/MLH3). The MutS heterodimers recognize mismatches and small insertions/deletions (indels). The MutL heterodimers form a MutS/MutL/DNA complex for exonuclease activity and termination of mismatch-provoked excision^{40,41}. MMR defects lead to genomic instability and the accumulation of secondary mutations, resulting in a strong mutator phenotype. Mutations occur especially in simple repetitive DNA sequences and microsatellites, resulting in microsatellite instability (MSI). MSI is a hallmark of MMRD cancers and is found in > 90% of LS colorectal cancers⁴²⁻⁴⁴. In up to 15% of sporadic CRCs, MSI is caused by somatic hypermethylation of the *MLH1* promoter and associated silencing of *MLH1*. These patients frequently also exhibit specific mutations in *BRAF* (V600E)³².

Familial colorectal cancer type X

In a fraction of families fulfilling the Amsterdam 1 criteria for HNPCC⁴⁵, CRCs are microsatellite stable and without MMR gene mutations. These families are defined as having familial colorectal cancer type X (FCCTX)^{46,47}. This heterogeneous group of families has an increased risk of developing CRC and other related tumors⁴⁸. Although the clinical identification of FCCTX has improved in recent years, its genetic etiology remains unknown^{47,49}. Some genes, such as *BMPR1A*⁵⁰, *BRCA2*⁵¹, *FAN1*⁵², *OGG1*⁵³, *RPS20*⁵⁴, *SEMA4A*⁵⁵ and *SETD6*⁵⁶, have already been reported to be potentially associated with FCCTX. In addition, a review suggested a possible association with *BCR*, *BMP4*, *CENPE*, *CDH18*, *GABBR2*, *GALNT12*, *GREM1*, *HABP4*, *KIF24* and *ZNF367*⁵⁷. Moreover, a review by Nejadtaghi et al.⁵⁸ identified *APC*, *BMPR1A*, *BRAF*, *BRCA2*, *KRAS*, *MGMT*, *RPS20*, *SEMA4A*, and hypermethylation of at least one gene of the MMR system as potentially related to FCCTX. Despite these studies, no defined set of genes is conclusively associated with FCCTX.

Polyposis syndromes

Familial adenomatous polyposis

Less than 1% of all CRCs occur due to familial adenomatous polyposis (FAP; MIM 175100). FAP represents the most common gastrointestinal polyposis syndrome

and the second most common cause of hereditary CRC^{59,60}, with an estimated incidence varying from 1:8000 to 1:37600⁶¹. FAP is an autosomal dominant precancerous condition characterized by the development of colorectal adenomas, which inevitably progress to colorectal carcinoma unless detected early¹⁰. In the classic form of FAP, patients develop hundreds to thousands of colorectal adenomas during adolescence or the third decade of life, and the lifetime risk of CRC is almost 100%. Attenuated FAP (AFAP) is a phenotypically distinct form of FAP in which patients have a milder manifestation than classic FAP. AFAP is characterized by fewer colorectal adenoma polyps (less than 100 polyps), a later age of adenoma development and a lower lifetime risk of CRC (70%)^{62,63}. FAP is caused by germline variants in the tumor suppressor gene *APC*⁶⁴⁻⁶⁷. *APC* is located on chromosome 5q21-q22 and consists of 15 exons encoding a protein of 2845 amino acids (310 kDa). *APC* plays a major role in the Wnt signaling pathway by negatively regulating the β -catenin oncoprotein⁶⁸⁻⁷⁰. Germline *APC* variants lead to the development of multiple adenomas as a result of inactivation of the remaining wild-type *APC* allele in the tumor, either through somatic mutations or through loss of heterozygosity of *APC*^{59,70,71}. Correlations between the FAP phenotype and the site of mutation in the *APC* gene have been reported; patients with AFAP generally have a mutation in the 5' or 3' region of the *APC* gene, whereas individuals with FAP carry mutations elsewhere in this gene⁷². De novo variants are responsible for approximately 25% of FAP cases who lack a family history of the disease, and approximately 20% of these have somatic mosaicism⁷³⁻⁷⁷.

MUTYH-associated polyposis

In 2002, Al Tassan et al. reported for the first time that inherited defects of the base excision repair gene *MUTYH* predispose patients to multiple colorectal adenomas and carcinoma⁷⁸, causing *MUTYH*-associated polyposis (MAP; MIM 608456)⁷⁸⁻⁸¹. MAP is an autosomal recessive inherited syndrome caused by biallelic germline variants in the base excision repair gene *MUTYH*, characterized by a greatly increased lifetime risk of CRC (80%)⁸² and accounting for less than 1% of CRC cases^{60,83}. An estimated 1 in every 20,000 European individuals have biallelic *MUTYH* variants⁶¹. *MUTYH* encodes a DNA glycosylase involved in oxidative DNA damage repair, is located on chromosome 1 between p32.1 and p34.3 and consists of 16 exons⁸⁴. The enzyme excises adenine bases from the DNA backbone at sites where adenine is inappropriately paired with guanine, cytosine, or 8-oxo-7,8-dihydroguanine, a major oxidatively damaged DNA lesion⁸⁵⁻⁸⁷. Consequently, tumors from MAP patients with dysfunctional *MUTYH* protein display an excess of somatic mutations with a strong bias toward C:G > to A:T

transversions at NpCpA or NpCpT sites in multiple genes, including *APC* and *KRAS*⁸⁸⁻⁹⁰. A molecular hallmark of cancers caused by MAP is the presence of the somatic *KRAS* c.34G>T mutation^{91,92}. MAP patients show substantial variability in clinical features but usually present with an attenuated polyposis phenotype, showing fewer than 100 adenomas, although a few MAP patients with CRC without polyps have also been reported. The evidence that monoallelic variants confer an elevated CRC risk is somewhat controversial. In a large population-based series, biallelic *MUTYH* variant carriers showed a 28-fold increased risk for CRC, while monoallelic *MUTYH* variants were not associated with an increased CRC risk⁹³. However, in other studies, a small increased risk for CRC was reported for *MUTYH* monoallelic variant carriers^{94,95}. Win et al. reported that the CRC risk for monoallelic variant carriers depends on family history and can be sufficiently high to warrant consideration of more intensive CRC screening than for the general population. CRC risk is higher for monoallelic carriers of Y179C than for G396D⁹⁶. A previous study reported that biallelic *MUTYH* carriers have an increased risk of bladder and ovarian cancers, while *MUTYH* monoallelic carriers have an increased risk of gastric, liver, breast and endometrial cancers⁹⁷.

Polymerase proofreading-associated polyposis

Germline pathogenic variants affecting the exonuclease domain of *POLE* and *POLD1* predispose patients to multiple colorectal adenomas and carcinomas, causing so-called polymerase proofreading-associated polyposis (PPAP; MIM 615083, 612591)⁹⁸⁻¹⁰². PPAP is an autosomal dominant disease with a high penetrance⁹⁸. In addition to multiple adenomas and CRC, variant carriers also present with extra colonic cancers, such as endometrial, ovarian, brain, pancreatic, and small intestinal cancer and melanoma¹⁰³⁻¹⁰⁶. A recent study indicated that PPAP constitutes 0.1-0.4% of familial cancer cases, reaching 0.3-0.7% when only CRC and polyposis are considered¹⁰⁷. Although the precise risk and mean age of CRC development are not clear, a study found patients with variants in *POLE* to have a 28% risk and patients with *POLD1* variants to have an 82% to 90% risk of CRC by the age of 70 years¹⁰⁸. *POLE* and *POLD1* encode the catalytic subunits of DNA polymerases epsilon and delta, respectively. Polymerase epsilon and delta are involved in DNA replication of the leading and lagging strands and possess an accurate proofreading domain that removes incorrectly inserted nucleotides during DNA replication¹⁰⁹. While the majority of CRCs from *POLE* or *POLD1* variant carriers are MMR proficient, a subset of CRCs in *POLE* variant carriers showed MMR deficiency without germline MMR gene variants¹¹⁰. De novo variants in *POLE* have been identified in several singletons⁹⁹, but the prevalence of de novo *POLE* variants remains to be determined. Tumors

from *POLE* and *POLD1* pathogenic variant carriers show an ultrahypermutated phenotype with the number of somatic mutations exceeding 100 mutations/Mb^{111, 112}. *POLE* defects are associated with signature SBS10 and show an excess of C:G>A:T and C:G>T:A^{113, 114}. Thus far, no clear signature has been described for *POLD1*-mutated CRCs.

NTHL1-associated tumor syndrome

In 2015, a rare recessive inherited form of polyposis and CRC syndrome that is caused by biallelic pathogenic variants in the base excision repair gene *NTHL1* was discovered¹¹⁵. After the discovery, several additional families from different ethnic groups with biallelic germline variants in *NTHL1* in a homozygous or compound heterozygous state were reported¹¹⁶⁻¹²². Different extracolonic malignancies were observed in individuals with biallelic germline *NTHL1* variants, including malignancies of the endometrium, breast and duodenum^{115, 116, 119}. Based on the frequency of loss-of-function (LoF) variants in the publicly available database, the incidence of *NTHL1* deficiency is estimated to be 1:114,770, approximately fivefold lower than the incidence of *MAP* (1:19,079)⁶¹. Endonuclease III-like protein 1, encoded by *NTHL1*, is a bifunctional glycosylase involved in base excision repair that recognizes and removes oxidized pyrimidines¹²³. Tumors from biallelic *NTHL1* LoF variant carriers show a bias toward C>T transitions at non-CpG sites^{115, 124} with a unique mutational signature referred to as signature SBS30¹²⁴. Signature 30 has previously been identified in one patient with breast cancer¹²⁵. Retrospective analysis of tumor and germline sequencing data of this breast cancer patient revealed a heterozygous germline *NTHL1* variant with loss of heterozygosity in the tumor¹²⁴.

MSH3-associated polyposis

Another polyposis syndrome with a recessive inheritance pattern is referred to as *MSH3*-associated polyposis (MIM 617100)¹²⁶. After whole-exome sequencing (WES) of leukocyte DNA from 102 unrelated individuals with unexplained adenomatous polyposis, two unrelated individuals with compound heterozygous LoF germline variants in *MSH3* were identified, suggesting that *MSH3* mutations represent an additional recessive subtype of colorectal adenomatous polyposis¹²⁶. The tumors from the carriers demonstrated high microsatellite instability of di- and tetranucleotides (Elevated Microsatellite Alterations at Selected Tetranucleotide repeats (EMAST)¹²⁷) and immunohistochemical loss of *MSH3* in normal and tumor tissues¹²⁶. The associated phenotype was characterized by the presence of colorectal and duodenal adenomas, CRC, gastric cancer and early-onset astrocytoma¹²⁶.

Constitutional MMR deficiency syndrome

Constitutional MMR deficiency (CMMRD; MIM 276300) syndrome is a rare autosomal recessive childhood cancer predisposition syndrome caused by biallelic pathogenic germline variants in one MMR gene (*MLH1*, *MSH2*, *MSH6* and *PMS2*). CMMRD is characterized by a high risk of developing a broad spectrum of malignancies during childhood and adolescence, including mainly T-cell non-Hodgkin lymphomas, high-grade gliomas and gastrointestinal tumors, mainly CRC tumors. Another characteristic of CMMRD is café-au-lait maculae (CALM)^{128, 129}. Remarkably, a large proportion of CMMRD patients develop multiple synchronous adenomas ranging from a few up to > 100 polyps, mimicking attenuated familial adenomatous polyposis¹³⁰⁻¹³². Polyps in CMMRD can also histologically resemble those in juvenile polyposis¹³¹.

Serrated polyposis syndrome

Serrated polyposis syndrome (SPS; MIM 617108) was previously known as hyperplastic polyposis syndrome (HPS). SPS is characterized by the presence of multiple serrated polyps throughout the colon and rectum and is associated with an increased risk of CRC for affected individuals and their first-degree relatives¹³³⁻¹³⁵. The prevalence of SPS is estimated to be 1:2000 in the general population¹³⁴. In 2014, Gala et al. reported the association between SPS and *RNF43* by identifying a novel germline variant in two individuals with multiple serrated polyps¹³⁶. Subsequently, the role of *RNF43* germline variants as the cause of multiple serrated polyps was supported by several other studies¹³⁷⁻¹³⁹. The study by Yan et al. showed loss of the remaining wild-type allele from carriers through somatic mutations or loss of heterozygosity, adding the potential role of *RNF43* in the development of colonic serrated neoplasia¹³⁸. Buchanan et al. proposed that mutations in *RNF43* might account for only a small proportion of SPS, and consequently, there is no need for routine germline testing of *RNF43* in individuals who meet the criteria for SPS¹⁴⁰.

Hereditary mixed polyposis syndrome

Hereditary mixed polyposis syndrome (HMPS1 MIM 601228) is a rare autosomal dominant disorder that is associated with an increased risk of CRC, characterized by polyps of multiple and mixed morphologies, including serrated lesions, Peutz-Jeghers polyps, juvenile polyps and conventional adenomas¹⁴¹⁻¹⁴⁴. The genetic etiology for HMPS1 was first described in 2012, when a 40-kb duplication in the 5' regulatory region of *GREM1* was identified as a causal mutation in families of Ashkenazi Jewish origin and was shown to lead to increased and ectopic expression of *GREM1* in the colonic mucosa¹⁴⁴. Excess GREM1 protein

levels suppress bone morphogenetic protein ¹⁴⁴, allowing epithelial cells to retain stem cell-like properties, form ectopic crypts and ultimately become neoplastic ¹⁴⁵. The 40-Kb duplication has been identified in 1:184 Ashkenazi Jewish individuals with a personal or familial history of polyposis or CRC ¹⁴⁶. In addition to the founder Ashkenazi duplication, several other *GREM1* variants were identified in families with polyposis and CRC ¹⁴⁷⁻¹⁴⁹.

Hamartomatous polyposis syndromes

Hamartomatous polyposis syndromes (HPSs) are a rare heterozygous group of disorders that are inherited in an autosomal-dominant manner and are characterized by the development of hamartomatous polyps of the gastrointestinal tract. Hamartomatous polyposis syndromes have malignant potential for the development of CRC as well as extracolonic cancers ⁶³. These conditions account for less than 1% of CRC cases and occur at approximately one-tenth of the frequency of adenomatous polyposis syndromes ^{150, 151}. The hamartomatous polyposis syndromes include juvenile polyposis syndrome (JPS), Peutz-Jegher's syndrome (PJS) and PTEN hamartoma tumor syndrome (PHTS).

Juvenile polyposis syndrome (JPS)

JPS is characterized by the development of multiple gastrointestinal polyps, the most common location of which is the colon (98%). Patients with JPS syndrome have a high risk of colon cancer, and there is also an increased risk of gastroduodenal cancer. Pathogenic germline variants in *SMAD4* or *BMPR1A* are found in approximately 20-60% of JPS patients ⁶³.

Peutz-Jeghers syndrome (PJS)

PJS is caused by germline variants in *STK11* (previously known as *LKB1*) and is characterized by multiple characteristic hamartomatous polyps in the gastrointestinal tract associated with mucocutaneous pigmentation. Patients with PJS have an increased risk for CRC and extra colonic cancers ⁶³.

PTEN hamartoma tumor syndrome (PHTS)

Germline variants in the tumor suppressor gene *PTEN* are responsible for a group of phenotypically diverse conditions, which have collectively been called PTEN hamartoma syndrome (PHTS) ^{63, 134, 152}. PHTS includes Cowden syndrome (CS) and Bannayan-Riley-Ruvalcaba syndrome (BRRS), both of which are inherited in an autosomal dominant pattern ^{151, 153, 154}. CS is rarely identified before adulthood and is characterized by multiple developmentally disorganized benign growths,

or hamartomas, with an increased risk of both benign and malignant tumors¹⁵⁵. Individuals with CS are at risk for developing breast, thyroid, endometrial, colon, skin and renal cancers¹⁵⁶. BRRS patients show gastrointestinal hamartomatous polyps, lipomas, macrocephaly and developmental delay¹⁵².

Missing heritable factors in CRC and polyposis

The exact contribution of heritable factors to CRC and polyposis is still not fully understood. Based on Nordic twin and family studies, it has been estimated that 12-35% of all CRCs are linked to genetic factors^{8, 157}. Later, estimates for heritability of CRC decreased to approximately 15% of all CRC cases^{158, 159}. The currently known high-penetrant Mendelian polyposis and/or CRC syndromes can only explain 5-10% of all CRC cases^{8, 60, 160, 161}. In the case of polyposis, the genetic causes remain unexplained in approximately 20% of polyposis cases¹⁶². In approximately 60% of MMRD CRCs without somatic *MLH1* promoter hypermethylation, no underlying germline MMR variants are known. These patients are referred to as having suspected Lynch syndrome (sLS) or Lynch-like syndrome (LLS)¹⁶³. Studies have shown that patients with double somatic MMR pathogenic variants can still have hereditary CRC caused by genes involved in DNA repair since they can lead to acquired pathogenic variants in the MMR genes¹⁶⁴⁻¹⁶⁶. The genetic background is unknown for 50-60% of hereditary nonpolyposis colorectal cancer (HNPCC) families who fulfil the Amsterdam criteria⁴⁵ but do not have a mutation in one of the MMR genes (MMRP), referred to as familial colorectal cancer type X (FCCTX)¹⁶⁷. In addition to the identification of rare high-penetrant risk genes contributing to the heredity of CRC, it is estimated that common variants may explain approximately 12% of the relative risk for CRC^{14, 16, 161, 168}. In more than approximately one-third of CRC patients with a suspected hereditary cause, the underlying genetic factors remain unexplained¹⁵⁷. It is important to resolve this issue with heritability, and the identification of genetic factors has important implications for the carriers and their families, as it helps risk assessment, directs clinical management, and guides preventive and therapeutic options^{10, 169}.

Novel candidate genes for CRC and polyposis

Recently, different candidate genes have been identified but require further evidence to be implemented in routine genetic testing. New candidate genes have been proposed for predisposition to hereditary CRC and polyposis, such as *BUB1*¹⁷⁰, *BUB3*¹⁷⁰, *FAN1*⁵², *LRP6*¹⁷¹, *RPS20*⁵⁴, *FOCAD*¹⁷², *PTPN12*¹⁷¹, *GALTN12*^{173, 174}, *MIA3*¹⁷⁵ and the constitutional epigenetic silencing of *PTPRJ*¹⁷⁶. Recently, *MCM8* was proposed for predisposition to CRC with a recessive pattern of

inheritance¹⁷⁷. In a systematic review performed to validate the association between variants in *RPS20*, *FANCM*, *FAN1*, *TP53*, *BUB1*, *BUB3*, *LRP6* and *PTPN12* and the development of CRC, the evidence supports the association between variants in *RPS20* and CRC but not in the other candidate genes¹⁷⁸.

Outline of this thesis

The aim of this thesis is to study the underlying genetic causes of unexplained polyposis and CRC. In particular, the role of *POLE*, *POLD1*, *APC* and *NTHL1* in unexplained cases was studied.

Chapter 2 describes the assessment of the prevalence of *POLE* p.(Leu424Val) and *POLD1* p.(Ser478Asn) in a Dutch series of index patients with unexplained familial early onset CRC and polyposis. In this study, we analyzed phenotypes and tumor characteristics in *POLE* variant carriers. We proposed that MMR deficiency in the tumors from *POLE* p.(Leu424Val) carriers is due to secondary MMR somatic mutation resulting from the hypermutation phenotype caused by the *POLE* variants.

In **Chapter 3**, the sequencing of the exonuclease domains of *POLE* and *POLD1* in unexplained index patients with multiple colorectal polyps is described in search for novel germline variants in these genes.

Chapter 4 focuses on screening of *APC* for mosaic and deep intronic variants in unexplained colorectal polyposis patients to study their role as predisposing factors for polyposis and CRC in this cohort.

Chapter 5 shows the molecular and clinical characterization of the tumor spectrum of individuals with biallelic LoF germline variants in *NTHL1*. To establish the disease phenotype of individuals with *NTHL1* deficiency, we identified individuals with biallelic LoF germline variants in *NTHL1* and performed mutational signature analysis on different tumor types from these individuals to determine the association between *NTHL1* deficiency and tumor development.

In **Chapter 6**, the role of monoallelic LoF germline variants in *NTHL1* in the risk of polyposis and/or CRC is investigated. Finally, **Chapter 7** provides a general discussion of the thesis and future perspectives.

References

1. Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394-424.
2. Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021;71:209-249.
3. Mattiuzzi C, Sanchis-Gomar F, Lippi G. Concise update on colorectal cancer epidemiology. *Ann Transl Med* 2019;7:609.
4. Ferlay J, Colombet M, Soerjomataram I, et al. Cancer statistics for the year 2020: An overview. *Int J Cancer* 2021.
5. Xi Y, Xu P. Global colorectal cancer burden in 2020 and projections to 2040. *Transl Oncol* 2021;14:101174.
6. Grady WM, Carethers JM. Genomic and epigenetic instability in colorectal cancer pathogenesis. *Gastroenterology* 2008;135:1079-99.
7. Goel A, Nagasaka T, Arnold CN, et al. The CpG island methylator phenotype and chromosomal instability are inversely correlated in sporadic colorectal cancer. *Gastroenterology* 2007;132:127-38.
8. Lichtenstein P, Holm NV, Verkasalo PK, et al. Environmental and heritable factors in the causation of cancer--analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 2000;343:78-85.
9. Monahan KJ, Bradshaw N, Dolwani S, et al. Guidelines for the management of hereditary colorectal cancer from the British Society of Gastroenterology (BSG)/Association of Coloproctology of Great Britain and Ireland (ACPGBI)/United Kingdom Cancer Genetics Group (UKCGG). *Gut* 2020;69:411-444.
10. Jasperson KW, Tuohy TM, Neklason DW, et al. Hereditary and familial colon cancer. *Gastroenterology* 2010;138:2044-58.
11. Broderick P, Carvajal-Carmona L, Pittman AM, et al. A genome-wide association study shows that common alleles of SMAD7 influence colorectal cancer risk. *Nat Genet* 2007;39:1315-7.
12. Dunlop MG, Dobbins SE, Farrington SM, et al. Common variation near CDKN1A, POLD3 and SHROOM2 influences colorectal cancer risk. *Nat Genet* 2012;44:770-6.
13. Houlston RS, Cheadle J, Dobbins SE, et al. Meta-analysis of three genome-wide association studies identifies susceptibility loci for colorectal cancer at 1q41, 3q26.2, 12q13.13 and 20q13.33. *Nat Genet* 2010;42:973-7.
14. Law PJ, Timofeeva M, Fernandez-Rozadilla C, et al. Association analyses identify 31 new risk loci for colorectal cancer susceptibility. *Nat Commun* 2019;10:2154.
15. Orlando G, Law PJ, Palin K, et al. Variation at 2q35 (PNKD and TMBIM1) influences colorectal cancer risk and identifies a pleiotropic effect with inflammatory bowel disease. *Hum Mol Genet* 2016;25:2349-2359.
16. Schmit SL, Edlund CK, Schumacher FR, et al. Novel Common Genetic Susceptibility Loci for Colorectal Cancer. *J Natl Cancer Inst* 2019;111:146-157.
17. Tomlinson IP, Webb E, Carvajal-Carmona L, et al. A genome-wide association study identifies colorectal cancer susceptibility loci on chromosomes 10p14 and 8q23.3. *Nat Genet* 2008;40:623-30.
18. Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;61:759-67.

19. Valle L, Vilar E, Tavtigian SV, et al. Genetic predisposition to colorectal cancer: syndromes, genes, classification of genetic variants and implications for precision medicine. *J Pathol* 2019;247:574-588.
20. Kastrinos F, Samadder NJ, Burt RW. Use of Family History and Genetic Testing to Determine Risk of Colorectal Cancer. *Gastroenterology* 2020;158:389-403.
21. Hampel H, Kalady MF, Pearlman R, et al. Hereditary Colorectal Cancer. *Hematol Oncol Clin North Am* 2022;36:429-447.
22. Lynch HT, Snyder CL, Shaw TG, et al. Milestones of Lynch syndrome: 1895-2015. *Nat Rev Cancer* 2015;15:181-94.
23. Terradas M, Munoz-Torres PM, Belhadj S, et al. Contribution to colonic polyposis of recently proposed predisposing genes and assessment of the prevalence of NTHL1- and MSH3-associated polyposes. *Hum Mutat* 2019;40:1910-1923.
24. Ma H, Brosens LAA, Offerhaus GJA, et al. Pathology and genetics of hereditary colorectal cancer. *Pathology* 2018;50:49-59.
25. Hampel H, Frankel WL, Martin E, et al. Feasibility of screening for Lynch syndrome among patients with colorectal cancer. *J Clin Oncol* 2008;26:5783-8.
26. Vasen HF, Offerhaus GJ, den Hartog Jager FC, et al. The tumour spectrum in hereditary non-polyposis colorectal cancer: a study of 24 kindreds in the Netherlands. *Int J Cancer* 1990;46:31-4.
27. Dowty JG, Win AK, Buchanan DD, et al. Cancer risks for MLH1 and MSH2 mutation carriers. *Hum Mutat* 2013;34:490-7.
28. Baglietto L, Lindor NM, Dowty JG, et al. Risks of Lynch syndrome cancers for MSH6 mutation carriers. *J Natl Cancer Inst* 2010;102:193-201.
29. Senter L, Clendenning M, Sotamaa K, et al. The clinical phenotype of Lynch syndrome due to germ-line PMS2 mutations. *Gastroenterology* 2008;135:419-28.
30. ten Broeke SW, Brohet RM, Tops CM, et al. Lynch syndrome caused by germline PMS2 mutations: delineating the cancer risk. *J Clin Oncol* 2015;33:319-25.
31. Win AK, Lindor NM, Young JP, et al. Risks of primary extracolonic cancers following colorectal cancer in lynch syndrome. *J Natl Cancer Inst* 2012;104:1363-72.
32. Cox VL, Saeed Bamashmos AA, Foo WC, et al. Lynch Syndrome: Genomics Update and Imaging Review. *Radiographics* 2018;38:483-499.
33. Ligtenberg MJ, Kuiper RP, Chan TL, et al. Heritable somatic methylation and inactivation of MSH2 in families with Lynch syndrome due to deletion of the 3' exons of TACSTD1. *Nat Genet* 2009;41:112-7.
34. Miyaki M, Konishi M, Tanaka K, et al. Germline mutation of MSH6 as the cause of hereditary nonpolyposis colorectal cancer. *Nat Genet* 1997;17:271-2.
35. Bronner CE, Baker SM, Morrison PT, et al. Mutation in the DNA mismatch repair gene homologue hMLH1 is associated with hereditary non-polyposis colon cancer. *Nature* 1994;368:258-61.
36. Nicolaides NC, Papadopoulos N, Liu B, et al. Mutations of two PMS homologues in hereditary nonpolyposis colon cancer. *Nature* 1994;371:75-80.
37. Fishel R, Lescoe MK, Rao MR, et al. The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer. *Cell* 1993;75:1027-38.
38. Leach FS, Nicolaides NC, Papadopoulos N, et al. Mutations of a mutS homolog in hereditary nonpolyposis colorectal cancer. *Cell* 1993;75:1215-25.
39. Giardiello FM, Allen JI, Axilbund JE, et al. Guidelines on genetic evaluation and management of Lynch syndrome: a consensus statement by the US Multi-Society Task Force on colorectal cancer. *Gastroenterology* 2014;147:502-26.
40. Jiricny J. The multifaceted mismatch-repair system. *Nat Rev Mol Cell Biol* 2006;7:335-46.
41. Li GM. Mechanisms and functions of DNA mismatch repair. *Cell Res* 2008;18:85-98.

42. Gausachs M, Mur P, Corral J, et al. MLH1 promoter hypermethylation in the analytical algorithm of Lynch syndrome: a cost-effectiveness study. *Eur J Hum Genet* 2012;20:762-8.
43. Loukola A, Eklin K, Laiho P, et al. Microsatellite marker analysis in screening for hereditary nonpolyposis colorectal cancer (HNPCC). *Cancer Res* 2001;61:4545-9.
44. Aaltonen LA, Salovaara R, Kristo P, et al. Incidence of hereditary nonpolyposis colorectal cancer and the feasibility of molecular screening for the disease. *N Engl J Med* 1998;338:1481-7.
45. Vasen HF, Mecklin JP, Khan PM, et al. The International Collaborative Group on Hereditary Non-Polyposis Colorectal Cancer (ICG-HNPCC). *Dis Colon Rectum* 1991;34:424-5.
46. Francisco I, Albuquerque C, Lage P, et al. Familial colorectal cancer type X syndrome: two distinct molecular entities? *Fam Cancer* 2011;10:623-31.
47. Garcia FAO, de Andrade ES, de Campos Reis Galvão H, et al. New insights on familial colorectal cancer type X syndrome. *Sci Rep* 2022;12:2846.
48. Martín-Morales L, Garre P, Lorca V, et al. BRIP1, a Gene Potentially Implicated in Familial Colorectal Cancer Type X. *Cancer Prev Res (Phila)* 2021;14:185-194.
49. Sánchez-Tomé E, Rivera B, Perea J, et al. Genome-wide linkage analysis and tumoral characterization reveal heterogeneity in familial colorectal cancer type X. *J Gastroenterol* 2015;50:657-66.
50. Nieminen TT, Abdel-Rahman WM, Ristimäki A, et al. BMPR1A mutations in hereditary nonpolyposis colorectal cancer without mismatch repair deficiency. *Gastroenterology* 2011;141:e23-6.
51. Garre P, Martín L, Sanz J, et al. BRCA2 gene: a candidate for clinical testing in familial colorectal cancer type X. *Clin Genet* 2015;87:582-7.
52. Seguí N, Mina LB, Lázaro C, et al. Germline Mutations in FAN1 Cause Hereditary Colorectal Cancer by Impairing DNA Repair. *Gastroenterology* 2015;149:563-6.
53. Kim IJ, Ku JL, Kang HC, et al. Mutational analysis of OGG1, MYH, MTH1 in FAP, HNPCC and sporadic colorectal cancer patients: R154H OGG1 polymorphism is associated with sporadic colorectal cancer patients. *Hum Genet* 2004;115:498-503.
54. Nieminen TT, O'Donohue MF, Wu Y, et al. Germline mutation of RPS20, encoding a ribosomal protein, causes predisposition to hereditary nonpolyposis colorectal carcinoma without DNA mismatch repair deficiency. *Gastroenterology* 2014;147:595-598.e5.
55. Schulz E, Klampfl P, Holzapfel S, et al. Germline variants in the SEMA4A gene predispose to familial colorectal cancer type X. *Nat Commun* 2014;5:5191.
56. Martín-Morales L, Feldman M, Vershinin Z, et al. SETD6 dominant negative mutation in familial colorectal cancer type X. *Hum Mol Genet* 2017;26:4481-4493.
57. Dominguez-Valentin M, Therkildsen C, Da Silva S, et al. Familial colorectal cancer type X: genetic profiles and phenotypic features. *Mod Pathol* 2015;28:30-6.
58. Nejadtaghi M, Jafari H, Farrokhi E, et al. Familial Colorectal Cancer Type X (FCCTX) and the correlation with various genes-A systematic review. *Curr Probl Cancer* 2017;41:388-397.
59. Leoz ML, Carballal S, Moreira L, et al. The genetic basis of familial adenomatous polyposis and its implications for clinical practice and risk management. *Appl Clin Genet* 2015;8:95-107.
60. Yurgelun MB, Kulke MH, Fuchs CS, et al. Cancer Susceptibility Gene Mutations in Individuals With Colorectal Cancer. *J Clin Oncol* 2017;35:1086-1095.
61. Weren RD, Ligtenberg MJ, Geurts van Kessel A, et al. NTHL1 and MUTYH polyposis syndromes: two sides of the same coin? *J Pathol* 2018;244:135-142.
62. Samadder NJ, Baffy N, Giridhar KV, et al. Hereditary Cancer Syndromes-A Primer on Diagnosis and Management, Part 2: Gastrointestinal Cancer Syndromes. *Mayo Clin Proc* 2019;94:1099-1116.

63. Kidambi TD, Kohli DR, Samadder NJ, et al. Hereditary Polyposis Syndromes. *Curr Treat Options Gastroenterol* 2019;17:650-665.
64. Bodmer WF, Bailey CJ, Bodmer J, et al. Localization of the gene for familial adenomatous polyposis on chromosome 5. *Nature* 1987;328:614-6.
65. Kinzler KW, Nilbert MC, Su LK, et al. Identification of FAP locus genes from chromosome 5q21. *Science* 1991;253:661-5.
66. Nishisho I, Nakamura Y, Miyoshi Y, et al. Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science* 1991;253:665-9.
67. Groden J, Thliveris A, Samowitz W, et al. Identification and characterization of the familial adenomatous polyposis coli gene. *Cell* 1991;66:589-600.
68. Fearnhead NS, Britton MP, Bodmer WF. The ABC of APC. *Hum Mol Genet* 2001;10:721-33.
69. Mishra N, Hall J. Identification of patients at risk for hereditary colorectal cancer. *Clin Colon Rectal Surg* 2012;25:67-82.
70. Fearnhead NS, Wilding JL, Bodmer WF. Genetics of colorectal cancer: hereditary aspects and overview of colorectal tumorigenesis. *Br Med Bull* 2002;64:27-43.
71. Syngal S, Brand RE, Church JM, et al. ACG clinical guideline: Genetic testing and management of hereditary gastrointestinal cancer syndromes. *Am J Gastroenterol* 2015;110:223-62; quiz 263.
72. Peters U, Bien S, Zubair N. Genetic architecture of colorectal cancer. *Gut* 2015;64:1623-36.
73. Bisgaard ML, Fenger K, Bülow S, et al. Familial adenomatous polyposis (FAP): frequency, penetrance, and mutation rate. *Hum Mutat* 1994;3:121-5.
74. Farrington SM, Dunlop MG. Mosaicism and sporadic familial adenomatous polyposis. *Am J Hum Genet* 1999;64:653-8.
75. Aretz S, Uhlhaas S, Caspari R, et al. Frequency and parental origin of de novo APC mutations in familial adenomatous polyposis. *Eur J Hum Genet* 2004;12:52-8.
76. Hes FJ, Nielsen M, Bik EC, et al. Somatic APC mosaicism: an underestimated cause of polyposis coli. *Gut* 2008;57:71-6.
77. Talseth-Palmer BA. The genetic basis of colonic adenomatous polyposis syndromes. *Hered Cancer Clin Pract* 2017;15:5.
78. Al-Tassan N, Chmiel NH, Maynard J, et al. Inherited variants of MYH associated with somatic G:C->T:A mutations in colorectal tumors. *Nat Genet* 2002;30:227-32.
79. Jones S, Emmerson P, Maynard J, et al. Biallelic germline mutations in MYH predispose to multiple colorectal adenoma and somatic G:C->T:A mutations. *Hum Mol Genet* 2002;11:2961-7.
80. Sieber OM, Lipton L, Crabtree M, et al. Multiple colorectal adenomas, classic adenomatous polyposis, and germ-line mutations in MYH. *N Engl J Med* 2003;348:791-9.
81. Sampson JR, Dolwani S, Jones S, et al. Autosomal recessive colorectal adenomatous polyposis due to inherited mutations of MYH. *Lancet* 2003;362:39-41.
82. Kastrinos F, Syngal S. Inherited colorectal cancer syndromes. *Cancer J* 2011;17:405-15.
83. Cleary SP, Cotterchio M, Jenkins MA, et al. Germline MutY human homologue mutations and colorectal cancer: a multisite case-control study. *Gastroenterology* 2009;136:1251-60.
84. Slupska MM, Baikalov C, Luther WM, et al. Cloning and sequencing a human homolog (hMYH) of the Escherichia coli mutY gene whose function is required for the repair of oxidative DNA damage. *J Bacteriol* 1996;178:3885-92.
85. Slupska MM, Luther WM, Chiang JH, et al. Functional expression of hMYH, a human homolog of the Escherichia coli MutY protein. *J Bacteriol* 1999;181:6210-3.
86. Cadet J, Davies KJA. Oxidative DNA damage & repair: An introduction. *Free Radic Biol Med* 2017;107:2-12.

87. Chen J, Huang Z, Wu X, et al. Oxidative stress induces different tissue dependent effects on Mutyh-deficient mice. *Free Radic Biol Med* 2019;143:482-493.
88. Rashid M, Fischer A, Wilson CH, et al. Adenoma development in familial adenomatous polyposis and MUTYH-associated polyposis: somatic landscape and driver genes. *J Pathol* 2016;238:98-108.
89. Pilati C, Shinde J, Alexandrov LB, et al. Mutational signature analysis identifies MUTYH deficiency in colorectal cancers and adrenocortical carcinomas. *J Pathol* 2017;242:10-15.
90. Viel A, Bruxelles A, Meccia E, et al. A Specific Mutational Signature Associated with DNA 8-Oxoguanine Persistence in MUTYH-defective Colorectal Cancer. *EBioMedicine* 2017;20:39-49.
91. Lipton L, Halford SE, Johnson V, et al. Carcinogenesis in MYH-associated polyposis follows a distinct genetic pathway. *Cancer Res* 2003;63:7595-9.
92. Aimé A, Coulet F, Lefevre JH, et al. Somatic c.34G>T KRAS mutation: a new prescreening test for MUTYH-associated polyposis? *Cancer Genet* 2015;208:390-5.
93. Lubbe SJ, Di Bernardo MC, Chandler IP, et al. Clinical implications of the colorectal cancer risk associated with MUTYH mutation. *J Clin Oncol* 2009;27:3975-80.
94. Win AK, Hopper JL, Jenkins MA. Association between monoallelic MUTYH mutation and colorectal cancer risk: a meta-regression analysis. *Fam Cancer* 2011;10:1-9.
95. Win AK, Cleary SP, Dowty JG, et al. Cancer risks for monoallelic MUTYH mutation carriers with a family history of colorectal cancer. *Int J Cancer* 2011;129:2256-62.
96. Win AK, Dowty JG, Cleary SP, et al. Risk of colorectal cancer for carriers of mutations in MUTYH, with and without a family history of cancer. *Gastroenterology* 2014;146:1208-11. e1-5.
97. Win AK, Reece JC, Dowty JG, et al. Risk of extracolonic cancers for people with biallelic and monoallelic mutations in MUTYH. *Int J Cancer* 2016;139:1557-63.
98. Palles C, Cazier JB, Howarth KM, et al. Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet* 2013;45:136-44.
99. Valle L, Hernandez-Illan E, Bellido F, et al. New insights into POLE and POLD1 germline mutations in familial colorectal cancer and polyposis. *Hum Mol Genet* 2014;23:3506-12.
100. Spier I, Holzapfel S, Altmüller J, et al. Frequency and phenotypic spectrum of germline mutations in POLE and seven other polymerase genes in 266 patients with colorectal adenomas and carcinomas. *Int J Cancer* 2015;137:320-31.
101. Chubb D, Broderick P, Frampton M, et al. Genetic diagnosis of high-penetrance susceptibility for colorectal cancer (CRC) is achievable for a high proportion of familial CRC by exome sequencing. *J Clin Oncol* 2015;33:426-32.
102. Bellido F, Pineda M, Aiza G, et al. POLE and POLD1 mutations in 529 kindred with familial colorectal cancer and/or polyposis: review of reported cases and recommendations for genetic testing and surveillance. *Genet Med* 2016;18:325-32.
103. Church DN, Briggs SE, Palles C, et al. DNA polymerase ϵ and δ exonuclease domain mutations in endometrial cancer. *Hum Mol Genet* 2013;22:2820-8.
104. Rohlin A, Zagoras T, Nilsson S, et al. A mutation in POLE predisposing to a multi-tumour phenotype. *Int J Oncol* 2014;45:77-81.
105. Hansen MF, Johansen J, Bjornevold I, et al. A novel POLE mutation associated with cancers of colon, pancreas, ovaries and small intestine. *Fam Cancer* 2015;14:437-48.
106. Aoude LG, Heitzer E, Johansson P, et al. POLE mutations in families predisposed to cutaneous melanoma. *Fam Cancer* 2015;14:621-8.
107. Mur P, García-Mulero S, Del Valle J, et al. Role of POLE and POLD1 in familial cancer. *Genet Med* 2020.

108. Buchanan DD, Stewart JR, Clendenning M, et al. Risk of colorectal cancer for carriers of a germ-line mutation in POLE or POLD1. *Genet Med* 2018;20:890-895.
109. Nick McElhinny SA, Gordenin DA, Stith CM, et al. Division of labor at the eukaryotic replication fork. *Mol Cell* 2008;30:137-44.
110. Jansen AM, van Wezel T, van den Akker BE, et al. Combined mismatch repair and POLE/POLD1 defects explain unresolved suspected Lynch syndrome cancers. *Eur J Hum Genet* 2016;24:1089-92.
111. Campbell BB, Light N, Fabrizio D, et al. Comprehensive Analysis of Hypermutation in Human Cancer. *Cell* 2017;171:1042-1056.e10.
112. Kane DP, Shcherbakova PV. A common cancer-associated DNA polymerase ϵ mutation causes an exceptionally strong mutator phenotype, indicating fidelity defects distinct from loss of proofreading. *Cancer Res* 2014;74:1895-901.
113. Alexandrov LB, Kim J, Haradhvala NJ, et al. The repertoire of mutational signatures in human cancer. *Nature* 2020;578:94-101.
114. Alexandrov LB, Nik-Zainal S, Wedge DC, et al. Signatures of mutational processes in human cancer. *Nature* 2013;500:415-21.
115. Weren RD, Ligtenberg MJ, Kets CM, et al. A germline homozygous mutation in the base-excision repair gene NTHL1 causes adenomatous polyposis and colorectal cancer. *Nat Genet* 2015;47:668-71.
116. Rivera B, Castellsague E, Bah I, et al. Biallelic NTHL1 Mutations in a Woman with Multiple Primary Tumors. *N Engl J Med* 2015;373:1985-6.
117. Chubb D, Broderick P, Dobbins SE, et al. Rare disruptive mutations and their contribution to the heritable risk of colorectal cancer. *Nat Commun* 2016;7:11883.
118. Belhadj S, Mur P, Navarro M, et al. Delineating the Phenotypic Spectrum of the NTHL1-Associated Polyposis. *Clin Gastroenterol Hepatol* 2017;15:461-462.
119. Fostira F, Kontopodis E, Apostolou P, et al. Extending the clinical phenotype associated with biallelic NTHL1 germline mutations. *Clin Genet* 2018;94:588-589.
120. Groves A, Gleeson M, Spigelman AD. NTHL1-associate polyposis: first Australian case report. *Fam Cancer* 2019.
121. Belhadj S, Quintana I, Mur P, et al. NTHL1 biallelic mutations seldom cause colorectal cancer, serrated polyposis or a multi-tumor phenotype, in absence of colorectal adenomas. *Sci Rep* 2019;9:9020.
122. Altaraihi M, Gerdes AM, Wadt K. A new family with a homozygous nonsense variant in NTHL1 further delineated the clinical phenotype of NTHL1-associated polyposis. *Hum Genome Var* 2019;6:46.
123. Krokan HE, Bjørås M. Base excision repair. *Cold Spring Harb Perspect Biol* 2013;5:a012583.
124. Drost J, van Boxtel R, Blokzijl F, et al. Use of CRISPR-modified human stem cell organoids to study the origin of mutational signatures in cancer. *Science* 2017;358:234-238.
125. Nik-Zainal S, Davies H, Staaf J, et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature* 2016;534:47-54.
126. Adam R, Spier I, Zhao B, et al. Exome Sequencing Identifies Biallelic MSH3 Germline Mutations as a Recessive Subtype of Colorectal Adenomatous Polyposis. *Am J Hum Genet* 2016;99:337-51.
127. Carethers JM, Koi M, Tseng-Rogenski SS. EMAST is a Form of Microsatellite Instability That is Initiated by Inflammation and Modulates Colorectal Cancer Progression. *Genes (Basel)* 2015;6:185-205.
128. Wimmer K, Etzler J. Constitutional mismatch repair-deficiency syndrome: have we so far seen only the tip of an iceberg? *Hum Genet* 2008;124:105-22.

129. Bakry D, Aronson M, Durno C, et al. Genetic and clinical determinants of constitutional mismatch repair deficiency syndrome: report from the constitutional mismatch repair deficiency consortium. *Eur J Cancer* 2014;50:987-96.
130. Jasperson KW, Samowitz WS, Burt RW. Constitutional mismatch repair-deficiency syndrome presenting as colonic adenomatous polyposis: clues from the skin. *Clin Genet* 2011;80:394-7.
131. Levi Z, Kariv R, Barnes-Kedar I, et al. The gastrointestinal manifestation of constitutional mismatch repair deficiency syndrome: from a single adenoma to polyposis-like phenotype and early onset cancer. *Clin Genet* 2015;88:474-8.
132. Aronson M, Gallinger S, Cohen Z, et al. Gastrointestinal Findings in the Largest Series of Patients With Hereditary Biallelic Mismatch Repair Deficiency Syndrome: Report from the International Consortium. *Am J Gastroenterol* 2016;111:275-84.
133. East JE, Vieth M, Rex DK. Serrated lesions in colorectal cancer screening: detection, resection, pathology and surveillance. *Gut* 2015;64:991-1000.
134. Patel R, Hyer W. Practical management of polyposis syndromes. *Frontline Gastroenterol* 2019;10:379-387.
135. Stanich PP, Pearlman R. Hereditary or Not? Understanding Serrated Polyposis Syndrome. *Curr Treat Options Gastroenterol* 2019;17:692-701.
136. Gala MK, Mizukami Y, Le LP, et al. Germline mutations in oncogene-induced senescence pathways are associated with multiple sessile serrated adenomas. *Gastroenterology* 2014;146:520-9.
137. Taupin D, Lam W, Rangiah D, et al. A deleterious RNF43 germline mutation in a severely affected serrated polyposis kindred. *Hum Genome Var* 2015;2:15013.
138. Yan HHN, Lai JCW, Ho SL, et al. RNF43 germline and somatic mutation in serrated neoplasia pathway and its association with BRAF mutation. *Gut* 2017;66:1645-1656.
139. Quintana I, Mejías-Luque R, Terradas M, et al. Evidence suggests that germline RNF43 mutations are a rare cause of serrated polyposis. *Gut* 2018;67:2230-2232.
140. Buchanan DD, Clendenning M, Zhuoer L, et al. Lack of evidence for germline RNF43 mutations in patients with serrated polyposis syndrome from a large multinational study. *Gut* 2017;66:1170-1172.
141. Thomas HJ, Whitelaw SC, Cottrell SE, et al. Genetic mapping of hereditary mixed polyposis syndrome to chromosome 6q. *Am J Hum Genet* 1996;58:770-6.
142. Whitelaw SC, Murday VA, Tomlinson IP, et al. Clinical and molecular features of the hereditary mixed polyposis syndrome. *Gastroenterology* 1997;112:327-34.
143. Rozen P, Samuel Z, Brazowski E. A prospective study of the clinical, genetic, screening, and pathologic features of a family with hereditary mixed polyposis syndrome. *Am J Gastroenterol* 2003;98:2317-20.
144. Jaeger E, Leedham S, Lewis A, et al. Hereditary mixed polyposis syndrome is caused by a 40-kb upstream duplication that leads to increased and ectopic expression of the BMP antagonist GREM1. *Nat Genet* 2012;44:699-703.
145. Davis H, Irshad S, Bansal M, et al. Aberrant epithelial GREM1 expression initiates colonic tumorigenesis from cells outside the stem cell niche. *Nat Med* 2015;21:62-70.
146. Lieberman S, Walsh T, Schechter M, et al. Features of Patients With Hereditary Mixed Polyposis Syndrome Caused by Duplication of GREM1 and Implications for Screening and Surveillance. *Gastroenterology* 2017;152:1876-1880.e1.
147. Rohlin A, Eiengård F, Lundstam U, et al. GREM1 and POLE variants in hereditary colorectal cancer syndromes. *Genes Chromosomes Cancer* 2016;55:95-106.
148. McKenna DB, Van Den Akker J, Zhou AY, et al. Identification of a novel GREM1 duplication in a patient with multiple colon polyps. *Fam Cancer* 2019;18:63-66.

149. Venkatachalam R, Verwiel ET, Kamping EJ, et al. Identification of candidate predisposing copy number variants in familial and early-onset colorectal cancer patients. *Int J Cancer* 2011;129:1635-42.
150. Gammon A, Jasperson K, Kohlmann W, et al. Hamartomatous polyposis syndromes. *Best Pract Res Clin Gastroenterol* 2009;23:219-31.
151. Manfredi M. Hereditary hamartomatous polyposis syndromes: understanding the disease risks as children reach adulthood. *Gastroenterol Hepatol (N Y)* 2010;6:185-96.
152. Macken WL, Tischkowitz M, Lachlan KL. PTEN Hamartoma tumor syndrome in childhood: A review of the clinical literature. *Am J Med Genet C Semin Med Genet* 2019;181:591-610.
153. Marsh DJ, Kum JB, Lunetta KL, et al. PTEN mutation spectrum and genotype-phenotype correlations in Bannayan-Riley-Ruvalcaba syndrome suggest a single entity with Cowden syndrome. *Hum Mol Genet* 1999;8:1461-72.
154. Smpokou P, Fox VL, Tan WH. PTEN hamartoma tumour syndrome: early tumour development in children. *Arch Dis Child* 2015;100:34-7.
155. Hobert JA, Eng C. PTEN hamartoma tumor syndrome: an overview. *Genet Med* 2009;11:687-94.
156. Gammon A, Jasperson K, Champine M. Genetic basis of Cowden syndrome and its implications for clinical practice and risk management. *Appl Clin Genet* 2016;9:83-92.
157. Czene K, Lichtenstein P, Hemminki K. Environmental and heritable causes of cancer among 9.6 million individuals in the Swedish Family-Cancer Database. *Int J Cancer* 2002;99:260-6.
158. Mucci LA, Hjelmborg JB, Harris JR, et al. Familial Risk and Heritability of Cancer Among Twins in Nordic Countries. *Jama* 2016;315:68-76.
159. Graff RE, Möller S, Passarelli MN, et al. Familial Risk and Heritability of Colorectal Cancer in the Nordic Twin Study of Cancer. *Clin Gastroenterol Hepatol* 2017;15:1256-1264.
160. Lynch HT, de la Chapelle A. Hereditary colorectal cancer. *N Engl J Med* 2003;348:919-32.
161. Huyghe JR, Bien SA, Harrison TA, et al. Discovery of common and rare genetic risk variants for colorectal cancer. *Nat Genet* 2019;51:76-87.
162. Mongin C, Coulet F, Lefevre JH, et al. Unexplained polyposis: a challenge for geneticists, pathologists and gastroenterologists. *Clin Genet* 2012;81:38-46.
163. Buchanan DD, Rosty C, Clendenning M, et al. Clinical problems of colorectal cancer and endometrial cancer cases with unknown cause of tumor mismatch repair deficiency (suspected Lynch syndrome). *Appl Clin Genet* 2014;7:183-93.
164. Morak M, Heidenreich B, Keller G, et al. Biallelic MUTYH mutations can mimic Lynch syndrome. *Eur J Hum Genet* 2014;22:1334-7.
165. Pearlman R, Frankel WL, Swanson B, et al. Prevalence and Spectrum of Germline Cancer Susceptibility Gene Mutations Among Patients With Early-Onset Colorectal Cancer. *JAMA Oncol* 2017;3:464-471.
166. Castillejo A, Vargas G, Castillejo MI, et al. Prevalence of germline MUTYH mutations among Lynch-like syndrome patients. *Eur J Cancer* 2014;50:2241-50.
167. Zetner DB, Bisgaard ML. Familial Colorectal Cancer Type X. *Curr Genomics* 2017;18:341-359.
168. Weigl K, Chang-Claude J, Hsu L, et al. Establishing a valid approach for estimating familial risk of cancer explained by common genetic variants. *Int J Cancer* 2020;146:68-75.
169. Patel SG, Boland CR. Colorectal Cancer in Persons Under Age 50: Seeking Causes and Solutions. *Gastrointest Endosc Clin N Am* 2020;30:441-455.
170. de Voer RM, Geurts van Kessel A, Weren RD, et al. Germline mutations in the spindle assembly checkpoint genes BUB1 and BUB3 are risk factors for colorectal cancer. *Gastroenterology* 2013;145:544-7.

171. de Voer RM, Hahn MM, Weren RD, et al. Identification of Novel Candidate Genes for Early-Onset Colorectal Cancer Susceptibility. *PLoS Genet* 2016;12:e1005880.
172. Weren RD, Venkatachalam R, Cazier JB, et al. Germline deletions in the tumour suppressor gene FOCAD are associated with polyposis and colorectal cancer development. *J Pathol* 2015;236:155-64.
173. Clarke E, Green RC, Green JS, et al. Inherited deleterious variants in GALNT12 are associated with CRC susceptibility. *Hum Mutat* 2012;33:1056-8.
174. Evans DR, Venkitachalam S, Revoredo L, et al. Evidence for GALNT12 as a moderate penetrance gene for colorectal cancer. *Hum Mutat* 2018;39:1092-1101.
175. Schubert SA, Ruano D, Elsayed FA, et al. Evidence for genetic association between chromosome 1q loci and predisposition to colorectal neoplasia. *Br J Cancer* 2017;117:1215-1223.
176. Venkatachalam R, Ligtenberg MJ, Hoogerbrugge N, et al. Germline epigenetic silencing of the tumor suppressor gene PTPRJ in early-onset familial colorectal cancer. *Gastroenterology* 2010;139:2221-4.
177. Golubicki M, Bonjoch L, Acuña-Ochoa JG, et al. Germline biallelic Mcm8 variants are associated with early-onset Lynch-like syndrome. *JCI Insight* 2020;5.
178. Broderick P, Dobbins SE, Chubb D, et al. Validation of Recently Proposed Colorectal Cancer Susceptibility Gene Variants in an Analysis of Families and Patients—a Systematic Review. *Gastroenterology* 2017;152:75-77.e4.

