



Universiteit
Leiden
The Netherlands

Mining the kinematics of discs to hunt for planets in formation

Izquierdo Cartagena, A.F.

Citation

Izquierdo Cartagena, A. F. (2023, December 1). *Mining the kinematics of discs to hunt for planets in formation*. Retrieved from <https://hdl.handle.net/1887/3665447>

Version: Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

License: <https://hdl.handle.net/1887/3665447>

Note: To cite this publication please use the final published version (if applicable).

Mining the kinematics of discs to hunt for planets in formation

Proefschrift

ter verkrijging van
de graad van doctor aan de Universiteit Leiden,
op gezag van rector magnificus prof. dr. ir. H. Bijl,
volgens besluit van het college voor promoties
te verdedigen op vrijdag 01 december 2023
klokke 11.15 uur
door

Andrés Felipe Izquierdo Cartagena

geboren te Palmira, Colombia
in 1996

Promotores:

Prof. dr. E. F. van Dishoeck

Prof. dr. L. Testi Università di Bologna

Co-promotor:

Dr. S. Facchini Università degli Studi di Milano

Promotiecommissie:

Prof. dr. I. A. G. Snellen

Prof. dr. M. R. Hogerheijde Universiteit Leiden

Universiteit van Amsterdam

Prof. dr. L. Hartmann University of Michigan

Prof. dr. C. P. Dullemond Ruprecht-Karls-Universität Heidelberg

Prof. dr. T. Birnstiel Ludwig-Maximilians-Universität München

Dr. P. Pinilla University College London

ISBN: 978-94-6419-981-9

Cover description: The coloured map represents azimuthal deviations from Keplerian rotation in a simulated disc perturbed by a planet like Jupiter. The grey lines follow a Voronoi mesh constructed so that its cells are more densely clustered around high values of the radial component of the velocity field.

Cover design: Andrés & Laila – Simulation credits: Giovanni Rosotti

Those who can imagine anything,
can create the impossible.
- *Alan Turing*

Table of contents

1	Introduction	1
1.1	Young Stellar Objects	2
1.1.1	Low-mass star formation	2
1.1.2	Physical and dynamical structure of discs	4
1.1.2.1	Density and temperature structure	4
1.1.2.2	Dynamical structure	6
1.1.3	Disc observations	8
1.1.3.1	Historical remarks	8
1.1.3.2	The ALMA era	9
1.2	The hunt for planets in formation	10
1.2.1	Planet signatures in gas discs: Theory	12
1.2.1.1	Analytical development	12
1.2.1.2	Semi-analytic theory and the need for numerical models	14
1.2.1.3	Numerical simulations and observational predictions	16
1.2.2	Planet signatures in gas discs: ALMA Observations	18
1.3	This thesis	21
1.4	Future prospects	24
2	The Disc Miner I.	
	A statistical framework to detect and quantify kinematical perturbations driven by young planets in discs	27
2.1	Introduction	29
2.2	Simulations and radiative transfer	30
2.3	Fitting channel maps	33
2.3.1	The Discminer package	33
2.3.2	Line intensity profile	35
2.3.3	Best-fit model of the simulation	36
2.4	Finding the kinematical footprints of planets	37
2.4.1	Intrinsic deviations from Keplerian rotation	37
2.4.2	Residual maps	39
2.4.3	Line centroid folding	41
2.4.4	Statistical methods to detect planets	43
2.5	Discussion	45

2.5.1	Comparison with other methods	45
2.5.2	Robustness of the spectral resolution of the method	49
2.5.3	Magnitude of observed planet-driven perturbations	50
2.5.4	Caveats	52
2.6	Conclusions	52
Appendices	53
2.A	Impact of the gap and emitting surfaces on the observed gas velocities and supporting figures	53
3	A new planet candidate detected in a dust gap of the disk around HD 163296 through localized kinematic signatures	59
3.1	Introduction	61
3.2	Line intensity model of the HD 163296 disc	62
3.2.1	Dataset	62
3.2.2	Discminer model setup	62
3.2.3	Parameter search with EMCEE	63
3.3	Results	66
3.3.1	Physical attributes of the ^{12}CO disc	66
3.3.1.1	Emission height	66
3.3.1.2	Line width and Brightness temperature	68
3.3.2	Residual maps	69
3.3.3	Gas gaps	71
3.3.3.1	Azimuthally averaged residuals	73
3.3.3.2	Non-axisymmetric gas substructure	74
3.3.4	Detection of planets	74
3.3.4.1	P94 perturbation	76
3.3.4.2	P261 perturbation	77
3.3.4.3	Detection significance of the localised perturbations	79
3.3.4.4	Non-detections	79
3.4	Conclusions	79
Appendices	81
3.A	Analytic propagation of errors	81
3.B	Supporting figures	85
4	The Disc Miner II.	
	Revealing gas substructures and kinematic signatures from planet-disc interaction through line profile analysis	89
4.1	Introduction	91
4.2	Observations and models	93
4.2.1	Discminer models	93
4.2.1.1	Model parameters and attributes	95
4.2.1.2	Data versus model channel maps	96
4.2.2	Line profile observables to compare data and models	98
4.2.2.1	Upper and lower emission surfaces	99
4.2.2.2	Decomposition method to extract azimuthal and meridional velocities	100

4.3	Kinematic signatures	102
4.3.1	Localised velocity and line width perturbations	105
4.3.1.1	Detections in the disc of HD 163296: Velocity	105
4.3.1.2	Detections in the disc of HD 163296: Line width	107
4.3.1.3	Non-detections	109
4.3.2	Extended velocity and line width perturbations	110
4.4	Gas substructures as sites to search for planets	118
4.4.1	Azimuthal and meridional velocity flows	119
4.4.2	Surface density gaps traced by azimuthal velocity modulations and line widths	124
4.4.3	Strong vertical dependence of radial pressure in the disc of HD 163296, around the planet candidate P94	128
4.4.4	Non-axisymmetric variations in the radial profiles of the disc around MWC 480	129
4.5	Discussion	131
4.5.1	Morphology of the P94 perturbation in HD 163296	131
4.5.2	Line broadening as a tracer of massive planets	131
4.5.3	Localised velocity perturbations are not so common	132
4.5.4	Correlation between pressure bumps and mm dust rings	132
4.6	Conclusions	133
4.6.1	Giant planets around HD 163296, MWC 480, AS 209	133
4.6.2	Non-detections in the discs of IM Lup and GM Aur	134
4.6.3	Gas gaps, and relationship between pressure bumps and dust substructures	135
Appendices	136
4.A	Supporting tables and figures	136
4.B	Azimuthal average of absolute velocity residuals	161
5	Line profile tomography of protoplanetary discs	163
5.1	Introduction	165
5.2	Modelling channel maps	166
5.3	Extraction of moment maps from line profile observables	169
5.3.1	Single-peaked sources	170
5.3.2	Double-peaked sources	173
5.3.3	Velocity and intensity profiles	176
5.4	Residual maps	179
5.4.1	Localised signatures in residual maps	180
5.4.2	Line width increments as signposts of planet-forming sites	186
5.4.3	Coherent large-scale signatures in residuals	189
5.5	Discussion	192
5.5.1	Planets in the disc of HD 135344B	192
5.5.2	Planet-driven spirals in MWC 758?	194
5.6	Conclusions	195
Appendices	197
5.A	Supporting figures	197

Bibliography	197
Nederlandse samenvatting	211
English summary	219
Resumen de la tesis	227
Publications	235
Curriculum Vitae	239
Acknowledgements	241