



Universiteit
Leiden
The Netherlands

Monitoring the Dutch Solid Start Program: developing an indicator set for municipalities to monitor their first thousand days approach

Molenaar, J.M.; Boesveld, I.C.; Kiefte-de Jong, J.C.; Struijs, J.N.

Citation

Molenaar, J. M., Boesveld, I. C., Kiefte-de Jong, J. C., & Struijs, J. N. (2022). Monitoring the Dutch Solid Start Program: developing an indicator set for municipalities to monitor their first thousand days approach. *International Journal Of Integrated Care*, 22(4).
doi:10.5334/ijic.6508

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3666115>

Note: To cite this publication please use the final published version (if applicable).



Monitoring the Dutch Solid Start Program: Developing an Indicator Set for Municipalities to Monitor their First Thousand Days-Approach

JOYCE M. MOLENAAR 

INGE C. BOESVELD 

JESSICA C. KIEFTE-DE JONG 

JEROEN N. STRUIJS 

*Author affiliations can be found in the back matter of this article

RESEARCH AND
THEORY

u[ubiquity press

ABSTRACT

Introduction: The Dutch Solid Start program aims to improve the collaboration between the medical and social sector to offer every child the best start in life. Municipalities form local coalitions of partners within the medical and social sector to support parents and children during the first thousand days. The aim of this study was to develop an indicator set for coalitions to monitor their local Solid Start program.

Methods: A modified Delphi study with three rounds was carried out among Dutch experts in Solid Start practice, policy and research (n = 39) to reach consensus.

Results: The indicator set included 19 indicators covering the three phases of the Solid Start program: preconception, pregnancy and after birth (up to two years). Prioritized indicators included both social and medical topics, among which poverty, psychological/psychiatric problems, stress, smoking, cumulation of risk factors, preconception care, low literacy, premature birth, and intellectual disability. Additionally, a development agenda was established with topics and indicators that lacked data or clear operationalization (e.g. stress, unintended pregnancy, loneliness).

Discussion and conclusion: The developed indicator set enhances the conversation between policymakers, managers, professionals and other stakeholders about the local situation and developments in order to prioritize interventions and policies. Next, the indicator set needs evaluation to assess its usefulness.

CORRESPONDING AUTHOR:

Joyce M. Molenaar

Centre for Nutrition, Prevention and Health Services,
Department of Quality of Care and Health Economics, National Institute for Public Health and the Environment, Bilthoven, The Netherlands; Health Campus The Hague/Department of Public Health and Primary Care, Leiden University Medical Center, The Hague, The Netherlands

joyce.molenaar@rivm.nl

KEYWORDS:

Delphi study; indicators; first thousand days; integration of care

TO CITE THIS ARTICLE:

Molenaar JM, Boesveld IC, Kieft-de Jong JC, Struijs JN. Monitoring the Dutch Solid Start Program: Developing an Indicator Set for Municipalities to Monitor their First Thousand Days-Approach. *International Journal of Integrated Care*, 2022; 22(4): 8, 1-12. DOI: <https://doi.org/10.5334/ijic.6508>

INTRODUCTION

Reducing perinatal health inequities and improving health outcomes for parents and children are high on the Dutch policy agenda since the early 2000s. Following alarming perinatal mortality and morbidity figures [1, 2], several policy measures were taken to improve maternity care, including the establishment of maternity care networks [3], experiments with bundled payment for maternity care [4] and the development of the 'Standard for Integrated Birth Care' [5]. Over the years, the focus of the programs shifted from the medical sector more towards the social and public health care sector, as perinatal and maternal health is strongly influenced by the wider social, economic and cultural contexts of families [6, 7]. For instance, a regional cross-sectoral approach to perinatal and maternal health, integrating the medical and social sector, was taken in the local 'Ready for a baby' program [8] and subsequent 'Healthy Pregnancy 4-All' programs [9-11]. These programs laid the foundation for the nationwide 'Solid Start' action program.

The Solid Start program was launched by the Dutch Ministry of Health, Welfare and Sport in September 2018 with the aim to give every child the best start in life by focusing on the first thousand days [12]. This period from preconception to the child's second birthday is crucial for children's further physical, mental and social development and is therefore regarded as a window of opportunity to improve population health [6, 13, 14]. The integrated approach of the Solid Start program combines medical and social services to offer better support during the first thousand days, specifically for parents in vulnerable situations. Consequently, the scope of integrated service delivery within the program is not limited to the health sector alone, but rather expanded to coordinate care and support also between the health and social sector (including public health) with its various organizations and providers (among which midwives, social workers, gynecologists, youth healthcare providers, debt counselors, and municipal officials). The Solid Start program is conceptualized and implemented over three phases: before pregnancy, during pregnancy and after birth (up to two years). Municipalities receive additional subsidies from the Ministry of Health to form local coalitions of partners within the medical and social sector, in order to tackle the region-specific challenges. Examples of region-specific challenges are unintended pregnancies, housing problems, domestic violence, and loneliness. This approach fits with the decentralization tendencies of social care in the Netherlands. Since 2015, the government has given municipalities new responsibilities in youth care, long-term care and income support, which cause local differences in policy implementation and outcomes [15]. Next to the subsidies, supportive methods were developed and offered to local coalitions. Examples include an analysis tool to map the current and desired

situation and an overview of effective interventions (e.g. prenatal home visits and 'Centering Pregnancy™': group care during pregnancy). Moreover, local coalitions receive support to develop and implement their local coalition and related programs by Pharos, which is the Dutch Centre of Expertise on Health Disparities.

The Ministry of Health commissioned the National Institute for Public Health and the Environment (Dutch abbreviation: RIVM) to monitor the implementation of the Solid Start program. To this end, an indicator set including fifteen indicators was developed in a Delphi study with experts in 2019 [16] and reported annually in order to monitor the implementation of the nationwide program and to identify whether health outcomes improve. The indicator set reflects both processes (e.g. percentage of municipalities in which youth healthcare offers prenatal home visits) and outcomes (e.g. percentage of children born prematurely or with a low birth weight). In addition, the RIVM conducted a process evaluation to collect the experiences of those involved in the Solid Start program in order to provide further insight into factors that promote and hinder the implementation. The Ministry of Health uses the results of the monitor in combination with other data sources and expert opinions to determine whether goals are being achieved and to timely adjust policies. The results of the national Solid Start monitor showed that local coalitions evolve and formalize and that the majority of them also plan to monitor their local program, or have started to do so [17-19]. However, the local coalitions generally experienced a lack of insight into which indicators to include in their local setting, where to find the data for their municipality and how to make optimal use of it. Because the national indicator set was considered less suitable for monitoring on a local level, they expressed a need for a uniform indicator set to use within their local coalition. In 2021, the RIVM started a support program that is focused on monitoring Solid Start on a local level (for additional information about the support program and its relation with the Solid Start program and national monitor see Appendix A). Key elements of this support program include learning from and with other stakeholders (both within and between local coalitions) and sharing best practices within learning communities. The local coalitions that participated in the monitoring support program considered the development of a suitable indicator set the essential first step to stimulate monitoring on a local level.

In this paper, we describe our approach in developing an indicator set to monitor the Solid Start program in Dutch local coalitions and we present this indicator set. The indicator set can be used by local coalitions to enhance the conversation between policymakers, managers, professionals and other stakeholders about the local situation and developments in order to prioritize interventions and policies. This can help to strengthen and promote integrated service delivery.

METHODS

DESIGN AND PROCEDURE

Within this mixed-methods study, we used a modified Delphi technique as a structured method to reach consensus on an indicator set to monitor Solid Start on a local level [20]. This commonly used approach in health research is suitable to synthesize knowledge from various experts with a different background or geographical location [21]. Our study had several iterative rounds of self-administered questionnaires and expert meetings (Figure 1). The study was conducted between March and June 2021.

PRIOR TO DELPHI STUDY

The study started with compiling a list of indicators originating from existing monitoring tools or documents from local coalitions, scientific and grey literature, and the indicator set used in the national Solid Start monitor [16, 22–26]. The list of possible indicators was long (in a first endeavor >350) because the scope of the first thousand days is comprehensive. As this was expected to be a burden to the participants, we decided to first select topics instead of indicators directly. One researcher (JM, health scientist) categorized and named the topics in line with existing monitoring tools and documents, and another researcher (IB, former midwife and advisor integrated maternity care organizations) cross-checked this. We categorized and named the topics based on the shared characteristics and common themes in indicators (e.g. indicators relating to a low household income, debts, receiving social benefits and stress due to finances were categorized into the topic ‘poverty’). Differences were discussed by three researchers (JM, IB and JS (expertise health economy)) until consensus was reached. We excluded topics that 1) did not have at least one operationalized indicator, or 2) exceeded the time period of the Solid Start program (i.e. beyond the first thousand days of life). Topics were classified in the three phases of Solid Start (preconception, pregnancy and after birth) with the reason to eventually get a sufficient number of indicators per phase. Some topics were relevant in more than one phase.

EXPERT PANEL

The expert panel consisted of a heterogeneous group of experts involved in Solid Start activities and experienced

with monitoring, geographically distributed over the Netherlands (i.e. both rural and urban areas in the northern, eastern, western and southern parts of the country). We aimed for a balanced representation of experts in practice, policy and research (purposive sampling), including managers of local coalitions, policy makers, policy advisors, epidemiologists, researchers, educators, primary and secondary healthcare providers (e.g. midwife, nurse, gynecologist, pediatrician) and social workers. We invited members of the monitoring support program (Appendix A) and their network (‘snowballing method’), and we recruited participants through social media, Solid Start newsletters and webpages, and personal invitation. Those interested received more information about the aim, design and voluntary nature of the study. The views of participants all received equal weight during the study.

DELPHI ROUND 1: QUESTIONNAIRE

In an online questionnaire, the Delphi panel was instructed to rate 121 topics based on relevance to monitor Solid Start on a local level on a nine-point Likert-scale (1 = not relevant at all, 9 = highly relevant). We gave an example of a possible indicator for each topic for comprehensibility. In addition, experts were invited to comment on the topics or to suggest additional topics for each of the three phases in the open spaces of the questionnaire.

All ratings were analyzed by calculating the median score and level of agreement between experts, following the RAND/UCLA Appropriateness Method user’s manual [27]. Based on the median scores, topics were classified as either inappropriate (median range 1 – 3), uncertain (median range 4 – 6) or appropriate (median range 7 – 9) (Appendix B). Level of agreement was assessed by the IPR-score (interpercentile range, difference between 30th and 70th percentile) and the IPRAS-score (interpercentile range adjusted for symmetry). If the IPRAS is larger than the IPR, there is agreement among experts and if the IPR is larger than the IPRAS, there is disagreement.

We planned to 1) accept topics with median score ≥ 7 with agreement, 2) reject topics with median score ≤ 3 with agreement, and 3) discuss all other topics (median score 4 – 6 or without agreement) in Delphi round 2. However, round 1 resulted in a large majority ‘accepted’ topics and well exceeded the number of intended indicators. We therefore decided to prioritize




Figure 1 Schematic representation of the development and establishment of the indicator set.

these ‘accepted’ topics in the second Delphi round and rejected all other topics.

The experts’ suggestions for new topics were read and discussed by the researchers (JM, IB, JS) until consensus was reached on additional topics. New topics were combined or reformulated if necessary and added to Delphi round 2.

DELPHI ROUND 2: EXPERT MEETING

The second Delphi round consisted of expert meetings to prioritize the topics using the cumulative voting method. Meetings were held online due to Dutch COVID policy restrictions and we organized three separate smaller meetings to encourage active participation during the online meetings. The meetings of +- 120 minutes were recorded. Experts were first informed about the results of Delphi round 1. Next, they were encouraged to prioritize topics by dividing 100 points at their own discretion. After the individual prioritization, experts entered their scores into an interactive program to aggregate scores of all participants in the meeting. We encouraged experts to reflect on these aggregated scores. After the discussion, experts were invited to reconsider their earlier individual scores again. This sequence was repeated for the three phases (preconception, pregnancy and after birth).

Subsequently, we aggregated all final scores and classified the topics from high to low sum scores. Within every phase (preconception, pregnancy and after birth) we searched for a sudden decline in sum scores as a natural cut-off point for prioritized topics. This led to a draft list of prioritized topics.

In addition, we transcribed the expert meetings verbatim and analyzed the data using MaxQDA. One of the researchers (JM) coded the data for considerations in the prioritization and requirements for the indicator set. Coding was checked by a second researcher (IB).

The researchers (JM, IB, JS) consequently checked the draft list of prioritized topics against the experts’ requirements for the indicator set. We checked whether the requirements were fulfilled or whether we should add lower prioritized topics to fulfill the requirements. At the end of the second Delphi round, we had a final list of prioritized topics.

INTERMEDIATE ROUND

Based on the final list of prioritized topics, we made a list of possible indicators for each topic. Indicators were derived from our previous list of possible indicators (prior to Delphi study) as well as suggestions made by experts during Delphi round 1 and 2. Indicators were reformulated or merged in case they were not clearly defined or overlapped, based on consensus between two researchers and in line with the other indicators (JM, IB). In the rare case that there was no indicator available in the mentioned sources for one of the topics, the researchers (JM, IB) formulated potential indicators

based on comparable indicators (e.g. indicators for the same topic in other phases). For each indicator, we described its numerator, denominator, data source, and data availability.

DELPHI ROUND 3: ONLINE QUESTIONNAIRE

The third Delphi round consisted of an online questionnaire to select and prioritize indicators. The experts received a list of possible indicators (including numerator and denominator) for each topic and were encouraged to 1) select a maximum of three indicators they considered suitable to monitor Solid Start on a local level, and 2) indicate their number one preference. In case only one possible indicator was presented, experts were asked whether or not they considered that indicator suitable. The experts were also invited to add comments.

For each indicator, we calculated the percentage of experts that selected the indicator within their top three or as their preference. The scores and comments were discussed by the researchers (JM, IB, JS) in order to select at least one indicator per topic. In this process, the following conditions were considered: 1) Is there a clear preference towards one indicator? 2) Is data available for this indicator in nationwide data sources for every municipality? 3) Is the indicator sufficiently operationalized? If all conditions were met, the preferred indicator was added to the draft indicator set. We additionally prepared a ‘development agenda’ for topics and indicators that were clearly preferred, but lacked data in nationwide data sources or a clear operationalization. In this case, a lower ranked indicator for this topic with data-availability and sufficient operationalization was added to the draft indicator set.

DISCUSSION – EXPERT MEETING

In a final two-hour online expert meeting we presented the draft indicator set (including the ‘development agenda’) and asked experts for feedback. Specifically, we checked whether the set covers the various elements to appropriately monitor Solid Start on a local level. Experts were encouraged to share their thoughts in the meetings’ chatbox or by e-mail afterwards. Pressing issues were discussed directly. Based on the meeting minutes and written feedback, we finalized the indicator set.

ETHICAL CONSIDERATIONS

Following the Dutch Medical Research Involving Human Subjects Act (WMO), ethical approval was not necessary for this study (<http://www.ccmo.nl>), as we did not conduct medical-scientific research and participants were not exposed to treatment or required to follow a certain behavioral strategy. All participants gave written informed consent. In an information letter and at the start of each round or meeting, we stressed that participation was voluntary and confidential, and that data were processed anonymously.

RESULTS

PARTICIPANTS

The expert panel consisted of 39 experts (Table 1). The full questionnaire to select topics (round 1) was completed by 39 experts and 28 experts joined the online expert

meeting to prioritize topics (round 2). A total of 28 experts participated in the questionnaire to select indicators (round 3) and 21 experts were present during the final expert meeting. 18 experts joined during the full study.

Figure 2 shows a flowchart of the selection of topics and indicators during the study.

	TOTAL **	ROUND 1 – QUESTIONNAIRE	ROUND 2 – EXPERT MEETING	ROUND 3 – QUESTIONNAIRE	DISCUSSION – EXPERT MEETING
Total number of participants	39	39	28	28	21
Field of expertise					
Policy*	22	22	16	16	14
Practice*	12	12	7	9	7
- Social sector	4	4	2	3	1
- Medical sector	3	3	1	1	1
- Both	5	5	4	5	5
Research*	9	9	7	6	4
Other* (e.g. providing support for collaboration and the formation of Solid Start coalitions in general)	3	3	3	3	2

Table 1 Characteristics of participants.

* More than one field of expertise is possible.

** The same pool of 39 experts was approached in each round (e.g. the discussion was attended by 21 of these 39 experts).




Figure 2 Flowchart of the selection of topics and indicators to monitor Solid Start on a local level.

ROUND 1 – QUESTIONNAIRE

The experts received 121 possible topics to rate. Out of these, 105 topics were selected (median score ≥ 7) and 16 topics were excluded (median score < 7) (Appendix B). These excluded topics mainly concerned complications or medical risks during pregnancy or after birth (e.g. gestational diabetes and caesarian-section). Based on the experts' suggestions, 34 topics were added. Some topics were completely new, but most were already mentioned in another one of the three phases (preconception, pregnancy, after birth). In total, 139 topics were selected for round 2.

ROUND 2 – EXPERT MEETING

Experts prioritized topics within each of the three phases (Appendix B). For the preconception phase, the topic 'poverty' received the highest sum score. The topic 'cumulation of risk factors' received the highest sum scores for the phases of pregnancy and after birth. A decline in sum scores was clear in the pregnancy-phase after 10 topics (from 112 points to 96 points), but less clear for the other phases. We selected the prioritized 10 topics within each phase (a total of 30 topics, Table 2). Most topics belonged to two or three phases.

Experts mentioned multiple requirements for the final indicator set (see Appendix C for a description of all requirements and corresponding quotes). The indicator set should include indicators regarding both processes and outcomes, and both parents and children. Experts moreover wanted to include indicators that have the potential to be influenced (to identify early effects of policy) as well as indicators that show prevalence rates (to be used in making policy). The total indicator

set should be balanced in terms of risk- and protective factors and in general it should provide a full picture of all relevant aspects. The indicator set should provide a starting point of the conversation within a cross-sector collaboration. Lastly, it was considered important that data are available for the indicators. No additional topics were added to the final indicator set based on these requirements, since the prioritized topics largely seemed to match these requirements.

INTERMEDIATE ROUND

For the 30 prioritized topics, 107 unique indicators were found by the research team in the different sources. The number of potential indicators per topic varied from 1 to 7.

ROUND 3 – QUESTIONNAIRE

Based on the experts' selection and prioritization, the preferred indicator was clear for 20 topics (Appendix B). 11 of these indicators lacked data and were added to the development agenda. As the 'second best' option, 5 lower prioritized indicators for the corresponding topics were added to the draft indicator set. The draft indicator set consisted of 16 indicators, the draft development agenda of 23 indicators.

DISCUSSION – EXPERT MEETING

In general, experts appreciated the draft indicator set. They mentioned a number of extra non-prioritized indicators, which were added to an additional 'choice set' in case data was available (Appendix B). This set complements the basic indicator set and allows local coalitions to use additional indicators (e.g. regarding educational level, single parent family, long-term low income) if they want to.

	PRECONCEPTION	PREGNANCY	AFTER BIRTH (UP TO TWO YEARS)
Topics in all three phases*	Poverty	Poverty	Poverty
	Early detection by healthcare provider	Early detection by healthcare provider	Early detection by healthcare provider
	Health: psychological/psychiatric problems	Health: psychological/psychiatric problems	Health: psychological/psychiatric problems parents
	Health: stress	Health: stress	Health: stress
Topics in two phases*	Domestic violence (including screening)	Domestic violence (including screening)	
	Substance use: smoking	Substance use: smoking	
		Social network	Social network
		Cumulation of risk factors	Cumulation of risk factors
Topics in one phase	Preconception care	Care: multidisciplinary collaboration	Health outcomes child: premature birth
	Interventions (process indicators)	Unintended and/or unwanted pregnancy	Relation parent – child
	Low literacy		Health: intellectual disability parent
	Client characteristics: socioeconomic status		Child abuse and neglect

Table 2 Overview of the prioritized topics (n = 30)*.

* The topics that occur in multiple phases are presented on the same row.

In reflecting on the indicator set, experts mentioned some conceptual considerations (e.g. indicators are often formulated as risks, while the reverse can be a protective factor). They also mentioned methodological considerations (e.g. indicators regarding children's health at age two are currently missing and should be added when more youth healthcare data is available). Experts gave their consent to the indicator set provided that the set will be piloted in practice. Based on the experts' feedback, the indicator set and development agenda were finalized.

FINAL INDICATOR SET

Finally, 19 indicators could be selected to monitor Solid Start on a local level (Table 3): 7 in the preconception phase, 5 during pregnancy and 7 after birth (up to two years). Some examples are debts, psychological or psychiatric problems, late antenatal care, smoking during pregnancy, vulnerability during pregnancy and after birth, not receiving postpartum care, and preterm birth and/or low birth weight for gestational age (SGA). Appendix B describes the selected indicators in more detail. Data is available in nationwide data sources for all these operationalized indicators and can be presented at local (municipality) level.

The development agenda consists of 21 indicators (Appendix B). These (preferred) indicators lacked data or a clear operationalization. Some examples are smoking before pregnancy, stress due to finances, unwanted or unplanned pregnancy, stress during pregnancy, loneliness among parents, secure bonding, abuse or neglect of children and stress with parenting.

DISCUSSION

In this paper, we present an indicator set to monitor the Solid Start program in Dutch local coalitions, and we describe how we used a modified Delphi technique to reach consensus. The final indicator set consists of 19 indicators, covering the three phases of the Solid Start program: preconception (n = 7), pregnancy (n = 5) and after birth (up to two years)(n = 7). These indicators are available in nationwide data sources and can be presented on local (municipality) level. The indicator set meets the requirements as mentioned by the experts; it contains indicators that cover both processes and outcomes, both parents and children, and both risk- and protective factors. Additionally, the indicator set reflects

PRECONCEPTION
Percentage of women and men in the reproductive age with debts
Percentage of women and men in the reproductive age with psychological or psychiatric problems
Percentage of women and men in the reproductive age with stress
Percentage of women in the reproductive age who smoke
Percentage of families reached with a preconception consultation (preconception care)
Percentage of low literacy among young people (<30 years) without partner and children
Percentage of women and men in the reproductive age living in a neighbourhood with a low liveability score
PREGNANCY
Percentage of pregnant women with debts *
Percentage of pregnant women who have their first antenatal care visit after the 10th week of pregnancy *
Percentage of pregnant women with psychological or psychiatric problems
Percentage of women who smoke at some point during pregnancy
Percentage of pregnant women in a potentially vulnerable situation (3 or more risk factors to vulnerability)
AFTER BIRTH (UP TO TWO YEARS)
Percentage of children born in a family with debts
Percentage of families not receiving postpartum care (at home) after birth *
Percentage of children aged 0 to 2 years of whom one or both parents have psychological or psychiatric problems
Percentage of children born in a family in a potentially vulnerable situation (3 or more risk factors to vulnerability) *
Percentage of children with a preterm birth or with a low birth weight for gestational age (SGA) *
Percentage of children born in a family of which one or both parents have a mild intellectual disability
Number of out-of-home placements for children before the age of 2 (per 1,000) *

Table 3 Selected indicators to monitor Solid Start on a local level (n = 19).

* These indicators are also included in the indicator set to monitor the national Solid Start program.

both medical and social factors. A development agenda was established with topics and indicators that were prioritized, but lacked data in nationwide data sources or a clear operationalization.

The indicator set covers the following topics: poverty, psychological/psychiatric problems, stress, smoking, cumulation of risk factors, preconception care, low literacy, socioeconomic status, premature birth, intellectual disability, and child abuse and neglect. The first four topics are presented in the indicator set for all three phases (preconception, pregnancy and after birth). In general, the social determinants of health [7, 28] are represented in the indicator set (e.g. debts, low literacy and living in a neighborhood with a low livability score). Specific clinical aspects that belong to one group of care providers (e.g. caesarean section, a child's hearing) are less present. Nonetheless, the indicator set reflects both medical and social care, which aligns with the aims of the Solid Start program. In comparison to the indicators used in the current national Solid Start monitor (Appendix D), there is some overlap (e.g. debts during pregnancy, preterm birth and low birth weight for gestational age) but also differences. For instance, the national monitor also includes indicators such as the percentage of municipalities that implemented the program 'Not Pregnant Now'. These differences are arguably caused by the different purposes of both indicator sets. The indicators in the national monitor can be used to monitor and evaluate the nationwide implementation of the program, and to monitor health outcomes of parents and children on a national level. As the implementation and health outcomes vary between municipalities, the indicator set of the local monitor aims to enhance the conversation between policymakers, managers, professionals and other stakeholders about the local situation and developments in order to prioritize interventions and policies at a local level.

A development agenda was made with indicators and topics that lacked data in nationwide data sources or a clear operationalization. Among others, the topics and indicators on the development agenda were related to stress, unwanted or unintended pregnancy, (quitting) smoking before pregnancy, loneliness, early detection, secure bonding, and child abuse or neglect. Multiple indicators related to stress were prioritized: stress due to finances, stress during pregnancy and stress with parenting. There is growing scientific evidence that stress during pregnancy or parenting has long- and short-term consequences for children's health and development [14, 29, 30]. The multidimensional concept of stress [31] may require different indicators. It seems, therefore, valuable to explore which topics of the development agenda should be prioritized to be incorporated in routine registries for the purpose of local monitoring.

There are, to the best of our knowledge, no other studies that used a Delphi technique to identify indicators for local monitoring of the full first thousand days (approach). There are, however, several previous studies that sought to describe indicators for aspects of the first thousand days, including antenatal care [32], obstetrical care [33], children's health [34], birth center care [35], and maternal and newborn health [36] or care [25] during pregnancy, childbirth and the postpartum period. Next to that, we found several programs in other countries that were focused to the first thousand days, but the aims, scope and key-design elements of the programs and their evaluation differ [37-41]. These programs were often not directly comparable to the Dutch Solid Start program and not (yet) focused on supporting monitoring on a local level. Consequently, a comparison between our indicator set and indicators in the aforementioned studies is hampered, with the exception of a study from Sweden [26]. In this Swedish study, the researchers developed indicators, sub-indices and a summary index in order to support municipalities with monitoring children's health. In comparison to our study, they also mentioned both risk- and protective factors and also selected indicators related to poverty, smoking and low birth weight.

STRENGTHS AND LIMITATIONS

A strength of this study is that the indicator set is developed based on the expertise of a heterogenic and balanced group of experts in policy, practice and research related to the first thousand days, who have an interest in using the set in daily practice [20]. The focus of the indicator set to the first thousand days, involving both the social and medical sector, is necessary for programs aimed at reducing health inequities as health outcomes are directly and indirectly influenced by both social and medical factors [6, 9, 42]. The experts exchanged information and expressed their views during two expert meetings, as done in previous Delphi studies [20]. We organized a meeting to discuss and prioritize topics (Delphi round 2) and a final expert meeting. We considered this final moment of reflection on the (draft) indicator set very important to increase the support and future uptake of the indicator set in practice.

However, this study also has several limitations. First, we selected indicators based on consensus without considering the scientific evidence for these indicators. This does not necessarily mean that indicators that were not prioritized are not valid and vice versa. For most indicators to monitor maternal and neonatal health, their level of evidence is not well described [25]. In general, the rare availability of evidence is one of the reasons to (partly) select indicators based on experts' opinions in a Delphi study [20]. Another limitation was that not all indicators in the final set were the preferred option by experts as

a consequence of limitations in data availability. Hence we included some ‘second best’ indicators and added the preferred indicators to the development agenda. Other limitations relate to the inclusion of experts. This depended on the availability and willingness of experts to participate within the study’s time period, and on the decisions of the researchers in how and who to invite. Moreover, we invited experts from practice, policy and research in both the social and medical sector. Making a clear distinction between and within those categories is not always possible, as multiple experts work at the intersection of the various fields of expertise (practice, policy and research) or in multiple sectors (medical and social). For example, managers of local coalitions can be categorized as working in both practice and policy, as well as within the medical and social sector. The inability to distinguish between the field of expertise and sector is however in line with the aims of the program (i.e. integrating service delivery across the medical and social sector). Therefore, we do not expect that this may have influenced the results. This is also reflected in our results, as the experts from different fields of expertise and sectors did not prioritize different topics and indicators. Additionally, some experts dropped out during the study period, but the three groups of experts from practice, policy and research were all well represented during the various rounds. In addition, we missed the perspective of parents themselves. Finally, due to the COVID-19 pandemic, we were unable to organize physical meetings. Our decision to organize three smaller online meetings hindered the exchange of information and considerations between all experts. However, since the results of each of the meetings were highly comparable, we expect little influence on the results.

FUTURE RESEARCH AND PRACTICE

Recently, the first indicators were quantified and presented to all municipalities in the Netherlands at www.regiobeeld.nl/kansrijke-start. In the future, we will further refine the website with additional indicators and new functionalities (among which maps with geospatial variation). In quantifying the indicators, we use nationwide observational data sources with routinely registered data, which are linked on individual level. In the last decade, the opportunities of linking observational data sources has increased at an enormous pace, which enhances the usefulness and applicability of the developed indicator set [43].

The indicator set has yet to be used and evaluated in practice, as we can only determine the feasibility through empirical testing. A previous systematic review concluded that not many published indicators for maternal and neonatal health are empirically tested for validity and feasibility [25]. Starting in 2022, we will evaluate and refine the indicator set in close

collaboration with the participants of the monitoring support program (Appendix A) in order to stimulate the uptake and adoption in daily practice. During this process, we expect to also discover which indicators are most often used and how, also for indicators that are similar across two or three phases (e.g. debts before pregnancy, during pregnancy and after birth). Using the indicator set should not be a one-time action, because the strength of using indicators for monitoring in municipalities is the comparison with previous comparable figures [26]. In the future, the indicator set will be refined because of new developments, changing demographics, new evidence and increased data-availability. In reflecting on the use of the indicator set, it is also important that we pay attention to questions about obtaining and presenting the data.

In the coming years, the topics on the development agenda will be prioritized and addressed in collaboration with national parties and local professionals. Central in this process is the formulation and operationalization of indicators and the expected increase of data-availability. Next to the indicator set and development agenda, the choice set with extra, non-prioritized indicators is also publicly shared (including where to find the data) for local coalitions to use.

RELEVANCY

We consider our study scientifically relevant as it increases our understanding of relevant indicators for Solid Start and of using a systematic approach in developing indicators for monitoring a cross-sectoral program. In addition, it is relevant for society, as we can directly benefit from the study results by using the indicator set in practice. In the Netherlands, the indicator set can be used by local coalitions in collaboration with local stakeholders to describe their population, to identify gaps in current processes, to make or adapt policies, to prioritize interventions, to monitor developments and to stress the importance of investing in the first thousand days. In this monitoring process, combining quantitative data with qualitative data about experiences, facilitators and barriers (in a mixed-methods approach) can help to interpret the quantitative data, gain more insight into processes and explore opportunities for improvement [44]. Using the indicator set in combination with qualitative data in a continuous learning cycle with local stakeholders can support an integrated approach that is adapted to the local context in Dutch municipalities. On an international level, the topics and indicators can potentially be a starting point for monitoring similar cross-sectoral programs into the first thousand days in other western countries [37–41]. Additionally, countries that aim to develop a supported and comprehensive indicator set to monitor a cross-sectoral program can learn from our systematic methodology of collaborating

with experts with varying backgrounds. Using a co-creative process can increase the support, relevancy and therewith impact of the research project [45, 46].

CONCLUSION

In this study we present an indicator set for monitoring the Dutch Solid Start program on a local level, which will be used and evaluated from 2022 onwards. The indicator set consists of 19 indicators that reflect both social and medical factors. The indicator set can be used by local coalitions to enhance the conversation between stakeholders about the local situation and developments in order to prioritize interventions and policies. Using the indicator set for monitoring is a continuous process that supports the optimisation and promotion of integrated service delivery across the medical and social sector at a local level. Ultimately, the indicator set contributes to the reduction of health inequities within the preconception period, during pregnancy and after birth in order to give each child a solid start.

ADDITIONAL FILES

The additional files for this article can be found as follows:

- **Appendix A.** RIVM monitoring support program – ‘Learning Local Monitor Solid Start’. DOI: <https://doi.org/10.5334/ijic.6508.s1>
- **Appendix B.** Results of Delphi round 1, 2, and 3, final local indicator set, choice set and development agenda. DOI: <https://doi.org/10.5334/ijic.6508.s2>
- **Appendix C.** Considerations in the prioritization and requirements for the final indicator set. DOI: <https://doi.org/10.5334/ijic.6508.s3>
- **Appendix D.** National indicator set. DOI: <https://doi.org/10.5334/ijic.6508.s4>

ACKNOWLEDGEMENTS

We are very grateful for the participation of all experts in the experts panel.

REVIEWERS

Prof. Dr. A.T.G. (Aggie) Paulus, Department of Health Services Research, Faculty of Health Medicine and Life Sciences, Care and Public Health Research Institute (Caphri), School of Health Professions Education (SHE), Maastricht University, Netherlands.

W. Douglas Tynan PhD, Department of Pediatrics, Sidney Kimmel Medical College, Thomas Jefferson University – Philadelphia, USA.

One anonymous reviewer.

FUNDING INFORMATION

The Dutch Ministry of Health, Welfare and Sport funded the monitor of the Dutch Solid Start program that was conducted by the National Institute for Public Health and Environment.

COMPETING INTERESTS

The authors have no competing interests to declare.

AUTHOR AFFILIATIONS

Joyce M. Molenaar  orcid.org/0000-0002-8025-040X

Centre for Nutrition, Prevention and Health Services, Department of Quality of Care and Health Economics, National Institute for Public Health and the Environment, Bilthoven, The Netherlands; Health Campus The Hague/Department of Public Health and Primary Care, Leiden University Medical Center, The Hague, The Netherlands

Inge C. Boesveld  orcid.org/0000-0002-1250-325X

Centre for Nutrition, Prevention and Health Services, Department of Quality of Care and Health Economics, National Institute for Public Health and the Environment, Bilthoven, The Netherlands

Jessica C. Kieft-de Jong  orcid.org/0000-0002-8136-0918

Health Campus The Hague/Department of Public Health and Primary Care, Leiden University Medical Center, The Hague, The Netherlands

Jeroen N. Struijs  orcid.org/0000-0002-7493-6720

Centre for Nutrition, Prevention and Health Services, Department of Quality of Care and Health Economics, National Institute for Public Health and the Environment, Bilthoven, The Netherlands; Health Campus The Hague/Department of Public Health and Primary Care, Leiden University Medical Center, The Hague, The Netherlands

REFERENCES

1. **Mohangoo AD, Buitendijk SE, Hukkelhouen C, Ravelli AC, Rijninks-van Driel GC, Tamminga P**, et al. Hoge perinatale sterfte in Nederland vergeleken met andere Europese landen: de Peristat-II-studie [Higher perinatal mortality in The Netherlands than in other European countries: the Peristat-II study]. *Ned Tijdschr Geneeskd.* 2008; 152(50): 2718.
2. **Euro-Peristat.** *European perinatal health report.* 2008.
3. **Stuurgroep Zwangerschap en Geboorte.** Een goed begin. Veilige zorg rond zwangerschap en geboorte. [A good start. Safe care for pregnancy and childbirth]. Utrecht, the Netherlands; 2009.
4. **Struijs JN, Hargreaves DS.** Turning a crisis into a policy opportunity: lessons learned so far and next steps in the Dutch early years strategy. *The Lancet Child & Adolescent Health.* 2019; 3(2): 66–8. DOI: [https://doi.org/10.1016/S2352-4642\(18\)30384-5](https://doi.org/10.1016/S2352-4642(18)30384-5)

5. **Zorginstituut Nederland.** Zorgstandaard Integrale Geboortezorg [Standard for Integrated Birth Care]; 2016.
6. **National Academies of Sciences E, Medicine.** Vibrant and healthy kids: Aligning science, practice, and policy to advance health equity; 2019.
7. **World Health Organization.** A conceptual framework for action on the social determinants of health; 2010.
8. **Denktas S, Bonsel GJ, Van der Weg EJ, Voorham AJ, Torij HW, De Graaf JP,** et al. An urban perinatal health programme of strategies to improve perinatal health. *Matern Child Health J.* 2012; 16(8): 1553–8 DOI: <https://doi.org/10.1007/s10995-011-0873-y>
9. **Waelput AJM, Sijkens MK, Lagendijk J, van Minde MRC, Raat H, Ernst-Smelt HE,** et al. Geographical differences in perinatal health and child welfare in the Netherlands: rationale for the healthy pregnancy 4 all-2 program. *BMC Pregnancy Childbirth.* 2017; 17(1): 254. DOI: <https://doi.org/10.1186/s12884-017-1425-2>
10. **Barsties LS, Daalderop LA, Lagendijk J, van Steenbergen F, Been JV, Bertens LCM,** et al. Addressing perinatal health inequities in Dutch municipalities: Protocol for the Healthy Pregnancy 4 All-3 programme. *Health Policy.* 2021; 125(3): 385–92 DOI: <https://doi.org/10.1016/j.healthpol.2020.12.013>
11. **Denktas S, Poeran J, van Voorst SF, Vos AA, de Jong-Potjer LC, Waelput AJ,** et al. Design and outline of the healthy pregnancy 4 all study. *BMC Pregnancy Childbirth.* 2014; 14(1): 1–11. DOI: <https://doi.org/10.1186/1471-2393-14-253>
12. **Ministry of Health, Welfare and Sport.** *Solid Start – the action programme.* The Hague; 2020. [<https://www.government.nl/documents/publications/2020/08/24/solid-start-the-action-programme>].
13. **Barker DJ.** The origins of the developmental origins theory. *J Intern Med.* 2007; 261(5): 412–7. DOI: <https://doi.org/10.1111/j.1365-2796.2007.01809.x>
14. **Wadhwa PD, Buss C, Entringer S, Swanson JM,** editors. Developmental origins of health and disease: brief history of the approach and current focus on epigenetic mechanisms. *Semin Reprod Med;* 2009: NIH Public Access. DOI: <https://doi.org/10.1055/s-0029-1237424>
15. **Vermeulen W.** Decentralisation of social policy in the Netherlands. *Decentralisation of education, health and social protection: issues and challenges.* 127.
16. **RIVM.** Indicatoren Kansrijke Start: Een Delphi-studie [Indicators Solid Start: a Delphi study]. Bilthoven, the Netherlands; 2019. [<https://www.rivm.nl/documenten/indicatoren-kansrijke-start-delphi-studie>].
17. **RIVM.** Monitor Kansrijke Start 2019. [Monitor Solid Start 2019]. Bilthoven, the Netherlands; 2019. [<https://www.rivm.nl/documenten/factsheet-kansrijke-start>].
18. **RIVM.** Monitor Kansrijke Start 2020 [Monitor Solid Start 2020]. Bilthoven, the Netherlands; 2020. [<https://www.rivm.nl/documenten/monitor-kansrijke-start-2020>].
19. **RIVM.** Monitor Kansrijke Start 2021 [Monitor Solid Start 2021]. Bilthoven, the Netherlands; 2021. [<https://www.rivm.nl/documenten/monitor-kansrijke-start-2021>].
20. **Boulkedid R, Abdoul H, Loustau M, Sibony O, Alberti C.** Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS One.* 2011; 6(6): e20476. DOI: <https://doi.org/10.1371/journal.pone.0020476>
21. **Keeney S, McKenna H, Hasson F.** The Delphi technique in nursing and health research. John Wiley & Sons; 2011. DOI: <https://doi.org/10.1002/9781444392029>
22. **Zorginstituut Nederland.** Indicatorenset Integrale Geboortezorg [Indicator set Integrated Birth Care]; 2020.
23. **Statistics Netherlands (CBS).** Kansrijk van Start in Heerlen, 2016 [Solid Start in Heerlen, 2016]; 2019.
24. **de Wolff M, Detmar S, Verkerk P, Reijneveld S, Geerdes-Maas T.** Verkenning JGZ Indicatoren; 2021.
25. **Saturno-Hernandez PJ, Martinez-Nicolas I, Moreno-Zegbe E, Fernandez-Elorriaga M, Poblano-Verastegui O.** Indicators for monitoring maternal and neonatal quality care: a systematic review. *BMC Pregnancy Childbirth.* 2019; 19(1): 25. DOI: <https://doi.org/10.1186/s12884-019-2173-2>
26. **Kohler L, Eriksson B.** A Child Health Index for Sweden's 290 Municipalities: A System of Indicators and Indices for Monitoring Children's Health on the Local Level. *Child Indic Res.* 2018; 11(6): 1889–906. DOI: <https://doi.org/10.1007/s12187-017-9515-2>
27. **Fitch K, Bernstein SJ, Aguilar MD, Burnand B, LaCalle JR.** The RAND/UCLA appropriateness method user's manual. Rand Corp Santa Monica, CA; 2001.
28. **Gómez CA, Kleinman DV, Pronk N, Gordon GLW, Ochiai E, Blakey C,** et al. Practice Full Report: Addressing Health Equity and Social Determinants of Health Through Healthy People 2030. *J Public Health Manag Pract.* 2021; 27(6): S249. DOI: <https://doi.org/10.1097/PHH.0000000000001297>
29. **Hattangadi N, Cost KT, Birken CS, Borkhoff CM, Maguire JL, Szatmari P,** et al. Parenting stress during infancy is a risk factor for mental health problems in 3-year-old children. *BMC Public Health.* 2020; 20(1): 1–7. DOI: <https://doi.org/10.1186/s12889-020-09861-5>
30. **Staneva A, Bogossian F, Pritchard M, Wittkowski A.** The effects of maternal depression, anxiety, and perceived stress during pregnancy on preterm birth: A systematic review. *Women Birth.* 2015; 28(3): 179–93. DOI: <https://doi.org/10.1016/j.wombi.2015.02.003>
31. **Alves AC, Cecatti JG, Souza RT.** Resilience and Stress during Pregnancy: A Comprehensive Multidimensional Approach in Maternal and Perinatal Health. *Scientific World Journal.* 2021; 2021: 9512854 DOI: <https://doi.org/10.1155/2021/9512854>
32. **Bollini P, Quack-Lotscher K.** Guidelines-based indicators to measure quality of antenatal care. *J Eval Clin Pract.* 2013; 19(6): 1060–6 DOI: <https://doi.org/10.1111/jep.12027>
33. **Boulkedid R, Sibony O, Goffinet F, Fauconnier A, Branger B, Alberti C.** Quality indicators for continuous monitoring to improve maternal and infant health in maternity departments: a modified Delphi survey of an international multidisciplinary panel. *PLoS One.* 2013; 8(4): e60663. DOI: <https://doi.org/10.1371/journal.pone.0060663>

34. **Selmani A, Coenen M, Voss S, Jung-Sievers C.** Health indices for the evaluation and monitoring of health in children and adolescents in prevention and health promotion: a scoping review. *BMC Public Health*. 2021; 21(1): 2309. DOI: <https://doi.org/10.1186/s12889-021-12335-x>
35. **Boesveld IC, Hermus MAA, de Graaf HJ, Hitzert M, van der Pal-de Bruin KM, de Vries RG,** et al. Developing quality indicators for assessing quality of birth centre care: a mixed- methods study. *BMC Pregnancy Childbirth*. 2017; 17(1): 259. DOI: <https://doi.org/10.1186/s12884-017-1439-9>
36. **Moller AB, Newby H, Hanson C, Morgan A, El Arifeen S, Chou D,** et al. Measures matter: A scoping review of maternal and newborn indicators. *PLoS One*. 2018; 13(10): e0204763. DOI: <https://doi.org/10.1371/journal.pone.0204763>
37. **Cattan S, Conti G, Farquharson C, Ginja R, Pecher M.** The health impacts of universal early childhood interventions: evidence from Sure Start. Institute for Fiscal Studies; 2021. DOI: <https://doi.org/10.1920/BN.IFS.2021.BN0332>
38. **Dubay L, Hill I, Garrett B, Blavin F, Johnston E, Howell E,** et al. Improving Birth Outcomes And Lowering Costs For Women On Medicaid: Impacts Of ‘Strong Start For Mothers And Newborns’ An evaluation of the federal Strong Start for Mothers and Newborns program’s impact on birth outcomes and costs for Medicaid-covered women. *Health Aff (Millwood)*. 2020; 39(6): 1042–50. DOI: <https://doi.org/10.1377/hlthaff.2019.01042>
39. **Harman-Smith Y, Wilson A, Brinkman S.** Strong Start – Northern Pilot: Evaluation report. Adelaide, Australia: Fraser Mustard Centre, Telethon Kids Institute and the South Australian Department for Education and Child Development; 2015.
40. **Blake-Lamb T, Boudreau AA, Matathia S, Tiburcio E, Perkins ME, Roche B,** et al. Strengthening integration of clinical and public health systems to prevent maternal-child obesity in the first 1,000 Days: a collective impact approach. *Contemp Clin Trials*. 2018; 65: 46–52. DOI: <https://doi.org/10.1016/j.cct.2017.12.001>
41. **U.S. Department of Health & Human Services.** Early Head Start Programs 2022 [<https://eclkc.ohs.acf.hhs.gov/programs/article/early-head-start-programs>].
42. **Timmermans S, Bonsel GJ, Steegers-Theunissen RP, Mackenbach JP, Steyerberg EW, Raat H,** et al. Individual accumulation of heterogeneous risks explains perinatal inequalities within deprived neighbourhoods. *Eur J Epidemiol*. 2011; 26(2): 165–80. DOI: <https://doi.org/10.1007/s10654-010-9542-5>
43. **Scheefhals TMZ, de Vries EF, Molenaar JM, Numans ME, Struijs JN.** Observational data for understanding maternity care and early life: experiences with DIAPER (Data-InfraStructure for ParEnts and childRen) [manuscript submitted for publication].
44. **Adatho M.** *Combining quantitative and qualitative methods for program monitoring and evaluation: Why are mixed-method designs best?*. World Bank; 2011.
45. **van Dijk-de Vries A, Stevens A, van der Weijden T, Beurskens AJ.** How to support a co-creative research approach in order to foster impact. The development of a Co-creation Impact Compass for healthcare researchers. *PLoS One*. 2020; 15(10): e0240543. DOI: <https://doi.org/10.1371/journal.pone.0240543>
46. **Greenhalgh T, Jackson C, Shaw S, Janamian T.** Achieving research impact through cocreation in communitybased health services: literature review and case study. *The Milbank Quarterly*. 2016; 94(2): 392–429. DOI: <https://doi.org/10.1111/1468-0009.12197>

TO CITE THIS ARTICLE:

Molenaar JM, Boesveld IC, Kieft-de Jong JC, Struijs JN. Monitoring the Dutch Solid Start Program: Developing an Indicator Set for Municipalities to Monitor their First Thousand Days-Approach. *International Journal of Integrated Care*, 2022; 22(4): 8, 1–12. DOI: <https://doi.org/10.5334/ijic.6508>

Submitted: 18 January 2022 **Accepted:** 06 October 2022 **Published:** 25 October 2022

COPYRIGHT:

© 2022 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

International Journal of Integrated Care is a peer-reviewed open access journal published by Ubiquity Press.