



Universiteit  
Leiden  
The Netherlands

## Temporal variability in organic amendment impacts on hydro-physical properties of sandy agricultural soils

Kok, D.D.; Scherer, L.A.; Vries W. de; Bodegom, P.M. van

### Citation

Kok, D. D., Scherer, L. A., & Bodegom, P. M. van. (2023). Temporal variability in organic amendment impacts on hydro-physical properties of sandy agricultural soils. *Soil Science Society Of America Journal*, 87(4), 963-984. doi:10.1002/saj2.20547

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3677266>

**Note:** To cite this publication please use the final published version (if applicable).

## ORIGINAL ARTICLE

## Soil &amp; Water Management &amp; Conservation

# Temporal variability in organic amendment impacts on hydro-physical properties of sandy agricultural soils

Dirk-Jan Daniel Kok<sup>1</sup>  | Laura Scherer<sup>1</sup>  | Wim de Vries<sup>2</sup> | Peter Michiel van Bodegom<sup>1</sup>

<sup>1</sup>Institute of Environmental Sciences (CML), Leiden University, Leiden, The Netherlands

<sup>2</sup>Wageningen University and Research, Wageningen, The Netherlands

## Correspondence

Dirk-Jan Kok, Institute of Environmental Sciences (CML), Leiden University, Leiden, The Netherlands.

Email: [d.d.kok@cml.leidenuniv.nl](mailto:d.d.kok@cml.leidenuniv.nl)

Assigned to Associate Editor Rongzhong Ye

## Funding information

Nederlandse Organisatie voor Wetenschappelijk Onderzoek, Grant/Award Number: NWO-GK-06

## Abstract

Organic amendments (OAs) can improve the hydro-physical properties of a soil and thereby potentially enhance the resilience of agricultural systems to droughts and floods. An OA's contribution to this resilience, however, depends on the timeliness of its impacts, as soil improvements should be achieved when droughts are most frequent or flood risks are greatest. Yet little is known regarding the temporal variability of OA impacts or the influence of OA quantity and quality thereupon. In this research, therefore, we investigated at two agricultural sites the temporal variability of improvements in soil bulk density, aggregate stability, infiltration capacity and water retention after the application of compost, farmyard manure, bokashi, a selection of organic residues from landscape maintenance, and a combination of these residues with manure. Results showed that, depending on management practices and soil type, OAs decrease bulk density by up to 9.8%, increase infiltration capacity by up to 108.1%, aggregate stability by up to 60.0%, and water retention by up to 77.8% relative to unamended controls within 3 years of repeated application. However, the magnitude of these improvements varies up to 96% between seasons, depending on the soil property and OA treatment. On average, for all treatments, impacts relative to the control varied between different seasons by 5% for bulk density, 47.1% for infiltration capacity, 22.6% for aggregate stability, and 26.3% for water retention. When offsetting OA nutrient differences with mineral fertilizers, this variability showed a stronger correlation to differences in OA application quantity than quality (i.e., chemical composition). Results suggest that disregarding temporal variability in OA impacts can result in an inaccurate valuation of OAs as either effective or ineffective in improving soil resilience, given that impacts may, instead of their frequently presumed persistency, actually be highly transient or lagged. Our findings highlight the importance of considering the potential intra-annual variability of OA impacts on soil hydro-physical properties when designing OA application strategies to ameliorate the effects of specific seasonal climatic challenges.

**Abbreviations:** EMM, estimated marginal mean; LM, linear regression model; OA, organic amendment; WWR, waterway weeds and reeds.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Soil Science Society of America Journal* published by Wiley Periodicals LLC on behalf of Soil Science Society of America.

## 1 | INTRODUCTION

Future agricultural production will be challenged by the intensification of seasonal precipitation extremes due to onsetting climate change (IPCC et al., 2021). Due to this, much of the world will experience decreases in the number of summer precipitation events and strong increases in winter precipitation intensity (Donat et al., 2013; Giorgi et al., 2014; IPCC et al., 2021). These climate change effects can hamper agricultural production by exposing it to flooding and droughts. Agricultural soils might be particularly vulnerable, as they often suffer from reduced infiltration capacity and water retention due to their degradation under conventional cultivation practices (Bai et al., 2008; Karlen & Rice, 2015; Lal, 2012). Rehabilitating agricultural soils and improving their hydro-physical properties may thus be necessary to improve the resilience of our food production systems to a changing climate (Cornelis et al., 2019).

Organic amendments (OAs) can improve soil hydro-physical properties and thereby potentially enhance the climate resilience of our crop production systems (Diacono & Montemurro, 2011; Larney & Angers, 2012; Montgomery, 2007). Different OAs have demonstrably improved soil hydraulic and physical properties such as bulk density (Cercioglu, 2017; García-Orenes et al., 2005; Khaleel et al., 1981; Reynolds et al., 2009; Tejada & Gonzalez, 2008), infiltration capacity and hydraulic conductivity (Eusufzai & Fujii, 2012; Meena et al., 2020; Wanniarachchi et al., 2019), penetration resistance (Celik et al., 2010; Negiş et al., 2020), pore size distribution (Luna et al., 2018), aggregate stability (Abiven et al., 2009; Albiach et al., 2001; Annabi et al., 2011; Metzger et al., 1987), and water retention (Ankenbauer & Loheide, 2017; Eden et al., 2017; Meena et al., 2020; Rawls et al., 2003; Zhou et al., 2020). Such improvements may support crop productivity as well as buffer climate change impacts. For instance, the roots of crops grown on soils of lower bulk densities more easily penetrate the soil and benefit from improved oxygen diffusion into the rhizosphere (Valentine et al., 2012); soils in which water readily infiltrates are less susceptible to ponding and land degradation by runoff (Hueso-González et al., 2015); soils capable of retaining more water provide crops with greater resilience to dry spells and lower irrigation requirements (Pandey & Shukla, 2006; Terleev et al., 2017; Zhang et al., 2006); and soils with high aggregate stability have increased nutrient retention through the occlusion of organic matter (Lutzow et al., 2006; Six et al., 2004; Tisdall & Oades, 1982).

The efficacy of OAs in improving soil resilience, however, depends not only on the magnitude of improvements but also on their intra-annual variation and persistence of impacts. Ideally, OAs improve relevant soil properties prior to the occurrence of a seasonal weather extreme. Yet little is known regarding the seasonal variability of OA impacts

### Core Ideas

- Hydro-physical properties improved within 3 years of repeated organic amendment (OA) application.
- Improvements were more closely related to OA application rates than to OA composition.
- The impact of OAs on hydro-physical properties showed significant temporal variability.
- Temporal variability in OA impacts was influenced by soil type and land management practices.
- No single OA quantity or qualitative property could consistently explain the observed temporal variability.

as most studies evaluate changes in soil properties at fixed yearly intervals (Cercioglu, 2017; Ferreras et al., 2006; Tejada & Gonzalez, 2008) or once at the end of an experiment (Li et al., 2018; Singh Brar et al., 2015; Yüksel & Kavdır, 2020; Zhao et al., 2009). Studies that do explore short-term OA impacts observe strong seasonal fluctuations in the response of soil hydro-physical properties to applications of OAs such as compost (Castellini et al., 2022; Felton, 1995), sewerage sludge (Delibacak et al., 2009), and biochar (Ouyang et al., 2013). These seasonal variations can furthermore differ strongly per soil hydro-physical property (Zare et al., 2010). Yet, the short-term temporal dynamics of soil hydro-physical properties—let alone OA impacts thereupon—remain generally poorly investigated (Alletto et al., 2015; Geris et al., 2021; Girei et al., 2016; Jirků et al., 2013).

OA impacts are influenced by the quantity (i.e., application rates) and qualitative properties (i.e., chemical composition) of the OAs applied (Kögel-Knabner, 2017; Tejada & Gonzalez, 2008; Yu et al., 2020; Zamani et al., 2016). Yet, even for intensively studied changes in aggregate stability, remarkably few studies attempt to correlate OA impacts with the initial biochemical characteristics of the OAs (Abiven et al., 2009). A dedicated study to assess the short-term impacts of different OAs in a dynamic agricultural setting will contribute to deepening our understanding of the functional relationship between OAs and soil hydro-physical properties, and will be valuable for the identification of OAs that are most effective in improving the seasonal resilience of our food production systems.

In this research, therefore, we explored the intra-annual variability in the response of four prominent soil hydro-physical properties to four annual OA treatments in a 3-year experiment on two agricultural soils (an anthrosol and a podzol). We monitored changes in bulk density, infiltration capacity, aggregate stability, and water retention after application of compost, farmyard manure, a locally available organic

residue, and a combination of manure and the above residues. Soil hydro-physical properties were measured twice per year, in the second and third year of the experiment, and analyzed for correlations with OA application rate (quantity) and composition (quality). Bulk density was only measured in the third year of the experiment. We hypothesized that, for each site, (i) OA treatments result in significant improvements in soil hydro-physical properties, (ii) these improvements significantly vary between seasons for both soil types, and (iii) these improvements and their seasonal differences are related to differences in OA quantity or quality (e.g., C:N, total N, organic matter content, etc.). Based on the significance of these improvements and seasonal differences therein, we classify the impact of each OA for each soil property as either persistent, lagged, transient, or enhanced (further defined in the materials and methods section) in order to facilitate a comparison of their temporal impacts.

## 2 | MATERIALS AND METHODS

### 2.1 | Field sites

The 3-year experiment was carried out in the period 2018–2020. In 2018, triplicate experimental blocks were laid out at two agricultural sites of contrasting soil types in a randomized complete block design. The first site is located near Heelsum, the Netherlands (51°58'42" N, 5°45'58" E), and consists of a coarse sandy Anthrosol (WRB-FAO classification; 74% sand (coarse), 20% silt, 2% clay, 3.7% organic matter, pH 5.4). The site lies elevated in its surroundings, resulting in a deep-water table (>2 m), making the crops vulnerable to drought. The second site is located near Harreveld (51°59'8" N, 6°29'53" E) and is characterized as a loamy-sand Podzol (WRB-FAO classification; 72% sand (fine), 19% silt, 3% clay, 6.2% organic matter, pH 5.6), an aeolian deposition of the Pleistocene. Both sites have a history of maize (*Zea mays* L.) cultivation with seasonal rotation of *Lolium multiflorum* Lam. as a winter catch crop. In contrast to the Heelsum site, however, the winter catch crop at Harreveld is sown together with the maize instead of after harvest and its maize cultivation was periodically alternated with meadow. Both sites lie within the same region and have an annual mean temperature of 10°C, with an average minimum of 0°C and maximum of 21°C, and an average annual precipitation of 825 mm for the period 1991–2020 (KNMI, The Royal Netherlands Meteorological Institute, 2022). An overview of the soil properties of each site is presented in Supporting Information A.

To create an experimental setting closest resembling agro-economic reality, the farmers at both sites had agreed to manage their fields according to their own conventional practices, but to stay consistent in these practices for the duration of the 3-year experiment. A resulting difference between the sites

is an extra tillage that occurs each year after harvest, before winter sampling, at the Anthrosol site. Its purpose was to integrate the corn stubble into the soil and plant the winter catch crop. At the Podzol site, this tillage was not necessary, as the catch crop was sown in together with the maize in spring. Other differences in site management are listed in Supporting Information B.

### 2.2 | Organic amendments

Four OAs were applied as treatment factors at each site: (i) compost, (ii) farmyard manure, (iii) Bokashi or a locally available organic residue, and (iv) the combination of (ii) and (iii). For the Anthrosol site, the organic residue (iii) consisted of grasses (G) recovered from public, rural, green areas such as roadsides and parks. For the Podzol site, the organic residue (iii) consisted of waterway weeds and reeds (WWR) in the first year (not included in this study), naturally fermented WWR in the second year, and a commercial, fermented Bokashi product in the third year—varying with local availability. We included the combination treatment of manure and organic residue as it reflects a Dutch agronomic reality where organic residues are unlikely to be applied alone given the excess of manure produced by the Dutch livestock industry. The four organic treatments at each site were compared to a control treatment that did not receive any organic input. Each treatment was replicated in three 10-by-10-m blocks.

Application rates were determined based on the nutrient properties (N, P, and K concentrations). Based on national fertilization recommendations and standard application norms (RVO, 2022; Schoonvelde et al., 2020), we limited the total nutrient input for each treatment to 120 kg ha<sup>-1</sup> available N, 50 kg ha<sup>-1</sup> P<sub>2</sub>O<sub>5</sub>, and 200 kg ha<sup>-1</sup> K<sub>2</sub>O. Each OA was applied in maximum until one of these nutrient limits was met. These rates varied slightly year-to-year due to the small variations in OA quality of each year. On average, application rates amounted to approximately 20 t ha<sup>-1</sup> fresh matter for compost, 30 t ha<sup>-1</sup> for manure, 20 t ha<sup>-1</sup> for grasses, 20 t ha<sup>-1</sup> for WWR, 20 t ha<sup>-1</sup> for Bokashi, and 50 t ha<sup>-1</sup> for the mixed treatment of combined grasses or WWR with manure. To isolate as much as possible the effect of the OAs and their organic chemical characteristics and thus to offset as much as possible the potential effect of differences in nutrient inputs on soil properties, each plot, including the control, was supplemented with mineral fertilizers (N, P, K). This furthermore reflects a Dutch agro-industrial reality where many farmers supplement OA applications with mineral fertilizers in order to meet the maximum fertilization limits for each nutrient. The resulting wide variations in both the quantity and quality of the applied OAs were regarded as an asset, allowing a general assessment

of the importance of both aspects in contributing to resulting changes in soil properties.

The OAs were mechanically integrated into the soil with a disc harrow to approximately 15-cm depth. An overview of the mineral and organic applications to each block, per year, and per site, is presented in Supporting Information C. Analysis of OA chemical properties was performed by the Nutrient Management Institute B.V., Wageningen, and Eurofins Agro, Wageningen following conventional methods using predominantly elemental combustion analysis. A complete overview of the OA characteristics is presented in Supporting Information D.

## 2.3 | Field sampling and measurements of hydro-physical soil properties

Organic amendments were applied after the harvest of the winter catch crop each year. Subsequently, maize was sown and mineral fertilizers were applied. A full overview of the field management and sampling dates is presented in Supporting Information B. Infiltration capacity, aggregate stability, and water retention were measured in late June, approximately 3 months after sowing, and in late November, approximately 3 months after harvest/ploughing, in years 2 (2019) and 3 (2020) of the experiment. Soil bulk density was only sampled in late May and late November, in the final year of the experiment (2020) when OA impacts were expected to be greatest.

### 2.3.1 | Bulk density

Soil cores of 5-cm length and 5.3-cm diameter (110.3 cm<sup>3</sup>) were inserted after carefully removing 10 cm of topsoil to recover an undisturbed soil sample spanning a depth of 10–15 cm. The contents of the soil cores were oven-dried at 70°C for 24 h and weighed. Soil bulk density ( $p_b$ ) was calculated by  $p_b = M_d/V$ , where  $M_d$  was the dried soil mass and  $V$  was the volume of the soil core.

### 2.3.2 | Infiltration capacity

Infiltration capacity was measured in situ using a standard, double-ring infiltrometer. Outer (ø55 cm) and inner (ø30 cm) galvanized steel rings were inserted into the soil up to a depth of 10 cm. Both rings were filled to ~10-cm head, after which the change in the water level of the inner ring was recorded over time. Throughout the infiltration process, the water level of the outer ring was kept equal to that of the inner ring. Measurements were repeated until we no longer observed a significant rate of change in water level. We commenced

each measurement by refilling both rings after the water level inside them had dropped to approximately 1-cm head.

### 2.3.3 | Aggregate stability

Changes in wet aggregate stability were determined following an adapted wet sieving method (International Maize & Wheat Improvement Center, 2013). Triplicate soil samples were collected spanning an approximate depth of 10–17 cm, transported in closed 70-mL polypropylene containers, and air-dried for 2 weeks. Samples were transferred to aluminum cups and oven-dried at 60°C for 24 h before sieving. The oven-dry samples were sieved at 8 mm to remove any roots, rocks and pebbles, and to break any large aggregates (>8 mm; along natural planes of weakness). A sieve stack of descending sizes (2 mm, 1 mm, 500 µm, 250 µm, and 125 µm) was then submerged into a column of water. Approximately 20–40 g of dry soil ( $M_T$ ) was weighed and slowly wetted with deionized water, using a wash bottle, and gently poured onto the top of the 2-mm sieve. The sample was sieved underwater in a vertical direction for 2 min at a frequency of 30 waves, of 3-cm amplitude, per minute. Each sieve was subsequently washed, breaking the aggregates, and recovering the dispersed aggregate material using a hand vacuum pump and a large funnel capped with a pre-weighed, rayon polyester filter cloth (pore size of 22–25 µm). The filter with sample ( $M_{Ai}$ ) was oven-dried at 60°C. Coarse material remaining on each sieve ( $M_{Ci}$ ) was pooled for each sample and dried accordingly. The fraction of water-stable aggregates ( $W_{Fi}$ ) was then calculated using the following equation:

$$W_{Fi} = \frac{M_{Ai}}{M_T - \sum_{i=1}^n M_{Ci}} \quad (1)$$

where  $i = 1, 2, \dots, n$  and corresponds to each aggregate size fraction. The mean weight diameter (MWD) [mm] was calculated as an index for aggregate stability (Equation 2):

$$\text{MWD} = \sum_{i=1}^n W_{Fi} \times x_i \quad (2)$$

where  $x_i$  is the average diameter of each size fraction (i.e., mean intersieve size) and  $W_{Fi}$  is as defined in Equation (1).

### 2.3.4 | Water retention

Changes in soil water retention were determined using pressure plates at the Soil Hydro-Physics lab of Wageningen University & Research. Two soil samples per treatment were extracted from 10-cm depth, one disturbed sample using a garden hand shovel and a second undisturbed sample using

a 110.3-cm<sup>3</sup> soil core. In the laboratory, the disturbed sample was turned into sludge with demineralized water. A 20-cm<sup>3</sup> rubber ring was placed on a porous ceramic pressure plate and filled with the soil sludge. The pressure plate was placed in a pressure pan and incubated at a relative pressure of 14,000 hPa for the Wilting Point estimate, which is more than sufficient pressure for sandy soils like the ones tested (Wiecheteck et al., 2020). The undisturbed soil core was saturated by capillary action and incubated in a pressure pan at 98 hPa for the Field Capacity estimate. After a 2-week incubation period, the samples were removed from the plate, weighed, oven-dried at 70°C, and weighed a second time to determine the difference in gravimetric water content ( $w$ ). Gravimetric water content was converted to volumetric water content ( $\theta$ ) by:  $\theta = w \times p_b / p_w$ ; where  $p_b$  is based on the measured dry mass of the undisturbed soil core sample and the known soil core volume; the density of water ( $p_w$ ) is assumed to be 1 g mL<sup>-1</sup>. Plant available water (PAW) was estimated by subtracting the field capacity measurement ( $\theta_{FC}$ ) from the wilting point ( $\theta_{WP}$ ) measurement:  $PAW = \theta_{FC} - \theta_{WP}$ .

## 2.4 | Statistical analyses

We assessed OA impacts on soil hydro-physical properties using repeated-measures generalized linear mixed-effects models (GLMEMs) of a gamma distribution family. To account for imbalances in the data while allowing an assessment of the impact of the site, two sets of models were fit; one set was fitted to all data at each site, and the other set was fitted only to the combined data of the mutually shared OA treatments at both sites (i.e., control, compost, and manure). The first set of models was designed to include additive and interactive effects between the factors *amendment* and sampling *season* (Equation 3). The second set of models, applied to data from the shared treatments, included the additive and interactive effects of the *site* (Equation 4). Both sets of models included random error effects for sampling year and treatment block:

$$y_{ijlm} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + \varepsilon_{l(i)} + \varepsilon_m \quad (3)$$

$$\begin{aligned} y_{ijklm} = & \mu + \alpha_i + \beta_j + \delta_k + (\alpha\delta)_{ik} + (\beta\delta)_{jk} + (\alpha\beta)_{ij} \\ & + (\alpha\beta\delta)_{ijk} + \varepsilon_{l(i,k)} + \varepsilon_{m(k)} \end{aligned} \quad (4)$$

where  $y_{ijlm} \in (0, \infty)$  is the measured hydro-physical property at each site for amendment  $i$ , season  $j$ , block  $l$ , and year  $m$ , and  $z_{ijklm} \in (0, \infty)$  is the measured hydro-physical property for shared treatments at both sites including, in addition to the terms of Equation (3), a site factor  $k$  and its interactive effects. Furthermore,  $\mu$  is the population mean;  $\alpha_i$ ,  $\beta_j$ , and  $\delta_k$  are the fixed effects of amendment, season, and site. For

the site-specific model,  $\varepsilon_{l(i)}$  is the random block error effect per treatment, and  $\varepsilon_m$  is the random year error effect. For the shared treatment model,  $\varepsilon_{l(i,k)}$  is the random block error effect per treatment and per site, and  $\varepsilon_{m(k)}$  is the random year error effect per site.

The models were fit to the acquired data through maximum likelihood estimation by Laplace approximation. The variance associated with each effect factor was evaluated through a multi-way analysis of deviance testing by Type III Wald  $\chi^2$  tests with power factor  $\alpha = 0.1$ . Subsequently, for each season, statistical significance of the improvements in soil properties of OA amended soils relative to the unamended control soil was determined through pairwise comparison of OA amended and control soils using single-tailed, Sidak post hoc tests. Sidak post hoc tests were then applied a second time to test for the statistical significance of seasonal differences in soil improvements for each OA through multiple pairwise comparisons of the improvement relative to the control of each season for each OA. The results of these tests are presented in bar graphs together with the estimated marginal means (EMMs) of seasonal improvements and seasonal differences per OA. Presenting the data as EMMs allows representation of OA effects while accounting for imbalances in the data (Lenth, 2020).

The water retention data, which were restricted in sample size due to the loss of data after using a leaky pressure plate, were scaled to the observations of the control soil ( $WR_{OA, impact} = \frac{WR_{OA} - WR_{Control}}{WR_{Control}}$ ). The results, which included negative values, were approximated by a Gaussian distribution and fit to a linear mixed-effects model (LMEM).

To evaluate the effect of organic matter application on hydro-physical improvements during each season, as well as the differences between seasons, simple linear regression models (LMs) were built. The LMs relate different forms of expression of OA application rates (i.e., as organic matter, fresh matter, or dry matter) as well as OA compositional properties (i.e., total carbon or nitrogen concentrations, organic N, P, or K concentrations and C:N, C:P, or N:P ratios) to the observed changes in soil hydro-physical properties. We identified the OA property with the strongest influence on each respective soil hydro-physical property by comparing the significance and size of the correlation coefficient for each model. All statistical analyses were performed using the emmeans (Lenth, 2020) and lme4 (Bates et al., 2015) packages in an RStudio programming environment (RStudio Team, 2020).

## 2.5 | Impact classification for temporal response

The temporal impact of OAs on each soil property was classified into one of four groups to simplify comparison:

**TABLE 1** Analysis of deviance  $\chi^2$  values for the generalized linear mixed-effects models (GLMEMs) for various soil hydro-physical properties.

Analysis of deviance Wald $\chi^2$ —Type III Per site (Equation 3)	Chi-square value							
	Bulk density		Infiltration capacity		Aggregate stability		Water retention	
	Phaeoz.	Podzol	Phaeoz.	Podzol	Phaeoz.	Podzol	Phaeoz.	Podzol
Amendment	8.25●	10.9*	28.2***	8.79	20.7***	15.6*	31.3***	23.6***
Season	0.72	3.96*	0.05	4.51*	0.61	0.11	1.00	0.01
Amendment:season	7.96●	7.70	16.1**	6.04	5.28	16.0*	48.2***	8.51●
<b>Shared treatments (Equation 4)</b>								
Amendment	14.6***		0.21		2.43		2.99	
Site	5.01*		0.94		5.28*		0.00	
Season	5.86*		7.18**		0.13		0.04	
Amendment:site	0.37		1.30		0.22		8.87*	
Amendment:season	9.95**		1.00		2.75		4.64●	
Site:season	0.63		1.84		0.01		0.01	
Amendment:site:season	4.41		1.82		1.61		13.3**	

Note: Higher  $\chi^2$  values indicate an increased likelihood that the detected variation in the data occurs due to the influence of the effect and not by chance. Site interaction effects are only representative of the shared treatments (i.e., control, compost, and manure treatments).

\*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; ● $p < 0.1$ .

1. *transient*, where a significant improvement was observed in an earlier but not a later sampling season;
2. *persistent*, where significant improvements were observed for both earlier and later sampling seasons, and/or sampling season did not affect the observed improvement;
3. *lagged*, where a significant improvement was not observed for an earlier sampling season but was observed for a later sampling season; and
4. *enhanced*, where a significant improvement was observed in an earlier season as well as a later season, and a significant enhancement of this improvement was observed between sampling seasons.

### 3 | RESULTS

#### 3.1 | Main and seasonal interaction effects of organic amendments on soil hydro-physical properties

Most of the evaluated soil hydro-physical properties were significantly affected by the *amendment* applied, except for infiltration capacity at the podzol site ( $p < 0.1$ ). This *amendment* effect depended on the *season* for bulk density and infiltration capacity at the anthrosol site, aggregate stability for the podzol site, and water retention at both sites ( $p < 0.1$ ; Table 1). We therefore conditionally accept the hypothesis (i) that improvements in soil hydro-physical properties are significantly affected by the OA treatments—except for infiltration capacity at the podzol site—and (ii) that these improvements are significantly different depending on sampling season for at least one of the sites. Moreover, we observe that—for the


treatments shared between sites—the effects of OAs on bulk density and aggregate stability varied strongly per site.

#### 3.2 | Bulk density

Sidak post hoc testing demonstrated that bulk density was significantly reduced for various OAs in the summer at both sites, but not in the winter at the podzol site ( $p < 0.1$ ; Figure 1, Supporting Information E). Greatest decreases in bulk density were observed for compost in the summer at the podzol site, reducing bulk density by  $0.12 \text{ g cm}^{-3}$  (−9.8%). Reductions in bulk density attained in summer had largely disappeared by winter for compost ( $\Delta 0.08 \text{ g cm}^{-3}$ ) at the anthrosol site and compost ( $\Delta 0.13 \text{ g cm}^{-3}$ ) and Bokashi ( $\Delta 0.06 \text{ g cm}^{-3}$ ) at the podzol site ( $p < 0.1$ ; Supporting Information E). The impact of different OAs on bulk density differed within a single season by 2.5%–9.8% relative to the unamended control (Supporting Information F).

##### 3.2.1 | Effect of amendment application rate

Amendment application rates, expressed as either organic matter, fresh matter or dry matter, were correlated to decreases in bulk density only in summer at the anthrosol site (Table 2). Strongest correlations were observed when amendment application rates were expressed in the form of dry matter ( $R_{\text{DM}} = 0.92^{***}$ ; Table 2). Differences between seasons for each OA treatment were also only significant at the anthrosol site, where the strongest correlation was again observed for the dry matter application rate ( $R_{\text{DM}} = 0.92^{***}$ ; Table 2; Figure 2).



**FIGURE 1** Estimated marginal means (EMMs) of bulk density after 3 years of repeated organic amendment (OA) application. Symbols at grey (summer) and black (winter) bars indicate the level of significance of the change in bulk density when compared to the control, while symbols for orange bars indicate the significance of the difference in decrease relative to the control between winter and summer for that OA treatment: \*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ .

**TABLE 2** Correlation coefficients for linear regression models (LMs) relating amendment application rates to seasonal improvements and differences in bulk density.

Correlation coefficients OA property versus bulk density	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
Organic matter	0.68*	0.11	0.81**	0.3	0.23	0.19
Fresh matter	0.76*	0.3	0.31	0.09	0.04	0.16
Dry matter	<b>0.92***</b>	0.01	<b>0.92***</b>	0.51	0.32	0.33

Note: Significant correlations are presented in bold.

Abbreviation: OA, organic amendment.


\*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ .

### 3.2.2 | Effect of amendment composition

Seasonal decreases and inter-seasonal differences in bulk density were correlated to OA compositional properties at both sites, except for summer at the podzol site (Table 3). At the anthrosol site, summer reductions in bulk density correlated most strongly to OAs phosphorus concentrations ( $R_P = 0.74^*$ ), winter reductions to carbon and active nitrogen concentrations ( $R_{C,R_{N-Active}} = 0.47^\bullet$ ), and differences between seasons to C:N ratio's ( $R_{C:N} = 0.72^*$ ). While neither reductions per season, nor differences between seasons, were significantly correlated to OM application rates at the podzol site (Supporting Information F1, Figure 2), we did observe correlations of winter improvements with OA total carbon ( $R_C = 0.59^*$ ) and nitrogen concentrations ( $R_{Nt} = 0.61^*$ ), and a correlation of inter-seasonal differences with potassium concentrations ( $R_K = 0.58^\bullet$ ).

### 3.3 | Infiltration capacity

Infiltration capacity was significantly increased in summer but not in winter for multiple OAs at the anthrosol site and both in summer and winter for several OAs at the podzol site ( $p < 0.1$ ; Figure 3, Supporting Information E). Greatest increases were observed in summer, where the grass and manure treatment increased infiltration capacity at the anthrosol site by  $7.2 \text{ mm min}^{-1}$  (108%) and WWR at the podzol site by  $5.52 \text{ mm min}^{-1}$  (99%). Summer improvements at both sites had significantly diminished by winter for all relevant treatments at the Anthrosol site but only for WWR at the podzol site ( $\Delta -4.82 \text{ mm min}^{-1}$ ;  $p < 0.1$ ; Supporting Information E). The impact of different OAs on infiltration capacity differed within a single season by 26.4%–100.8% relative to the unamended control (Supporting Information F).



**FIGURE 2** Bulk density response to different rates of total dry matter application in the experimental period of 3 years with the different organic amendments (OAs). Symbols indicate the significance of the regression trend: \*\*\* $p < 0.001$ ; n.s.  $p > 0.1$ . n.s., not significant.

**TABLE 3** Correlation coefficients for linear regression models (LMs) relating organic amendment (OA) composition (concentrations and ratios) to seasonal improvements (reductions) and differences in bulk density.

Correlation coefficients OA property versus bulk density	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
C <sub>Total</sub>	0.17	<b>0.47●</b>	0.05	0.13	0.59*	0.25
N <sub>Total</sub>	0.16	0.46●	0.23	0.12	<b>0.61*</b>	0.28
N <sub>Active</sub>	0.13	<b>0.47●</b>	0.17	0.08	0.38	0.18
P <sub>Organic</sub>	<b>0.74*</b>	0.24	0.3	0.54	0.2	0.46
K <sub>Organic</sub>	0.64*	0.03	0.37	0.45	0.21	<b>0.58●</b>
C:N	0.54	0.27	<b>0.72*</b>	0.31	0.04	0.34
C:P	0.35	0.21	0.67*	0.13	0.2	0.3
N:P	0.55	0.31	0.64*	0.2	0.28	0.41

Note: The maximum of each column is presented in bold.


\* $p < 0.05$ ; ● $p < 0.1$ .

### 3.3.1 | Effect of amendment application rate

Amendment application rates in the form of organic matter (OM) correlated with increases in infiltration capacity in summer ( $R_{OM} = 0.59^{**}$ ) and winter ( $R_{OM} = 0.42^{*}$ ) at the anthrosol site, whereas, for the podzol site, only a weak correlation for winter increases was observed with fresh matter application rate ( $R_{FM} = 0.32^{*}$ ; Table 4). Also, the differences between seasons for each OA treatment correlated only to OM application rates and only at the anthrosol site ( $R_{OM} = 0.47^{*}$ ; Table 4, Figure 4).

### 3.3.2 | Effect of organic amendment composition

Seasonal increases and inter-seasonal differences demonstrated *weaker* correlations to amendment composition than OM application rates at both sites (Table 5 and Table 2). At the anthrosol site, summer improvements were significant but less strongly related to C:P ratios than to organic matter application rates ( $R_{C:P} = 0.52^{**}$  vs.  $R_{OM} = 0.59^{**}$ ), while no compositional property was related to winter improvements. Differences between seasons at the anthrosol site also were slightly less correlated with C:P and N:P ratios than with OM



**FIGURE 3** Estimated marginal means (EMMs) of infiltration capacity response to repeated organic amendment (OA) application. Symbols at grey (summer) and black (winter) bars indicate the level of significance of the increase when compared to the control for each season, while symbols at orange bars indicate the significance of the difference between winter and summer for that OA treatment: \*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ . WWR, waterway weeds and reeds.

**TABLE 4** Correlation coefficients for linear regression models (LMs) relating amendment application rates to seasonal improvements and differences in infiltration capacity.

Correlation coefficients OA property versus infiltration capacity	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
Organic matter	<b>0.59**</b>	<b>0.42*</b>	<b>0.47*</b>	0.08	0.25	0.06
Fresh matter	0.31	0.06	0.31	0.05	<b>0.32•</b>	0.16
Dry matter	0.26	0.28	0.28	0.23	0.11	0.21

Note: Significant correlations are presented in bold.

Abbreviation: OA, organic amendment.


\*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ .

**TABLE 5** Correlation coefficients for linear regression models (LMs) relating organic amendment (OA) composition (as concentrations and ratios) to seasonal improvements and differences in infiltration capacity.

Correlation coefficients OA property versus infiltration capacity	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
C <sub>Total</sub>	0.28	0.1	0.27	0.23	0.09	0.05
N <sub>Total</sub>	0.15	0.2	0.17	0.24	0.08	0.05
N <sub>Active</sub>	0.22	0.16	0.23	0.22	0.09	0.06
P <sub>Organic</sub>	0.14	0.13	0.04	0.09	0.15	0.02
K <sub>Organic</sub>	0.28	0.09	0.12	0.09	0.12	0.02
C:N	<b>0.42*</b>	0.17	<b>0.37•</b>	0.17	0.17	0.08
C:P	<b>0.52**</b>	0.3	<b>0.41*</b>	0.18	0.14	0.11
N:P	<b>0.50**</b>	0.19	<b>0.41*</b>	0.16	0.14	0.12

Note: The maximum of each column is presented in bold.

\*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ .



**FIGURE 4** Infiltration capacity response to different rates of organic matter application. Symbols indicate significance of the regression trend: \*\* $p < 0.01$ ; \* $p < 0.05$ ; n.s., not significant.

application rates ( $R_{C:P}$ ,  $R_{N:P} = 0.41^*$  vs.  $R_{OM} = 0.47^*$ ). At the podzol, OA composition properties showed no correlation to seasonal improvements nor inter-seasonal differences.

### 3.4 | Aggregate stability

Aggregate stability was significantly increased for various OAs in both summer and winter at both sites ( $p < 0.1$ ; Supporting Information E, Figure 5). Greatest increases were observed in the winter, where the grass and manure treatment improved aggregate stability at the anthrosol site by 0.75 mm (60.0%) and bokashi at the podzol site by 0.73 mm (40.5%). Aggregate stability was significantly improved between summer and winter by compost ( $\Delta 0.19$  mm) and manure ( $\Delta 0.35$  mm) at the anthrosol site and by bokashi ( $\Delta 1.21$  mm) at the podzol site ( $p < 1$ ; Supporting Information E; Figure 5). The impact of OAs on aggregate stability differed within a single season by 8.2%–46.3% relative to the unamended control (Supporting Information F).

#### 3.4.1 | Effect of amendment application rate

Amendment application rates as fresh matter correlated strongly to increases in aggregate stability in summer ( $R_{FM} = 0.61^{***}$ ) and winter ( $R_{FM} = 0.42^*$ ) at the anthrosol site, whereas application rates expressed as dry matter correlated most strongly to winter increases at the podzol site ( $R_{DM} = 0.53^{**}$ ; Table 6). The seasonal differences at the anthrosol site were not related to the OA application rate,


whereas at the podzol site, they correlated again to the dry matter application rate ( $R_{DM} = 0.48^*$ ; Table 6; Figure 6).

### 3.4.2 | Effect of amendment composition

Seasonal increases and inter-seasonal differences in aggregate stability demonstrated a *weaker* correlation to OA compositional properties than to OM application rates at both sites (Tables 6 and 7). At the anthrosol site, summer increases were most strongly correlated to N:P ratios ( $R_{N:P} = 0.61^{***}$  vs.  $R_{FM} = 0.69^{***}$ ), and winter increases were most strongly correlated to C:N and N:P ratios ( $R_{C:N}$ ,  $R_{N:P} = 0.46^*$  vs.  $R_{FM} = 0.57^{**}$ ). Seasonal differences correlated only with phosphorus ( $R_P = 0.42^*$ ) and potassium concentrations ( $R_K = 0.39^*$ ). At the podzol site, OA application rates did not correlate to summer increases, whereas a weak correlation was observed with total nitrogen concentrations ( $R_{N-Total} = 0.37^\bullet$ ). Weaker correlations with OA composition than OA application rates were observed for winter increases, which correlated with C:N ratios ( $R_{C:N} = 0.38^*$  vs.  $R_{DM} = 0.53^{**}$ ), and the difference between seasons, which correlated with phosphorus concentrations ( $R_P = 0.39^\bullet$  vs.  $R_{DM} = 0.48^{**}$ ).

### 3.5 | Water retention

Water retention was significantly increased at the anthrosol site in both seasons for the grass and manure treatments and



**FIGURE 5** Estimated marginal means (EMMs) of aggregate stability response to repeated organic amendment (OA) application. Symbols at grey (summer) and black (winter) bars indicate the level of significance of the increase when compared to the control for each season, while symbols at orange bars indicate the significance of the difference between winter and summer for that OA treatment: \*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ . WWR, waterway weeds and reeds.

**TABLE 6** Correlation coefficients for linear regression models (LMs) relating amendment application rates to seasonal improvements and differences in aggregate stability.

Correlation coefficients OA property versus aggregate stability	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
Organic matter	0.61***	0.42*	0.05	0.02	0.33•	0.25
Fresh matter	<b>0.69***</b>	<b>0.57**</b>	0.14	0.19	0.26	0.02
Dry matter	0.37•	0.39*	0.2	0.13	<b>0.53**</b>	<b>0.48*</b>

Note: Significant correlations are presented in bold.

Abbreviation: OA, organic amendment.


\*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ .

**TABLE 7** Correlation coefficients for linear regression models (LMs) relating organic amendment (OA) composition (as concentrations and ratios) to seasonal increases and differences in aggregate stability.


Correlation coefficients OA property versus aggregate stability	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
C <sub>Total</sub>	0.41*	0.44*	0.2	0.34•	0.03	0.26
N <sub>Total</sub>	0.32	0.41*	0.22	<b>0.37•</b>	0.04	0.29
N <sub>Active</sub>	0.46*	0.44*	0.16	0.24	0.09	0.08
P <sub>Organic</sub>	0.12	0.45*	<b>0.42*</b>	0.16	0.34•	<b>0.39•</b>
K <sub>Organic</sub>	0.14	0.24	0.39*	0.07	0.08	0.00
C:N	<b>0.60***</b>	<b>0.46*</b>	0.12	0.04	<b>0.38*</b>	<b>0.35•</b>
C:P	0.53**	0.35•	0.06	0.16	0.3	0.13
N:P	<b>0.61***</b>	<b>0.46**</b>	0.12	0.24	0.32•	0.07

Note: The maximum of each column is presented in bold.

\*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; • $p < 0.1$ .



**FIGURE 6** Aggregate stability response to different amendment application rates expressed in the form of fresh matter for the anthrosol site and dry matter for the podzol site because of their respective greatest significance. Symbols indicate significance of the regression trend: \*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; n.s., not significant.



**FIGURE 7** Estimated marginal means (EMMs) of water retention response to repeated organic amendment (OA) application. Symbols above grey (summer) and black (winter) bars indicate the level of significance of the improvement when compared to the control for each season, while symbols above orange bars indicate the significance of the difference between summer and winter for that OA treatment: \*\* $p < 0.01$ ; \* $p < 0.05$ ; \* $p < 0.1$ . WWR, waterway weeds and reeds.

in winter additionally for the grasses treatments. At the podzol site, it was significantly improved in summer for WWR and manure and bokashi and manure, and in winter for bokashi and bokashi and manure ( $p < 0.1$ ; Figure 7; Supporting Information E). Greatest increases were observed in the winter,

where water retention was improved for the grasses treatment at the anthrosol site by 57% and for the bokashi and manure treatment at the podzol site by 77%. Water retention at both sites had significantly improved between summer and winter only for the grasses treatment ( $\Delta 25\%$ ) at the


**TABLE 8** Correlation coefficients for linear regression models (LMs) relating amendment application rates to seasonal improvements and differences in water retention.

Correlation coefficients OA property versus water retention	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
Organic matter	<b>0.94***</b>	<b>0.82●</b>	0.54	0.55●	<b>0.83●</b>	<b>0.95*</b>
Fresh matter	0.5	0.13	0.23	0.58●	0.76	0.77
Dry matter	0.71*	0.5	0.14	0.01	0.45	0.79

Note: Significant correlations are presented in bold.

Abbreviation: OA, organic amendment.

\*\*\* $p < 0.001$ ; \* $p < 0.05$ ; ● $p < 0.1$ .



**FIGURE 8** Water retention response to different rates of organic matter application. Symbols indicate significance of the regression trend: \*\*\* $p < 0.001$ ; \* $p < 0.05$ ; ● $p < 0.1$ ; n.s., not significant.

anthrosol site ( $p < 1$ ; Supporting Information E). The impact of OAs on water retention differed within a single season by 11.1%–77.8% relative to the unamended control (Supporting Information F).

### 3.5.1 | Effect of amendment application rate

Amendment application rates correlated to increases in water retention in both seasons at both sites when expressed as organic matter (Table 8). At the anthrosol site, these correlations were relatively stronger (summer:  $R_{OM} = 0.94^{***}$ ; winter  $R_{OM} = 0.82●$ ) than at the podzol site (summer:  $R_{OM} = 0.55●$ ; winter  $R_{OM} = 0.83●$ ; Supporting Information F). The differences between seasons, however, were linearly related to OM application rates only at the podzol site ( $R_{OM} = 0.95^*$ ; Table 8; Figure 8). Due to the loss of

1 year of winter data, the standard error of winter and season difference trends are much higher than for other soil properties.

### 3.5.2 | Effect of amendment composition

For water retention, the seasonal improvements and inter-seasonal differences demonstrated a *weaker* correlation to OA compositional properties than to OM application rates at both sites (Tables 8 and 9). At the anthrosol site, only summer increases were significantly correlated with OA compositional properties, namely C:N and C:P ( $R_{C:N} = 0.58●$ ,  $R_{C:P} = 0.63●$  vs.  $R_{OM} = 0.94^{***}$ ). At the podzol site, only the differences between seasons were significantly correlated to OA compositional properties ( $R_P = 0.89^*$ ,  $R_{C:N} = 0.87●$  vs.  $R_{OM} = 0.95^*$ ).

**TABLE 9** Correlation coefficients for linear regression models (LMs) relating organic amendment (OA) composition (as concentrations and ratios) to seasonal improvements and differences in water retention.

Correlation coefficients OA property versus water retention	Anthrosol			Podzol		
	Summer	Winter	Difference	Summer	Winter	Difference
C <sub>Total</sub>	0.00	0.02	0.07	0.21	0.04	0.03
N <sub>Total</sub>	0.17	0.3	0.35	0.16	0.01	0.01
N <sub>Active</sub>	0.04	0.11	0.16	0.43	0.42	0.36
P <sub>Organic</sub>	0.08	0.38	0.66	0.11	0.52	<b>0.89*</b>
K <sub>Organic</sub>	0.01	0.44	0.67	0.21	0.05	0.41
C:N	0.58●	0.6	0.43	0.39	0.7	0.87●
C:P	<b>0.63●</b>	0.76	0.67	0.4	0.6	0.67
N:P	0.54	0.57	0.38	0.27	0.48	0.64

Note: The maximum of each column is presented in bold.

\* $p < 0.05$ ; ● $p < 0.1$ .

**TABLE 10** Seasonal differences in organic amendment (OA) impacts as percent relative to control (%winter–%summer) for different OA treatments.

Seasonal difference OA impact relative to control	Site	Bulk density (%W–%S)	Infiltration capacity (%W–%S)	Aggregate stability (%W–%S)	Water retention (%W–%S)	Average absolute difference (%W–%S)
Compost	Anthrosol	6.1●	9.7	27.8●	–29.4	18.3
	Podzol	10.7*	41.6†	24.5†	20.2	24.3
Manure (M)	Anthrosol	–3.2●	–45.5●	28.7●	–23.2	25.2
	Podzol	8.6	18.4	–1.6	18.1	11.7
Grasses (G)	Anthrosol	3.0	–68.9*	6.2	10.1●	22.1
G&M	Anthrosol	2.2	–96.2**	13.7	25.6	34.4
WWR	Podzol	n.d.	–73.0●	21.5	n.d.	47.3
WWR&M	Podzol	n.d.	46.9	16.0	n.d.	31.5
Bokashi (B)	Podzol	4.7	–12.4	66.4***	39.5†	30.8
B&M	Podzol	1.5	58.5	19.2†	44.3	30.9
Average absolute difference		5.0	47.1	22.6	26.3	27.6
Standard deviation ( $\sigma$ )		2.9	25.5	16.0	9.9	8.8

Note: Original data are presented in Supporting Information E.

Abbreviations: n.d., no data; WWR, waterway weeds and reeds.

Symbols indicate the significance of the difference between winter and summer impacts for each OA treatment: \*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$ ; ● $p < 0.1$ , where “†” indicates that an OA impact was significant for one season but not another while the difference between seasons was not statistically significant.

### 3.6 | Seasonal variability of OA impacts

OAs varied strongly in their seasonal impacts on soil properties relative to the control (Table 10). Greatest seasonal impacts were observed for the WWR treatment (on average 47% relative to the control). Least impact was observed for manure at the podzol site (11.7%). Infiltration capacity had the greatest seasonal variation in OA impacts, showing an average 47% difference between summer and winter OA impacts relative to the control. Least variation was observed in bulk density, showing average seasonal differences of 5%.

The classification of the temporal variability in OAs impacts shows how some OAs have a lagged effect on one

property and a persistent, transient or enhanced effect on another, and how one soil property can demonstrate a transient response to one type of OA, but a lagged response to another type of OA (Table 11). It furthermore illustrates differences between sites. In general, bulk density appeared to be transiently improved at the podzol site regardless of the OA treatment, and transiently or persistently improved at the anthrosol site depending on the OA treatment. Infiltration capacity was mostly only transiently improved, especially at the anthrosol site, while for water retention the improvements were either lagged or persistent depending again on the OA treatment. For aggregate stability, most OAs had a persistent, lagged or enhanced effect. Furthermore, the classification

**TABLE 11** Classification of amendment effects per soil property and site.

	Amendment impact classification			
	Bulk density	Infiltration capacity	Aggregate stability	Water retention
<b>Anthrosol</b>				
Compost	Transient	–	Lagged	–
Manure	Lagged	Transient	Enhanced	–
Grasses	Persistent	Transient	Persistent	Lagged
Grasses and manure	Persistent	Transient	Persistent	–
<i>Amendment app.</i>	Lagged	Transient	Persistent	Persistent
<b>Podzol</b>				
Compost	Transient	Lagged	Lagged	–
Manure	Transient	–	Persistent	–
WWR <sup>a</sup>	n.d.	Transient	–	n.d.
WWR and manure <sup>a</sup>	n.d.	Persistent	–	n.d.
Bokashi <sup>b</sup>	Transient	–	Lagged	Lagged
Bokashi and manure <sup>b</sup>	–	–	Lagged	Persistent
<i>Amendment app.</i>	–	–	Lagged	Enhanced

Note: ‘–’ indicates no effect.

Abbreviations: n.d., no data; WWR, waterway weeds and reeds.

<sup>a</sup>Sampled only in year 2 of the experiment.

<sup>b</sup>Sampled only in year 3 of the experiment.

shows that no specific OA treatment consistently outperforms the others across all soil hydro-physical properties and sites. The limited significance of some of the OA treatments at the podzol site (with footnotes in Table 11) might be explained by their lower data count (1 year instead of 2 years of sampling).

## 4 | DISCUSSION

### 4.1 | Improvements following amendment application

Results showed substantial improvements in bulk density (up to –9.8% decrease), infiltration capacity (up to 108.1% increase), aggregate stability (up to 60.0% increase), and water retention (up to 77.8% increase) for most of the investigated OAs within 3 years of repeated application (Supporting Information F). While OA application is often regarded as a long-term strategy for soil improvement, as soil properties are typically slow to evolve in response to OA treatments (Diacono & Montemurro, 2011), the current experiment corroborates findings that significant soil improvements can already be achieved within a few years (Carron et al., 2015; Castellini et al., 2022; Delibacak et al., 2009; Ouyang et al., 2013).

OA application generally improved soil hydro-physical properties, but the magnitude of these improvements differed significantly depending on the OA treatment. Differences between OA impacts (relative to the control) were as large

as 9.6% for bulk density, 135% for infiltration capacity, 43% for aggregate stability, and 85% for water retention, depending on the season (Supporting Information F). Up to 140% differences for infiltration capacity have also been observed in other studies (Zamani et al., 2016). This suggests that not all OAs are equally effective in contributing to building soil resilience, as their impacts may not have developed yet or already have dissipated by the time that season-specific challenges occur.

Temporal effects of OAs were significant for all soil hydro-physical properties and were frequently larger than spatial, site-dependent differences (Table 2). Similar greater effects of temporal than (field-scale) spatial variability on soil hydro-physical properties have also been observed for unamended soils (van Es et al., 1999). Temporal influences such as climate and OAs thus appear to exert greater control on soil changes than differences that arise due to field-scale heterogeneity or differences between sites. Highest temporal variability was observed for the combined grasses and manure treatment, which showed nearly a 100% difference between summer and winter seasons for infiltration capacity. On average, OA treatments varied in their seasonal impacts relative to the control by 5% for bulk density, 47.1% for infiltration capacity, 22.6% for aggregate stability, and 26.3% for water retention. Field studies on *unamended* soils have also noted short-term temporal variation in soil hydro-physical properties such as bulk density, porosity, aggregate stability, and hydraulic conductivity, of which the latter furthermore demonstrated large differences in its temporal patterns for different soils (Jirků et al., 2013; Scott et al., 1994). To explain this variation,

Popolizio et al. (2022) identified soil moisture as a critical factor. Different soil moisture concentrations are believed to significantly influence changes in bulk density and hydraulic conductivity through effects such as swelling and slaking. Such differences could explain contrasting variations between the anthrosol and podzol sites but cannot explain differences in seasonal variability of OA impacts within each site, as seasonal water balances are the same for all treatments at a site. It would be interesting for future studies to measure seasonal changes in biological properties known to affect soil structure (Kinsbursky et al., 1989; Oades, 1984, 1993), and evaluate whether and how these relate to short-term changes in hydro-physical soil properties.

## 4.2 | Influence of quantity versus quality

Diversity in the composition and application rates of OAs allowed assessing the relative importance of OA quantity versus soil hydro-physical properties. Unfortunately, no single OA quantity or quality indicator could consistently explain the observed temporal trends for any soil property at either site. However, seasonal improvements and differences between seasons generally correlate better with OA quantity indicators (i.e., application rates) than with OA quality indicators (i.e., compositional properties). Other OA studies without combined mineral fertilizer applications have similarly observed the importance of OA application quantity over differences in OA type (Barzegar et al., 2002; Castellini et al., 2022; Delibacak et al., 2009; Tejada & Gonzalez, 2008).

Of the OA quality indicators, the C:N and N:P ratios, as well as P concentrations, generally demonstrated more and/or stronger correlations to both seasonal improvements and differences between seasons. The recurrence of nitrogen and phosphorus as a quality indicator is not surprising as the stoichiometric balance of these with other elements is recognized to be an important regulator of organic matter decomposition (Manzoni et al., 2010; Mooshammer et al., 2017). Since no single OA quality indicator performed best at both sites for any soil hydro-physical property, the impact of OA composition likely also depends on the conditions of the local soil environment. A complex interaction effect with the soil biotic and abiotic properties prior to OA application can often explain subsequent changes in the soil (Chen et al., 2022; Hamer & Marschner, 2005; Kok et al., 2022; Lloyd et al., 2016). Potentially, OA quality may have a more consistent impact on soil improvements in agricultural settings where the total nutrient inputs are not supplemented with mineral fertilizers and are, instead, completely dependent on the nutrients provided by the OAs applied.

While OA quantity correlated better than OA quality to observed changes in soil hydro-physical properties, the strength of these correlations varied significantly depending

on the expression of application quantity in either the form of fresh, dry or organic matter. Manure, for instance, due to its high water content, was applied at rates of around 30 kg *fresh matter* ha<sup>-1</sup> but only 2.5 kg ha<sup>-1</sup> in terms of *dry matter*. Compost, instead, was applied at much lower rates of 20 kg ha<sup>-1</sup> *fresh matter* and higher rates of 13 kg ha<sup>-1</sup> *dry matter*. Also, *organic matter* application rates varied significantly (i.e., ~3 kg ha<sup>-1</sup> for compost and manure, 4–14 kg ha<sup>-1</sup> for grasses, etc.; Supporting Information C). Differences in the strength of correlation between these metrics suggest that factors such as the initial water content of OAs (fresh vs. dry matter) or the ratio of organic to non-organic components (organic matter vs. dry or fresh matter) are potentially relevant in explaining differences in OA impacts.

Aggregate stability, for instance, showed stronger correlations than organic matter to application rates in terms of fresh matter, particularly at the anthrosol site (Table 6). Such site-dependent effects of manure applications have been observed before (Paré et al., 1999; Roldán et al., 1996; Whalen & Chang, 2002) and have been linked to differences in the leaching rates of aggregate dispersal agents (Guo et al., 2019). Such differences in leaching may also explain our results for the anthrosol site, given its coarser texture and higher infiltration capacity. Alternatively, the strong correlation of aggregate stability with high fresh matter application rates may relate to the slurry-like state of manure, allowing it to infiltrate soil pores and encapsulate soil particles resulting in its more efficient colonization and subsequent stimulation of aggregate stabilization. In general, however, for all soil properties, *organic matter* application rates demonstrated the most consistent correlations to observed improvements at both sites (Tables 2, 4, 6, and 8). This underscores the well-established importance of external carbon input as a stimulant of soil biological processes driving changes in the soil (Bronick & Lal, 2005; Tang et al., 2022; Ward et al., 2007).

## 4.3 | Influence of land management practices

Seasonal variability in OA impacts on hydro-physical properties is likely also influenced by land management practices, as these often involve considerable disturbance of a soil at different spatial and temporal scales (Hu et al., 2018). The occurrence of a disturbance event can affect improvements achieved by OAs, which might be reverted, remain unaffected, or potentially be amplified depending on the soil property and disturbance intensity. At the anthrosol site, for example, OAs with significant impacts all demonstrated a transient improvement in infiltration capacity, where improvements during the summer had completely dissipated by winter. A possible explanation for this consistent response for all OAs may be found in the effect of tillage between seasons at this site. Infiltration capacity measurements are sensitive to the

presence of preferential flow paths; thus, OAs potentially improved soil infiltration capacity by stimulating preferential flow path development (Ali et al., 2018). Such soil macrostructures, however, are also sensitive to physical disturbance events (Andreini & Steenhuis, 1990; Shipitalo et al., 2000), and so it is likely that the post-harvest tillage at the anthrosol site destroyed many of the developed preferential flow paths, resulting in the reversal of infiltration capacity improvements by winter. This explanation is furthermore supported by data from the podzol site, which was not tilled post-harvest, and where soils retained some of their improvements providing lagged or persistent effects by winter. Studies on unamended soils has similarly observed greater improvements in soil hydraulic properties when fields were not tilled by conventional means (Arshad et al., 1999; Pagliai et al., 2004; Pires et al., 2017). Additional tillage at the podzol site thus potentially contributed to generally reduced OA impacts on bulk density, infiltration capacity, and aggregate stability, as well as the greater seasonal differences observed for properties at this site.

Parallel to the challenge of uncovering universal relationships between OA properties with chemical and biological changes in soils (Kögel-Knabner, 2017), it also appears that linking OA properties to changes in soil hydro-physical properties is similarly not straightforward given the many possible interaction effects with land management. Moreover, some of the contradicting findings published in studies regarding variation in aggregate stability (Abiven et al., 2009; Albiach et al., 2001) and water retention (Minasny & McBratney, 2018) may be partially explained by the combination of temporal variability in these soil hydro-physical properties and differences in study sampling times (Carron et al., 2015).

#### 4.4 | Design of organic amendment application strategies

Though impacts varied, the application of OAs led to improvements in soil hydro-physical properties for nearly every OA type, at nearly every sampling moment, at both anthrosol and podzol sites. The application of compost, manure, grasses, grasses and manure, WWR, WWR and manure, bokashi, and bokashi and manure to dynamic agricultural soils thus appears substantially more likely to improve soil water management and crop growth rather than harm them (Abiven et al., 2009; Lal, 2020). However, not all investigated OAs appear equally effective for the purpose of improving soil resilience to specific seasonal weather extremes, given the differences in their temporal variability of impacts and the potential interaction of these effects with soil type and management practices. We have demonstrated that these temporal effects were significant for all soil hydro-physical properties, and that they frequently explained

similar or greater variation in the data than spatial, site-dependent differences (Table 2). This implies that designs of OA application strategies should account for temporal variability in their impacts and not rely on single or annual measurements, as is the current practice (e.g., Cercioğlu, 2017; Ferreras et al., 2006; Li et al., 2018; Singh Brar et al., 2015; Tejada & Gonzalez, 2008; Yüksel & Kavdır, 2020; Zhao et al., 2009). Our findings furthermore underscore that the parameterization of soil hydro-physical properties based solely on soil type may not be appropriate for agricultural lands given their significantly high short-term temporal variability due to soil-management factors such as OA treatments or tillage practices (van Es et al., 1999). Finally, for cultivation soils with fertilizer restriction and soils where OAs are supplemented with synthetic fertilizers, the design of OA application strategies would benefit from increasing the quantity of OA applied rather than the quality, as organic matter application rates demonstrated more consistent and stronger correlations to changes in soil hydro-physical properties than OA compositional properties did.

Most of the OAs investigated in this study are waste products derived from landscape maintenance, that is, clearing of waterways and mowing of road-verges, in treated or untreated forms. With little to no apparent negative consequence, for at least soil hydro-physical properties, the application of these organic wastes to immediately surrounding agricultural soils may be an efficient means of waste disposal (Sharma et al., 2019). Naturally, soil chemical impacts as well as toxicological, pathological, and biological hazards, would require further investigation before the widespread adoption of the less familiar OAs, for example waterway residues, in soil improvement practices (Ros et al., 2012).

Of the OAs investigated, compost generally produced temporally favorable impacts on hydro-physical properties. Comparing compost impacts is difficult as more than 600 studies on its role in agriculture have been published over the past 11 years alone (Rivier et al., 2022), although the findings of the current study correspond well with the ranges observed elsewhere (Supporting Information G).

Manure only produced temporally beneficial effects on aggregate stability at the anthrosol site and on bulk density and aggregate stability at the podzol site. Comparatively, the grass & manure combination treatment, which was only applied to the anthrosol soil, seems a more promising alternative to applying manure alone as it produced temporally favorable effects for more soil properties, namely, aggregate stability, bulk density, and water retention. The application of manure is a traditional soil improvement strategy that dates back to 6000 B.C. (Bogaard et al., 2013), and consequentially its effects have been extensively studied and reported (Rayne & Aula, 2020). Our observed manure impacts are at the lower end of observations in other studies for bulk density (2.1–14%; Celik et al., 2010) and hydraulic

conductivity (up to 234%; Fares et al., 2008). Changes in aggregate stability are in a similar range as reported elsewhere (from no significant effects (Karami et al., 2012) to 75% (Celik et al., 2010).

While OAs like compost and manure have been extensively studied, few experiments have evaluated the potential soil benefits resulting from the application of grasses recovered from public green areas and waterway residues recovered from public waterways and/or their combination with manure. An estimated 770,000 t of dry matter is produced each year in the regular landscape maintenance of public green areas in the Netherlands, indicating an enormous potential for their application as a soil improver (Elbersen & Spijker, 2014). Although only tested at one of the sites, our results showed that, when compared to compost, grasses resulted in a similar improvement in bulk density but greater improvements in infiltration capacity, aggregate stability, and water retention, though again the large improvement in the latter was not statistically significant. The impact of waterway residues (WWR) was difficult to assess due to the limited data (only sampled in 1 year, after 2 years of application) but appears to outperform compost only in improvements in infiltration capacity. Despite the excellent performance of grasses, their application to the soil posed practical issues. The grasses were difficult to integrate as they stuck to the tillage and harrow blades. For the purpose of the experiment, therefore, we manually facilitated soil integration. An alternative solution that is more easily implementable at an agro-industrial level could be to shred the grasses before application – although this may change the soil response. Nevertheless, grasses recovered from public green areas demonstrated a significant and favorable impact on a soil's hydro-physical properties which warrants further investigation of their utilization potential in agricultural settings.

## 5 | CONCLUSION

Our findings show that the application of OAs can improve soil hydro-physical properties up to 108%, but that differences in these improvements between seasons could vary up to 96% depending on the OA and soil property. While no OA property could consistently explain this temporal variability, results demonstrate a greater influence of OA application quantity and a lesser influence of OA composition (i.e., its quality), when OA nutrient differences are offset with mineral fertilizers. Of the OAs investigated, compost generally produced a temporally most favorable impact on the soil hydro-properties, though the improvements in some seasons were sometimes greater for the landscape residues (i.e., grasses, waterway residues, and bokashi fermented waterway residues) and/or their combined application with manure. Overall, findings indicate that not all OAs are equally effective

for combating season-specific climatic challenges, given that short-term OA effects may not have developed yet or already have dissipated by the time that an anticipated weather extreme occurs. We, therefore, stress the need to consider the temporal variability of OA impacts when comparing the performance of OAs in existing and future field studies and when designing OA application strategies aimed at improving soil resilience to specific climatic challenges.

## AUTHOR CONTRIBUTIONS

**Dirk-Jan D. Kok:** Conceptualization; data curation; formal analysis; investigation; methodology; visualization; writing—original draft; writing—review and editing. **Laura Scherer:** Conceptualization; supervision; writing—review and editing. **Wim de Vries:** Conceptualization; funding acquisition; project administration; supervision; writing—review and editing. **Peter M. van Bodegom:** Conceptualization; formal analysis; funding acquisition; project administration; supervision; writing—review and editing.

## ACKNOWLEDGMENTS

This work is part of the research project Social Innovation for Nutrient re-Cycling and Resource Efficiency (SIN-CERE) with project number NWO-GK-06, financed by the Netherlands Organization for Scientific Research (NWO) and co-financed by the Province of Gelderland (Provincie Gelderland), Vitens, Agricultural Landscape Collective Achterhoek (Vereniging Agrarisch Landschap Achterhoek), Foundation for Sustainable Agricultural Innovation Haarloseveld and Olden Eibergen (Stichting HOEduurzaam), the Directorate-General for Public Works and Water Management of the Netherlands (Rijkswaterstaat), Dutch Waste Management Association (DWMA), and the Netherlands Society for the Preservation of Nature Monuments (Natuurmonumenten). We thank also our fieldwork partners, Romke Postma, Dirk Thijssen and others involved at NMI Agro, as well as the field owners Jan Gerritsen and Alex ten Have. Finally, we would also like to acknowledge the valuable contribution of two anonymous reviewers to the improvement of this manuscript.

## CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

## DATA AVAILABILITY STATEMENT

The presented data has been made publicly accessible in the DANS-Easy data repository under (DANS: 10.17026/dans-znp-8pw2).

## ORCID

**Dirk-Jan Daniel Kok**  <https://orcid.org/0000-0003-0962-180X>

**Laura Scherer**  <https://orcid.org/0000-0002-0194-9942>

## REFERENCES

- Abiven, S., Menasseri, S., & Chenu, C. (2009). The effects of organic inputs over time on soil aggregate stability - A literature analysis. *Soil Biology and Biochemistry*, 41, 1–12. <https://doi.org/10.1016/j.soilbio.2008.09.015>
- Albiach, R., Canet, R., Pomares, F., & Ingelmo, F. (2001). Organic matter components and aggregate stability after the application of different amendments to a horticultural soil. *Bioresource Technology*, 76, 125–129. [https://doi.org/10.1016/S0960-8524\(00\)00090-0](https://doi.org/10.1016/S0960-8524(00)00090-0)
- Ali, G., Macrae, M., Walker, M., Laing, J., & Lobb, D. (2018). Preferential flow in vertisolic soils with and without organic amendments. *Agricultural & Environmental Letters*, 3, 180018. <https://doi.org/10.2134/aer2018.04.0018>
- Alletto, L., Pot, V., Giuliano, S., Costes, M., Perdrieux, F., & Justes, E. (2015). Temporal variation in soil physical properties improves the water dynamics modeling in a conventionally-tilled soil. *Geoderma*, 243–244, 18–28. <https://doi.org/10.1016/j.geoderma.2014.12.006>
- Andreini, M. S., & Steenhuis, T. S. (1990). Preferential paths of flow under conventional and conservation tillage. *Geoderma*, 46, 85–102.
- Ankenbauer, K. J., & Loheide, S. P. (2017). The effects of soil organic matter on soil water retention and plant water use in a meadow of the Sierra Nevada, CA. *Hydrological Processes*, 31, 891–901. <https://doi.org/10.1002/hyp.11070>
- Annabi, M., Le Bissonnais, Y., Le Villio-Poitrenaud, M., & Houot, S. (2011). Improvement of soil aggregate stability by repeated applications of organic amendments to a cultivated silty loam soil. *Agriculture Ecosystems and Environment*, 144, 382–389. <https://doi.org/10.1016/j.agee.2011.07.005>
- Arshad, M. A., Franzluebbers, A. J., & Azooz, R. H. (1999). Components of surface soil structure under conventional and no-tillage in northwestern Canada. *Soil and Tillage Research*, 53, 41–47. [https://doi.org/10.1016/S0167-1987\(99\)00075-6](https://doi.org/10.1016/S0167-1987(99)00075-6)
- Bai, Z., Dent, D. L., Olsson, L., & Schaepman, M. E. (2008). *Global assessment of land degradation and improvement: 1. Identification by remote sensing* (Report 2008/01). ISRIC-World Soil Information. [https://www.isric.org/sites/default/files/isric\\_report\\_2008\\_01.pdf](https://www.isric.org/sites/default/files/isric_report_2008_01.pdf)
- Barzegar, A. R., Yousefi, A., & Daryashenas, A. (2002). The effect of addition of different amounts and types of organic materials on soil physical properties and yield of wheat. *Plant and Soil*, 247, 295–301. <https://doi.org/10.1023/A:1021561628045>
- Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Bogaard, A., Fraser, R., Heaton, T. H., Wallace, M., Vaiglova, P., Charles, M., Jones, G., Evershed, R. P., Styring, A. K., Andersen, N. H., Arbogast, R.-M., Bartosiewicz, L., Gardeisen, A., Kanstrup, M., Maier, ... Stephan, E. (2013). Crop manuring and intensive land management by Europe's first farmers. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 12589–12594.
- Bronick, C. J., & Lal, R. (2005). Soil structure and management: A review. *Geoderma*, 124, 3–22. <https://doi.org/10.1016/j.geoderma.2004.03.005>
- Carron, M. P., Pierrat, M., Snoeck, D., Villenave, C., Ribeyre, F., Suhardi Marichal, R., & Caliman, J. P. (2015). Temporal variability in soil quality after organic residue application in mature oil palm plantations. *Soil Research*, 53, 205–215. <https://doi.org/10.1071/SR14249>
- Castellini, M., Diacono, M., Preite, A., & Montemurro, F. (2022). Short- and medium-term effects of on-farm compost addition on the physical and hydraulic properties of a clay soil. *Agronomy*, 12, 1446. <https://doi.org/10.3390/agronomy12061446>
- Celik, I., Gunal, H., Budak, M., & Akpinar, C. (2010). Effects of long-term organic and mineral fertilizers on bulk density and penetration resistance in semi-arid Mediterranean soil conditions. *Geoderma*, 160, 236–243. <https://doi.org/10.1016/j.geoderma.2010.09.028>
- Cercioglu, M. (2017). The role of organic soil amendments on soil physical properties and yield of maize (*Zea mays* L.). *Communications in Soil Science and Plant Analysis*, 48, 683–691. <https://doi.org/10.1080/00103624.2017.1298787>
- Chen, Y., Li, W., You, Y., Ye, C., Shu, X., Zhang, Q., & Zhang, K. (2022). Soil properties and substrate quality determine the priming of soil organic carbon during vegetation succession. *Plant and Soil*, 471, 559–575. <https://doi.org/10.1007/s11104-021-05241-z>
- Cornelis, W., Waweru, G., & Araya, T. (2019). Building resilience against drought and floods: The soil-water management perspective. In R. Lal, & R. Francaviglia (Eds.), *Sustainable agriculture reviews 29: Sustainable soil management: Preventive and ameliorative strategies* (pp. 125–142). Springer International Publishing. [https://doi.org/10.1007/978-3-030-26265-5\\_6](https://doi.org/10.1007/978-3-030-26265-5_6)
- Delibacak, S., Okur, B., & Ongun, A. R. (2009). Effects of treated sewage sludge levels on temporal variations of some soil properties of a Typic Xerofluvent soil in Menemen Plain, Western Anatolia, Turkey. *Environmental Monitoring and Assessment*, 148, 85–95. <https://doi.org/10.1007/s10661-007-0141-1>
- Diacono, M., & Montemurro, F. (2011). Long-term effects of organic amendments on soil fertility. In E. Lichtfouse, M. Hamelin, M. Navarrete, & P. Debaeke (Eds.), *Sustainable agriculture volume 2* (pp. 761–786). Springer. [https://doi.org/10.1007/978-94-007-0394-0\\_34](https://doi.org/10.1007/978-94-007-0394-0_34)
- Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., Dunn, R. J. H., Willett, K. M., Aguilar, E., Brunet, M., Caesar, J., Hewitson, B., Jack, C., Klein Tank, A. M. G., Kruger, A. C., Marengo, J., Peterson, T. C., Renom, M., Oria Rojas, C., Rusticucci, M., ... Kitching, S. (2013). Updated analyses of temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset. *Journal of Geophysical Research: Atmospheres*, 118, 2098–2118. <https://doi.org/10.1002/jgrd.50150>
- Eden, M., Gerke, H. H., & Houot, S. (2017). Organic waste recycling in agriculture and related effects on soil water retention and plant available water: A review. *Agronomy for Sustainable Development*, 37, 11. <https://doi.org/10.1007/s13593-017-0419-9>
- Elbersen, H. W., & Spijker, J. H. (2014). *Biomassapotentie Rijkswaterstaat: Analyse van hoeveelheden en huidige toepassing*. Wageningen UR - Food & Biobased Research.
- Eusufzai, M. K., & Fujii, K. (2012). Effect of organic matter amendment on hydraulic and pore characteristics of a clay loam soil. *Open Journal of Soil Science*, 02, 372–381. <https://doi.org/10.4236/ojss.2012.24044>
- Fares, A., Abbas, F., Ahmad, A., Deenik, J. L., & Safeeq, M. (2008). Response of selected soil physical and hydrologic properties to manure amendment rates, levels, and types. *Soil Science*, 173, 522–533. <https://doi.org/10.1097/SS.0b013e318182b063>
- Felton, G. K. (1995). Temporal variation of soil hydraulic properties on municipal solid waste amended mine soils. *Transactions of the ASAE*, 38, 775–782.
- Ferreras, L., Gomez, E., Toresani, S., Firpo, I., & Rotondo, R. (2006). Effect of organic amendments on some physical, chemical and

- biological properties in a horticultural soil. *Bioresource Technology*, 97, 635–640. <https://doi.org/10.1016/j.biortech.2005.03.018>
- García-Orenes, F., Guerrero, C., Mataix-Solera, J., Navarro-Pedreño, J., Gómez, I., & Mataix-Beneyto, J. (2005). Factors controlling the aggregate stability and bulk density in two different degraded soils amended with biosolids. *Soil and Tillage Research*, 82, 65–76. <https://doi.org/10.1016/j.still.2004.06.004>
- Geris, J., Verrot, L., Gao, L., Peng, X., Oyesiku-Blakemore, J., Smith, J. U., Hodson, M. E., McKenzie, B. M., Zhang, G., & Hallett, P. D. (2021). Importance of short-term temporal variability in soil physical properties for soil water modelling under different tillage practices. *Soil and Tillage Research*, 213, 105132. <https://doi.org/10.1016/j.still.2021.105132>
- Giorgi, F., Coppola, E., & Raffaele, F. (2014). A consistent picture of the hydroclimatic response to global warming from multiple indices: Models and observations. *Journal of Geophysical Research: Atmospheres*, 119, 11527–11891. <https://doi.org/10.1002/2014JD022238>
- Girei, A., Abdu, N., & Abdulkadir, A. (2016). Temporal variability of soil physico-chemical properties under a long-term fertilizer trial at Samaru, Northern Guinea Savanna of Nigeria. *International Journal of Plant & Soil Science*, 9, 1–10. <https://doi.org/10.9734/IJPSS/2016/22161>
- Guo, Z., Zhang, J., Fan, J., Yang, X., Yi, Y., Han, X., Wang, D., Zhu, P., & Peng, X. (2019). Does animal manure application improve soil aggregation? Insights from nine long-term fertilization experiments. *Science of the Total Environment*, 660, 1029–1037. <https://doi.org/10.1016/j.scitotenv.2019.01.051>
- Hamer, U., & Marschner, B. (2005). Priming effects in different soil types induced by fructose, alanine, oxalic acid and catechol additions. *Soil Biology & Biochemistry*, 37, 445–454. <https://doi.org/10.1016/j.soilbio.2004.07.037>
- Hu, W., Tabley, F., Beare, M., Tregurtha, C., Gillespie, R., Qiu, W., & Gosden, P. (2018). Short-term dynamics of soil physical properties as affected by compaction and tillage in a silt loam soil. *Vadose Zone Journal*, 17, 180115. <https://doi.org/10.2136/vzj2018.06.0115>
- Hueso-González, P., Ruiz-Sinoga, J. D., Martínez-Murillo, J. F., & Lavee, H. (2015). Overland flow generation mechanisms affected by topsoil treatment: Application to soil conservation. *Geomorphology*, 228, 796–804. <https://doi.org/10.1016/j.geomorph.2014.10.033>
- International Maize and Wheat Improvement Center. (2013). *Soil aggregate stability by wet sieving: A practical guide for comparing crop management practices*. CIMMYT.
- Intergovernmental Panel on Climate Change (IPCC), Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., & Zhou, B. (2021). *Climate change 2021: The physical science basis. (Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change)*. IPCC, Cambridge University Press.
- Jirků, V., Kodešová, R., Nikodem, A., Mühlhanslová, M., & Žigová, A. (2013). Temporal variability of structure and hydraulic properties of topsoil of three soil types. *Geoderma*, 204–205, 43–58. <https://doi.org/10.1016/j.geoderma.2013.03.024>
- Kögel-Knabner, I. (2017). The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter: Fourteen years on. *Soil Biology & Biochemistry*, 105, A3–A8. <https://doi.org/10.1016/j.soilbio.2016.08.011>
- Karami, A., Homae, M., Afzalnia, S., Ruhipour, H., & Basirat, S. (2012). Organic resource management: Impacts on soil aggregate stability and other soil physico-chemical properties. *Agriculture Ecosystems and Environment*, 148, 22–28. <https://doi.org/10.1016/j.agee.2011.10.021>
- Karlen, D. L., & Rice, C. W. (2015). Soil degradation: Will humankind ever learn? *Sustainability*, 7, 12490–12501. <https://doi.org/10.3390/su70912490>
- Khaleel, R., Reddy, K. R., & Overcash, M. R. (1981). Changes in soil physical properties due to organic waste applications: A review. *Journal of Environmental Quality*, 10, 133–141.
- Kinsbursky, R. S., Levanon, D., & Yaron, B. (1989). Role of fungi in stabilizing aggregates of sewage sludge amended soils. *Soil Science Society of America Journal*, 53, 1086–1091. <https://doi.org/10.2136/sssaj1989.03615995005300040017x>
- KNMI, The Royal Netherlands Meteorological Institute. (2022). *Klimaatviewer*. <https://www.knmi.nl/klimaat-viewer/kaarten/neerslag-verdamping>
- Kok, D. D., Scherer, L., de Vries, W., Trimbos, K., & van Bodegom, P. M. (2022). Relationships of priming effects with organic amendment composition and soil microbial properties. *Geoderma*, 422, 115951. <https://doi.org/10.1016/j.geoderma.2022.115951>
- Lal, R. (2012). Climate change and soil degradation mitigation by sustainable management of soils and other natural resources. *Agricultural Research*, 1, 199–212. <https://doi.org/10.1007/s40003-012-0031-9>
- Lal, R. (2020). Soil organic matter and water retention. *Agronomy Journal*, 112, 3265–3277. <https://doi.org/10.1002/agj2.20282>
- Larney, F. J., & Angers, D. A. (2012). The role of organic amendments in soil reclamation: A review. *Canadian Journal of Soil Science*, <https://doi.org/10.4141/CJSS2010-064>
- Lenth, R. (2020). *emmeans: Estimated marginal means, aka least-squares means*. R package version 1.5.2-1. <https://mran.microsoft.com/snapshot/2020-11-10/web/packages/emmeans/index.html>
- Li, Z., Schneider, R. L., Morreale, S. J., Xie, Y., Li, C., & Li, J. (2018). Woody organic amendments for retaining soil water, improving soil properties and enhancing plant growth in desertified soils of Ningxia, China. *Geoderma*, 310, 143–152. <https://doi.org/10.1016/j.geoderma.2017.09.009>
- Lloyd, D. A., Ritz, K., Paterson, E., & Kirk, G. J. D. (2016). Effects of soil type and composition of rhizodeposits on rhizosphere priming phenomena. *Soil Biology & Biochemistry*, 103, 512–521. <https://doi.org/10.1016/j.soilbio.2016.10.002>
- Luna, L., Vignozzi, N., Miralles, I., & Solé-Benet, A. (2018). Organic amendments and mulches modify soil porosity and infiltration in semiarid mine soils. *Land Degradation and Development*, 29, 1019–1030. <https://doi.org/10.1002/ldr.2830>
- Lutzow, M. v., Kogel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., & Flessa, H. (2006). Stabilization of organic matter in temperate soils: Mechanisms and their relevance under different soil conditions - a review. *European Journal of Soil Science*, 57, 426–445. <https://doi.org/10.1111/j.1365-2389.2006.00809.x>
- Manzoni, S., Trofymow, J. A., Jackson, R. B., & Porporato, A. (2010). Stoichiometric controls on carbon, nitrogen, and phosphorus dynamics in decomposing litter. *Ecological Monographs*, 80, 89–106. <https://doi.org/10.1890/09-0179.1>
- Meena, R. S., Lal, R., & Yadav, G. S. (2020). Long-term impacts of topsoil depth and amendments on soil physical and hydrological

- properties of an Alfisol in central Ohio, USA. *Geoderma*, 363, 114164. <https://doi.org/10.1016/j.geoderma.2019.114164>
- Metzger, L., Levanon, D., & Mingelgrin, U. (1987). The effect of sewage sludge on soil structural stability: Microbiological aspects. *Soil Science Society of America Journal*, 51, 346–351. <https://doi.org/10.2136/sssaj1987.03615995005100020016x>
- Minasny, B., & McBratney, A. B. (2018). Limited effect of organic matter on soil available water capacity. *European Journal of Soil Science*, 69, 39–47. <https://doi.org/10.1111/ejss.12475>
- Montgomery, D. R. (2007). Soil erosion and agricultural sustainability. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 13268–13272. <https://doi.org/10.1073/pnas.0611508104>
- Mooshammer, M., Hofhansl, F., Frank, A. H., Wanek, W., Hämmerle, I., Leitner, S., Schneckner, J., Wild, B., Watzka, M., Keiblinger, K. M., Zechmeister-Boltenstern, S., & Richter, A. (2017). Decoupling of microbial carbon, nitrogen, and phosphorus cycling in response to extreme temperature events. *Science Advances*, 3, e1602781. <https://doi.org/10.1126/sciadv.1602781>
- Negiş, H., Şeker, C., Gümüş, I., Manirakiza, N., & Mücevher, O. (2020). Effects of biochar and compost applications on penetration resistance and physical quality of a sandy clay loam soil. *Communications in Soil Science and Plant Analysis*, 51, 38–44. <https://doi.org/10.1080/00103624.2019.1695819>
- Oades, J. M. (1984). Soil organic matter and structural stability: Mechanisms and implications for management. *Plant and Soil*, 76, 319–337. <https://doi.org/10.1007/BF02205590>
- Oades, J. M. (1993). The role of biology in the formation, stabilization and degradation of soil structure. In L. Brussaard, & M. J. Kooistra (Eds.), *Soil structure/soil biota interrelationships* (pp. 377–400). Elsevier. <https://doi.org/10.1016/B978-0-444-81490-6.50033-9>
- Ouyang, L., Wang, F., Tang, J., Yu, L., & Zhang, R. (2013). Effects of biochar amendment on soil aggregates and hydraulic properties. *Journal of Soil Science and Plant Nutrition*, 13, 991–1002. <https://doi.org/10.4067/S0718-95162013005000078>
- Pagliai, M., Vignozzi, N., & Pellegrini, S. (2004). Soil structure and the effect of management practices. *Soil and Tillage Research*, 79, 131–143. <https://doi.org/10.1016/j.still.2004.07.002>
- Pandey, C., & Shukla, S. (2006). Effects of soil organic amendment on water and nutrient movement in a sandy soil. *American Society of Agricultural and Biological Engineers*, 2006, 062007. <https://doi.org/10.13031/2013.20667>
- Paré, T., Dinel, H., Moulin, A. P., & Townley-Smith, L. (1999). Organic matter quality and structural stability of a black Chernozemic soil under different manure and tillage practices. *Geoderma*, 91, 311–326. [https://doi.org/10.1016/S0016-7061\(99\)00011-7](https://doi.org/10.1016/S0016-7061(99)00011-7)
- Pires, L. F., Borges, J. A. R., Rosa, J. A., Cooper, M., Heck, R. J., Passoni, S., & Roque, W. L. (2017). Soil structure changes induced by tillage systems. *Soil and Tillage Research*, 165, 66–79. <https://doi.org/10.1016/j.still.2016.07.010>
- Popolizio, S., Stellacci, A. M., Giglio, L., Barca, E., Spagnuolo, M., & Castellini, M. (2022). Seasonal and soil use dependent variability of physical and hydraulic properties: An assessment under minimum tillage and no-tillage in a long-term experiment in southern Italy. *Agronomy*, 12, 3142. <https://doi.org/10.3390/agronomy12123142>
- Rawls, W. J., Pachepsky, Y. A., Ritchie, J. C., Sobecki, T. M., & Bloodworth, H. (2003). Effect of soil organic carbon on soil water retention. *Geoderma*, 116, 61–76. [https://doi.org/10.1016/S0016-7061\(03\)00094-6](https://doi.org/10.1016/S0016-7061(03)00094-6)
- Rayne, N., & Aula, L. (2020). Livestock manure and the impacts on soil health: A review. *Soil Systems*, 4, 64. <https://doi.org/10.3390/soilsystems4040064>
- Reynolds, W. D., Drury, C. F., Tan, C. S., Fox, C. A., & Yang, X. M. (2009). Use of indicators and pore volume-function characteristics to quantify soil physical quality. *Geoderma*, 152, 252–263. <https://doi.org/10.1016/j.geoderma.2009.06.009>
- Rivier, P.-A., Jamniczky, D., Nemes, A., Makó, A., Barna, G., Uzing, N., Rékási, M., & Farkas, C. (2022). Short-term effects of compost amendments to soil on soil structure, hydraulic properties, and water regime. *Journal of Hydrology and Hydromechanics*, 70, 74–88. <https://doi.org/10.2478/johh-2022-0004>
- Roldán, A., Albaladejo, J., & Thornes, J. (1996). Aggregate stability changes in a semiarid soil after treatment with different organic amendments. *Arid Land Research and Management*, 10, 139–148.
- Ros, G. H., Termorshuizen, A. J., & van Dijk, T. A. (2012). *Risico's van diffuse verspreiding van groenafvalstromen*. <https://www.nmi-agro.nl/wp-content/uploads/2020/08/Rapport-1474-def.pdf>
- RStudio Team. (2020). *RStudio: Integrated development for R*.
- RVO. (2022). *Tabellen mest rijksdienst voor ondernemend nederland*. <https://www.rvo.nl/onderwerpen/mest/tabellen>
- Schoonvelde, H., van Middelkoop, J. C., Philipsen, A. P., van Dongen, C., Bussink, D. W., Bos, A. J., Velthof, G. L., de Haan, J. J., Schröder, J. J., Reijneveld, J. A., & Eekeren, N. V., CBGV. (2020). *Bemestingsadvies commissie bemesting grasland en voedergrassen*. Commissie Bemesting Grasland en Voedergrassen.
- Scott, H. D., Mauromoustakos, A., Handayani, I. P., & Miller, D. M. (1994). Temporal Variability of Selected Properties of Loessial Soil as Affected by Cropping. *Soil Science Society of America Journal*, 58, 1531–1538. <https://doi.org/10.2136/sssaj1994.03615995005800050037x>
- Sharma, B., Vaish, B., Monika Singh, U. K., Singh, P., & Singh, R. P. (2019). Recycling of organic wastes in agriculture: An environmental perspective. *International Journal of Environmental Research*, 13, 409–429. <https://doi.org/10.1007/s41742-019-00175-y>
- Shipitalo, M. J., Dick, W. A., & Edwards, W. M. (2000). Conservation tillage and macropore factors that affect water movement and the fate of chemicals. *Soil and Tillage Research*, 53, 167–183. [https://doi.org/10.1016/S0167-1987\(99\)00104-X](https://doi.org/10.1016/S0167-1987(99)00104-X)
- Singh Brar, B., Singh, J., Singh, G., & Kaur, G. (2015). Effects of long term application of inorganic and organic fertilizers on soil organic carbon and physical properties in maize–wheat rotation. *Agronomy*, 5, 220–238. <https://doi.org/10.3390/agronomy5020220>
- Six, J., Bossuyt, H., Degryze, S., & Denef, K. (2004). A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research*, 79, 7–31. <https://doi.org/10.1016/J.STILL.2004.03.008>
- Tang, C., Yang, F., & Antonietti, M. (2022). Carbon materials advancing microorganisms in driving soil organic carbon regulation. *Research*, 2022, 9857374.
- Tejada, M., & Gonzalez, J. L. (2008). Influence of two organic amendments on the soil physical properties, soil losses, sediments and runoff water quality. *Geoderma*, 145, 325–334. <https://doi.org/10.1016/j.geoderma.2008.03.020>
- Terleev, V., Ginevsky, R., Lazarev, V., Nikonov, A., Togo, I., Topaj, A., Moiseev, K., Abakumov, E., Melnichuk, A., & Dunaieva, I. (2017). Predicting the scanning branches of hysteretic soil water-retention capacity with use of the method of mathematical modeling. *IOP*

- Conference Series: Earth and Environmental Science*, 90, 012105. <https://doi.org/10.1088/1755-1315/90/1/012105>
- Tisdall, J. M., & Oades, J. M. (1982). Organic matter and water-stable aggregates in soils. *European Journal of Soil Science*, 33, 141–163. <https://doi.org/10.1111/j.1365-2389.1982.tb01755.x>
- Valentine, T. A., Hallett, P. D., Binnie, K., Young, M. W., Squire, G. R., Hawes, C., & Bengough, A. G. (2012). Soil strength and macropore volume limit root elongation rates in many UK agricultural soils. *Annals of botany*, 110, 259–270. <https://doi.org/10.1093/aob/mcs118>
- van Es, H. M., Ogden, C. B., Hill, R. L., Schindelfbeck, R. R., & Tsegaye, T. (1999). Integrated assessment of space, time, and management-related variability of soil hydraulic properties. *Soil Science Society of America Journal*, 63, 1599–1608. <https://doi.org/10.2136/sssaj1999.6361599x>
- Wanniarachchi, D., Cheema, M., Thomas, R., Kavanagh, V., & Galagedara, L. (2019). Impact of soil amendments on the hydraulic conductivity of boreal agricultural Podzols. *Agriculture*, 9, 133. <https://doi.org/10.3390/agriculture9060133>
- Ward, B. B., Capone, D. G., & Zehr, J. P. (2007). What's new in the nitrogen cycle? *Oceanography*, 20, 101–109.
- Whalen, J. K., & Chang, C. (2002). Macroaggregate characteristics in cultivated soils after 25 annual manure applications. *Soil Science Society of America Journal*, 66, 1637–1647. <https://doi.org/10.2136/sssaj2002.1637>
- Wiecheteck, L. H., Giarola, N. F. B., de Lima, R. P., Tormena, C. A., Torres, L. C., & de Paula, A. L. (2020). Comparing the classical permanent wilting point concept of soil (−15,000 hPa) to biological wilting of wheat and barley plants under contrasting soil textures. *Agricultural Water Management*, 230, 105965. <https://doi.org/10.1016/j.agwat.2019.105965>
- Yüksel, O., & Kavdır, Y. (2020). Improvement of soil quality parameters by municipal solid waste compost application in clay-loam soil. *Turkish Journal of Agriculture - Food Science and Technology*, 8, 603–609. <https://doi.org/10.24925/turjaf.v8i3.603-609.3062>
- Yu, G.-H., Chen, C.-M., He, X.-H., Zhang, X.-Z., & Li, L.-N. (2020). Unexpected bulk density and microstructures response to long-term pig manure application in a ferralic Cambisol soil: Implications for rebuilding a healthy soil. *Soil and Tillage Research*, 203, 104668. <https://doi.org/10.1016/j.still.2020.104668>
- Zamani, J., Afyuni, M., Sepehrnia, N., & Schulin, R. (2016). Opposite effects of two organic wastes on the physical quality of an agricultural soil. *Archives of Agronomy and Soil Science*, 62, 413–427. <https://doi.org/10.1080/03650340.2015.1060321>
- Zare, M., Abbaspour, M., & Aafyuni, K. C. (2010). Effects of biosolids application on temporal variations in soil physical and unsaturated hydraulic properties. *Journal of Residuals Science & Technology*, 7, 227–235.
- Zhang, S., Yang, X., Wiss, M., Grip, H., & Lövdahl, L. (2006). Changes in physical properties of a loess soil in China following two long-term fertilization regimes. *Geoderma*, 136, 579–587. <https://doi.org/10.1016/j.geoderma.2006.04.015>
- Zhao, Y., Wang, P., Li, J., Chen, Y., Ying, X., & Liu, S. (2009). The effects of two organic manures on soil properties and crop yields on a temperate calcareous soil under a wheat–maize cropping system. *European Journal of Agronomy*, 31, 36–42. <https://doi.org/10.1016/j.eja.2009.03.001>
- Zhou, H., Chen, C., Wang, D., Arthur, E., Zhang, Z., Guo, Z., Peng, X., & Mooney, S. J. (2020). Effect of long-term organic amendments on the full-range soil water retention characteristics of a Vertisol. *Soil and Tillage Research*, 202, 104663. <https://doi.org/10.1016/j.still.2020.104663>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Kok, D. D., Scherer, L., de Vries, W., & van Bodegom, P. M. (2023). Temporal variability in organic amendment impacts on hydro-physical properties of sandy agricultural soils. *Soil Science Society of America Journal*, 87, 963–984. <https://doi.org/10.1002/saj2.20547>