



Universiteit
Leiden
The Netherlands

Developmental effects of polystyrene nanoparticles in the chicken embryo

Wang, M.

Citation

Wang, M. (2024, January 16). *Developmental effects of polystyrene nanoparticles in the chicken embryo*. Retrieved from <https://hdl.handle.net/1887/3704678>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3704678>

Note: To cite this publication please use the final published version (if applicable).

Chapter 1. General introduction

Problem statement

A new category of potentially harmful substances is attracting the attention of researchers. These substances are so-called nanomaterials, which are various substances in particulate form, and with a predominant particle size on the nanoscale. One group of nanomaterials are the nanoplastics (NPs). These are either intentionally manufactured for example for use as research reagents; or arise naturally in the environment from the breakdown of plastic waste (Hernandez et al., 2017; Sangkham et al., 2022). The breakdown of plastics first releases particles called microplastics (MPs) which are in the size range ≤ 5 mm (Ray et al., 2022) and these in turn can break down into smaller fragments including NPs.

NPs are widely present in the environment as pollutants in water bodies, soil or the air, and this provides a potential route for wildlife and humans (Sangkham et al., 2022). NPs are being considered as potential drug-delivery agents in the field of nanomedicine (Boehnke et al., 2022). This could open up an additional route through which humans could be exposed. The toxicity of NPs has mainly been studied using aquatic invertebrates and fish model species. In this thesis, I will use a warm-blooded vertebrate model, the chicken embryo, to investigate NP toxicity.

The origins and fate of nanoplastics

Origins of nanoplastics

Nanoplastics are small fragments of plastic — that is, polymer-based materials synthesized from petroleum products (Desai and Galage, 2015; Geyer et al., 2017). Plastics have a broad range of applications in manufacturing, the electronics, packaging and food industries, and in agriculture and medicine and surgery (Gourmelon, 2015). This is because they are durable, cheap, water-resistant, have

relatively low energy requirements for their manufacture, and are relatively light in weight (Gourmelon, 2015; Panda et al., 2010). Moreover, plastics can be significantly altered by heating (Becker and Locascio, 2002; Levine and Berman, 1995), allowing them to be molded into required shapes. Since the 1950s, the annual global growth rate in plastics production has remained at an average of 8.5% (Platt, 2003). By 2018, global plastics production reached 359 million tonnes. According to some predictions, this value will have doubled by 2025 (Gibb, 2019). This thesis is concerned with a particular form of plastics called small plastic particles. In a sense, plastics, and small plastic particles, are typical representative 'fossils' of human activity (Ng et al., 2018), that is to say, they are enduring artefacts in the environment that are **anthropogenic** (uniquely created by human activity).

Classification of small plastic particles

Small plastic particles include microplastics (MPs) whose particle size is ≤ 5 mm (Arthur et al., 2008); and nanoplastics (NPs), whose particle size is either ≤ 100 nm (Zhang and Webster, 2009) or ≤ 1000 nm (Gigault et al., 2018) depending on the opinion of the author.

Other than classifying them by their size, small plastic particles can be classified according to their origins. Thus, MPs and NPs can be classified as being primary or secondary (Ziajahromi et al., 2017). **Primary** MPs and NPs are small plastic particles manufactured for a specific purpose, in the case of MPs: personal-use products including toothpaste, cosmetic, shampoo (Hernandez et al., 2017); and in the case of NPs as potential drug-delivery vehicles in the emerging field of nanomedicine (Galafassi et al., 2019; Patel et al., 2009). The potential route of administration of NPs to human patients might include intravenous injection or inhalation (Chai et al., 2020).

By contrast, **secondary** MPs and NPs are fragments coming from larger pieces of plastic such as plastic bags, bottles and fishing nets (Boucher and Friot, 2017). They are generated through the natural processes of physical, biological, and chemical

degradation in the environment, for example, the action of UV light, ocean waves, and wind abrasion (Stevens, 2015). Another source of MPs and NPs is the microfibers generated by laundering of synthetic textiles (Center and Wash, 2017), and the accidental discarding of plastic products during manufacturing or transport (Rezania et al., 2018). Surprisingly, even plastic teabags can release nanoplastics (Hernandez et al., 2019).

Some cosmetic products contain both primary and secondary small plastic particles. One study (Hernandez et al., 2017) analyzed commercially-available facial scrubs, and found 'microbeads' of around 200 μm diameter which were ingredients added intentionally by the manufacturer — along with nanoplastic particles of 24 ± 6 to 52 ± 14 nm. It is not clear why nanoplastics were present, because they were not part of the product formulation. It is possible that they arose from the breakdown of the larger microbeads.

Small plastic particles are also distinguished by their composition; for example they may be made of polystyrene (PS), polypropylene, polyethylene, low-density polyethylene, polyacrylates, polyvinylchloride, polyamide, polyethylene terephthalate and polyvinyl alcohol (Anderson et al., 2016; Avio et al., 2017). Finally, MPs and NPs can be categorized according to their shapes which include: beads or spheres, fibers, fragments, film, pellets and foam (Burns and Boxall, 2018).

Plastic waste as a source of small plastic particles

We mentioned above that secondary plastic particles are released from manufactured plastics when they break down (Stevens, 2015; Zhang et al., 2021). For example, at their end of usable life, plastics end up forming a large component of domestic garbage and industrial waste (Subramanian, 2000). The great majority of this plastic waste is dumped into land-fills, while some is burned and a smaller amount is recycled (Geyer et al., 2017). Dumped plastic waste can be found in freshwater bodies, oceans, soil, air, and even in remote, uninhabited islands (Bergmann et al., 2019). For these and other reasons, plastic pollution to our

environment is a matter for great concern (Chauhan and Wani, 2019). Plastic waste can persist for years in the environment due to the low speed at which plastics undergo degradation (Webb et al., 2013). One reason so much plastic is dumped rather than being recycled is that recycling technology is not currently optimal. For example, it is common practice to collect all types of recyclable plastic into one mixed batch, making it difficult to produce a high-quality recycled product (Geyer et al., 2017). Furthermore, plastics can only be recycled twice (Geyer et al., 2017). Because of the size of nanoplastics it is not practical to remove them from water by filtration, nor is it practical to try to sort them into different plastic types for recycling (Nguyen et al., 2019). In any case, no matter how plastic wastes are disposed, they can potentially break down and release small plastic particles into the natural environment.

Distribution of MPs and NPs in the environment

After the breakdown of waste plastic, and its release of small plastic particles into the natural environment, MPs and NPs become widely distributed in freshwater bodies, sediments, air and the sea, from the equator to the polar regions (Browne et al., 2011; Wan et al., 2019a). Furthermore, MPs and NPs persist in aquatic systems because the waste-water technology to deal with these kinds of materials is lacking. Even though membrane bioreactors have been created by scientists to remove the particles, the high cost of implementation results in feasibility problems for large-scale, public application (Westphalen and Abdelrasoul, 2017). As a result, a large amount of MPs and NPs enter the ecosystem (Browne et al., 2007; Fendall and Sewell, 2009).

Trophic transfer of MPs and NPs

MPs and NPs are transferred between species that are part of food chains (Shruti and Kutralam-Muniasamy, 2019; Vom Sqaal et al., 2008). This is called trophic transfer, and is one of the most important uptake routes of MPs and NPs (Toussaint et al., 2019). MPs and NPs can be taken up by wildlife passively or actively (Vom Sqaal et al.,

2008). For instance, fish can take up MPs and NPs while feeding or drinking (passively), or when foraging by accidental ingestion (actively; (Roch et al., 2020).

Animals tend to accidentally ingest plastic items in their food (de Sá et al., 2018). In Gall & Thompson's (2015) report, large pieces of plastic have been found in the digestive tract of 208 species of marine organisms worldwide including 86% (6/7) sea turtle species, 39% (122/312) of seabird species, 0.3% (50/16,754) fish species and 26% (30/115) marine mammals (Gall and Thompson, 2015). Such large pieces of plastic (plastic bottles, plastic bags etc.) can cause adverse effect to wild animals, resulting in starvation and even death (Nkwachukwu et al., 2013). In addition, animals (including sea turtles, seals and sharks) can become entangled in plastic fishing nets and fishing lines and this may impair the movement of the animals, and even cause asphyxia (Butterworth et al., 2012).

MPs and NPs are difficult to detect in terrestrial environments owing to the complex composition of soils. Fortunately, analytical methods for measuring MPs in soils have been developed (He et al., 2018). Large amounts of MPs and NPs have been found in agricultural soils and in industrial regions (Galloway et al., 2017; Zhang and Liu, 2018). The accumulation of MPs and NPs in the soil cause negative effects through altering its biophysical properties (Navarro et al., 2008; Wan et al., 2019b).

Toxicity of small plastic particles to living organisms

MPs and NPs pose a number of health threats to wildlife and humans. For example, once they have entered the natural environment, they can adsorb further toxic chemicals such as polycyclic aromatic hydrocarbons (PAHs) (Maes et al., 2017), polychlorinated biphenyls (PCBs) (Velzeboer et al., 2014) and dichlorodiphenyltrichloroethane (DDT) (Bakir et al., 2014), which can potentiate any intrinsic toxicity they may have (Souza et al., 2022; Ziccardi et al., 2016).

Any of these toxins may later get leached out and released into the environment when the plastic is degraded into particles (Burns and Boxall, 2018; de Souza Machado et al., 2018; Horton et al., 2017; Smith et al., 2018). Adsorption and leaching, respectively, are influenced by internal factors, namely: particle composition, size and shape, and external factors such as temperature, pressure and pH in the environment (Bakir et al., 2014; Frere et al., 2017). In this context, it is important to note that NPs have a very high surface-area-to-volume ratio.

Like all plastics, NPs may contain toxic plasticizers and pigments from manufacture (Unar et al., 2010). Toxic chemicals used in the manufacture of some plastics include nonylphenol, bisphenol A, and vinyl chloride, and these are able to leach from the plastics when they degrade in the environment (Gallo et al., 2018; Koelmans et al., 2014). Also, MPs and NPs can affect the food chain by transferring biocide (pesticide and herbicide) residues in agro-ecosystems (Ng et al., 2018). Other pollutants such as heavy metals and organic chemicals can also be adsorbed to the surface of the MPs and NPs (Zeng, 2018). In summary, MPs and NPs are widely distributed in biotic and abiotic matrices (Karlsson et al., 2017), and can show enhanced toxicity due to physical and chemical changes.

Toxic effects of MPs and NPs in experimental animals

The biological effects of MPs and NPs have been widely studied in aquatic organisms including crustaceans such as *Daphnia* sp. (Brun et al., 2017), and teleost fish such as the zebrafish (*Danio rerio*) (Bashirova et al., 2023), its adults or embryos/larvae (Veneman et al., 2017), reviewed by (Jiang et al., 2020). When MPs or NPs enter into the body of animals, they can accumulate and cause significant effects. These effects may be on **higher levels of the biological hierarchy**, such as changes to the populations, communities and ecosystems of animals (Eerkes-Medrano and Thompson, 2018; Galloway et al., 2017). Pollution by MPs and NPs can affect a community if a an invasive species contaminated by MPs and NPs is introduced; this

can result in alterations to the composition of species in ecosystem (Gall and Thompson, 2015; Lusher, 2015).

Effects of MPs and NPs on **lower levels in the biological hierarchy**, include deleterious effects at the molecular, cellular and organismal levels such as reproductive and development toxicity (Wang et al., 2019; Zhang et al., 2019). Such organismal effects have been seen in the common goby (*Pomatoschistus microps*) (Ferreira et al., 2016), the Pacific mole crab (*Emerita analoga*) (Horn et al., 2020) and the water flea (*Daphnia magna*) (Aljaibachi and Callaghan, 2018). Furthermore, polyethylene MPs accumulated in gill and intestine of the adult zebrafish can lead to abnormal behavior and movements, including seizures (Brun et al., 2019; Mak et al., 2019). More seriously, it has been reported that 40 nm and 200 nm polystyrene NPs can increase the risk of cardiovascular disease by disrupting the cellular components and extracellular matrix in human induced pluripotent cells (Bojic et al., 2020; Prata, 2018).

In the rainbow trout (*Oncorhynchus mykiss*), PS-Pd NPs (Palladium-doped polystyrene nanoplastics) can adhere to the gut epithelium, and can be taken up into the liver, kidney, gills and muscles (Clark et al., 2023). In another study it was found that NPs of 51 nm can be transferred into the gallbladder, liver, pancreas, heart and brain of zebrafish larvae (Pitt et al., 2018). Moreover, it has been found that MPs and NPs can cause inflammation in the liver, heart, lungs, kidney and brain in the Medaka fish (*Oryzias latipes*) (Kashiwada, 2006).

It is not clear whether they then enter the blood stream or the lymphatic system, both of which drain the gut. It is possible that MPs and NPs might be cleared from the circulation by the spleen, and some might be secreted in the urine reviewed by (Bouwmeester et al., 2015; Li et al., 2022; Zhu et al., 2022). Zhu *et al.* (2020) also found that MPs of 10 µm diameter accumulated in the gills and gut of the medaka (*Oryzias latipes*) (Zhu et al., 2020). The distribution of the NPs *in vivo* appeared to be influenced the size of the NPs. Thus, when zebrafish were exposed to 70 nm, 5 µm,

and 20 µm diameter polystyrene particles, the 5 µm particles could be found in the gills, gut, and liver whereas the 70 nm and 20 µm particles were not taken up (Lu et al., 2016). In this kind of study, it is important to realize that, with increased particle size, the number of particles decreases, for any given concentration of the plastic in the carrier vehicle (Wang et al., 2023).

A recent study in zebrafish larvae suggests that fluorescein-tagged 20 nm polystyrene NPs are able to cross the blood-brain barrier, and are subsequently accumulated in the brain itself, of the zebrafish larvae (Sökmen et al., 2020). The blood-brain barrier appears to develop in the normal zebrafish sometime between 2- and 10-days post fertilization, depending on the molecular size of the marker studied (Quiñonez-Silvero et al., 2020). The uptake, accumulation and transportation of MPs and NPs can have deleterious effects on the immune system, nervous system, and on metabolism in mammals (reviewed by (Yong et al., 2020). Another zebrafish study found that 5 µm polystyrene particles can induce inflammation and oxidative stress in the gut and changes in the metabolome and microbiome of the gut (Qiao et al., 2019).

Potential risks of MPs and NPs to humans

Relatively little has been published about the biological effects of small plastic particles on humans, not least because of the ethical difficulties in doing research on human subjects. Among their possible toxic effects on humans, are those affecting the embryo or fetus (reviewed by (Hougaard et al., 2015). Little is known about the possibility of placental transfer in any mammalian species, but one study used the BeWo cell line (which is used to model placental cells), and found that they could take up 50 nm polystyrene NPs (Dusza et al., 2022).

In addition, Ragusa *et al.* (2021) found a few fragments of MPs in human placentas including on the maternal side, the fetal side and in the chorioamniotic membrane (Ragusa et al., 2021). As a result, it has been speculated that MPs and NPs might be able to cause decreased birth weight, autoimmune lung disease, and a series of central nervous system abnormalities (Bates, 2019). Another recent study using the

perfused human placenta found that the NPs affect the gene expression related to inflammation and iron homeostasis (Chortarea et al., 2023).

MPs and NPs potentially cause effects on humans *via* various other exposure routes (Galloway, 2015; Powell et al., 2007). For example, MPs and NPs can be transferred to humans from food species (Chang et al., 2020). Thus, they have been widely found in fish, shellfish and shrimps (Garrido Gamarro et al., 2020; Li et al., 2015; Smith et al., 2018); in honey, salt and sugar (Liebezeit and Liebezeit, 2013; Yang et al., 2015); and in food and beverage packages such as teabags and soup cups (Du et al., 2020; Hernandez et al., 2019). Cox *et al.* revealed that people eat at least 50,000 MPs particles per year on average (Cox et al., 2019). Hence, there is at least the potential for humans to suffer harmful effects from exposure to MPs and NPs.

Even though the skin functions as a biological barrier, it is still another potential route of human exposure (Prata *et al.*, 2020) because MPs and NPs are commonly contained in personal products such as sun cream, toothpaste and shower gel (Sharma & Chatterjee, 2017) in many countries (except the EU, Canada and USA, where personal care products containing such plastic 'microbeads' are banned (Hernandez et al., 2017). It has been shown that other classes of nanomaterials, namely the so-called 'quantum dot' nanoparticles, can penetrate the skin of mice (Mortensen et al., 2008). There is evidence that 40 nm polystyrene NPs can enter through hair follicles in human skin (Vogt et al., 2006). Both 40 nm and 200 nm polystyrene NPs were found to be able to penetrate into mouse skin (Mahe et al., 2009). In addition, poly(l-lactide-co-glycolide) nanoparticles of 70 nm and 300 nm diameter were found to accumulate more, and penetrate deeper, in inflamed skin compared to healthy skin, in mouse and pig models (Try et al., 2016). Moreover, the degree of penetration of NPs into the skin was influenced by their size (Try et al., 2016).

The penetration and accumulation of NPs in skin can, in principle, result in the exposure of epidermal and dermal cells to the NPs. However, the epidermis does not

contain blood vessels or lymphatic vessels (Lund et al., 2016). Therefore, in order for the NPs to spread to other parts of the body, from the skin, they will have to cross the epidermal basement membrane and enter the blood vessels or lymphatic vessels in the dermis. Furthermore, it has been shown that exposure to NPs can lead to oxidative stress in human cerebral and epithelial cells *in vitro* (Schirinzi et al., 2017).

While it is difficult or impossible to perform the necessary experiments in humans with MPs and NPs, we can get insight from other warm-blooded vertebrates such as rats and birds (De Jong et al., 2008; Wang et al., 2023). We can also get indirect evidence about plastic nanoparticles from experiments with metal or other sorts of nanoparticles (De Jong et al., 2008; Yan et al., 2020; Yan et al., 2021), although great care should be used in extrapolating from one nanomaterial to the other. In summary, there is growing evidence that MPs and NPs might be a potential threat to human health.

The potential advantages of using chicken embryos in nanoplastic toxicity research

The aim of this thesis was to use a model closer to humans than the fish and invertebrates that have often been used for testing the toxicity of MPs and NPs. This model is the embryo of the chicken (*Gallus gallus*). Birds are warm-blooded, 'higher' vertebrates with a physiology quite similar to humans (Pozio, 2005). Birds share a most recent common ancestor with humans that lived approximately 319 million years ago (Pardo et al., 2020; Rezanian et al., 2018; Sánchez-Villagra, 2012; St John et al., 2012). By contrast, the most recent common ancestor of humans and the zebrafish (*Danio rerio*) lived 450 million years ago (Hedges and Kumar, 2009). Furthermore, the chicken embryo can be directly exposed experimentally to NPs *in ovo*, and then the egg returned to the incubator (Tickle, 1993). Such experiments are

not possible to perform in mammalian model species because of the placental barrier.

Other reasons for using the chicken embryo (often called the chick embryo) are that there is a genome sequenced and the chicken embryo has been for decades an important model in developmental and toxicology studies (Bryda, 2013; Davey and Tickle, 2007; Hamburger and Hamilton, 1951; Korhonen et al., 1982; Rashidi and Sottile, 2009; Romanoff, 1960). Another argument in favor of using the chick embryo model is that the chicken is a bird, and there is a need to understand the effects of pollutants on wild birds, as shown by Rachel Carson in *Silent Spring* (Carson, 1962).

Before hatching, the chicken embryo is not subject to the European Laws on animal welfare licensing (European Union (EU) directive no. 2010/63/EU). At Leiden University, we have chosen to use the embryos no further than day 14, which is $\frac{2}{3}$ of the incubation period. Additionally, the chicken embryo is a relatively cost-effective model, and at the time of writing, in our laboratory, we buy fertilized chicken eggs costing approximately €1.00 each (including transport costs) from a commercial hatchery.

A standard series of embryonic stages has been described for the chicken (Hamburger and Hamilton, 1951) from laying to hatching (21 d). These stages allow for the standardization of research between laboratories. Duman *et al.* found that the chicken embryo model can be used to mimic human tissues and can be considered as a platform for the study of teratogen-induced malformations (Duman et al., 2019). For example, exposure of the human embryo to valproic acid (a teratogenic drug used to treat epilepsy) can harm the developing embryo, producing fetal valproate syndrome (Ornoy, 2009). The same exposure has also been shown to be teratogenic in chicken embryos, producing a pattern of defects similar to those in humans (Nanau and Neuman, 2013; Tanoshima et al., 2015). Thus, the chicken embryo is likely to act as a practical model with relevance to humans.

The chicken embryo model has been used for examining the effects of nanomaterials in a small number of studies (reviewed by (Ghimire et al., 2022)). These include studies of nanoparticles made out of polystyrene (Nie et al., 2021; Wang et al., 2023), zinc oxide (Yan et al., 2020; Yan et al., 2021), titanium oxide (Patel et al., 2018), carbon (Kurantowicz et al., 2017; Samak et al., 2020), gold (Zielinska et al., 2011), platinum (Prasek et al., 2013), silver (Grodzik and Sawosz, 2006), magnetic iron oxide (Patel et al., 2019). We do not claim that the chicken embryo is a model for environmental exposure, nor do we make any claims about the uptake of particles into the bird egg in the field. We are simply using it as a convenient experimental model to unravel the cellular and molecular effects of nanoplastics in a warm-blooded, higher vertebrate model.

Outline of this thesis

The main objective of this thesis is to explore the developmental toxicity of nanoplastics and the underlying mechanisms of that toxicity. Another objective is to determine whether the chicken embryo a good model for ecotoxicology. My specific research questions are these:

- Do MPs and NPs cause developmental toxicity in chicken embryos?
- If so, what are the cellular and molecular mechanisms underlying the toxic effects?

I use a multi-technique approach in this doctoral work to: (i) unravel the effects of surface embryonic exposure to different concentration of 25 nm nanoplastics in the stage 8 chicken embryo using experimental embryological techniques; (ii) explore the teratogenic effects induced by nanoplastics in the chicken embryo using MicroCT and synchrotron x-ray tomography; (iii) determine the cellular and molecular mechanisms of toxicity of nanoplastics exposure by *in situ* hybridization in the chicken embryo.

In **Chapter 1**, I review the relevant research literature relating to nanoplastics, their origins and harmful effects, including their potential (but poorly understood) risks to humans. I also explore the literature justifying the my use of the chicken embryo as a model for this research.

In **Chapter 2**, I describe the research in which I exposed chick embryos to nanoplastics and looked at their dose-dependent and size-dependent developmental toxicity (and teratogenicity). I found that the the nanoplastics have a dose-dependent harmful effect on the neural tube, eye, tail and other organs. I could not make any conclusions about the possible size-dependency of these effects because of confounding variables (surface area, number of particles) which also change with particle size.

Chapter 3 consists of a detailed analysis of the effects of polystyrene nanoparticles (PS-NPs) on the cardiovascular system of the chicken embryo. Using synchrotron scanning, I showed that 25 nm PS-NPs cause ventricular septal defects and abnormal numbers of aortic arches. Video recordings of live, exposed chicken embryos showed that PS-NPs caused bradycardia. Gene expression studies and immunochemistry showed that malformations in the cardiovascular system might be understood in terms of abnormal migration of the cardiac neural crest.

In **Chapter 4**, I conducted transcriptional profiling of neural crest marker-genes in early chick embryos. I found that exposure of chick embryos to 25 nm PS-NPs caused disrupted migration of the trunk and cranial neural crest. Using fluorescein-tagged PS-NPs I showed that the nanoparticles co-localised to the neural crest (dorsal midline of the neural tube). TUNEL staining showed that exposure to PS-NPs caused a wave of cell death in the dorsal cells of the embryo.

Chapter 5 presents a summary, discussion and perspective of my research. I conclude that PS-NPs pose a potential health risk because they bind to, and disrupt the development of, neural crest cells. I argue that more work is needed to understand the mechanism underlying these effects, and to map the biodistribution of PS-NPs in the body of animals and humans.

References

- Aljaibachi, R. and Callaghan, A.** (2018). Impact of Polystyrene Microplastics on *Daphnia Magna* Mortality and Reproduction in Relation to Food Availability. *PeerJ* **6**, e4601-e4601.
- Anderson, J. C., Park, B. J. and Palace, V. P.** (2016). Microplastics in Aquatic Environments: Implications for Canadian Ecosystems. *Environmental Pollution* **218**, 269-280.
- Arthur, C., Baker, J. and Bamford, H.** (2008). International Research Workshop on the Occurrence, Effects, and Fate of Microplastic Marine Debris. In *Conference Proceedings. Sept*, pp. 9-11.
- Avio, C. G., Gorbi, S. and Regoli, F.** (2017). Plastics and Microplastics in the Oceans: From Emerging Pollutants to Emerged Threat. *Marine Environmental Research* **128**, 2-11.
- Bakir, A., Rowland, S. J. and Thompson, R. C.** (2014). Enhanced Desorption of Persistent Organic Pollutants from Microplastics under Simulated Physiological Conditions. *Environmental Pollution* **185**, 16-23.
- Bashirova, N., Poppitz, D., Klüver, N., Scholz, S., Matysik, J. and Alia, A.** (2023). A Mechanistic Understanding of the Effects of Polyethylene Terephthalate Nanoplastics in the Zebrafish (*Danio Rerio*) Embryo. *Scientific Reports* **13**, 1891.
- Bates, A.** (2019). *Transforming Plastic: From Pollution to Evolution*: GroundSwell Books.
- Becker, H. and Locascio, L. E.** (2002). Polymer Microfluidic Devices. *Talanta* **56**, 267-287.
- Bergmann, M., Mützel, S., Primpke, S., Tekman, M. B., Trachsel, J. and Gerdt, G.** (2019). White and Wonderful? Microplastics Prevail in Snow from the Alps to the Arctic. *Science Advances* **5**, eaax1157- eaax1157.
- Boehnke, N., Straehla, J. P., Safford, H. C., Kocak, M., Rees, M. G., Ronan, M., Rosenberg, D., Adelman, C. H., Chivukula, R. R., Nabar, N., et al.** (2022). Massively Parallel Pooled Screening Reveals Genomic Determinants of Nanoparticle Delivery. *Science* **377**, eabm5551.
- Bojic, S., Falco, M. M., Stojkovic, P., Ljujic, B., Gazdic Jankovic, M., Armstrong, L., Markovic, N., Dopazo, J., Lako, M., Bauer, R., et al.** (2020). Platform to Study Intracellular Polystyrene Nanoplastic Pollution and Clinical Outcomes. *Stem Cells* **38**, 1321-1325.
- Boucher, J. and Friot, D.** (2017). *Primary Microplastics in the Oceans: A Global Evaluation of Sources*: IUCN Gland, Switzerland.
- Bouwmeester, H., Hollman, P. C. H. and Peters, R. J. B.** (2015). Potential Health Impact of Environmentally Released Micro-and Nanoplastics in the Human Food Production Chain: Experiences from Nanotoxicology. *Environmental Science & Technology* **49**, 8932-8947.
- Browne, M. A., Crump, P., Niven, S. J., Teuten, E., Tonkin, A., Galloway, T. and Thompson, R.** (2011). Accumulation of Microplastic on Shorelines Worldwide: Sources and Sinks. *Environmental Science & Technology* **45**, 9175-9179.
- Browne, M. A., Galloway, T. and Thompson, R.** (2007). Microplastic—an Emerging Contaminant of Potential Concern? *Integrated Environmental Assessment and Management: An International Journal* **3**, 559-561.
- Brun, N. R., Beenakker, M. M. T., Hunting, E. R., Ebert, D. and Vijver, M. G.** (2017). Brood Pouch-Mediated Polystyrene Nanoparticle Uptake During *Daphnia Magna* Embryogenesis. *Nanotoxicology* **11**, 1059-1069.
- Brun, N. R., van Hage, P., Hunting, E. R., Haramis, A. G., Vink, S. C., Vijver, M. G., Schaaf, M. J. M. and Tudorache, C.** (2019). Polystyrene Nanoplastics Disrupt Glucose Metabolism and Cortisol Levels with a Possible Link to Behavioural Changes in Larval Zebrafish. *Commun Biol* **2**, 382.
- Bryda, E. C.** (2013). The Mighty Mouse: The Impact of Rodents on Advances in Biomedical Research. *Missouri Medicine* **110**, 207-207.
- Burns, E. E. and Boxall, A. B. A.** (2018). Microplastics in the Aquatic Environment: Evidence for or against Adverse Impacts and Major Knowledge Gaps. *Environmental Toxicology and Chemistry* **37**, 2776-2796.
- Butterworth, A., Clegg, I. and Bass, C.** (2012). Marine Debris: A Global Picture of the Impact on Animal Welfare and of Animal-Focused Solutions. *WSPA International* **222**.
- Carson, R.** (1962). *Silent Spring* *iii*. New Yorker **23**.
- Center, L. T. and Wash, O. C.** (2017). Microfiber Release from Clothes after Washing: Hard Facts, Figures and Promising Solutions.
- Chai, G., Hassan, A., Meng, T., Lou, L., Ma, J., Simmers, R., Zhou, L., Rubin, B. K., Zhou, Q. T., Longest, P. W., et al.** (2020). Dry Powder Aerosol Containing Muco-Inert Particles for Excipient Enhanced Growth Pulmonary Drug Delivery. *Nanomedicine* **29**, 102262.

- Chang, X., Xue, Y., Li, J., Zou, L. and Tang, M.** (2020). Potential Health Impact of Environmental Micro- and Nanoplastics Pollution. *Journal of Applied Toxicology* **40**, 4-15.
- Chauhan, G. S. and Wani, S.** (2019). Plastic Pollution: A Major Environmental Threat. *Int. J. Innov. Res. Technol* **6**, 43-46.
- Chortarea, S., Gupta, G., Saarimäki, L. A., Netkueakul, W., Manser, P., Aengenheister, L., Wichser, A., Fortino, V., Wick, P., Greco, D., et al.** (2023). Transcriptomic Profiling Reveals Differential Cellular Response to Copper Oxide Nanoparticles and Polystyrene Nanoplastics in Perfused Human Placenta. *Environment International* **177**, 108015.
- Clark, N. J., Khan, F. R., Crowther, C., Mitrano, D. M. and Thompson, R. C.** (2023). Uptake, Distribution and Elimination of Palladium-Doped Polystyrene Nanoplastics in Rainbow Trout (*Oncorhynchus Mykiss*) Following Dietary Exposure. *Science of The Total Environment* **854**, 158765.
- Cox, K. D., Covernton, G. A., Davies, H. L., Dower, J. F., Juanes, F. and Dudas, S. E.** (2019). Human Consumption of Microplastics. *Environmental Science & Technology* **53**, 7068-7074.
- Davey, M. G. and Tickle, C.** (2007). The Chicken as a Model for Embryonic Development. *Cytogenetic and Genome Research* **117**, 231-239.
- De Jong, W. H., Hagens, W. I., Krystek, P., Burger, M. C., Sips, A. J. and Geertsma, R. E.** (2008). Particle Size-Dependent Organ Distribution of Gold Nanoparticles after Intravenous Administration. *Biomaterials* **29**, 1912-1919.
- de Sá, L. C., Oliveira, M., Ribeiro, F., Rocha, T. L. and Futter, M. N.** (2018). Studies of the Effects of Microplastics on Aquatic Organisms: What Do We Know and Where Should We Focus Our Efforts in the Future? *Science of The Total environment* **645**, 1029-1039.
- de Souza Machado, A. A., Kloas, W., Zarfl, C., Hempel, S. and Rillig, M. C.** (2018). Microplastics as an Emerging Threat to Terrestrial Ecosystems. *Global Change Biology* **24**, 1405-1416.
- Desai, S. B. and Galage, C. K.** (2015). Production and Analysis of Pyrolysis Oil from Waste Plastic in Kolhapur City. *International Journal of Engineering Research and General Science* **3**, 590-595.
- Du, F., Cai, H., Zhang, Q., Chen, Q. and Shi, H.** (2020). Microplastics in Take-out Food Containers. *Journal of Hazardous Materials*, 122969-122969.
- Duman, R., Ertekin, T., Duman, R., Aslan, E., Sabaner, M. C. and Cetinkaya, E.** (2019). The Novel Model: Experimental Optical Coherence Tomography-Guided Anterior Segment Imaging Chick Embryo Model. *Indian J Ophthalmol* **67**, 54-58.
- Dusza, H. M., Katrukha, E. A., Nijmeijer, S. M., Akhmanova, A., Vethaak, A. D., Walker, D. I. and Legler, J.** (2022). Uptake, Transport, and Toxicity of Pristine and Weathered Micro- and Nanoplastics in Human Placenta Cells. *Environmental Health Perspectives* **130**, 097006.
- Eerkes-Medrano, D. and Thompson, R.** (2018). Occurrence, Fate, and Effect of Microplastics in Freshwater Systems. In *Microplastic Contamination in Aquatic Environments*, pp. 95-132: Elsevier.
- Fendall, L. S. and Sewell, M. A.** (2009). Contributing to Marine Pollution by Washing Your Face: Microplastics in Facial Cleansers. *Marine Pollution Bulletin* **58**, 1225-1228.
- Ferreira, P., Fonte, E., Soares, M. E., Carvalho, F. and Guilhermino, L.** (2016). Effects of Multi-Stressors on Juveniles of the Marine Fish *Pomatoschistus Microps*: Gold Nanoparticles, Microplastics and Temperature. *Aquatic Toxicology* **170**, 89-103.
- Frere, L., Paul-Pont, I., Rinnert, E., Petton, S., Jaffré, J., Bihannic, I., Soudant, P., Lambert, C. and Huvet, A.** (2017). Influence of Environmental and Anthropogenic Factors on the Composition, Concentration and Spatial Distribution of Microplastics: A Case Study of the Bay of Brest (Brittany, France). *Environmental Pollution* **225**, 211-222.
- Galafassi, S., Nizzetto, L. and Volta, P.** (2019). Plastic Sources: A Survey across Scientific and Grey Literature for Their Inventory and Relative Contribution to Microplastics Pollution in Natural Environments, with an Emphasis on Surface Water. *Sci. Total Environ* **693**, 133499-133499.
- Gall, S. C. and Thompson, R. C.** (2015). The Impact of Debris on Marine Life. *Marine Pollution Bulletin* **92**, 170-179.
- Gallo, F., Fossi, C., Weber, R., Santillo, D., Sousa, J., Ingram, I., Nadal, A. and Romano, D.** (2018). Marine Litter Plastics and Microplastics and Their Toxic Chemicals Components: The Need for Urgent Preventive Measures. *Environmental Sciences Europe* **30**, 13-13.
- Galloway, T. S.** (2015). *Micro- and Nano-Plastics and Human Health*. pp. 343-366: Springer, Cham.
- Galloway, T. S., Cole, M. and Lewis, C.** (2017). Interactions of Microplastic Debris Throughout the Marine Ecosystem. *Nature Ecology & Evolution* **1**, 1-8.
- Garrido Gamarro, E., Ryder, J., Elvevoll, E. O. and Olsen, R. L.** (2020). Microplastics in Fish and Shellfish—a Threat to Seafood Safety? *Journal of Aquatic Food Product Technology* **29**, 417-425.

- Geyer, R., Jambeck, J. R. and Law, K. L.** (2017). Production, Use, and Fate of All Plastics Ever Made. *Sci Adv* **3**, e1700782.
- Ghimire, S., Zhang, X., Zhang, J. and Wu, C.** (2022). Use of Chicken Embryo Model in Toxicity Studies of Endocrine-Disrupting Chemicals and Nanoparticles. *Chemical Research in Toxicology* **35**, 550-568.
- Gibb, B. C.** (2019). Plastics Are Forever. *Nat Chem* **11**, 394-395.
- Gigault, J., Ter Halle, A., Baudrimont, M., Pascal, P.-Y., Gauffre, F., Phi, T.-L., El Hadri, H., Grassl, B. and Reynaud, S.** (2018). Current Opinion: What Is a Nanoplastic? *Environmental Pollution* **235**, 1030-1034.
- Gourmelon, G.** (2015). Global Plastic Production Rises, Recycling Lags. *Vital Signs* **22**, 91-95.
- Grodzik, M. and Sawosz, E.** (2006). The Influence of Silver Nanoparticles on Chick Embryo Development and Bursa of Fabricius Morphology. *Journal of Animal and Feed Sciences* **15**, 111.
- Hamburger, V. and Hamilton, H. L.** (1951). A Series of Normal Stages in the Development of the Chick Embryo. *J Morphol* **88**, 49-92.
- He, D., Luo, Y., Lu, S., Liu, M., Song, Y. and Lei, L.** (2018). Microplastics in Soils: Analytical Methods, Pollution Characteristics and Ecological Risks. *TrAC Trends in Analytical Chemistry* **109**, 163-172.
- Hedges, S. B. and Kumar, S.** (2009). *The Timetree of Life*: OUP Oxford.
- Hernandez, L. M., Xu, E. G., Larsson, H. C. E., Tahara, R., Maisuria, V. B. and Tufenkji, N.** (2019). Plastic Teabags Release Billions of Microparticles and Nanoparticles into Tea. *Environmental Science & Technology* **53**, 12300-12310.
- Hernandez, L. M., Yousefi, N. and Tufenkji, N.** (2017). Are There Nanoplastics in Your Personal Care Products? *Environmental Science & Technology Letters* **4**, 280-285.
- Horn, D. A., Granek, E. F. and Steele, C. L.** (2020). Effects of Environmentally Relevant Concentrations of Microplastic Fibers on Pacific Mole Crab (*Emerita Analoga*) Mortality and Reproduction. *Limnology and Oceanography Letters* **5**, 74-83.
- Horton, A. A., Walton, A., Spurgeon, D. J., Lahive, E. and Svendsen, C.** (2017). Microplastics in Freshwater and Terrestrial Environments: Evaluating the Current Understanding to Identify the Knowledge Gaps and Future Research Priorities. *Science of the Total Environment* **586**, 127-141.
- Hougaard, K. S., Campagnolo, L., Chavatte-Palmer, P., Tarrade, A., Rousseau-Ralliard, D., Valentino, S., Park, M. V. D. Z., de Jong, W. H., Wolterink, G. and Piersma, A. H.** (2015). A Perspective on the Developmental Toxicity of Inhaled Nanoparticles. *Reproductive Toxicology* **56**, 118-140.
- Jiang, B., Kauffman, A. E., Li, L., McFee, W., Cai, B., Weinstein, J., Lead, J. R., Chatterjee, S., Scott, G. I. and Xiao, S.** (2020). Health Impacts of Environmental Contamination of Micro- and Nanoplastics: A Review. *Environmental Health and Preventive Medicine* **25**, 1-15.
- Karlsson, T. M., Vethaak, A. D., Almroth, B. C., Ariese, F., van Velzen, M., Hassellöv, M. and Leslie, H. A.** (2017). Screening for Microplastics in Sediment, Water, Marine Invertebrates and Fish: Method Development and Microplastic Accumulation. *Marine Pollution Bulletin* **122**, 403-408.
- Kashiwada, S.** (2006). Distribution of Nanoparticles in the See-through Medaka (*Oryzias Latipes*). *Environmental Health Perspectives* **114**, 1697-1702.
- Koelmans, A. A., Besseling, E. and Foekema, E. M.** (2014). Leaching of Plastic Additives to Marine Organisms. *Environmental Pollution* **187**, 49-54.
- Korhonen, A., Hemminki, K. and Vainio, H.** (1982). Application of the Chicken Embryo in Testing for Embryotoxicity: Thiurams. *Scand J Work Environ Health* **8**, 63-69.
- Kurantowicz, N., Sawosz, E., Halik, G., Strojny, B., Hotowy, A., Grodzik, M., Piast, R., Pasanphan, W. and Chwalibog, A.** (2017). Toxicity Studies of Six Types of Carbon Nanoparticles in a Chicken-Embryo Model. *Int J Nanomedicine* **12**, 2887-2898.
- Levine, B. and Berman, W. E.** (1995). The Current Status of Expanded Polytetrafluoroethylene (Gore-Tex) in Facial Plastic Surgery. *Ear, Nose & Throat Journal* **74**, 681-684.
- Li, J., Chen, C. and Xia, T.** (2022). Understanding Nanomaterial–Liver Interactions to Facilitate the Development of Safer Nanoapplications. *Advanced Materials* **34**, 2106456.
- Li, J., Yang, D., Li, L., Jabeen, K. and Shi, H.** (2015). Microplastics in Commercial Bivalves from China. *Environmental Pollution* **207**, 190-195.
- Liebezeit, G. and Liebezeit, E.** (2013). Non-Pollen Particulates in Honey and Sugar. *Food Additives & Contaminants: Part A* **30**, 2136-2140.
- Lu, Y., Zhang, Y., Deng, Y., Jiang, W., Zhao, Y., Geng, J., Ding, L. and Ren, H.** (2016). Uptake and Accumulation of Polystyrene Microplastics in Zebrafish (*Danio Rerio*) and Toxic Effects in Liver. *Environmental Science & Technology* **50**, 4054-4060.
- Lund, A. W., Medler, T. R., Leachman, S. A. and Coussens, L. M.** (2016). Lymphatic Vessels, Inflammation, and Immunity in Skin Cancer. *Cancer Discov* **6**, 22-35.

- Lusher, A.** (2015). Microplastics in the Marine Environment: Distribution, Interactions and Effects, Pp. 245e307.
- Maes, T., Jessop, R., Wellner, N., Haupt, K. and Mayes, A. G.** (2017). A Rapid-Screening Approach to Detect and Quantify Microplastics Based on Fluorescent Tagging with Nile Red. *Scientific Reports* **7**, 44501-44501.
- Mahe, B., Vogt, A., Liard, C., Duffy, D., Abadie, V., Bonduelle, O., Boissonnas, A., Sterry, W., Verrier, B., Blume-Peytavi, U., et al.** (2009). Nanoparticle-Based Targeting of Vaccine Compounds to Skin Antigen-Presenting Cells by Hair Follicles and Their Transport in Mice. *Journal of Investigative Dermatology* **129**, 1156-1164.
- Mak, C. W., Yeung, K. C.-F. and Chan, K. M.** (2019). Acute Toxic Effects of Polyethylene Microplastic on Adult Zebrafish. *Ecotoxicology and Environmental Safety* **182**, 109442-109442.
- Mortensen, L. J., Oberdörster, G., Pentland, A. P. and DeLouise, L. A.** (2008). In Vivo Skin Penetration of Quantum Dot Nanoparticles in the Murine Model: The Effect of Uvr. *Nano letters* **8**, 2779-2787.
- Nanau, R. M. and Neuman, M. G.** (2013). Adverse Drug Reactions Induced by Valproic Acid. *Clinical Biochemistry* **46**, 1323-1338.
- Navarro, E., Baun, A., Behra, R., Hartmann, N. B., Filser, J., Miao, A.-J., Quigg, A., Santschi, P. H. and Sigg, L.** (2008). Environmental Behavior and Ecotoxicity of Engineered Nanoparticles to Algae, Plants, and Fungi. *Ecotoxicology* **17**, 372-386.
- Ng, E.-L., Lwanga, E. H., Eldridge, S. M., Johnston, P., Hu, H.-W., Geissen, V. and Chen, D.** (2018). An Overview of Microplastic and Nanoplastic Pollution in Agroecosystems. *Science of The Total Environment* **627**, 1377-1388.
- Nguyen, B., Claveau-Mallet, D., Hernandez, L. M., Xu, E. G., Farner, J. M. and Tufenkji, N.** (2019). Separation and Analysis of Microplastics and Nanoplastics in Complex Environmental Samples. *Accounts of Chemical Research* **52**, 858-866.
- Nie, J.-h., Shen, Y., Roshdy, M., Cheng, X., Wang, G. and Yang, X.** (2021). Polystyrene Nanoplastics Exposure Caused Defective Neural Tube Morphogenesis through Caveolae-Mediated Endocytosis and Faulty Apoptosis. *Nanotoxicology*, 1-20.
- Nkwachukwu, O. I., Chima, C. H., Ikenna, A. O. and Albert, L.** (2013). Focus on Potential Environmental Issues on Plastic World Towards a Sustainable Plastic Recycling in Developing Countries. *International Journal of Industrial Chemistry* **4**, 34-34.
- Ornoy, A.** (2009). Valproic Acid in Pregnancy: How Much Are We Endangering the Embryo and Fetus? *Reproductive Toxicology* **28**, 1-10.
- Panda, A. K., Singh, R. K. and Mishra, D. K.** (2010). Thermolysis of Waste Plastics to Liquid Fuel: A Suitable Method for Plastic Waste Management and Manufacture of Value Added Products—a World Prospective. *Renewable and Sustainable Energy Reviews* **14**, 233-248.
- Pardo, J. D., Lennie, K. and Anderson, J. S.** (2020). Can We Reliably Calibrate Deep Nodes in the Tetrapod Tree? Case Studies in Deep Tetrapod Divergences. *Frontiers in Genetics* **11**.
- Patel, M. M., Goyal, B. R., Bhadada, S. V., Bhatt, J. S. and Amin, A. F.** (2009). Getting into the Brain. *CNS Drugs* **23**, 35-58.
- Patel, S., Jana, S., Chetty, R., Thakore, S., Singh, M. and Devkar, R.** (2018). Tio2 Nanoparticles Induce Omphalocele in Chicken Embryo by Disrupting Wnt Signaling Pathway. *Scientific Reports* **8**, 4756.
- Patel, S., Jana, S., Chetty, R., Thakore, S., Singh, M. and Devkar, R.** (2019). Toxicity Evaluation of Magnetic Iron Oxide Nanoparticles Reveals Neuronal Loss in Chicken Embryo. *Drug Chem Toxicol* **42**, 1-8.
- Pitt, J. A., Kozal, J. S., Jayasundara, N., Massarsky, A., Trevisan, R., Geitner, N., Wiesner, M., Levin, E. D. and Di Giulio, R. T.** (2018). Uptake, Tissue Distribution, and Toxicity of Polystyrene Nanoparticles in Developing Zebrafish (*Danio Rerio*). *Aquatic Toxicology* **194**, 185-194.
- Platt, D. K.** (2003). *Engineering and High Performance Plastics Market Report: A Rapra Market Report*: iSmithers Rapra Publishing.
- Powell, J. J., Thoree, V. and Pele, L. C.** (2007). Dietary Microparticles and Their Impact on Tolerance and Immune Responsiveness of the Gastrointestinal Tract. *British Journal of Nutrition* **98**, S59-S63.
- Pozio, E.** (2005). The Broad Spectrum of *Trichinella* Hosts: From Cold-to Warm-Blooded Animals. *Veterinary Parasitology* **132**, 3-11.
- Prasek, M., Sawosz, E., Jaworski, S., Grodzik, M., Ostaszewska, T., Kamaszewski, M., Wierzbicki, M. and Chwalibog, A.** (2013). Influence of Nanoparticles of Platinum on Chicken Embryo Development and Brain Morphology. *Nanoscale Res Lett* **8**, 251.
- Prata, J. C.** (2018). Airborne Microplastics: Consequences to Human Health? *Environmental Pollution* **234**, 115-126.

- Qiao, R., Sheng, C., Lu, Y., Zhang, Y., Ren, H. and Lemos, B.** (2019). Microplastics Induce Intestinal Inflammation, Oxidative Stress, and Disorders of Metabolome and Microbiome in Zebrafish. *Science of the Total Environment* **662**, 246-253.
- Quiñonez-Silvero, C., Hübner, K. and Herzog, W.** (2020). Development of the Brain Vasculature and the Blood-Brain Barrier in Zebrafish. *Dev Biol* **457**, 181-190.
- Ragusa, A., Svelato, A., Santacroce, C., Catalano, P., Notarstefano, V., Carnevali, O., Papa, F., Rongioletti, M. C. A., Baiocco, F., Draghi, S., et al.** (2021). Plasticenta: First Evidence of Microplastics in Human Placenta. *Environ Int* **146**, 106274.
- Rashidi, H. and Sottile, V.** (2009). The Chick Embryo: Hatching a Model for Contemporary Biomedical Research. *Bioessays* **31**, 459-465.
- Ray, S. S., Lee, H. K., Huyen, D. T. T., Chen, S.-S. and Kwon, Y.-N.** (2022). Microplastics Waste in Environment: A Perspective on Recycling Issues from Ppe Kits and Face Masks During the Covid-19 Pandemic. *Environmental Technology & Innovation*, 102290.
- Rezania, S., Park, J., Din, M. F. M., Taib, S. M., Talaiekhosani, A., Yadav, K. K. and Kamyab, H.** (2018). Microplastics Pollution in Different Aquatic Environments and Biota: A Review of Recent Studies. *Marine Pollution Bulletin* **133**, 191-208.
- Roch, S., Friedrich, C. and Brinker, A.** (2020). Uptake Routes of Microplastics in Fishes: Practical and Theoretical Approaches to Test Existing Theories. *Scientific Reports* **10**, 1-12.
- Romanoff, A. L.** (1960). *The Avian Embryo : Structural and Functional Development*. New York: Macmillan.
- Samak, D. H., El-Sayed, Y. S., Shaheen, H. M., El-Far, A. H., Abd El-Hack, M. E., Noreldin, A. E., El-Naggar, K., Abdelnour, S. A., Saied, E. M., El-Seedi, H. R., et al.** (2020). Developmental Toxicity of Carbon Nanoparticles During Embryogenesis in Chicken. *Environ Sci Pollut Res Int* **27**, 19058-19072.
- Sánchez-Villagra, M.** (2012). *Embryos in Deep Time: The Rock Record of Biological Development*: Univ of California Press.
- Sangkham, S., Faikhaw, O., Munkong, N., Sakunkoo, P., Arunlertaree, C., Chavali, M., Mousazadeh, M. and Tiwari, A.** (2022). A Review on Microplastics and Nanoplastics in the Environment: Their Occurrence, Exposure Routes, Toxic Studies, and Potential Effects on Human Health. *Marine Pollution Bulletin* **181**, 113832.
- Schirinzi, G. F., Pérez-Pomeda, I., Sanchís, J., Rossini, C., Farré, M. and Barceló, D.** (2017). Cytotoxic Effects of Commonly Used Nanomaterials and Microplastics on Cerebral and Epithelial Human Cells. *Environmental Research* **159**, 579-587.
- Shruti, V. C. and Kutralam-Muniasamy, G.** (2019). Bioplastics: Missing Link in the Era of Microplastics. *Science of The Total Environment* **697**, 134139-134139.
- Smith, M., Love, D. C., Rochman, C. M. and Neff, R. A.** (2018). Microplastics in Seafood and the Implications for Human Health. *Current Environmental Health Reports* **5**, 375-386.
- Sökmen, T. Ö., Sulukan, E., Türkoğlu, M., Baran, A., Özkaraca, M. and Ceyhun, S. B.** (2020). Polystyrene Nanoplastics (20 Nm) Are Able to Bioaccumulate and Cause Oxidative DNA Damages in the Brain Tissue of Zebrafish Embryo (Danio Rerio). *NeuroToxicology* **77**, 51-59.
- Souza, A. M. d., Santos, A. L., Araújo, D. S., Magalhães, R. R. d. B. and Rocha, T. L.** (2022). Micro(Nano)Plastics as a Vector of Pharmaceuticals in Aquatic Ecosystem: Historical Review and Future Trends. *Journal of Hazardous Materials Advances* **6**, 100068.
- St John, J. A., Braun, E. L., Isberg, S. R., Miles, L. G., Chong, A. Y., Gongora, J., Dalzell, P., Moran, C., Bed'Hom, B., Abzhanov, A., et al.** (2012). Sequencing Three Crocodylian Genomes to Illuminate the Evolution of Archosaurs and Amniotes. *Genome Biology* **13**, 415.
- Stevens, A. P.** (2015). Tiny Plastic, Big Problem. *Science News for Students*.
- Subramanian, P. M.** (2000). Plastics Recycling and Waste Management in the Us. *Resources, Conservation and Recycling* **28**, 253-263.
- Tanoshima, M., Kobayashi, T., Tanoshima, R., Beyene, J., Koren, G. and Ito, S.** (2015). Risks of Congenital Malformations in Offspring Exposed to Valproic Acid in Utero: A Systematic Review and Cumulative Meta-Analysis. *Clinical Pharmacology & Therapeutics* **98**, 417-441.
- Tickle, C.** (1993). Chick Limb Buds. In *Essential Developmental Biology: A Practical Approach*, pp. 119-125: IRL Press at Oxford University Press.
- Toussaint, B., Raffael, B., Angers-Loustau, A., Gilliland, D., Kestens, V., Petrillo, M., Rio-Echevarria, I. M. and Van den Eede, G.** (2019). Review of Micro-and Nanoplastic Contamination in the Food Chain. *Food Additives & Contaminants: Part A* **36**, 639-673.

- Try, C., Moulari, B., Béduneau, A., Fantini, O., Pin, D., Pellequer, Y. and Lamprecht, A. (2016). Size Dependent Skin Penetration of Nanoparticles in Murine and Porcine Dermatitis Models. *European Journal of Pharmaceutics and Biopharmaceutics* **100**, 101-108.
- Unar, I. N., Soomro, S. A. and Aziz, S. (2010). Effect of Various Additives on the Physical Properties of Polyvinylchloride Resin. *Pakistan Journal of Analytical & Environmental Chemistry* **11**, 7-7.
- Velzeboer, I., Kwadijk, C. and Koelmans, A. A. (2014). Strong Sorption of Pcb's to Nanoplastics, Microplastics, Carbon Nanotubes, and Fullerenes. *Environmental Science & Technology* **48**, 4869-4876.
- Veneman, W. J., Spaijk, H. P., Brun, N. R., Bosker, T. and Vijver, M. G. (2017). Pathway Analysis of Systemic Transcriptome Responses to Injected Polystyrene Particles in Zebrafish Larvae. *Aquatic Toxicology* **190**, 112-120.
- Vogt, A., Combadiere, B., Hadam, S., Stieler, K. M., Lademann, J., Schaefer, H., Autran, B., Sterry, W. and Blume-Peytavi, U. (2006). 40nm, but Not 750 or 1,500nm, Nanoparticles Enter Epidermal Cd1a+ Cells after Transcutaneous Application on Human Skin. *Journal of Investigative Dermatology* **126**, 1316-1322.
- Vom Sqaal, F., Parmigiani, S., Palanza, P. L., Everett, L. G. and Ragaini, R. (2008). The Plastic World: Sources, Amounts, Ecological Impacts and Effects on Development Reproduction, Brain and Behavior in Aquatic and Terrestrial Animals and Humans. *Environmental Research (New York, NY: Print)* **108**, 127-184.
- Wan, J.-K., Chu, W.-L., Kok, Y.-Y. and Lee, C.-S. (2019a). Distribution of Microplastics and Nanoplastics in Aquatic Ecosystems and Their Impacts on Aquatic Organisms, with Emphasis on Microalgae. *Reviews of Environmental Contamination and Toxicology Volume 246*, 133-158.
- Wan, Y., Wu, C., Xue, Q. and Hui, X. (2019b). Effects of Plastic Contamination on Water Evaporation and Desiccation Cracking in Soil. *Science of The Total Environment* **654**, 576-582.
- Wang, M., Rücklin, M., Poelmann, R. E., de Mooij, C. L., Fokkema, M., Lamers, G. E. M., de Bakker, M. A. G., Chin, E., Bakos, L. J., Marone, F., et al. (2023). Nanoplastics Causes Extensive Congenital Malformations During Embryonic Development by Passively Targeting Neural Crest Cells. *Environment International* **173**, 107865.
- Wang, W., Gao, H., Jin, S., Li, R. and Na, G. (2019). The Ecotoxicological Effects of Microplastics on Aquatic Food Web, from Primary Producer to Human: A Review. *Ecotoxicology and Environmental Safety* **173**, 110-117.
- Webb, H. K., Arnott, J., Crawford, R. J. and Ivanova, E. P. (2013). Plastic Degradation and Its Environmental Implications with Special Reference to Poly (Ethylene Terephthalate). *Polymers* **5**, 1-18.
- Westphalen, H. and Abdelrasoul, A. (2017). Challenges and Treatment of Microplastics in Water. IntechOpen.
- Yan, Y., Wang, G., Huang, J., Zhang, Y., Cheng, X., Chuai, M., Brand-Saberi, B., Chen, G., Jiang, X. and Yang, X. (2020). Zinc Oxide Nanoparticles Exposure-Induced Oxidative Stress Restricts Cranial Neural Crest Development During Chicken Embryogenesis. *Ecotoxicology and Environmental Safety* **194**, 110415.
- Yan, Y., Wang, G., Luo, X., Zhang, P., Peng, S., Cheng, X., Wang, M. and Yang, X. (2021). Endoplasmic Reticulum Stress-Related Calcium Imbalance Plays an Important Role on Zinc Oxide Nanoparticles-Induced Failure of Neural Tube Closure During Embryogenesis. *Environment International* **152**, 106495.
- Yang, D., Shi, H., Li, L., Li, J., Jabeen, K. and Kolandhasamy, P. (2015). Microplastic Pollution in Table Salts from China. *Environmental Science & Technology* **49**, 13622-13627.
- Yong, C. Q. Y., Valiyaveetill, S. and Tang, B. L. (2020). Toxicity of Microplastics and Nanoplastics in Mammalian Systems. *International Journal of Environmental Research and Public Health* **17**, 1509-1509.
- Zeng, E. Y. (2018). *Microplastic Contamination in Aquatic Environments: An Emerging Matter of Environmental Urgency*: Elsevier.
- Zhang, G. S. and Liu, Y. F. (2018). The Distribution of Microplastics in Soil Aggregate Fractions in Southwestern China. *Science of the Total Environment* **642**, 12-20.
- Zhang, K., Hamidian, A. H., Tubić, A., Zhang, Y., Fang, J. K. H., Wu, C. and Lam, P. K. S. (2021). Understanding Plastic Degradation and Microplastic Formation in the Environment: A Review. *Environmental Pollution* **274**, 116554.
- Zhang, L. and Webster, T. J. (2009). Nanotechnology and Nanomaterials: Promises for Improved Tissue Regeneration. *Nano Today* **4**, 66-80.
- Zhang, S., Wang, J., Liu, X., Qu, F., Wang, X., Wang, X., Li, Y. and Sun, Y. (2019). Microplastics in the Environment: A Review of Analytical Methods, Distribution, and Biological Effects. *TrAC Trends in Analytical Chemistry* **111**, 62-72.
- Zhu, G. H., Gray, A. B. C. and Patra, H. K. (2022). Nanomedicine: Controlling Nanoparticle Clearance for Translational Success. *Trends in Pharmacological Sciences* **43**, 709-711.

- Zhu, M., Chernick, M., Rittschof, D. and Hinton, D. E.** (2020). Chronic Dietary Exposure to Polystyrene Microplastics in Maturing Japanese Medaka (*Oryzias Latipes*). *Aquatic Toxicology* **220**, 105396-105396.
- Ziajahromi, S., Neale, P. A., Rintoul, L. and Leusch, F. D. L.** (2017). Wastewater Treatment Plants as a Pathway for Microplastics: Development of a New Approach to Sample Wastewater-Based Microplastics. *Water Research* **112**, 93-99.
- Ziccardi, L. M., Edgington, A., Hentz, K., Kulacki, K. J. and Driscoll, S. K.** (2016). Microplastics as Vectors for Bioaccumulation of Hydrophobic Organic Chemicals in the Marine Environment: A State-of-the-Science Review. *Environmental Toxicology and Chemistry* **35**, 1667-1676.
- Zielinska, M., Sawosz, E., Grodzik, M., Wierzbicki, M., Gromadka, M., Hotowy, A., Sawosz, F., Lozicki, A. and Chwalibog, A.** (2011). Effect of Heparan Sulfate and Gold Nanoparticles on Muscle Development During Embryogenesis. *Int J Nanomedicine* **6**, 3163-3172.