



Universiteit  
Leiden  
The Netherlands

## Special issue adaptive tools for resilient bones: biostatistical approaches to past physical activity in osteoarchaeology

Schrader S.A., Carballo Pérez J.

### Citation

Schrader S.A., C. P. J. (2022). Special issue adaptive tools for resilient bones: biostatistical approaches to past physical activity in osteoarchaeology. *International Journal Of Osteoarchaeology*, 33(3), 381-388. doi:10.1002/oa.3177

Version: Publisher's Version



License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3716702>

**Note:** To cite this publication please use the final published version (if applicable).

**SPECIAL ISSUE PAPER**

# Special issue adaptive tools for resilient bones: Biostatistical approaches to past physical activity in osteoarchaeology

Sarah A. Schrader<sup>1</sup>  | Jared Carballo Pérez<sup>2</sup> <sup>1</sup>Faculty of Archaeology, Leiden University, Leiden, The Netherlands<sup>2</sup>UDI de Prehistoria, Arqueología e Historia Antigua, Facultad de Humanidades, Universidad de La Laguna, Tenerife, Canary Islands, Spain**Correspondence**Sarah A. Schrader, Faculty of Archaeology, Leiden University, Leiden, The Netherlands.  
Email: [s.a.schrader@arch.leidenuniv.nl](mailto:s.a.schrader@arch.leidenuniv.nl)**Funding information**

This workshop was funded by Leiden University Fund (W211115-6)

**Abstract**

In this introduction to the special issue, *Adaptive Tools for Resilient Bones: Biostatistical Approaches to Past Physical Activity in Osteoarchaeology*, we discuss the outcome of the workshop held in Leiden (the Netherlands; November 18–19, 2021). We review statistical approaches to enthesal changes and present a series of new contributions to this field. These research, commentary, and review articles present different statistical approaches to enthesal changes and reflect the current state of research in the field.

**KEYWORDS**

activity, enthesal changes, statistics

## 1 | INTRODUCTION

Investigations into physical activities of past individuals and populations have been an enduring goal of bioarchaeology. Jurmain and colleagues referred to activity reconstruction as the Holy Grail of bioarchaeology, simultaneously highlighting the value and potential impact of such research and, equally, the difficulty of attaining such an ambitious objective (Jurmain et al., 2012). With Angel, Kennedy, Merbs, and Dutour publishing some of the earliest case studies of musculoskeletal modifications to the skeleton, this line of research has been continually studied from an osteoarchaeological perspective for more than five decades (Angel, 1966; Dutour, 1986; Kennedy, 1983; Merbs, 1983).

The *International Journal of Osteoarchaeology* has a long history with activity-related research. In 1998, the journal published a special issue on “Stress Markers,” which focused on methodological approaches to quantifying and standardizing skeletal indicators of activity (Peterson & Hawkey, 1998). More than a decade later, the *International Journal of Osteoarchaeology* published another special issue related to activity, which focused on technical and theoretical advances in enthesal changes research (Henderson & Alves Cardoso, 2013). In addition to these transformative special issues, there have also been numerous impactful publications relating to

interdisciplinary perspectives, methodological advancements, experimental studies, terminology, questions regarding contributing factors, and critiques (see Acosta et al., 2017; Djukić et al., 2015, 2020; Godde et al., 2018; He & de Almeida Prado, 2021; Henderson et al., 2016; Karakostis et al., 2018; Karakostis, Jeffery, et al., 2019; Karakostis, Wallace, et al., 2019; Karakostis, Haeufle, et al., 2021; Karakostis, Reyes-Centeno, et al., 2021; Michopoulou et al., 2015, 2017; Milella et al., 2015; Niinimäki & Salmi, 2016, 2021; Nikita et al., 2019; Nolte & Wilczak, 2013; Salmi et al., 2020; Villotte & Knüsel, 2013; Villotte et al., 2016; Wallace et al., 2017).

However, one crucial point seemed to be lacking from this list—statistical clarity. Although this topic was discussed in a previous special issue, a lack of statistical transparency persisted (Robb, 1998). For this reason, the authors organized a two-day workshop entitled, *Adaptive Tools for Resilient Bones: Biostatistical Approaches to Past Physical Activity in Osteoarchaeology* held at the University of Leiden, the Netherlands (November 18–19, 2021). The objective of the conference was not to select the single best statistical approach, nor define an all-encompassing set of statistical guidelines. Rather, the goal was to discuss the advantages and disadvantages of various statistical methods currently being applied to enthesal changes and activity-related research more broadly. We consider this workshop to be a success, with more than 300 registered attendees

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *International Journal of Osteoarchaeology* published by John Wiley & Sons Ltd.

from 44 countries. We present here a brief discussion on the history of statistical approaches to activity related studies in bioarchaeology as well as a series of articles, which illustrate new research and review perspectives, that are a product of the Adaptive Tools for Resilient Bones workshop.

## 2 | STATISTICAL INQUIRY

As mentioned above, statistical approaches to enthesal changes have been diverse (Table 1). It is important to note that the statistical test of choice is often directly related to the how enthesal changes data has been recorded (e.g., continuous, ordinal, and binary dichotomy). At a most basic level, descriptive statistics are frequently used. This often involves reporting measures of central tendency (e.g., mean) and sometimes includes measures of variability (e.g., standard deviation, variance, minimum and maximum variables, kurtosis, and skewness). While these data are certainly informative, particularly if complete (i.e., measures of variability and measures of variance), many studies have adopted statistical approaches that allow for comparisons (e.g., between populations, time periods, and sexes).

The chi-square test has been used to assess whether two variables are associated. Bivariate correlations have been employed to examine the strength and direction of a linear relationship between two continuous or ordinal variables. This usually involves questions pertaining to the degree to which two variables are related (e.g., enthesal changes and age). Some authors opt for Pearson's correlation coefficient, while others argue that the more conservative nonparametric Spearman's rank correlation coefficient is more appropriate for enthesal data.

If researchers want to compare the means of two populations (e.g., population groups and time periods), a *t*-test or nonparametric Mann-Whitney *U* have been used. Multivariate correlations are also applied in scenarios where multiple variables are being compared ANOVA/ANCOVA/Factorial ANOVA, as well as the nonparametric Kruskal-Wallis, have been utilized. The Wilcoxon signed-rank test (nonparametric version of paired *t*-test) has been used, typically to test bilateral asymmetry within individuals.

More recently, various forms of regression have been applied to enthesal changes data. Generalized linear models (GLM) are a type of linear regression that allows for the study of the impact of multiple independent predictor variables and their interactions. Generalized estimating equations (GEE) are similar to GLM but allow for the analysis of longitudinal data. This brief summary is not exhaustive but highlights some of the major trends in statistical approaches in enthesal changes analysis in the past few decades.

## 3 | A BRIEF OVERVIEW OF PAPERS IN THIS SPECIAL ISSUE

In addition to this introduction, this special issue consists of seven contributions, including research articles, review articles, and short

reports. Here, we will briefly highlight these papers and their unique additions to the field.

In a compelling contribution, Villotte and Santos discuss the relationship between age-at-death and enthesal changes and, in doing so, question the “muscle-use” hypothesis that osteoarchaeologists often rely upon in their interpretations. Thirty fibrous and fibrocartilaginous attachment sites were examined in a large sample ( $n = 721$ ), originating from four collections (all European, 18th–20th centuries), where sex and age have been identified. Spearman's rank correlation coefficients and Fisher's exact tests were used to examine the relationship between age-at-death and enthesal changes. Logistic regression and log-linear models were applied to examine the interaction between age, sex, and enthesal severity. Adjusted standardized residuals were then utilized to identify which contributing factors impacted enthesal changes the most. Lastly, generalized odds ratios were employed to assess if the distribution of enthesal changes scores differed across age groups. Villotte and Santos found that the majority of fibrous points of attachment prior to the fourth decade were stage B (slight modifications affecting the margin and/or inner part of the enthesis), with stage C enthesal changes (major modifications affecting the margin and inner part of the enthesis) increasing with age. The authors note that, while these results seem to support the “muscle use” hypothesis, this should be used with caution as more histological and experimental analyses are required. Fibrocartilaginous entheses were found to increase across age classes, with the most major changes occurring in the fifth and sixth decade of life. These findings suggest that enthesal changes in young or middle adult skeletons may actually reflect microtrauma during life; however, this is increasingly unreliable with age.

Cheverko and colleagues present new enthesal data from a context of emergent inequality in the San Pedro de Atacama oases. Through the analysis of four contemporary sites (Middle Period AD 400–1000;  $n = 210$ ) of varying status, the authors examine the intersection of physical activity and increasing social and economic disparity. Thirty-four entheses were examined (17 per side) and data were then combined into joint categories (shoulder, elbow, wrist, hip, knee, ankle). ANCOVA and Factorial ANOVA were used to compare enthesal scores between sites. Both approaches allow for direct comparisons between multiple groups, while controlling for covariates—in this case, the most significant potential covariate was deemed to be age. Factorial ANOVAs were used to test the dependent variable (enthesal changes) against two or more categorical variables (i.e., linear interaction), while ANCOVAs were suited for numeric and ordinal variables as predictors. Cheverko and colleagues found that the elite site of Casa Parroquial had lower enthesal scores in most joints. These data suggest a varied lived experience, manifesting in embodied labour practices, between high-status and low-status groups in San Pedro de Atacama during a period of population expansion, trade network development, and socioeconomic inequality institutionalization.

Nikita and Radini elucidate how labour, activity, and daily life changed from the Anglo-Saxon period (6th–9th centuries CE) to the Medieval period (11th–16th centuries CE), in the East Midlands (England;  $n = 151$ ). This is a critical time period in history, as it marks

**TABLE 1** Examples of statistical approaches applied to enthesal changes research

Associations	Chi-square	Alves Cardoso & Henderson, 2013
		Bakirci et al., 2020
		Campanacho & Santos, 2013
		Godde & Taylor, 2013
		Listi, 2016
	Pearson's Spearman's	Milella et al., 2012
		Nolte & Wilczak, 2013
		Bakirci et al., 2020
		Lieverse et al., 2013
		Milella et al., 2012; Milella, 2014
Odds ratio	Niinimäki, 2012	
	Nikita et al., 2019	
	Palmer et al., 2016, 2019	
	Schrader, 2012, 2015	
	Acosta et al., 2017	
Differences between groups	t-test	Laffranchi et al., 2020
		Karakostis, Jeffery, & Harvati, 2019
	Mann–Whitney <i>U</i>	Nolte & Wilczak, 2013
		Alves Cardoso & Henderson, 2013
		Bakirci et al., 2020
		Carballo-Pérez et al., 2021
		He & de Almeida Prado, 2021
		Laffranchi et al., 2020
		Niinimäki & Baiges Sotos, 2013
		Palmer et al., 2016, 2019
Refai, 2019		
ANOVA/ANCOVA	Santana-Cabrera et al., 2015	
	Schrader, 2012, 2015	
	Yonemoto, 2016	
MANOVA	Godde et al., 2018; Godde & Taylor, 2013	
	Niinimäki, 2012; Niinimäki & Salmi, 2021	
	Schrader & Buzon, 2017	
Kruskal–Wallis	Bousquié et al., 2022	
	Milella et al., 2015	
	Thomas, 2014	
Differences within group	Wilcoxon	Alves Cardoso & Henderson, 2013
		Lieverse et al., 2013
		Schrader, 2012
		Yonemoto, 2016
		Listi, 2016
Regression	Logistic	Milella, 2014; Milella et al., 2012
		Niinimäki, 2012
		Refai, 2019
		Santana-Cabrera et al., 2015
		Alves Cardoso & Henderson, 2013
		Campanacho & Santos, 2013
		Godde & Taylor, 2013

(Continues)

TABLE 1 (Continued)

		Myszka et al., 2020
		Salega & Grosskopf, 2022
	Linear	Bakirci et al., 2020
		Listi, 2016
	Least squares	Niinimäki, 2011
	Ordinal	Henderson et al., 2013, 2017
	GLM/GEE	Alonso-Llamazares et al., 2022
		Becker, 2019
		Castro et al., 2022
		Henderson & Nikita, 2016
		Kubicka & Myszka, 2020
		Laffranchi et al., 2020
		Mazza, 2019
		Michopoulou et al., 2015, 2017
		Nikita, 2014
		Villotte et al., 2010
Other	Principal components analysis (PCA)	Karakostis et al., 2017
		Kubicka & Myszka, 2020
		Santana-Cabrera et al., 2015
		Yonemoto, 2016
	Nonlinear PCA	Milella et al., 2015
	Bayesian	Alonso-Llamazares et al., 2022
	3D landmark-based geometric morphometrics	Casado et al., 2019; Karakostis et al., 2018; Karakostis, Haeufle, et al., 2021; Karakostis, Reyes-Centeno, et al., 2021

the foundations of urbanization, increasing population size, and socio-economic stratification. The authors examined seven fibrocartilaginous entheses and cross-sectional geometric properties of the upper limb long bones. They innovatively used the Mean Measure of Divergence (for binary data, i.e., enthesal changes) and Mahalanobis distances (for continuous data, i.e., cross-sectional geometry) to examine and visualize similarities in activity patterns within and between assemblages. Additionally, the authors utilized the Shapiro–Wilk test to assess normality; ANOVA for inter-assemblage comparisons; Kruskal–Wallis, followed by pairwise Mann–Whitney tests, for those datasets lacking normal distribution; independent *t*-tests and Mann–Whitney tests for sexual dimorphism assessment, as well as GLM for exploring the effect the assemblage, sex, age, and body size on EC expression. Nikita and Radini found a general lack of sexual dimorphism, with the exception of cross sectional geometry for the medieval St. Peter's collection. For most assemblages, this suggests no pronounced gender-based division of labour was in place during either the Anglo-Saxon or medieval period. One other assemblage stood out in terms of enthesal changes; individuals at Austin Friars, a medieval monastery, had markedly gracile enthesal changes, which aligns well with the theorized limited physical activities they would have been engaged in. In conclusion, Nikita and Radini suggest activity during the Anglo-Saxon period may have been less complex and less

diversified than the later Medieval period; however, there were notable similarities between the assemblages of these periods.

Becker et al. problematize current approaches to enthesal changes research and provide an innovative alternative to how these data can be interpreted going forward. Oftentimes, enthesal changes research is conducted at the population level, involving numerous entheses, which can result in overwhelming and unclear findings. The authors propose an individual-level approach to enthesal analysis using a large collection of skeletal data ( $n = 1203$ ) from the Tiwanaku culture (500–1100 CE; Bolivia). A total of 37 entheses are examined and grouped into use areas (upper arm, forearm, mid-body, lower body, foot). By applying GEE, Becker and colleagues are able to examine a dependent variable's relationship to both within-subject and between-subject variables (e.g., location, time period, sex, and age). The authors also discuss database design and advocate for the use of one burial or specimen number per individual, for which all associated data can be “anchored,” thus facilitating queries, exploratory analysis, as well as, but not limited to, multiresearch use. Using these tools, Becker and colleagues found significant differences between the Tiwanaku core and colony communities, with modelled frequencies often higher in the core than the colony. Females in the Tiwanaku core also had higher enthesal scores than in the colony (upper arm and lower body). Similarly, core males had higher enthesal scores

than colony males (upper arm, forearm, mid-body, lower body). Additionally, differences between colonial neighbourhoods were identified, suggesting there may have been specific community function and specialization.

Carballo Pérez and Schrader examine enthesal changes in the Nile Valley during the state formation process (ca. 4800–1750 BCE). This period was a critical phase as agricultural practices were dramatically intensified, more and more people settled into permanent communities, and socioeconomic status became increasingly divided. Multiple skeletal collections from predynastic and early-dynastic Egypt and Nubia ( $n = 259$ ) were examined. Nonparametric tests, namely, Spearman's correlation coefficient, Mann–Whitney  $U$ , and Kruskal–Wallis, are applied to compare enthesal changes between sexes, sites, time periods, and left and right sides. Results indicate a decrease in enthesal changes early in the predynastic period, which the authors argue may be linked to increasing specialization and sedentism. This is followed by an increase in enthesal changes, which aligns well with the archaeological data that suggests an institutionalization of inequalities around this same time. There is a great deal of variability in enthesal changes between sites, with high-status sites exhibiting lower enthesal changes scores. Lastly, there also appear to be differences in enthesal changes between sexes, potentially indicating gendered labour activities. Women have higher levels of forearm supination and rotation and could have been in repetitive and strenuous activities involving handcrafting, pottery production, as well as planting, collecting, and processing agricultural resources.

In a review article, Karakostis provides a current summary of enthesal research using the Validated Entheses-based Reconstruction of Activity (VERA) method. Several authors have argued that visual assessment of enthesal severity is unreliable, with high inter- and intraobserver error. Karakostis has provided evidence that the VERA method, employing three-dimensional bone models, more accurately defines and quantifies points of muscle and ligament attachment. Additionally, the validity of the VERA approach has been confirmed using experimental studies (Bousquie et al., 2022; Castro et al., 2022; Karakostis et al., 2017; Karakostis, Jeffery, et al., 2019; Karakostis, Wallace, et al., 2019). Karakostis applies size adjustment and multivariate statistical analyses to identify habitual and coordinated muscle groups. Here, Karakostis provides the first step-by-step protocol for univariate and multivariate analysis of VERA quantified data and presents a novel plotting technique, the Group Enthesal Patterning Index. Additionally, techniques for size adjustment, analysis of single entheses, multivariate statistics, biomechanical efficiency, as well as limitations and future objectives, are also discussed in this highly informative and thorough contribution.

The Van der Pas and Schrader contribution reflects a collaboration between a statistician and an osteoarchaeologist. In this commentary, the authors argue that standardizing statistical methods for enthesal changes is not realistic or even desired. Data structure and research questions differ greatly and, therefore, there cannot be a single recommendation for a statistical application. Additionally, examples are provided for how standardization of statistical testing can actually block innovation rather than foster reliability. However, what

can be standardized is how statistical analysis with enthesal data are reported within osteoarchaeological studies. The authors use STROBE (STrengthening the Reporting of OBservational studies in Epidemiology) as an example of how guidelines can be put in place to structure data reporting and statistical transparency. While such a STROBE list has not yet been developed for enthesal changes, a list of considerations is provided to help guide authors in their statistical analysis reporting.

## 4 | CONCLUSIONS

We are very pleased with the outcome of Adaptive Tools for Bones and view this special issue as a step forward in statistical communication within biological anthropology. The contributions in this volume highlight the variety of statistical tests that can and have been used when examining enthesal changes. We suggest that the type of statistical test employed is highly dependent upon the data and research question(s). We hope that this series of publications provide examples and directions for those interested in enthesal changes and activity-related osteological research. Each contribution has aimed to discuss the advantages, limitations, and assumptions of their statistical method of choice. We argue that the first and next step to enthesal changes data handling should be clarity of statistical approach, including how missing data were handled, how statistical assumptions were met (or not), and a complete statistical output, moving beyond just reporting a  $p$  value (Smith, 2020; Valeggia & Fernández-Duque, 2022).

## ACKNOWLEDGMENTS

The authors would like to thank all participants of the Adaptive Tools for Resilient Bones workshop. This workshop would not have been possible without the support of Leiden University Fund (W211115-6) and the Laboratory for Human Osteoarchaeology, Leiden University.

## CONFLICT OF INTEREST

Authors of this article have no conflicts of interest to declare.

## DATA AVAILABILITY STATEMENT

No data were used in this article.

## ORCID

Sarah A. Schrader  <https://orcid.org/0000-0003-0424-6748>

Jared Carballo Pérez  <https://orcid.org/0000-0001-6188-3183>

## REFERENCES

- Acosta, M. A., Henderson, C. Y., & Cunha, E. (2017). The effect of terrain on enthesal changes in the lower limbs. *International Journal of Osteoarchaeology*, 27, 828–838. <https://doi.org/10.1002/oa.2597>
- Alonso-Llamazares, C., Lopez, B., & Pardiñas, A. (2022). Sex differences in the distribution of enthesal changes: Meta-analysis of published evidence and its use in Bayesian paleopathological modeling. *American Journal of Biological Anthropology*, 177, 249–265. <https://doi.org/10.1002/ajpa.24425>

- Alves Cardoso, F. A., & Henderson, C. (2013). The categorisation of occupation in identified skeletal collections: A source of bias? *International Journal of Osteoarchaeology*, 23, 186–196. <https://doi.org/10.1002/oa.2285>
- Angel, J. L. (1966). *Early skeletons from tranquility California*. (Vol. 2) Smithsonian Press.
- Bakirci, S., Solmaz, D., Stephenson, W., Eder, L., Roth, J., & Aydin, S. Z. (2020). Enteseal changes in response to age, body mass index, and physical activity: An ultrasound study in healthy people. *The Journal of Rheumatology*, 47, 968–972. <https://doi.org/10.3899/jrheum.190540>
- Becker, S. K. (2019). Evaluating elbow osteoarthritis within the prehistoric Tiwanaku state using generalized estimating equations (GEE). *American Journal of Physical Anthropology*, 169, 186–196. <https://doi.org/10.1002/ajpa.23806>
- Bousquié, L., Karakostis, F. A., Crevecoeur, I., & Villotte, S. (2022). Technical note: Investigating activity-induced 3d hand enteseal variation in a documented South African sample. *Archaeological and Anthropological Sciences*, 14, 213. <https://doi.org/10.1007/s12520-022-01677-1>
- Campanacho, V., & Santos, A. L. (2013). Comparison of the enteseal changes of the os coxae of Portuguese males (19th–20th centuries) with known occupation. *International Journal of Osteoarchaeology*, 23, 229–236. <https://doi.org/10.1002/oa.2297>
- Carballo-Pérez, J., Sánchez-Cañadillas, E., Arnay-de-la-Rosa, M., Hernández-Marrero, J. C., & González-Reimers, E. (2021). Quotidian lives on isolated bodies: Enteseal changes and cross-sectional geometry among the aboriginal population of La Gomera (ca. 200–1500 AD, Canary Islands). *International Journal of Osteoarchaeology*, 31, 366–381. <https://doi.org/10.1002/oa.2956>
- Casado, A., Punsola, V., Gómez, M., de Diego, M., Barbosa, M., de Paz, F. J., Pastor, J. F., & Potau, J. M. (2019). Three-dimensional geometric morphometric analysis of the distal radius insertion sites of the palmar radiocarpal ligaments in hominoid primates. *American Journal of Physical Anthropology*, 170, 24–36. <https://doi.org/10.1002/ajpa.23885>
- Castro, A. A., Karakostis, F. A., Copes, L. E., McClendon, H. E., Trivedi, A. P., Schwartz, N. E., & Garland, T. Jr. (2022). Effects of selective breeding for voluntary exercise, chronic exercise, and their interaction on muscle attachment site morphology in house mice. *Journal of Anatomy*, 240, 279–295. <https://doi.org/10.1111/joa.13547>
- Djukic, K., Milovanovic, P., Hahn, M., Busse, B., Amling, M., & Djuric, M. (2015). Bone microarchitecture at muscle attachment sites: The relationship between macroscopic scores of entheses and their cortical and trabecular microstructural design. *American Journal of Physical Anthropology*, 157, 81–93. <https://doi.org/10.1002/ajpa.22691>
- Djukić, K., Milovanović, P., Milenković, P., & Djurić, M. (2020). A micro-architectural assessment of the gluteal tuberosity suggests two possible patterns in enteseal changes. *American Journal of Physical Anthropology*, 172, 291–299. <https://doi.org/10.1002/ajpa.24038>
- Dutour, O. (1986). Enthesopathies (lesions of muscular insertions) as indicators of the activities of Neolithic Saharan populations. *American Journal of Physical Anthropology*, 71, 221–224. <https://doi.org/10.1002/ajpa.1330710209>
- Godde, K., & Taylor, R. W. (2013). Distinguishing body mass and activity level from the lower limb: Can entheses diagnose obesity? *Forensic Science International*, 226, 303.e1–303.e7. <https://doi.org/10.1016/j.forsciint.2013.01.027>
- Godde, K., Wilson Taylor, R. W., & Gutierrez, C. (2018). Enteseal changes and demographic/health indicators in the upper extremity of modern Americans: Associations with age and physical activity. *International Journal of Osteoarchaeology*, 28, 285–293. <https://doi.org/10.1002/oa.2653>
- He, L. R., & de Almeida Prado, P. S. (2021). An evaluation of the relationship between the degree of enteseal changes and the severity of osteodegenerative processes at fibrocartilaginous entheses. *The Anatomical Record*, 304, 1255–1265. <https://doi.org/10.1002/ar.24541>
- Henderson, C. Y., & Alves Cardoso, F. (2013). Special issue enteseal changes and occupation: Technical and theoretical advances and their applications. *International Journal of Osteoarchaeology*, 23, 127–134. <https://doi.org/10.1002/oa.2298>
- Henderson, C. Y., & Nikita, E. (2016). Accounting for multiple effects and the problem of small sample sizes in osteology: A case study focussing on enteseal changes. *Archaeological and Anthropological Sciences*, 8, 805–817. <https://doi.org/10.1007/s12520-015-0256-1>
- Henderson, C. Y., Mariotti, V., Pany-Kucera, D., Villotte, S., & Wilczak, C. (2013). Recording specific enteseal changes of fibrocartilaginous entheses: Initial tests using the Coimbra method. *International Journal of Osteoarchaeology*, 23, 152–162. <https://doi.org/10.1002/oa.2287>
- Henderson, C. Y., Mariotti, V., Pany-Kucera, D., Villotte, S., & Wilczak, C. (2016). The new 'Coimbra method': A biologically appropriate method for recording specific features of fibrocartilaginous enteseal changes. *International Journal of Osteoarchaeology*, 26, 925–932. <https://doi.org/10.1002/oa.2477>
- Henderson, C. Y., Mariotti, V., Santos, F., Villotte, S., & Wilczak, C. A. (2017). The new Coimbra method for recording enteseal changes and the effect of age-at-death. *Bulletins et Mémoires de la Société d'Anthropologie de Paris*, 29, 140–149. <https://doi.org/10.1007/s13219-017-0185-x>
- Jurmain, R., Cardoso, F. A., Henderson, C., & Villotte, S. (2012). Bioarchaeology's Holy Grail: The reconstruction of activity. In A. L. Grauer (Ed.), *A companion to paleopathology*. Wiley-Blackwell. <https://doi.org/10.1002/97811444345940.ch29>
- Karakostis, F. A., Haeufle, D., Anastopoulou, I., Moraitis, K., Hotz, G., Tourloukis, V., & Harvati, K. (2021). Biomechanics of the human thumb and the evolution of dexterity. *Current Biology*, 31, 1317–1325.e8. <https://doi.org/10.1016/j.cub.2020.12.041>
- Karakostis, F. A., Hotz, G., Scherf, H., Wahl, J., & Harvati, K. (2017). Occupational manual activity is reflected on the patterns among hand entheses. *American Journal of Physical Anthropology*, 164, 30–40. <https://doi.org/10.1002/ajpa.23253>
- Karakostis, F. A., Hotz, G., Scherf, H., Wahl, J., & Harvati, K. (2018). A repeatable geometric morphometric approach to the analysis of hand enteseal three-dimensional form. *American Journal of Physical Anthropology*, 166, 246–260. <https://doi.org/10.1002/ajpa.23421>
- Karakostis, F. A., Jeffery, N., & Harvati, K. (2019). Experimental proof that multivariate patterns among muscle attachments (entheses) can reflect repetitive muscle use. *Scientific Reports*, 9, 16577. <https://doi.org/10.1038/s41598-019-53021-8>
- Karakostis, F. A., Reyes-Centeno, H., Franken, M., Hotz, G., Rademaker, K., & Harvati, K. (2021). Biocultural evidence of precise manual activities in an Early Holocene individual of the high-altitude Peruvian Andes. *American Journal of Physical Anthropology*, 174, 35–48. <https://doi.org/10.1002/ajpa.24160>
- Karakostis, F. A., Wallace, I. J., Konow, N., & Harvati, K. (2019). Experimental evidence that physical activity affects the multivariate associations among muscle attachments (entheses). *Journal of Experimental Biology*, 222, jeb213058. <https://doi.org/10.1242/jeb.213058>
- Kennedy, K. A. R. (1983). Morphological variations in ulnar supinator crests and fossae as identifying markers of occupational stress. *Journal of Forensic Science*, 28, 871–876.
- Kubicka, A. M., & Mysza, A. (2020). Are enteseal changes and cross-sectional properties associated with the shape of the upper limb? *American Journal of Physical Anthropology*, 173, 293–306. <https://doi.org/10.1002/ajpa.24096>
- Laffranchi, Z., Charisi, D., Jiménez-Brobeil, S. A., & Milella, M. (2020). Gendered division of labor in a Celtic community? A comparison of sex differences in enteseal changes and long bone shape and robusticity in the pre-Roman population of Verona (Italy, third–first century BC). *American Journal of Physical Anthropology*, 173, 568–588. <https://doi.org/10.1002/ajpa.24111>

- Lieverse, A. R., Bazaliiskii, V. I., Goriunova, O. I., & Weber, A. W. (2013). Lower limb activity in the Cis-Baikal: Enteseal changes among middle holocene siberian foragers. *American Journal of Physical Anthropology*, 150, 421–432. <https://doi.org/10.1002/ajpa.22217>
- Listi, G. A. (2016). The use of enteseal changes in the femur and os coxa for age assessment. *Journal of Forensic Sciences*, 61, 12–18. <https://doi.org/10.1111/1556-4029.12905>
- Mazza, B. (2019). Enteseal changes among late Holocene hunter-gatherers from the southern extreme of La Plata basin (Argentina). *Archaeological and Anthropological Sciences*, 11, 1865–1885. <https://doi.org/10.1007/s12520-018-0638-2>
- Merbs, C. (1983). *Patterns of activity-induced pathology in a Canadian Inuit population (Vol. Archaeological Survey of Canada Paper No. 119)*. National Museums of Canada. <https://doi.org/10.2307/j.ctv173qw>
- Michopoulou, E., Nikita, E., & Henderson, C. Y. (2017). A test of the effectiveness of the Coimbra method in capturing activity-induced enteseal changes. *International Journal of Osteoarchaeology*, 27, 409–417. <https://doi.org/10.1002/oa.2564>
- Michopoulou, E., Nikita, E., & Valakos, E. D. (2015). Evaluating the efficiency of different recording protocols for enteseal changes in regards to expressing activity patterns using archival data and cross-sectional geometric properties. *American Journal of Physical Anthropology*, 158, 557–568. <https://doi.org/10.1002/ajpa.22822>
- Milella, M. (2014). The influence of life history and sexual dimorphism on enteseal changes in modern humans and African great apes. *PLoS ONE*, 9, e107963. <https://doi.org/10.1371/journal.pone.0107963>
- Milella, M., Cardoso, F. A., Assis, S., Lopreno, G. P., & Speith, N. (2015). Exploring the relationship between enteseal changes and physical activity: A multivariate study. *American Journal of Physical Anthropology*, 156, 215–223. <https://doi.org/10.1002/ajpa.22640>
- Milella, M., Giovanna Belcastro, M., Zollikofer, C. P. E., & Mariotti, V. (2012). The effect of age, sex, and physical activity on enteseal morphology in a contemporary Italian skeletal collection. *American Journal of Physical Anthropology*, 148, 379–388. <https://doi.org/10.1002/ajpa.22060>
- Myszka, A., Krenz-Niedbala, M., Tomczyk, J., & Zalewska, M. (2020). Osteoarthritis: A problematic disease in past human populations. A dependence between enteseal changes, body size, age, sex, and osteoarthritic changes development. *The Anatomical Record*, 303, 2357–2371. <https://doi.org/10.1002/ar.24316>
- Niinimäki, S. (2011). What do muscle marker ruggedness scores actually tell us? *International Journal of Osteoarchaeology*, 21, 292–299. <https://doi.org/10.1002/oa.1134>
- Niinimäki, S. (2012). The relationship between musculoskeletal stress markers and biomechanical properties of the humeral diaphysis. *American Journal of Physical Anthropology*, 147, 618–628. <https://doi.org/10.1002/ajpa.22023>
- Niinimäki, S., & Baiges Sotos, L. (2013). The relationship between intensity of physical activity and enteseal changes on the lower limb. *International Journal of Osteoarchaeology*, 23, 221–228. <https://doi.org/10.1002/oa.2295>
- Niinimäki, S., & Salmi, A.-K. (2016). Enteseal changes in free-ranging versus zoo reindeer—Observing activity status of reindeer. *International Journal of Osteoarchaeology*, 26, 314–323. <https://doi.org/10.1002/oa.2423>
- Niinimäki, S., & Salmi, A.-K. (2021). Covariation between enteseal changes and cross-sectional properties of reindeer long bones—Considering bone functional adaptation as partial contributing factor. *Journal of Archaeological Science: Reports*, 36, 102840. <https://doi.org/10.1016/j.jasrep.2021.102840>
- Nikita, E. (2014). The use of generalized linear models and generalized estimating equations in bioarchaeological studies. *American Journal of Physical Anthropology*, 153, 473–483. <https://doi.org/10.1002/ajpa.22448>
- Nikita, E., Xanthopoulou, P., Bertatos, A., Chovalopoulou, M.-E., & Hafez, I. (2019). A three-dimensional digital microscopic investigation of enteseal changes as skeletal activity markers. *American Journal of Physical Anthropology*, 169, 704–713. <https://doi.org/10.1002/ajpa.23850>
- Nolte, M., & Wilczak, C. (2013). Three-dimensional surface area of the distal biceps enthesis, relationship to body size, sex, age and secular changes in a 20th century American sample. *International Journal of Osteoarchaeology*, 23, 163–174. <https://doi.org/10.1002/oa.2292>
- Palmer, J. L. A., Hoogland, M. H. L., & Waters-Rist, A. L. (2016). Activity reconstruction of post-medieval Dutch rural villagers from upper limb osteoarthritis and enteseal changes. *International Journal of Osteoarchaeology*, 26, 78–92. <https://doi.org/10.1002/oa.2397>
- Palmer, J. L. A., Quintelier, K., Inskip, S., & Waters-Rist, A. L. (2019). A comparison of two methods for recording enteseal change on a post-medieval urban skeletal collection from Aalst (Belgium). *Archaeometry*, 61, 211–225. <https://doi.org/10.1111/arc.12409>
- Peterson, J., & Hawkey, D. E. (1998). Preface. *International Journal of Osteoarchaeology*, 8, 303–304. [https://doi.org/10.1002/\(SICI\)1099-1212\(199809\)8:5%3C303::AID-OA450%3E3.0.CO;2-N](https://doi.org/10.1002/(SICI)1099-1212(199809)8:5%3C303::AID-OA450%3E3.0.CO;2-N)
- Refai, O. (2019). Enteseal changes in ancient Egyptians from the pyramid builders of Giza—Old kingdom. *International Journal of Osteoarchaeology*, 29, 513–524. <https://doi.org/10.1002/oa.2748>
- Robb, J. E. (1998). The interpretation of skeletal muscle sites: A statistical approach. *International Journal of Osteoarchaeology*, 8, 363–377. [https://doi.org/10.1002/\(SICI\)1099-1212\(199809\)8:5<363::AID-OA438>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1099-1212(199809)8:5<363::AID-OA438>3.0.CO;2-K)
- Salega, S., & Grosskopf, B. (2022). Evaluation of enteseal changes in a modern identified skeletal collection from Inden (Germany). *International Journal of Osteoarchaeology*, 32, 86–99. <https://doi.org/10.1002/oa.3046>
- Salmi, A.-K., Niinimäki, S., & Pudas, T. (2020). Identification of working reindeer using palaeopathology and enteseal changes. *International Journal of Paleopathology*, 30, 57–67. <https://doi.org/10.1016/j.ijpp.2020.02.001>
- Santana-Cabrera, J., Velasco-Vázquez, J., & Rodríguez-Rodríguez, A. (2015). Enteseal changes and sexual division of labor in a North-African population: The case of the pre-Hispanic period of the Gran Canaria Island (11th–15th c. CE). *Homo*, 66, 118–138. <https://doi.org/10.1016/j.jchb.2014.10.005>
- Schrader, S. A. (2012). Activity patterns in New Kingdom Nubia: An examination of enteseal remodeling and osteoarthritis at Tombos. *American Journal of Physical Anthropology*, 149, 60–70. <https://doi.org/10.1002/ajpa.22094>
- Schrader, S. A. (2015). Elucidating inequality in Nubia: An examination of enteseal changes at Kerma (Sudan). *American Journal of Physical Anthropology*, 156, 192–202. <https://doi.org/10.1002/ajpa.22637>
- Schrader, S. A., & Buzon, M. R. (2017). Everyday life after collapse: A bioarchaeological examination of enteseal change and accidental injury in postcolonial Nubia. *Bioarchaeology International*, 1, 19–34. <https://doi.org/10.5744/bi.2017.1000>
- Smith, R. J. (2020). P>.05: The incorrect interpretation of “not significant” results is a significant problem. *American Journal of Biological Anthropology*, 172, 521–527. <https://doi.org/10.1002/ajpa.24092>
- Thomas, A. (2014). Bioarchaeology of the middle Neolithic: Evidence for archery among early European farmers. *American Journal of Physical Anthropology*, 154, 279–290. <https://doi.org/10.1002/ajpa.22504>
- Valeggia, C. R., & Fernández-Duque, E. (2022). Moving biological anthropology research beyond p < 0.05. *American Journal of Biological Anthropology*, 177, 193–195. <https://doi.org/10.1002/ajpa.24444>
- Villotte, S., & Knüsel, C. J. (2013). Understanding enteseal changes: Definition and life course changes. *International Journal of Osteoarchaeology*, 23, 135–146. <https://doi.org/10.1002/oa.2289>
- Villotte, S., Assis, S., Cardoso, F. A., Henderson, C. Y., Mariotti, V., Milella, M., Pany-Kucera, D., Speith, N., Wilczak, C. A., & Jurmain, R.



- (2016). In search of consensus: Terminology for enthesal changes (EC). *International Journal of Paleopathology*, 13, 49–55. <https://doi.org/10.1016/j.ijpp.2016.01.003>
- Villotte, S., Castex, D., Couallier, V., Dutour, O., Knüsel, C. J., & Henry-Gambier, D. (2010). Enthesopathies as occupational stress markers: Evidence from the upper limb. *American Journal of Physical Anthropology*, 142, 224–234. <https://doi.org/10.1002/ajpa.21217>
- Wallace, I. J., Winchester, J. M., Su, A., Boyer, D. M., & Konow, N. (2017). Physical activity alters limb bone structure but not enthesal morphology. *Journal of Human Evolution*, 107, 14–18. <https://doi.org/10.1016/j.jhevol.2017.02.001>
- Yonemoto, S. (2016). Differences in the effects of age on the development of enthesal changes among historical Japanese populations. *American Journal of Physical Anthropology*, 159, 267–283. <https://doi.org/10.1002/ajpa.22870>

**How to cite this article:** Schrader, S. A., & Carballo Pérez, J. (2023). Special issue adaptive tools for resilient bones: Biostatistical approaches to past physical activity in osteoarchaeology. *International Journal of Osteoarchaeology*, 33(3), 381–388. <https://doi.org/10.1002/oa.3177>