



Universiteit
Leiden
The Netherlands

Cardiometabolic determinants of cognitive function in later life: unravelling the roles of risk factors

Zonneveld, M.H.

Citation

Zonneveld, M. H. (2024, February 15). *Cardiometabolic determinants of cognitive function in later life: unravelling the roles of risk factors*. Retrieved from <https://hdl.handle.net/1887/3717601>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3717601>

Note: To cite this publication please use the final published version (if applicable).



6

Weight loss, visit-to-visit body weight variability and cognitive function in older individuals

Michelle H. Zonneveld, Raymond Noordam, Behnam Sabayan, David J. Stott, Simon P. Mooijaart, Gerard J. Blauw, J. Wouter Jukema, Naveed Sattar, Stella Trompet

Age and ageing, vol. 52,1 (2023): afac312. doi:10.1093/ageing/afac312

Abstract

Objective

To investigate the association between variability and loss of body weight with subsequent cognitive performance and activities of daily living in older individuals.

Design

Cross-sectional cohort study.

Setting

PROspective Study of Pravastatin in the Elderly at Risk (PROSPER), multicentre trial with participants from Scotland, Ireland and the Netherlands.

Subjects

4,309 participants without severe cognitive dysfunction (mean age 75.1 years, SD=3.3), at higher risk for cardiovascular disease.

Methods

Body weight was measured every three months for 2.5 years. Weight loss is defined as an average slope across all weight measurements and as $\geq 5\%$ decrease in baseline body weight during follow-up. Visit-to-visit variability is defined as the SD of weight measurements (kg) between visits. Four tests of cognitive function were examined: Stroop test, Letter-Digit Coding test, immediate and delayed Picture-Word learning tests. Two measures of daily living activities: Barthel Index (BI) and instrumental activities of daily living (IADL). We examined all tests at month 30.

Results

Both larger body weight variability and loss of $\geq 5\%$ of baseline weight were independently associated with worse scores on all cognitive tests, but minimally with BI and IADL. Compared to participants with stable weight, participants with significant weight loss scored 5.83 seconds (95%CI 3.74; 7.92) slower on the Stroop test, coded 1.72 digits less (95%CI -2.21; -1.13) on the Letter-Digit Coding test, and remembered 0.71 less pictures (95%CI -0.93; -0.48) on the delayed Picture-Word Learning test.

Conclusion

In older people at higher risk for CVD, weight loss and variability are independent risk-factors for worse cognitive function.

1. Introduction

The process of ageing is accompanied by fluctuations in homeostatic processes, resulting in intrinsic intraindividual variability in physiological parameters. For example, variability in weight, including both gaining and losing weight, is associated with significant increases in mortality [1, 2]. Several observational studies have demonstrated that *unintentional* weight loss in older adults is related to increased frailty and functional decline [2-5]. The origin of unintentional weight loss in older adults is often linked to the manifestation of malignancies but numerous social, behavioural and health factors may also be instrumental [6, 7]. Over 27% of frail older individuals above the age of 65 years' experience unintentional weight loss [8], where a specific cause cannot be found in as many as 25% of cases [2].

Recent evidence indicates that unintentional weight loss in older individuals associates with brain atrophy and cognitive impairment, which are known to associate with Alzheimer's Disease [3, 6, 9]. Other markers of unstable homeostasis and intraindividual variability, including variability in systolic blood pressure and low-density lipoprotein cholesterol (LDL-C), have also been associated with worsened cognitive function [10, 11]. We hypothesized that larger variation in body weight, as well as loss of weight, are independently associated with lower cognitive performance, independent of baseline body mass index (BMI) and traditional risk factors. Therefore, in the present study, we investigated the association of 2.5-years body weight loss and variation in body weight with subsequent cognitive performance and activities of daily living in a cohort of older individuals at increased risk of cardiovascular disease (CVD) but without severe cognitive dysfunction at baseline.

2. Methods

Study design and participants

The data employed in the present study originates from the PROspective Study of Pravastatin in the Elderly at Risk (PROSPER). 5,804 men and women aged 70-82 years from three countries (the Netherlands, Scotland, Ireland) were enrolled between December 1997 and May 1999 in a prospective, multicentre randomized trial in order to assess the safety and efficacy of pravastatin in reducing the risk of major vascular events. Participants were eligible for enrolment if they had pre-existing vascular disease or increased risk due to smoking, hypertension or diabetes.

During recruitment, the following exclusion criteria were: cognitive impairment (Mini-Mental Score Examination score <24); history of malignancy within the past 5 years except localized basal cell carcinoma; recent stroke, transient ischemic attack, myocardial infarction, surgery or amputation for vascular disease ≤ 6 months before study entry. More details regarding exclusion criteria of PROSPER have been described elsewhere [12, 13]. The PROSPER study was approved by the Medical Ethics Committees of the three collaborating centres and complied with the Declaration of Helsinki. All participants gave written informed consent.

In the present study, the following inclusion criteria were applied: ≥ 1 out of the 4 cognitive tests scores at month 30 of follow-up; ≥ 2 weight measurements recorded between baseline and month 30 of follow-up with a maximum of 11 repetitive measurements.

Data collection

Exposure variables

Participants in the PROSPER study were reviewed every three months and body weight was measured at each visit, resulting in a maximum of 11 repetitive weight measurements.

From the weight collected at baseline and follow-up visits, we computed two determinants: weight loss and weight variability. Visit-to-visit weight variability was calculated by means of the intraindividual standard deviation (S.D.) over each individual's measurements between baseline and every 3 months up to 30 months of follow-up. Weight loss was defined using two methods: first by calculating an average slope across all weight measurements, followed by calculating a delta incorporating only the first and last weight measurements during follow-up. The average slope incorporating all weight assessments during follow-up was computed using a linear mixed model capable of handling missing data during the visits. By fitting a linear-mixed model over the various measurements during follow-up, an average slope of weight change was calculated per participant. Then, the delta of weight change from baseline to month 30 was calculated and defined according to the following classification: weight loss as $\geq 5\%$ of body weight decreased from baseline to month 30; weight gain as $\geq 5\%$ of weight increased; and stability if within $< 5\%$ weight variation between baseline and month 30 of follow-up [2, 6]. Although no universally accepted definition of clinically relevant weight loss exists, previous observational studies have employed this definition [2, 6].

Cognitive function measurements

Serving as our outcome measurements, cognitive function was assessed face-to-face using four neuropsychological performance tests. More details regarding the tests are described in detail elsewhere [14]. The Stroop colour-word inference test (selective attention) and the Letter-Digit Coding Test (LDCT) (processing speed) were used to measure executive functioning [14]. The outcome parameter for the Stroop test was the total number of seconds to complete the third Stroop card. The outcome variable for the LDCT was the total number of correct entries in 60 seconds. Visual episodic memory was assessed with the 15-Picture Learning test (PLT) testing immediate and delayed recall. The main outcome was the accumulated number of recalled pictures over the three learning trials and the number of pictures recalled after 20 minutes. Functional status was assessed using the BI (Barthel Index) and IADL (Instrumental activities of daily living). BI assesses self-care activities of daily living using 10 items (such as dressing) where a higher score indicates higher independence, with a maximum of 20 points [15]. Similarly, IADL measures activities of daily living using 7 items with a maximum of 14 points, but in addition includes the interaction with the social and physical environment [16]. Likewise, here a higher score also higher functional capacity

and independence. All outcomes were assessed at month 30 to maximize the availability of outcomes after the measurement of weight variability and weight change.

Covariates

Covariates were obtained from an extensive medical history using routine care data during a 10-week screening period. The participant's general practitioner reported on history of various clinical diseases, including CVD, diabetes mellitus, transient ischemic attack, stroke and myocardial infarction. The following were evaluated using a medical inventory: number of medications used, use of diuretics and antidepressants, years of education, smoking status, alcohol intake. Systolic and diastolic blood pressure were measured every 3 months [10]. Data on diseases developed during follow-up were ascertained using hospital records and general practitioners' records, and included: diabetes; non-fatal cardiovascular and coronary events including myocardial infarction, stroke and transient ischemic attack; hospitalization due to heart failure; serious non-fatal cancer [13]. These endpoints were classified by an independent committee. Data on these baseline covariates were complete for all participants.

Statistical analyses

Baseline characteristics of the study participants are reported as mean (standard deviation) or median (interquartile range) for continuous variables and number (percentage) for categorical variables. To investigate whether slope and variability were two independent phenotypes, we assessed the correlation with a Pearson's correlation coefficient. We repeated the analyses with both slope of weight change and weight variability in the same multivariable-adjusted regression model to study whether these phenotypes have independent effects.

The associations of visit-to-visit variability in total weight (in SD), slope of weight change (in kg), and weight loss (in categories) with measures of cognitive function at month 30 were assessed using multivariable-adjusted linear regression models. Weight variability and slope of weight change were analysed as a continuous variable and in equally-sized thirds, where the lower third of weight variability, and the middle third of the slope of weight change, were defined as reference categories. For reasons of clinical interpretation, we presented the results on slope of weight change as the difference in cognitive function at month 30 per extra 0.1 kg/month weight loss.

The linear regression analyses were adjusted for covariates based on their biological plausibility as potential confounders. Therefore, in the minimally adjusted model, we included sex, age, country as a 3-level variable, mean weight during follow-up, height at baseline, and years of education. The fully adjusted model additionally included smoking status, alcohol intake, number of medications used, use of diuretics and antidepressants, and history of CVD, diabetes, and myocardial infarction.

Data is reported as the mean multivariable-adjusted difference in outcome between the second and third thirds of weight change in comparison with the reference group, with accompanying 95% confidence interval. For example, "weight lost" and "weight gained" are

compared to the reference group “stable weight”. All analyses were performed using SPSS Windows version 26 (IBM Corp., 2019).

Sensitivity analyses

In addition to these overall analyses, we performed sensitivity analyses to investigate the robustness of the associations considering different subgroups of the population. First, we performed separate analyses for placebo and pravastatin treatment groups. Next, it has been shown that high blood pressure variability was associated with worse cognitive function [10]. As variability in weight and blood pressure may have a common cause, we additionally adjusted our models for systolic blood pressure variability and mean systolic blood pressure from baseline to month 30. This allows for the separation of effects of weight variability from those originating from blood pressure variability. Systolic blood pressure variability was defined as the intraindividual standard deviation from baseline to month 30, where blood pressure was measured every 3 months, as previously done [10, 17]. Furthermore, we included both weight loss (slope) and visit-to-visit body weight variables in the same multivariable-adjusted linear regression model to test independence of the two phenotypes. Last, we performed analyses excluding individuals who developed any of the following diseases during follow-up, to ensure weight loss did not follow as a result: incident diabetes; non-fatal cancer; non-fatal stroke or transient ischemic attack; hospitalization due to heart failure; non-fatal coronary or cardiovascular events.

3. Results

After excluding participants with <2 measurements (n=225) and participants without cognitive test scores at month 30 (n=1,270), 4,309 participants were eligible for inclusion (Table 1). The mean age was 75.1 years (SD=3.3) and more than half of the study population was female (n=2,222, 51.6%). Large majority of participants had a history of hypertension (n=2,706, 62.8%). The mean weight during follow-up was 72.9 kilograms (SD=13.3) and participants had a median weight loss of 0.01 kilograms per month (IQR -0.07; 0.06). Baseline weight characteristics per third of weight loss and weight variability are reported in supplementary table 1.

The correlation between continuous weight variability and the continuous slope of weight change was negligible (Pearson's $r=0.22$).

Table 1. Demographics and clinical characteristics of study population

Sociodemographics	All (N=4309)
Age, y, mean (SD)	75.1 (3.3)
Female, n (%)	2222 (51.6)
Age left school, y, mean (SD)	15.2 (2.1)
Current smoker, n (%)	1079 (25.0)
Alcohol intake, unit intake per week, mean (SD)	5.30 (9.2)
Cardiovascular risk factors	
History of CVD, n (%)	1878 (43.6)
History of hypertension, n (%)	2706 (62.8)
History of stroke or TIA, n (%)	465 (10.8)
History of myocardial infarction, n (%)	560 (13.0)
Diabetes mellitus, n (%)	446 (10.4)
Serum cholesterol, mmol/L, mean (SD)	5.68 (0.9)
Weight during follow-up, kilograms, mean (SD)	72.9 (13.3)
Weight change, kg/month, mean (SD)	-0.01 (0.1)
Lost more than 5% of baseline body weight during follow-up, n (%)	802 (18.6)
Gained more than 5% of baseline body weight during follow-up, n (%)	580 (13.5)
Number of weight measurements, median (IQR)	10 (10; 10)
Body mass index, kg/m ² , mean (SD)	26.9 (4.1)
Pravastatin treatment, n (%)	2146 (49.8)
Number of medications, median (IQR)	3 (2; 5)
Use of diuretics, n (%)	1804 (41.9)
Cognitive function at month 30 of follow-up	
Stroop test, seconds, mean (SD) ¹	64.5 (26.1)
LDCT, digits coded, mean (SD) ²	22.9 (7.8)
PLTi, pictures remembered, mean (SD) ³	9.5 (2.0)
PLTd, pictures remembered, mean (SD) ³	10.2 (2.9)
Barthel, index, mean (SD) ⁴	19.7 (0.9)
IADL, points, mean (SD) ⁴	13.5 (1.3)

Abbreviations: CVD = cardiovascular disease; TIA = transient ischemic attack; SD = standard deviation; IQR = interquartile range; LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed. ¹=performed by 4122 participants; ²=performed by 4264 participants; ³=performed by 4309 participants; ⁴=performed by 4428 participants.

Association between weight loss and cognitive function

Table 2 displays the association of the slope of weight change and cognitive function. After full adjustments, in comparison to the middle third, the lower third of the slope of weight change was associated with a worse performance on all cognitive and functional domains except on the BI (-0.04 points, 95% CI -0.10; 0.03). To illustrate; at month 30 of follow-up, the lower third coded 1.42 (95% CI -1.98; -0.86) digits less on the LDCT and performed 4.39 (95% CI 2.42; 6.37) seconds slower on the Stroop test. On the other hand, in comparison to the middle third, the upper third of the slope of weight change was not significantly associated with cognitive performance. Continuously, the slope of weight change was also associated with worse cognitive function on all tests. Per 0.10 kg/month additional average weight loss, the score on the Stroop test was 1.82 (95% CI 1.13; 2.49) seconds slower, 0.70 less (95% CI -0.90; -0.51) digits were coded on the LDCT, and 0.21 less (95% CI -0.29; -0.14) pictures were remembered on the delayed PLT.

Loss of $\geq 5\%$ of weight during follow-up was associated with worse performance on all domains, but did not show association with BI and IADL (Table 3 and Figure 1). After full adjustments, in comparison to maintaining stable weight during follow-up, participants who lost $\geq 5\%$ of baseline body weight performed 5.83 (95% CI 3.74; 7.92) seconds slower on the Stroop test. Furthermore, weight loss was also associated with a worse performance on the LDCT (Beta -1.72 digits coded, 95% CI -2.21; -1.13) and the PLT, both immediate (Beta -0.48 pictures remembered, 95% CI -0.64; -0.33) and delayed (Beta -0.71 pictures remembered, 95% CI -0.93; -0.48). In comparison to individuals who maintained stable weight, we did not find evidence of a significant association between weight gain during follow-up and cognitive function.

Table 2. Associations of weight change (slope) during follow-up time and cognitive function at month 30 of follow-up

Cognitive test	Weight change (slope)				Continuous ³ All (N _{max} = 4309) Beta (95% CI)
	Low third	Middle third	Upper third		
	(N _{max} = 1436) Beta (95% CI)	(N _{max} = 1436) Beta (95% CI)	(N _{max} = 1437) Beta (95% CI)		
Minimally adjusted¹					
Stroop, seconds	4.75 (2.84; 6.67)	Ref	0.34 (-1.52; 2.19)		1.91 (1.25; 2.57)
LDCT, digits coded	-1.52 (-2.07; -0.97)	Ref	0.32 (-0.21; 0.86)		-0.69 (-0.88; -0.50)
PLTi, pictures remembered	-0.37 (-0.52; -0.23)	Ref	0.01 (-0.13; 0.15)		-0.15 (-0.20; -0.10)
PLTd, pictures remembered	-0.56 (-0.77; -0.36)	Ref	-0.06 (-0.26; 0.14)		-0.21 (-0.28; -0.14)
Barthel, index	-0.04 (-0.10; 0.03)	Ref	0.00 (-0.06; 0.07)		-0.02 (-0.04; 0.00)
IADL, points	-0.11 (-0.20; -0.02)	Ref	-0.07 (-0.16; 0.02)		-0.04 (-0.07; 0.01)
Fully adjusted²					
Stroop, seconds	4.39 (2.42; 6.37)	Ref	0.15 (-1.78; 2.06)		1.82 (1.13; 2.49)
LDCT, digits coded	-1.42 (-1.98; -0.86)	Ref	0.43 (-0.12; 0.97)		-0.70 (-0.90; -0.51)
PLTi, pictures remembered	-0.37 (-0.52; -0.22)	Ref	0.02 (-0.12; 0.17)		-0.15 (-0.20; -0.10)
PLTd, pictures remembered	-0.52 (-0.73; -0.31)	Ref	-0.01 (-0.22; 0.20)		-0.21 (-0.29; -0.14)
Barthel, index	-0.04 (-0.10; 0.03)	Ref	0.00 (-0.06; 0.07)		-0.02 (-0.05; 0.00)
IADL, points	-0.11 (-0.21; -0.02)	Ref	-0.07 (-0.16; 0.02)		-0.04 (-0.07; -0.01)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, mean weight during follow-up, height, education. ²=sex, age, country, education level, height, mean weight during follow-up time, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, number of medications, use of diuretics, use of antidepressants. ³ Presented results can be interpreted as the difference is cognitive performance at month 30 per 0.1 kg/month additional weight loss. Stroop test was performed by 4122 participants, Letter-Digit Coding test was performed by 4264 participants, and the Picture-Word Learning immediate and delayed tests were performed by 4309 participants. Items written in bold indicate that the 95% confidence interval does not contain zero.

Table 3. Associations of weight change during follow-up time and cognitive function at month 30 of follow-up

Cognitive test	Weight change		
	Weight lost	Weight stable	Weight gained
	(N _{max} =802)	(N _{max} =2927)	(N _{max} =644)
	Beta (95% CI)	Beta (95% CI)	Beta (95% CI)
Minimally adjusted¹			
Stroop, seconds	6.34 (4.31; 8.37)	Ref	-0.74 (-3.02; 1.54)
LDCT, digits coded	-1.85 (-2.43; -1.27)	Ref	0.52 (-0.14; 1.17)
PLTi, pictures remembered	-0.48 (-0.64; -0.33)	Ref	0.02 (-0.15; 0.20)
PLTd, pictures remembered	-0.74 (-0.96; -0.52)	Ref	-0.08 (-0.32; 0.17)
Barthel, index	-0.02 (-0.09; 0.04)	Ref	-0.02 (-0.09; 0.06)
IADL, points	-0.09 (-0.19; 0.01)	Ref	-0.07 (-0.18; 0.04)
Fully adjusted²			
Stroop, seconds	5.83 (3.74; 7.92)	Ref	-1.28 (-3.62; 1.06)
LDCT, digits coded	-1.72 (-2.21; -1.13)	Ref	0.78 (0.12; 1.45)
PLTi, pictures remembered	-0.48 (-0.64; -0.33)	Ref	0.06 (-0.12; 0.23)
PLTd, pictures remembered	-0.71 (-0.93; -0.48)	Ref	0.01 (-0.24; 0.26)
Barthel, index	-0.02 (-0.09; 0.05)	Ref	-0.01 (-0.09; 0.07)
IADL, points	-0.08 (-0.18; 0.02)	Ref	-0.05 (-0.16; 0.07)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, mean weight during follow-up, height, education. ²=sex, age, country, education level, height, mean weight during follow-up time, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, number of medications, use of diuretics, use of antidepressants. Weight loss was defined as $\geq 5\%$ of body weight decreased from baseline to month 30; weight gain as $\geq 5\%$ of weight increased; and stability if within $< 5\%$ weight variation between baseline and month 30 of follow-up. Stroop test was performed by 4122 participants, Letter-Digit Coding test was performed by 4264 participants, and the Picture-Word Learning immediate and delayed tests were performed by 4309 participants. Items written in bold indicate that the 95% confidence interval does not contain zero.




Figure 1. Fully adjusted associations of weight change during follow-up time and cognitive function at month 30, where weight loss was defined as $\geq 5\%$ of body weight decreased from baseline to month 30; weight gain as $\geq 5\%$ of weight increased; and stability if within $< 5\%$ weight variation between baseline and month 30 of follow-up.

Association between visit-to-visit body weight variability and cognitive function

The associations of visit-to-visit weight variability and cognitive function are presented in Table 4. After full adjustments, both middle and upper thirds of weight variability, in comparison to the lower third, performed worse on all tests. For example, the middle third (moderate variability) of weight variability was associated with 1.93 seconds (95% CI 0.01; 3.86) slower performance on the Stroop test, whilst the upper third (high variability) was associated with 4.52 (95% CI 2.52; 6.51) seconds slower performance, in comparison to the lower third (low variability). Likewise, higher weight variability as a continuous variable was associated with worse cognitive function on all domains. Here, a one-SD higher weight variability was associated with 1.46 seconds (95% CI 0.82; 2.09) slower performance on the Stroop test. There were no associations with functional capacity.

Sensitivity analyses

Adjusting the analyses between weight variability and cognitive function for systolic blood pressure variability did not materially change the results (Table 4). The results were essentially unchanged when stratifying for the treatment groups (Supplementary tables 2-4). Repeating the analyses after excluding participants with incident disease-states during follow-up also did not materially change the associations (Supplementary tables 5-7). The associations between continuous weight variability and continuous slope of weight change with cognitive function remained significant when including the two determinants in the same model (Supplementary table 8).

Table 4. Associations of weight variability (SD) and cognitive function at month 30 of follow-up

Weight variability, SD					
	Low third	Middle third	Upper third	Continuous	
	(N _{max} = 1436)	(N _{max} = 1436)	(N _{max} = 1437)	All (N _{max} = 4309)	
Cognitive test	Beta (95% CI)	Beta (95% CI)	Beta (95% CI)	Beta (95% CI)	
Minimally adjusted¹					
Stroop, s	Ref	2.30 (0.42; 4.18)	6.01 (4.07; 7.95)	1.98 (1.35; 2.60)	
LDCT, digits coded	Ref	-1.42 (-1.96; -0.88)	-2.29 (-2.85; -1.74)	-0.73 (-0.91; -0.55)	
PLTi, pictures remembered	Ref	-0.21 (-0.35; -0.06)	-0.63 (-0.78; -0.49)	-0.19 (-0.24; -0.14)	
PLTd, pictures remembered	Ref	-0.21 (-0.35; -0.06)	-0.96 (-1.16; -0.75)	-0.29 (-0.35; -0.22)	
Barthel, index	Ref	-0.04 (-0.10; 0.03)	-0.09 (-0.15; -0.02)	-0.06 (-0.08; -0.04)	
IADL, points	Ref	-0.04 (-0.13; 0.05)	-0.19 (-0.28; -0.10)	-0.10 (-0.13; -0.07)	
Fully adjusted²					
Stroop, s	Ref	2.25 (0.31; 4.18)	5.32 (0.31; 4.18)	1.72 (1.08; 2.35)	
LDCT, digits coded	Ref	-1.36 (-1.92; -0.81)	-2.16 (-2.72; -1.59)	-0.67 (-0.86; -0.49)	
PLTi, pictures remembered	Ref	-0.19 (-0.33; -0.04)	-0.61 (-0.76; -0.46)	-0.18 (-0.23; -0.13)	
PLTd, pictures remembered	Ref	-0.20 (-0.41; 0.01)	-0.91 (-1.12; -0.69)	-0.27 (-0.33; -0.20)	
Barthel, index	Ref	-0.04 (-0.11; 0.02)	-0.08 (-0.15; -0.02)	-0.05 (-0.08; -0.03)	
IADL, points	Ref	-0.05 (-0.14; 0.04)	-0.17 (-0.27; -0.08)	-0.09 (-0.12; -0.06)	

Table 4. Associations of weight variability (SD) and cognitive function at month 30 of follow-up (*continued*)

Weight variability, SD			
	Low third	Middle third	Upper third
	(N _{max} = 1436)	(N _{max} = 1436)	(N _{max} = 1437)
Cognitive test	Beta (95% CI)	Beta (95% CI)	Beta (95% CI)
Fully adjusted with systolic blood pressure variability³			
Stroop, s	Ref	1.93 (0.01; 3.86)	4.52 (2.52; 6.51)
LDCT, digits coded	Ref	-1.28 (-1.83; -0.73)	-1.95 (-2.51; -1.39)
PLTi, pictures remembered	Ref	-0.17 (-0.31; -0.03)	-0.58 (-0.73; -0.43)
PLTd, pictures remembered	Ref	-0.18 (-0.39; 0.02)	-0.86 (-1.08; -0.65)
Barthel, index	Ref	-0.04 (-0.10; 0.03)	-0.08 (-0.14; -0.01)
IADL, points	Ref	-0.04 (-0.14; 0.05)	-0.17 (-0.25; -0.06)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, mean weight during follow-up, height, education. ²=sex, age, country, education level, height, mean weight during follow-up time, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, number of medications, use of diuretics, use of antidiuretics. ³=sex, age, country, education level, height, mean weight during follow-up time, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, number of medications, use of diuretics, use of antidiuretics, systolic blood pressure variability, mean systolic blood pressure. Stroop test was performed by 4122 participants, Letter-Digit Coding test was performed by 4264 participants, and the Picture-Word Learning immediate and delayed tests were performed by 4309 participants. Items written in bold indicate that the 95% confidence interval does not contain zero.

4. Discussion

In the present study, we investigated the association of 2.5-years variation in weight and loss of weight with subsequent cognitive performance in a cohort of older individuals at increased risk of CVD. Loss of weight and a higher weight variability were independently significantly associated with worse cognition. These findings were consistent in all tested cognitive domains and independent of incident disease-states, use of diuretics or antidepressants, cardiovascular risk-factors. This study found no associations with activities of daily living.

Major strengths of this study are its size with >4,300 older participants, use of multiple consecutive weight measurements and the ability to investigate various cognitive domains. Furthermore, participants were free of dementia at baseline due to the PROSPER exclusion criteria of MMSE scores <24. A limitation is the lack of information regarding physical activity and intentional weight loss. At the onset of data collection, study participants received health counselling which may have led some participants to intentionally lose weight. Intentional loss of weight has been showed to result in improved cognitive function [18], whereas the present study could not corroborate this observation. Furthermore, reverse causation may also play a role as cognitive impairment can also lead to raised energy expenditure or changes in eating habits [2]. Lack of associations of weight loss or variability with BI and IADL may be due to the ceiling effects.

Current evidence on the association between cognitive decline and variation of weight mainly comes from studies that collected fewer weights measurements (≤ 3) than the present study (median of 11 measurements, IQR 11; 11) [1, 6, 19, 20]. Furthermore, these studies did not calculate an average slope of weight change during follow-up as done in the present study. In line with our results, these studies demonstrate that greater variation in weight is associated with higher risk of dementia. In the present study, we used a battery of cognitive tests to examine various domains as opposed to solely the diagnosis of dementia [1, 19-21], adding nuance to our findings. Consistent with our findings, these studies suggest that larger variation in weight may function as a marker of risk of early cognitive impairment.

On the other hand, some studies present mixed results. Improved cognitive function following intended weight loss has also been observed among older individuals [18, 22, 23] However, these studies were designed to *intentionally* induce weight loss in participants whereas in the present study, it is believed that weight loss in the vast majority was *unintentional* and therefore a consequence of other subclinical processes.

Incident disease-states such as cancer are often thought to be the underlying cause of weight loss [2]. In the present study, participants with a recent history of malignancies (<5 years) and cardiovascular events (<6 months) were excluded during recruitment of the original PROSPER trial, and we found that associations did not change after repeating the analyses excluding participants with incident disease-states during follow-up. It is therefore more likely that weight loss in the present study may have resulted from unstable homeostasis rather than due to major disease during the follow-up. We also demonstrated that higher weight variability was associated with worse cognitive function, independent of systolic blood pressure variability. This may suggest that variability in body weight and

systolic blood pressure do not share a common cause, and that the current findings are different from what we have previously published [10, 11]. In addition, the two variability variables are more likely to symbolize different biological pathways are involved in regulating homeostasis.

The biological mechanisms by which weight loss and most notably weight variability is associated with cognitive function are not fully understood. Weight loss can result as a downstream effect of normal ageing as metabolic needs may change [2]. In addition, polypharmacy can cause dentition and absorption issues, altered gastric signals, causing early satiation, or loss of appetite. Reverse causality, where weight loss is an early manifestation of dementia [24], could also contribute to our findings. However, weight loss has been shown to precede symptoms of cognitive decline, implying that pathological processes of weight loss could contribute, perhaps indirectly, to cognitive decline [25]. Future long-term investigations are warranted to examine whether maintaining weight stability can effectively decrease the risk of cognitive decline.

In conclusion, we found that in older participants at increased risk for vascular disease, steeper decline in weight and a higher variability in body weight, were strongly associated with lower cognitive function in multiple domains. These findings were independent of cardiovascular risk-factors, comorbidities, incident disease-states, and independent of each other. Although we did not produce evidence favouring weight change to cause decreased cognitive function, it may represent an early manifestation or signal of cognitive decline.

5. References

1. Ravona-Springer R, Schnaider-Beeri M, Goldbourt U. Body weight variability in midlife and risk for dementia in old age. *Neurology*. 2013;80(18):1677-83.
2. McMinn J, Steel C, Bowman A. Investigation and management of unintentional weight loss in older adults. *BMJ*. 2011;342:d1732.
3. Cova I, Clerici F, Rossi A, Cucumo V, Ghirelli R, Maggiore L, et al. Weight Loss Predicts Progression of Mild Cognitive Impairment to Alzheimer's Disease. *PLoS One*. 2016;11(3):e0151710.
4. Strandberg TE, Stenholm S, Strandberg AY, Salomaa VV, Pitkälä KH, Tilvis RS. The "obesity paradox," frailty, disability, and mortality in older men: a prospective, longitudinal cohort study. *Am J Epidemiol*. 2013;178(9):1452-60.
5. Strandberg TE, Sirola J, Pitkälä KH, Tilvis RS, Strandberg AY, Stenholm S. Association of midlife obesity and cardiovascular risk with old age frailty: a 26-year follow-up of initially healthy men. *Int J Obes (Lond)*. 2012;36(9):1153-7.
6. Giudici KV, Guyonnet S, Rolland Y, Vellas B, de Souto Barreto P, Nourhashemi F. Body Weight Variation Patterns as Predictors of Cognitive Decline over a 5 Year Follow-Up among Community-Dwelling Elderly (MAPT Study). *Nutrients*. 2019;11(6).
7. Sattar N, Welsh P. The obesity paradox in secondary prevention: a weighty intervention or a wait for more evidence? *Eur Heart J*. 2020.
8. Alibhai SM, Greenwood C, Payette H. An approach to the management of unintentional weight loss in elderly people. *Cmaj*. 2005;172(6):773-80.
9. Jimenez A, Pegueroles J, Carmona-Iragui M, Vilaplana E, Montal V, Alcolea D, et al. Weight loss in the healthy elderly might be a non-cognitive sign of preclinical Alzheimer's disease. *Oncotarget*. 2017;8(62):104706-16.
10. Sabayan B, Wijsman LW, Foster-Dingley JC, Stott DJ, Ford I, Buckley BM, et al. Association of visit-to-visit variability in blood pressure with cognitive function in old age: prospective cohort study. *BMJ*. 2013;347:f4600.
11. Smit RA, Trompet S, Sabayan B, le Cessie S, van der Grond J, van Buchem MA, et al. Higher Visit-to-Visit Low-Density Lipoprotein Cholesterol Variability Is Associated With Lower Cognitive Performance, Lower Cerebral Blood Flow, and Greater White Matter Hyperintensity Load in Older Subjects. *Circulation*. 2016;134(3):212-21.
12. Shepherd J, Blauw GJ, Murphy MB, Bollen EL, Buckley BM, Cobbe SM, et al. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet*. 2002;360(9346):1623-30.
13. Shepherd J, Blauw GJ, Murphy MB, Cobbe SM, Bollen EL, Buckley BM, et al. The design of a prospective study of Pravastatin in the Elderly at Risk (PROSPER). PROSPER Study Group. PROspective Study of Pravastatin in the Elderly at Risk. *Am J Cardiol*. 1999;84(10):1192-7.
14. Houx PJ, Shepherd J, Blauw GJ, Murphy MB, Ford I, Bollen EL, et al. Testing cognitive function in elderly populations: the PROSPER study. PROspective Study of Pravastatin in the Elderly at Risk. *Journal of neurology, neurosurgery, and psychiatry*. 2002;73(4):385-9.
15. Mahoney FI, Barthel DW. FUNCTIONAL EVALUATION: THE BARTHEL INDEX. *Md State Med J*. 1965;14:61-5.
16. Lawton MP. The functional assessment of elderly people. *J Am Geriatr Soc*. 1971;19(6):465-81.
17. Wijsman LW, de Craen AJ, Muller M, Sabayan B, Stott D, Ford I, et al. Blood Pressure Lowering Medication, Visit-to-Visit Blood Pressure Variability, and Cognitive Function in Old Age. *Am J Hypertens*. 2016;29(3):311-8.
18. Horie NC, Serrao VT, Simon SS, Gascon MR, Dos Santos AX, Zambone MA, et al. Cognitive Effects of Intentional Weight Loss in Elderly Obese Individuals With Mild Cognitive Impairment. *The Journal of clinical endocrinology and metabolism*. 2016;101(3):1104-12.
19. Nam GE, Park YG, Han K, Kim MK, Koh ES, Kim ES, et al. BMI, Weight Change, and Dementia Risk in Patients With New-Onset Type 2 Diabetes: A Nationwide Cohort Study. *Diabetes Care*. 2019;42(7):1217-24.

20. Park S, Jeon SM, Jung SY, Hwang J, Kwon JW. Effect of late-life weight change on dementia incidence: a 10-year cohort study using claim data in Korea. *BMJ Open*. 2019;9(5):e021739.
21. Stewart R, Masaki K, Xue QL, Peila R, Petrovitch H, White LR, et al. A 32-year prospective study of change in body weight and incident dementia: the Honolulu-Asia Aging Study. *Arch Neurol*. 2005;62(1):55-60.
22. Siervo M, Nasti G, Stephan BC, Papa A, Muscariello E, Wells JC, et al. Effects of intentional weight loss on physical and cognitive function in middle-aged and older obese participants: a pilot study. *J Am Coll Nutr*. 2012;31(2):79-86.
23. Witte AV, Fobker M, Gellner R, Knecht S, Flöel A. Caloric restriction improves memory in elderly humans. *Proc Natl Acad Sci U S A*. 2009;106(4):1255-60.
24. Burns JM, Johnson DK, Watts A, Swerdlow RH, Brooks WM. Reduced lean mass in early Alzheimer disease and its association with brain atrophy. *Arch Neurol*. 2010;67(4):428-33.
25. Buchman AS, Wilson RS, Bienias JL, Shah RC, Evans DA, Bennett DA. Change in body mass index and risk of incident Alzheimer disease. *Neurology*. 2005;65(6):892-7.

