



Universiteit
Leiden
The Netherlands

BASS-XXIII: a new mid-infrared diagnostic for absorption in AGN

Pfeifle, R.; Ricci, C.; Boorman, P.; Stalevski, M.; Asmus, D.; Trakhtenbrot, B.; ... ; Harrison, F.

Citation

Pfeifle, R., Ricci, C., Boorman, P., Stalevski, M., Asmus, D., Trakhtenbrot, B., ... Harrison, F. (2023). BASS-XXIII: a new mid-infrared diagnostic for absorption in AGN. *Bulletin Of The American Astronomical Society*, 254.04. Retrieved from <https://hdl.handle.net/1887/3718880>

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)
Downloaded from: <https://hdl.handle.net/1887/3718880>

Note: To cite this publication please use the final published version (if applicable).

Bulletin of the AAS • Vol. 55, Issue 2 (AAS241 Abstracts)

BASS-XXIII: A New Mid-Infrared Diagnostic for Absorption in AGN

**Ryan Pfeifle¹ Claudio Ricci² Peter Boorman³ Marko Stalevski⁴
Daniel Asmus⁵ Benny Trakhtenbrot⁶ Michael Koss⁷ Daniel Stern⁸
Federica Ricci⁹ Shobita Satyapal¹⁰ Kohei Ichikawa¹¹ David Rosario¹²
Turgay Caglar¹³ Ezequiel Treister⁹ Meredith Powell¹⁴ Kyuseok Oh¹⁵
C. Urry¹⁶ Fiona Harrison³**

¹NASA Goddard Space Flight Center, ²Universidad Diego Portales,

³California Institute of Technology, ⁴Universiteit Gent, ⁵University of Southampton,

⁶Tel Aviv University, ⁷Eureka Scientific, Inc., ⁸Jet Propulsion Laboratory,

⁹Pontificia Universidad Catolica de Chile, ¹⁰George Mason University, ¹¹Tohoku University,

¹²Newcastle University, ¹³Leiden Observatory, ¹⁴Stanford University,

¹⁵Korea Astronomy and Space Science Institute, ¹⁶Yale University

Published on: Jan 31, 2023

URL: <https://baas.aas.org/pub/2023n2i254p04>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

In this study, we use the *Swift*/BAT AGN sample, which has received extensive multiwavelength follow-up analysis as a result of the BAT AGN Spectroscopic Survey (BASS), to develop a diagnostic for nuclear obscuration by examining the relationship between the line-of-sight column densities (N_{H}), the 2-10 keV-to-12 μm luminosity ratio, and *WISE* mid-infrared colors. We demonstrate that heavily obscured AGNs tend to exhibit both preferentially “redder” mid-infrared colors and lower values of $L_{2-10 \text{ keV, Obs.}}/L_{12\mu\text{m}}$ than less obscured AGNs, and we derive expressions relating N_{H} to the $L_{2-10 \text{ keV, Obs.}}/L_{12\mu\text{m}}$ and $L_{22\mu\text{m}}/L_{4.6\mu\text{m}}$ luminosity ratios as well as develop diagnostic criteria using these ratios. Our diagnostic regions yield samples that are $\geq 80\%$ complete and $\geq 60\%$ pure for AGNs with $\log(N_{\text{H}}/\text{cm}^{-2}) \geq 24$, as well as $\geq 85\%$ pure for AGNs with $\log(N_{\text{H}}/\text{cm}^{-2}) \geq 23.5$. We find that these diagnostics cannot be used to differentiate between optically star forming galaxies and active galaxies. Further, mid-IR contributions from host galaxies that dominate the observed 12 μm emission can lead to larger apparent X-ray deficits and redder mid-IR colors than the AGNs would intrinsically exhibit, though this effect helps to better separate less obscured and more obscured AGNs. Finally, we test our diagnostics on two catalogs of AGNs and infrared galaxies, including the XNN-Newton XXL-N field, and we identify several known Compton-thick AGNs as well as a handful of candidate heavily obscured AGNs based upon our proposed obscuration diagnostics.