



Universiteit
Leiden
The Netherlands

The photodissociation and ionization fronts in M17-SW localized with FIFI-LS on board SOFIA

Klein, R.; Reedy, A.; Fischer, C.; Looney, L.; Colditz, S.; Fadda, D.; Tielens, A.G.G.M.

Citation

Klein, R., Reedy, A., Fischer, C., Looney, L., Colditz, S., Fadda, D., & Tielens, A. G. G. M. (2023). The photodissociation and ionization fronts in M17-SW localized with FIFI-LS on board SOFIA. *Bulletin Of The American Astronomical Society*, 210.04. Retrieved from <https://hdl.handle.net/1887/3719216>

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3719216>

Note: To cite this publication please use the final published version (if applicable).

The Photodissociation and Ionization Fronts in M17-SW Localized with FIFI-LS on Board SOFIA

**Randolf Klein¹ Alexander Reedy² Christian Fischer³ Leslie Looney⁴
Sebastian Colditz³ Dario Fadda¹ Alexander Tielens⁵**

¹SOFIA/USRA, ²Rutgers University, ³University of Stuttgart,

⁴University of Illinois, Urbana-Champaign, ⁵Leiden Observatory

Published on: Jul 01, 2023

URL: <https://baas.aas.org/pub/2023n6i210p04>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

To understand star formation rates, studying feedback mechanisms that regulate star formation is necessary. The radiation emitted by nascent massive stars play a significant role in feedback by photodissociating and ionizing their parental molecular clouds. To gain a detailed picture of the physical processes, we mapped the photodissociation region (PDR) M17-SW in several fine-structure and high-J CO lines with FIFI-LS, the far-infrared imaging spectrometer aboard SOFIA. An analysis of the CO and [O I]146 μm line intensities, combined with the far-infrared intensity, allows us to create a density and UV intensity map using a one-dimensional model. The density map reveals a sudden change in the gas density crossing the PDR. The strengths and limits of the model and the locations of the ionization and photodissociation front of the edge-on PDR are discussed.