



Universiteit
Leiden
The Netherlands

Black hole demographics in low-mass galaxies: improving tidal disruption event rate estimations

Hannah, C.; Seth, A.; Stone, N.; Velzen, S. van

Citation

Hannah, C., Seth, A., Stone, N., & Velzen, S. van. (2023). Black hole demographics in low-mass galaxies: improving tidal disruption event rate estimations. *Bulletin Of The American Astronomical Society*, 214.03. Retrieved from <https://hdl.handle.net/1887/3719262>

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3719262>

Note: To cite this publication please use the final published version (if applicable).

Bulletin of the AAS • Vol. 55, Issue 2 (AAS241 Abstracts)

Black Hole Demographics in Low-Mass Galaxies: Improving Tidal Disruption Event Rate Estimations

Christian Hannah¹ Anil Seth¹ Nicholas Stone² Sjoert van Velzen³

¹University of Utah, ²The Hebrew University of Jerusalem, ³Leiden University

Published on: Jan 31, 2023

URL: <https://baas.aas.org/pub/2023n2i214p03>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Measuring massive black hole (MBH) demographics in low-mass galaxies provides a promising route toward constraining the different theoretical seeding mechanisms for MBHs. The advent of the Rubin Observatory (LSST) will allow tidal disruption events (TDEs) to be used in measuring MBH demographics. This measurement requires accurate TDE rate predictions. The current, widely used TDE rate estimates of Stone & Metzger (2016) do not account for stellar density components resulting from the presence of nuclear star clusters (NSCs). Due to the prevalence of NSCs in galaxies with masses between 10^8 and $10^{10} M_{\odot}$ paired with the large enhancement in TDE rates caused by the presence of NSCs (Pfister et al. 2020), the rates of TDEs in these galaxies are determined by the density profiles of their NSCs. In this work, we are constructing the largest currently available set of high-resolution 3-D stellar density profiles of galaxies with NSCs. We are using these stellar density profiles to develop accurate relations between the stellar mass/type of galaxies and their central density profiles. We then combine these relations with realistic model galaxy distributions to create new loss-cone models and improve TDE rate estimates. With these estimates, we will predict the galaxy demographics of TDE hosts for ZTF and Rubin with a wide range MBH demographic scenarios.