



Universiteit
Leiden
The Netherlands

Extragalactic science goals and predictions for the SPICE Far-IR Probe

Bonato, M.; Leisawitz, D.; Urry, C.; Shivaiei, I.; Aalto, S.; Eales, S.; ... ; Wilson, G.

Citation

Bonato, M., Leisawitz, D., Urry, C., Shivaiei, I., Aalto, S., Eales, S., ... Wilson, G. (2023). Extragalactic science goals and predictions for the SPICE Far-IR Probe. *Bulletin Of The American Astronomical Society*, 160.05. Retrieved from <https://hdl.handle.net/1887/3719298>

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3719298>

Note: To cite this publication please use the final published version (if applicable).

Extragalactic science goals and predictions for the SPICE Far-IR Probe

**Matteo Bonato¹ David Leisawitz² C. Urry³ Irene Shivaie⁴
Susanne Aalto⁵ Stephen Eales⁶ Duncan Farrah⁷ Alan Kogut⁸
David Sanders⁷ Nicholas Scoville⁹ Jessica Sutter¹⁰ Serena Viti¹¹
Grant Wilson¹²**

¹INAF-IRA, ²NASA GSFC, ³Yale University, ⁴University of California, Riverside,
⁵Chalmers University of Technology, ⁶Cardiff University, ⁷University of Hawaii,
⁸NASA Goddard Space Flight Center, ⁹Caltech, ¹⁰SOFIA Science Center, ¹¹Leiden Observatory,
¹²University of Massachusetts, Amherst

Published on: Jan 31, 2023

URL: <https://baas.aas.org/pub/2023n2i160p05>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

The reference extragalactic survey of $\sim 0.83 \text{ deg}^2$ with The Space Interferometer for Cosmic Evolution (SPICE), exploiting its excellent sensitivity and its $1' \times 1'$ field of view will detect tens of thousands of star-forming galaxies and thousands of AGNs (in line and in continuum), thus probing the co-evolution of galaxies and AGNs, up to high redshifts ($z \sim 6$). Moreover, SPICE's excellent (sub-arcsecond) angular resolution will allow us to resolve star formation in nearby galaxies. It will also permit us to study the evolution of the morphology of star-forming galaxies up to quite high redshifts (discriminating among galaxy formation and evolution models). Its high spectral resolution ($R \sim 3000$), line sensitivity and broad spectral coverage will allow us to learn about the physical conditions (temperature, density, metallicity) of galaxies over the past ~ 12 billion years. By resolving the rich variety of far-IR spectral lines, SPICE will provide information on streaming motions due to feedback processes and on the relationships among the kinematics of different atomic and molecular gas phases. We present quantitative predictions for blind (photometric+spectroscopic) surveys and pointed observations and discuss the optimization of the observing strategy.